

Developed at the request of:



Research conducted by:



# Climate: Observations, projections and impacts: France

Met Office

Simon N. Gosling, University of Nottingham

Robert Dunn, Met Office

Fiona Carrol, Met Office

Nikos Christidis, Met Office

John Fullwood, Met Office

Diogo de Gusmao, Met Office

Nicola Golding, Met Office

Lizzie Good, Met Office

Trish Hall, Met Office

Lizzie Kendon, Met Office

John Kennedy, Met Office

Kirsty Lewis, Met Office

Rachel McCarthy, Met Office

Carol McSweeney, Met Office

Colin Morice, Met Office

David Parker, Met Office

Matthew Perry, Met Office

Peter Stott, Met Office

Kate Willett, Met Office

Myles Allen, University of Oxford

Nigel Arnell, Walker Institute, University of Reading

Dan Bernie, Met Office

Richard Betts, Met Office

Niel Bowerman, Centre for Ecology and Hydrology

Bastiaan Brak, University of Leeds

John Caesar, Met Office

Andy Challinor, University of Leeds

Rutger Dankers, Met Office

Fiona Hewer, Fiona's Red Kite

Chris Huntingford, Centre for Ecology and Hydrology

Alan Jenkins, Centre for Ecology and Hydrology

Nick Klingaman, Walker Institute, University of Reading

Kirsty Lewis, Met Office

Ben Lloyd-Hughes, Walker Institute, University of Reading

Jason Lowe, Met Office

Rachel McCarthy, Met Office

James Miller, Centre for Ecology and Hydrology

Robert Nicholls, University of Southampton

Maria Noguera, Walker Institute, University of Reading

Friedreike Otto, Centre for Ecology and Hydrology

Paul van der Linden, Met Office

Rachel Warren, University of East Anglia

**The country reports were written by a range of climate researchers, chosen for their subject expertise, who were drawn from institutes across the UK. Authors from the Met Office and the University of Nottingham collated the contributions in to a coherent narrative which was then reviewed. The authors and contributors of the reports are as above.**

Developed at the request of:



Research conducted by:



# Climate: Observations, projections and impacts

France



We have reached a critical year in our response to climate change. The decisions that we made in Cancún put the UNFCCC process back on track, saw us agree to limit temperature rise to 2 °C and set us in the right direction for reaching a climate change deal to achieve this. However, we still have considerable work to do and I believe that key economies and major emitters have a leadership role in ensuring a successful outcome in Durban and beyond.

To help us articulate a meaningful response to climate change, I believe that it is important to have a robust scientific assessment of the likely impacts on individual countries across the globe. This report demonstrates that the risks of a changing climate are wide-ranging and that no country will be left untouched by climate change.

I thank the UK's Met Office Hadley Centre for their hard work in putting together such a comprehensive piece of work. I also thank the scientists and officials from the countries included in this project for their interest and valuable advice in putting it together. I hope this report will inform this key debate on one of the greatest threats to humanity.

**The Rt Hon. Chris Huhne MP, Secretary of State for Energy and Climate Change**



There is already strong scientific evidence that the climate has changed and will continue to change in future in response to human activities. Across the world, this is already being felt as changes to the local weather that people experience every day.

Our ability to provide useful information to help everyone understand how their environment has changed, and plan for future, is improving all the time. But there is still a long way to go. These reports – led by the Met Office Hadley Centre in collaboration with many institutes and scientists around the world – aim to provide useful, up to date and impartial information, based on the best climate science now available. This new scientific material will also contribute to the next assessment from the Intergovernmental Panel on Climate Change.

However, we must also remember that while we can provide a lot of useful information, a great many uncertainties remain. That's why I have put in place a long-term strategy at the Met Office to work ever more closely with scientists across the world. Together, we'll look for ways to combine more and better observations of the real world with improved computer models of the weather and climate; which, over time, will lead to even more detailed and confident advice being issued.

**Julia Slingo, Met Office Chief Scientist**

# Introduction

Understanding the potential impacts of climate change is essential for informing both adaptation strategies and actions to avoid dangerous levels of climate change. A range of valuable national studies have been carried out and published, and the Intergovernmental Panel on Climate Change (IPCC) has collated and reported impacts at the global and regional scales. But assessing the impacts is scientifically challenging and has, until now, been fragmented. To date, only a limited amount of information about past climate change and its future impacts has been available at national level, while approaches to the science itself have varied between countries.

In April 2011, the Met Office Hadley Centre was asked by the United Kingdom's Secretary of State for Energy and Climate Change to compile scientifically robust and impartial information on the physical impacts of climate change for more than 20 countries. This was done using a consistent set of scenarios and as a pilot to a more comprehensive study of climate impacts. A report on the observations, projections and impacts of climate change has been prepared for each country. These provide up to date science on how the climate has already changed and the potential consequences of future changes. These reports complement those published by the IPCC as well as the more detailed climate change and impact studies published nationally.

Each report contains:

- A description of key features of national weather and climate, including an analysis of new data on extreme events.
- An assessment of the extent to which increases in greenhouse gases and aerosols in the atmosphere have altered the probability of particular seasonal temperatures compared to pre-industrial times, using a technique called 'fraction of attributable risk.'
- A prediction of future climate conditions, based on the climate model projections used in the Fourth Assessment Report from the IPCC.
- The potential impacts of climate change, based on results from the UK's Avoiding Dangerous Climate Change programme (AVOID) and supporting literature.  
For details visit: <http://www.avoid.uk.net>

The assessment of impacts at the national level, both for the AVOID programme results and the cited supporting literature, were mostly based on global studies. This was to ensure consistency, whilst recognising that this might not always provide enough focus on impacts of most relevance to a particular country. Although time available for the project was short, generally all the material available to the researchers in the project was used, unless there were good scientific reasons for not doing so. For example, some impacts areas were omitted, such as many of those associated with human health. In this case, these impacts are strongly dependant on local factors and do not easily lend themselves to the globally consistent framework used. No attempt was made to include the effect of future adaptation actions in the assessment of potential impacts. Typically, some, but not all, of the impacts are avoided by limiting global average warming to no more than 2 °C.

The Met Office Hadley Centre gratefully acknowledges the input that organisations and individuals from these countries have contributed to this study. Many nations contributed references to the literature analysis component of the project and helped to review earlier versions of these reports.

We welcome feedback and expect these reports to evolve over time. For the latest version of this report, details of how to reference it, and to provide feedback to the project team, please see the website at [www.metoffice.gov.uk/climate-change/policy-relevant/obs-projections-impacts](http://www.metoffice.gov.uk/climate-change/policy-relevant/obs-projections-impacts)

In the longer term, we would welcome the opportunity to explore with other countries and organisations options for taking forward assessments of national level climate change impacts through international cooperation.





# Summary

## Climate observations

- Over the period 1960 to 2010 there was a spatially consistent warming trend in summer over France.
- There is a clear trend to fewer cool nights and more warm nights since 1960, and fewer cool days and more hot days with a very coherent pattern of changes over the whole of France.
- There has been a general increase in summer temperatures averaged over the country as a result of human influence on climate, making the occurrence of warm summer temperatures more frequent and cool summer temperatures less frequent.

## Climate change projections

- For the A1B emissions scenario projected increases in temperature over France are generally between 2.5°C and 3.5°C. Increases are lower in the north, and higher in southern regions. The agreement between the CMIP3 models is generally moderate over most of the country.
- Europe shows a strong contrast in projected precipitation changes, with large decreases in the south and large increases in the north. France falls towards the southern region with decreasing precipitation, with projected decreases of up to 20% in the southwest, but with smaller decreases of up to 5% further north. There is generally moderate agreement between ensemble members over central and southern France, and low agreement in the north, indicating uncertainty in the position of the transition zone between increasing and decreasing precipitation over Europe.

# Climate change impacts projections

## Crop yields

- Global- and regional-scale studies generally project yield gains for wheat and yield losses for maize, two of France's major crops.
- National- and sub-national-scale studies included here also concur with the results from global-scale studies. In addition they note that repeated dry spells during the growing season may have a large impact on yields, although how the frequency of such events may be affected by climate change remains poorly understood.

## Food security

- France is currently a country with extremely low levels of undernourishment. Global-scale studies included here generally project that France will not face serious food security issues over the next 40 years as a consequence of climate change.
- However, recent research by the AVOID programme and others demonstrates that adaptive measures will be needed to ensuring France's food security under climate change.

## Water stress and drought

- The majority of global and regional assessments included here project that drought could become more frequent in France with climate change, and that water stress could increase.
- National-scale studies included here suggest that agricultural drought could increase in duration and intensity with climate change, to an extent never observed in the historical record. However there remains uncertainty in the magnitude to which such changes may be realised.
- Recent simulations by the AVOID programme project a median increase of 12% in the population may be exposed to increased water stress by 2100 under the A1B emissions scenario.

## Pluvial flooding and rainfall

- Recent modelling studies confirm conclusions from the IPCC AR4 that heavy winter precipitation and flooding could increase with climate change for continental France,

although these projections are based upon large-scale climate modelling experiments across the European domain.

- However, recent regional climate model simulations for France suggest that extreme precipitation may not change under climate change.
- To this end, large uncertainty remains in quantifying the degree to which pluvial flooding could be affected by climate change.

### **Fluvial flooding**

- Projected changes in flood hazard in France are subject to large uncertainties owing to large natural variability and uncertainties in the simulated climate change signal from models.
- European-scale simulations have projected an increase in extreme flow levels in the main French rivers by the end of the century.
- Recent national-scale studies mostly found little change in flooding in France with climate change, however.
- Simulations by the AVOID programme support this, with the majority of the projections showing a greater tendency towards decreasing flood risk at first. Later in the century the models become more evenly divided between increases and decreases in flood risk, especially under the A1B scenario.

### **Coastal regions**

- Continental France's coasts and port cities could be affected by sea level rise (SLR) but to a lesser extent than many other countries across the globe.
- Results from the limited number of studies available indicate that by the 2080s under a high SLR scenario and without adaptation, the average annual number of people flooded in France could be around 463,000. This is greatly reduced with adaptation, to around 2,500. Under a low SLR scenario, 3,000 people could be flooded annually without adaptation and 1,800 could be flooded with adaptation.
- Some French districts in the West Indies, South America, Indian Ocean and Pacific Ocean could be affected by SLR, but these were not considered in this review.





# Table of Contents

?

<b>Chapter 1 – Climate Observations</b> .....	<b>9?</b>
<b>Rationale</b> .....	<b>10?</b>
<b>Climate overview</b> .....	<b>12?</b>
Analysis of long-term features in the mean temperature .....	13?
<b>Temperature extremes</b> .....	<b>15?</b>
Recent extreme temperature events .....	15?
Heat wave, summer 2003 .....	15?
Heat wave, summer 2006 .....	16?
Analysis of long-term features in moderate temperature extremes .....	16?
Attribution of changes in likelihood of occurrence of seasonal mean temperatures .....	22?
Summer 2003.....	22?
Summer 2006.....	23?
<b>Precipitation extremes</b> .....	<b>24?</b>
Recent extreme precipitation events .....	25?
Severe Flooding, June 2010 .....	25?
Analysis of long-term features in precipitation .....	25?
<b>Storms</b> .....	<b>28?</b>
Recent extreme storm events.....	28?
Winter storm Xynthia, February 2010 .....	28?
<b>Summary</b> .....	<b>30?</b>
<b>Methodology annex</b> .....	<b>31?</b>
Recent, notable extremes.....	31?
Observational record .....	32?
Analysis of seasonal mean temperature .....	32?
Analysis of temperature and precipitation extremes using indices .....	33?
Presentation of extremes of temperature and precipitation .....	43?
Attribution.....	47?
<b>References</b> .....	<b>50?</b>
<b>Acknowledgements</b> .....	<b>54?</b>
<b>Chapter 2 – Climate Change Projections</b> .....	<b>55?</b>
<b>Introduction</b> .....	<b>56?</b>
<b>Climate projections</b> .....	<b>58?</b>
Summary of temperature change in France .....	59?
Summary of precipitation change in France .....	59?
<b>Chapter 3 – Climate Change Impact Projections</b> .....	<b>61?</b>
<b>Introduction</b> .....	<b>62?</b>
Aims and approach.....	62?
Impact sectors considered and methods .....	62?
Supporting literature .....	63?

AVOID programme results.....	63
Uncertainty in climate change impact assessment.....	64
<b>Summary of findings for each sector .....</b>	<b>71</b>
<b>Crop yields .....</b>	<b>74</b>
Headline.....	74
Supporting literature .....	74
Introduction .....	74
Assessments that include a global or regional perspective .....	76
National-scale or sub-national scale assessments .....	84
AVOID programme results.....	85
Methodology.....	86
Results .....	87
<b>Food security .....</b>	<b>89</b>
Headline.....	89
Supporting literature .....	89
Introduction .....	89
Assessments that include a global or regional perspective .....	89
National-scale or sub-national scale assessments .....	96
<b>Water stress and drought .....</b>	<b>97</b>
Headline.....	97
Supporting literature .....	97
Introduction .....	97
Assessments that include a global or regional perspective .....	98
National-scale or sub-national scale assessments .....	106
AVOID Programme Results.....	107
Methodology.....	107
Results .....	108
<b>Pluvial flooding and rainfall .....</b>	<b>110</b>
Headline.....	110
Supporting literature .....	110
Introduction .....	110
Assessments that include a global or regional perspective .....	111
National-scale or sub-national scale assessments .....	112
<b>Fluvial flooding .....</b>	<b>115</b>
Headline.....	115
Supporting literature .....	115
Introduction .....	115
Assessments that include a global or regional perspective .....	116
National-scale or sub-national scale assessments .....	117
AVOID programme results.....	118
Methodology.....	118
Results .....	119
<b>Tropical cyclones.....</b>	<b>121</b>
<b>Coastal regions .....</b>	<b>122</b>

Headline.....	122
Supporting literature .....	122
Assessments that include a global or regional perspective .....	122
National-scale or sub-national scale assessments .....	131
<b>References.....</b>	<b>133</b>

?

?



?

# Chapter 1 – Climate Observations

?

?

?


?

?



## Rationale

Present day weather and climate play a fundamental role in the day to day running of society. Seasonal phenomena may be advantageous and depended upon for sectors such as farming or tourism. Other events, especially extreme ones, can sometimes have serious negative impacts posing risks to life and infrastructure, and significant cost to the economy. Understanding the frequency and magnitude of these phenomena, when they pose risks or when they can be advantageous and for which sectors of society, can significantly improve societal resilience. In a changing climate it is highly valuable to understand possible future changes in both potentially hazardous events and those reoccurring seasonal events that are depended upon by sectors such as agriculture and tourism. However, in order to put potential future changes in context, the present day must first be well understood both in terms of common seasonal phenomena and extremes.



**Figure 1.** Location of boxes for the regional average time series (red dashed box) in Figures 3 and 5 and the attribution region (grey box) in Figure 4.

The purpose of this chapter is to summarise the weather and climate from 1960 to present day. This begins with a general climate overview including an up to date analysis of changes in surface mean temperature. These changes may be the result of a number of factors including climate change, natural variability and changes in land use. There is then a focus on extremes of temperature, precipitation and storms selected from 2000 onwards, reported in the World Meteorological Organization (WMO) Annual Statement on the Status of the Global Climate and/or the Bulletin of the American Meteorological Society (BAMS) State of the Climate reports. This is followed by a discussion of changes in moderate extremes from 1960 onwards using an updated version of the HadEX extremes database (Alexander et al., 2006) which categorises extremes of temperature and precipitation. These are core climate variables which have received significant effort from the climate research community in terms of data acquisition and processing and for which it is possible to produce long high quality records for monitoring. No new analysis is included for storms (see the methodology section that follows for background). For seasonal temperature extremes, an attribution

analysis then puts the seasons with highlighted extreme events into context of the recent climate versus a hypothetical climate in the absence of anthropogenic emissions (Christidis et al., 2011). It is important to note that we carry out our attribution analyses on seasonal mean temperatures over the entire country. Therefore these analyses do not attempt to attribute the changed likelihood of individual extreme events. The relationship between extreme events and the large scale mean temperature is likely to be complex, potentially being influenced by *inter alia* circulation changes, a greater expression of natural internal variability at smaller scales, and local processes and feedbacks. Attribution of individual extreme events is an area of developing science. The work presented here is the foundation of future plans to systematically address the region's present and projected future weather and climate, and the associated impacts.

The methodology section provides details of the data shown here and of the scientific analyses underlying the discussions of changes in the mean temperature and in temperature and precipitation extremes. It also explains the methods used to attribute the likelihood of occurrence of seasonal mean temperatures.

## Climate overview


France is located between the latitudes of 43° and 51°N and has a mainly temperate climate. Northern and western France are mainly low-lying, but there are some high mountain regions with the Alps and Jura in the east and the Pyrenees in the south, as well as the Massif Central in central southern France. Consequently, there are considerable variations in climate. The mountainous areas are the coldest regions of France, especially in winter, with temperature decreasing with altitude. Northern and north-western France has a maritime climate with mild winters and warm summers, for example Brest on the north-west coast has a mean annual temperature of 11°C, and a seasonal range of ±5°C. Central northern France has rather colder winters and slightly warmer summers, with Paris having a mean annual temperature of 13°C and a seasonal range of ±8°C. Bordeaux in the south-west has very similar temperature climatology to Paris. Lyon, towards the south-east, has a more continental climate with colder winters and a greater chance of frost and snow; here, the annual mean temperature is 12°C with a seasonal range of ±9°C. The Mediterranean coast and the island of Corsica have a Mediterranean climate with hot summers and mild, sunny winters. The pleasant winter and spring weather is often interrupted, however, by cold and blustery weather brought by a northerly wind called the Mistral. Nice has an annual mean temperature of 16°C, with a seasonal range of ±7°C.

Most of France receives a moderate amount of precipitation throughout the year. The exception to this is the Mediterranean climate area which has dry summers as the track of eastward-moving Atlantic weather systems migrates away to the north. Nice has average annual rainfall of 800 mm, which mostly falls in the autumn and winter, although occasional thunderstorms do occur in summer. Central areas have little seasonal variation in rainfall, although the mechanisms change with summer rainfall associated with convective thunderstorms and winter rainfall with frontal Atlantic disturbances. Lyon has an annual average rainfall of 840 mm and Paris 670 mm. Western France is more exposed to storms coming from the Atlantic, so has more seasonal variation with increased rainfall amounts in winter. Bordeaux has an annual average amount of 980 mm and Brest 1140 mm. The high mountain areas have heavier precipitation, much of it falling as snow in winter.

Summer thunderstorms can lead to flooding. Droughts and heat waves are also a potential hazard in the summer.

## **Analysis of long-term features in the mean temperature**

CRUTEM3 data (Brohan et al., 2006) have been used to provide an analysis of mean temperatures from 1960 to 2010 over France using the median of pairwise slopes method to fit the trend (Sen, 1968; Lanzante, 1996). The methods are fully described in the methodology section. In agreement with increasing global average temperatures (Sánchez-Lugo et al., 2011), over the period 1960 to 2010 there is a spatially consistent warming signal for temperature over France as shown in Figure 2. Grid boxes in which the 5<sup>th</sup> to 95<sup>th</sup> percentiles of the slopes are of the same sign can be more confidently regarded as showing this signal and they are geographically widespread for the summer months (June to August). For the winter months (December to February) confidence in this signal is lower in all but one grid-box in the south. Regionally averaged trends (over grid boxes included in the red dashed box in Figure 1) calculated by the median of pairwise slopes show clear warming trends but with high confidence only for summer. This trend for summer is 0.35°C per decade (5<sup>th</sup> to 95<sup>th</sup> percentile of slopes: 0.24 to 0.48°C per decade) and for winter is 0.20°C per decade (5<sup>th</sup> to 95<sup>th</sup> percentile of slopes: -0.02 to 0.40°C per decade).



**Figure 2.** Decadal trends in seasonally averaged temperatures for France and the surrounding regions over the period 1960 to 2010. Monthly mean anomalies from CRUTEM3 (Brohan et al., 2006) are averaged over each 3 month season (June-July-August – JJA and December-January-February – DJF). Trends are fitted using the median of pairwise slopes method (Sen, 1968; Lanzante, 1996). There is high confidence in the trends shown if the 5<sup>th</sup> to 95<sup>th</sup> percentiles of the pairwise slopes do not encompass zero because here the trend is considered to be significantly different from a zero trend (no change). This is shown by a black dot in the centre of the respective grid box.



# Temperature extremes

Both hot and cold temperature extremes can place many demands on society. While seasonal changes in temperature are normal and indeed important for a number of societal sectors (e.g. tourism, farming etc.), extreme heat or cold can have serious negative impacts. Importantly, what is 'normal' for one region may be extreme for another region that is less well adapted to such temperatures.

Table 1 shows selected extreme events since 2000 that are reported in WMO Statements on Status of the Global Climate and/or BAMS State of the Climate reports. The heat waves of 2003 and 2006 are highlighted below as examples of extreme temperature events in France.

Year	Month	Event	Details	Reference
2003	Jul-Aug	Heat wave	Likely the warmest summer in parts of central Europe since at least 1540.	BAMS (Levinson & Waple, 2004)
2006	Jul	Heat wave	June-July temperatures significantly above average. July was the second hottest month on record.	BAMS (Obregon et al., 2007)
2009	Aug	Heat wave	15-20 August temperatures reached 36°C in the North and 40 °C in the South	BAMS (Obregon et al., 2010)
2009	Dec	Cold spell	15-22 December, temperatures fell to -20 °C across France.	BAMS (Obregon et al., 2010)

**Table 1.** Selected extreme temperature events reported in WMO Statements on Status of the Global Climate and/or BAMS State of the Climate reports since 2000.

## Recent extreme temperature events

### Heat wave, summer 2003

The summer of 2003 was one of the warmest on record across parts of Europe, and in parts of Central Europe was likely the warmest since 1540 (Levinson and Waple, 2004). Two distinct periods of exceptional heat occurred during the summer season— the first in June and the second during the first half of August. The heat waves resulted from strong high pressure over western Europe. Such “blocking highs” can persist for many days in Europe

during summer (Bell & Eichler, 2004). In 2003, heated air from the south reinforced the strength and persistence of the heat wave, and nearly all the sun's radiation was converted to heat because of the soil and vegetation dryness (WMO, 2004). The August heat wave was the more serious of the two, because it coincided with the normal peak in summer temperatures and was accompanied by an almost complete absence of rainfall. During June-August 2003 daily maximum temperatures averaged 36°–38°C across southern and central France (Bell & Eichler, 2004).

One of impacts of the high temperatures and dry conditions was the exacerbation of forest fires that burned across southern France in July (Alexander & Parker, 2004). Many people were affected by the heat wave, in France, 11,000 heat-related deaths were reported between late July and mid-August (Levinson and Waple, 2004). These deaths resulted not only from the extreme temperatures, but also particularly from the duration that these conditions persisted in areas not accustomed to such conditions (Bell & Eichler, 2004).

### **Heat wave, summer 2006**

In June and July 2006 summer temperatures were considerably above average. In France it was the hottest July on record since 1949 and the second hottest month overall behind August 2003 (Obregón et al., 2007). USA Today reported that 64 people died in France as a result of the high temperatures, with the majority of those affected being elderly. The causes of death ranged from severe dehydration to heat stroke (USA Today, 2006).

## **Analysis of long-term features in moderate temperature extremes**


ECA&D data (Klein Tank et al., 2002) have been used to update the HadEX extremes analysis for France from 1960 to 2010 using daily maximum and minimum temperatures. Here we discuss changes in the frequency of cool days and nights and warm days and nights which are moderate extremes. Cool days/nights are defined as being below the 10<sup>th</sup> percentile of daily maximum/minimum temperature and warm days/nights are defined as being above the 90<sup>th</sup> percentile of the daily maximum/minimum temperature. The methods are fully described in the methodology section.

Over the period analysed (1960-2009) there is a clear trend to fewer cool nights and more warm nights, and also to fewer cool days and more hot days (Figure 3). This matches the


trend in the mean temperatures outlined in Section 2. This implies that there is a shift in the temperature distribution, with very little change in its shape. The warming signal is very coherent over the whole of France making it difficult to comment on regional differences.

□



a)




b)




Monthly: -1.01% per decade (-1.30 to -0.74)  
Total change of -5.06% from 1960 to 2011 (-6.51% to -3.69%)  
Annual: -1.15% per decade (-1.58 to -0.69)  
Total change of -5.74% from 1960 to 2010 (-7.90% to -3.43%)




e)



f)



Monthly: -0.98% per decade (-1.28 to -0.69)  
Total change of -4.91% from 1960 to 2011 (-6.39% to -3.47%)  
Annual: -1.15% per decade (-1.68 to -0.62)  
Total change of -5.74% from 1960 to 2010 (-8.39% to -3.10%)



**Figure 3.** Change in cool nights (a,b), warm nights (c,d), cool days (e,f) and warm days (g,h) for France over the period 1960 to 2010 relative to 1961-1990 from the ECA&D dataset (Klein Tank et al., 2002). a,c,e,g) Grid-box decadal trends. Grid-boxes outlined in solid black contain at least 3 stations and so are likely to be more representative of the wider grid-box. Trends are fitted using the median of pairwise slopes method (Sen, 1968; Lanzante, 1996). Higher confidence in a long-term trend is shown by a black dot if the 5th to 95th percentile slopes are of the same sign. Differences in spatial coverage occur because each index has its own decorrelation length scale (see methodology section); b,d,f,h) Area averaged annual time series for 5.625° W to 9.375° E and 41.25° to 56.25° N as shown by the green box on the map and red box in Figure 1. Thin and thick black lines show the monthly and annual variation respectively. Monthly (orange) and annual (blue) trends are fitted as described above. The decadal trend and its 5th to 95th percentile confidence intervals are stated along with the change over the period for which there are data available. All the trends have higher confidence that they are different from zero as their 5th to 95th percentile slopes are of the same sign. The green vertical lines show the locations of the heat waves in 2003 and 2006.

The time series of the temperature indicators show a large amount of natural variability. However, the heat wave for 2003 stands out as exceptional in the number of warm days and nights. The heat wave of 2006 was climatologically weaker than the one in 2003 with a strong signal only in the number of warm nights. All four time-series show a clear trend linking into the warming signal seen in the average temperatures (Figure 2) with high confidence that the trend is different from zero.

## **Attribution of changes in likelihood of occurrence of seasonal mean temperatures**

Today's climate covers a range of likely extremes. Recent research has shown that the temperature distribution of seasonal means would likely be different in the absence of anthropogenic emissions (Christidis et al., 2011). Here we discuss the seasonal means, within which the highlighted extreme temperature events occur, in the context of recent climate and the influence of anthropogenic emissions on that climate. The methods are fully described in the methodology section.


### **Summer 2003**

The distributions of the summer mean regional temperature in recent years in the presence and absence of anthropogenic forcings are shown in Figure 4. Analyses with both models suggest that human influences on the climate have shifted the distribution to higher temperatures. Considering the average over the entire region, the 2003 summer is exceptionally hot, as it lies at the far end of the warm tail of the temperature distributions for the climate influenced by anthropogenic forcings (red distributions) and is the hottest since 1900 in the CRUTEM3 dataset. In the absence of human influences on the climate (green distributions), the season would be even more extreme. It should be noted that the attribution results shown here refer to temperature anomalies over the entire region and over an entire season, whereas the actual extreme event had a shorter duration and affected a smaller region.



## Summer 2006

The observed summer mean regional temperature in 2006 is also shown in Figure 4. Considering the average over the entire region, the 2006 summer is warm, as it lies at the warm tail of the temperature distributions for the climate influenced by anthropogenic forcings (red distributions), but not as extreme as the summer of 2003. In the absence of human influences on the climate, the season would be more extreme, as it lies further into the tails of the distributions plotted in green. It should again be noted that the attribution results shown here refer to temperature anomalies over the entire region and over an entire season, whereas the actual extreme event had a shorter duration and affected a smaller region.



**Figure 4.** Distributions of the June-July-August mean temperature anomalies (relative to 1961-1990) averaged over a Northern European region that encompasses France (10W-20E, 40-60N – as shown in Figure 1) including (red lines) and excluding (green lines) the influence of anthropogenic forcings. The distributions describe the seasonal mean temperatures expected in recent years (2000-2009) and are based on analyses with the HadGEM1 (solid lines) and MIROC (dotted lines) models. The vertical black line marks the observed anomaly in 2006 and the vertical orange and blue lines correspond to the maximum and minimum anomaly in the CRUTEM3 dataset since 1900 respectively.

## Precipitation extremes

Precipitation extremes, either excess or deficit, can be hazardous to human health, societal infrastructure, and livestock and agriculture. While seasonal fluctuations in precipitation are normal and indeed important for a number of societal sectors (e.g. tourism, farming etc.), flooding or drought can have serious negative impacts. These are complex phenomena and often the result of accumulated excesses or deficits or other compounding factors such as spring snow-melt, high tides/storm surges or changes in land use. The analysis section below deals purely with precipitation amounts.

Table 2 shows selected extreme events since 2000 that are reported in WMO Statements on Status of the Global Climate and/or BAMS State of the Climate reports. The severe flooding which occurred in the département of Var in 2010 is highlighted below as an example of an extreme precipitation event for France.

Year	Month	Event	Details	Source
2000	Oct-Nov	Heavy rainfall	Records of 2-month rainfall totals broken in many areas e.g. Paris-Montsouris, Boulogne, La-Rochelle, Bordeaux and Nice.	WMO (2001)
2001	Oct-Mar	Heavy rainfall	Bretagne region had exceptional precipitation in the 6- month period ending in March with numerous cities setting 6-month rainfall records.	BAMS (Waple et al., 2002)
2005		Drought	Drought throughout the year	BAMS (Trigo et al., 2006)
2008	Oct-Nov	Heavy rainfall/flooding	3 day rainfall totals of 500mm in places. Severe flooding/flash floods in central and east-central parts.	WMO (2009)
2009	Jan	Snow	20-40cm snow fell in southeastern France for first time since 1987.	BAMS (Obregón et al., 2010).
2010	Jun	Flooding	Heavy rain from storms leading to worst floods since 1827.	WMO (2011)

**Table 2.** Selected extreme precipitation events reported in WMO Statements on Status of the Global Climate and/or BAMS State of the Climate reports since 2000.

## **Recent extreme precipitation events**



### **Severe Flooding, June 2010**

In June storms hit the south-east of France and the large amounts of heavy rain led to localised flash flooding (WMO, 2011). The rainfall amounts were exceptional for the time of year. 400 mm of rain fell in less than 2 days in Provence, an amount that had not been seen over the last 50 years (Météo France, 2010). These were the worst floods since 1827 and caused severe damage and loss of life in southern France (WMO, 2011). A number of towns in the department of Var were affected, with hundreds of homes flooded. According to BBC reports, at least 19 people were killed by flash floods and several others reported missing (BBC, 2010).



## **Analysis of long-term features in precipitation**

ECA&D data (Klein Tank et al., 2002) have been used to update the HadEX extremes analysis for France from 1960 to 2010 for daily precipitation totals. Here we discuss changes in the annual total precipitation, and in the frequency of prolonged (greater than 6 days) wet and dry spells. The methods are fully described in the methodology section.


□



Annual: -5.75mm per decade (-23.42 to 13.97)  
Total change of -28.76mm from 1960 to 2010 (-117.11mm to 69.87mm)



Annual: -0.03days per decade (-0.65 to 0.60)  
Total change of -0.17days from 1960 to 2010 (-3.26days to 2.98days)



**Figure 5.** The change in annual total rainfall (a,b), the annual number of continuous dry days (c,d) and the annual number of continuous wet days (e,f) over the period 1960-2010. The maps and time series have been created in exactly the same way as Figure 3. The green vertical lines show the date of the flooding of 2010. Only annual regional averages are shown in b,d,f. All the trends have lower confidence as they are different from zero, as their 5th to 95th percentile slopes are of different signs, and hence are marked with dotted lines.

There is no strong signal in the precipitation indices across France, neither in the maps, nor in the time series (which have been created from a regional average of all the grid boxes contained within the red box, Figure 5). There is an indication for a north to south change in the total precipitation, with a drying in the south and an increase in rainfall in the north, but no grid box is significant. The number of consecutive dry and wet days both appear to decrease, but with low confidence for all the grid boxes. The dates of the floods of June 2010 are marked in the time series. There is no clear signal from the flooding as this was a short, intense and very localised event not well captured by these indices and the subsequent gridding.

## Storms

Storms can be very hazardous to all sectors of society. They can be small with localised impacts or spread across wide regions. There is no systematic observational analysis included for storms because, despite recent progress (Peterson et al., 2011; Cornes and Jones, 2011), land surface wind data are not yet adequate for worldwide robust analysis (see methodology section). Further progress awaits studies of the more reliable barometric pressure data through the new 20<sup>th</sup> Century Reanalysis (Compo et al., 2011) and its planned successors.

Table 3 shows selected extreme events since 2000 that are reported in WMO Statements on Status of the Global Climate and/or BAMS State of the Climate reports. The European windstorm Xynthia which crossed Europe in 2010 is highlighted below as an example of a storm event for France.

Year	Month	Event	Details	Source
2010	Feb	Storm	Atlantic storm Xynthia, 27-28 February.	BAMS (Maier et al., 2011)

**Table 3.** Selected extreme storm events reported in WMO Statements on Status of the Global Climate and/or BAMS State of the Climate reports since 2000.

## Recent extreme storm events

### Winter storm Xynthia, February 2010

The violent extratropical storm Xynthia hit coastal Western Europe on the 27–28 February (Maier et al., 2011). Bringing hurricane force winds and heavy rain in what was the worst storm in the region since 1999. Gusts of 120–140 km/h were common on low ground in France, and at Pic du Midi in the French Pyrenees a wind gust of 238 km/h was recorded (WMO, 2011).

More than 60 people were killed across Western Europe. The majority of deaths were in France where storm surges reached 1.5 m at La Rochelle and caused sea walls to break in L'Agillon-sur-Mer, Vendee (Maier et al., 2011; WMO, 2011). The widespread wind and storm

surge damage resulted in insured losses in France and Germany exceeding US\$ 4 billion (WMO, 2011).

# Summary

**The main features seen in observed climate over France from this analysis are:**

- Over the period 1960 to 2010 there is a spatially consistent warming trend in summer over France.
- There is a clear trend to fewer cool nights and more warm nights since 1960, and fewer cool days and more hot days with a very coherent pattern of changes over the whole of France.
- There has been a general increase in summer temperatures averaged over the country as a result of human influence on climate, making the occurrence of warm summer temperatures more frequent and cool summer temperatures less frequent.



## Methodology annex

### Recent, notable extremes

In order to identify what is meant by 'recent' events the authors have used the period since 1994, when WMO Status of the Global Climate statements were available to the authors. However, where possible, the most notable events during the last 10 years have been chosen as these are most widely reported in the media, remain closest to the forefront of the memory of the country affected, and provide an example likely to be most relevant to today's society. By 'notable' the authors mean any event which has had significant impact either in terms of cost to the economy, loss of life, or displacement and long term impact on the population. In most cases the events of largest impact on the population have been chosen, however this is not always the case.

Tables of recent, notable extreme events have been provided for each country. These have been compiled using data from the World Meteorological Organisation (WMO) Annual Statements on the Status of the Global Climate. This is a yearly report which includes contributions from all the member countries, and therefore represents a global overview of events that have had importance on a national scale. The report does not claim to capture all events of significance, and consistency across the years of records available is variable. However, this database provides a concise yet broad account of extreme events per country. This data is then supplemented with accounts from the monthly National Oceanic and Atmospheric Administration (NOAA) State of the Climate reports which outline global extreme events of meteorological significance.

We give detailed examples of heat, precipitation and storm extremes for each country where these have had significant impact. Where a country is primarily affected by precipitation or heat extremes this is where our focus has remained. An account of the impact on human life, property and the economy has been given, based largely on media reporting of events, and official reports from aid agencies, governments and meteorological organisations. Some data has also been acquired from the Centre for Research on Epidemiological Disasters (CRED) database on global extreme events. Although media reports are unlikely to be

completely accurate, they do give an indication as to the perceived impact of an extreme event, and so are useful in highlighting the events which remain in the national psyche.

Our search for data has not been exhaustive given the number of countries and events included. Although there are a wide variety of sources available, for many events, an official account is not available. Therefore figures given are illustrative of the magnitude of impact only (references are included for further information on sources). It is also apparent that the reporting of extreme events varies widely by region, and we have, where possible, engaged with local scientists to better understand the impact of such events.

The aim of the narrative for each country is to provide a picture of the social and economic vulnerability to the current climate. Examples given may illustrate the impact that any given extreme event may have and the recovery of a country from such an event. This will be important when considering the current trends in climate extremes, and also when examining projected trends in climate over the next century.

## **Observational record**

In this section we outline the data sources which were incorporated into the analysis, the quality control procedure used, and the choices made in the data presentation. As this report is global in scope, including 23 countries, it is important to maintain consistency of methodological approach across the board. For this reason, although detailed datasets of extreme temperatures, precipitation and storm events exist for various countries, it was not possible to obtain and incorporate such a varied mix of data within the timeframe of this project. Attempts were made to obtain regional daily temperature and precipitation data from known contacts within various countries with which to update existing global extremes databases. No analysis of changes in storminess is included as there is no robust historical analysis of global land surface winds or storminess currently available.

### **Analysis of seasonal mean temperature**

Mean temperatures analysed are obtained from the CRUTEM3 global land-based surface-temperature data-product (Brohan et al. 2006), jointly created by the Met Office Hadley Centre and Climatic Research Unit at the University of East Anglia. CRUTEM3 comprises of more than 4000 weather station records from around the world. These have been averaged together to create 5° by 5° gridded fields with no interpolation over grid boxes that do not

contain stations. Seasonal averages were calculated for each grid box for the 1960 to 2010 period and linear trends fitted using the median of pairwise slopes (Sen 1968; Lanzante 1996). This method finds the slopes for all possible pairs of points in the data, and takes their median. This is a robust estimator of the slope which is not sensitive to outlying points. High confidence is assigned to any trend value for which the 5<sup>th</sup> to 95<sup>th</sup> percentiles of the pairwise slopes are of the same sign as the trend value and thus inconsistent with a zero trend.

### **Analysis of temperature and precipitation extremes using indices**

In order to study extremes of climate a number of indices have been created to highlight different aspects of severe weather. The set of indices used are those from the World Climate Research Programme (WCRP) Climate Variability and Predictability (CLIVAR) Expert Team on Climate Change Detection and Indices (ETCCDI). These 27 indices use daily rainfall and maximum and minimum temperature data to find the annual (and for a subset of the indices, monthly) values for, e.g., the 'warm' days where daily maximum temperature exceeds the 90<sup>th</sup> percentile maximum temperature as defined over a 1961 to 1990 base period. For a full list of the indices we refer to the website of the ETCCDI (<http://cccma.seos.uvic.ca/ETCCDI/index.shtml>).

Index	Description	Shortname	Notes
Cool night frequency	Daily minimum temperatures lower than the 10 <sup>th</sup> percentile daily minimum temperature using the base reference period 1961-1990	TN10p	---
Warm night frequency	Daily minimum temperatures higher than the 90 <sup>th</sup> percentile daily minimum temperature using the base reference period 1961-1990	TN90p	---
Cool day frequency	Daily maximum temperatures lower than the 10 <sup>th</sup> percentile daily maximum temperature using the base reference period 1961-1990	TX10p	---
Warm day frequency	Daily maximum temperatures higher than the 90 <sup>th</sup> percentile daily maximum temperature using the base reference period 1961-1990	TX90p	---
Dry spell duration	Maximum duration of continuous days within a year with rainfall <1mm	CDD	Lower data coverage due to the requirement for a 'dry spell' to be at least 6 days long resulting in intermittent temporal coverage
Wet spell duration	Maximum duration of continuous days with rainfall >1mm for a given year	CWD	Lower data coverage due to the requirement for a 'wet spell' to be at least 6 days long resulting in intermittent temporal coverage
Total annual precipitation	Total rainfall per year	PRCPTOT	---

**Table 4.** Description of ETCCDI indices used in this document

A previous global study of the change in these indices, containing data from 1951-2003 can be found in Alexander et al. 2006, (HadEX; see <http://www.metoffice.gov.uk/hadobs/hadex/>). In this work we aimed to update this analysis to the present day where possible, using the most recently available data. A subset of the indices is used here because they are most easily related to extreme climate events (Table 4).

#### **Use of HadEX for analysis of extremes**

The HadEX dataset comprises all 27 ETCCDI indices calculated from station data and then smoothed and gridded onto a 2.5° x 3.75° grid, chosen to match the output from the Hadley Centre suite of climate models. To update the dataset to the present day, indices are calculated from the individual station data using the RClimDex/FClimDex software; developed and maintained on behalf of the ETCCDI by the Climate Research Branch of the Meteorological Service of Canada. Given the timeframe of this project it was not possible to

obtain sufficient station data to create updated HadEX indices to present day for a number of countries: Brazil; Egypt; Indonesia; Japan (precipitation only); South Africa; Saudi Arabia; Peru; Turkey; and Kenya. Indices from the original HadEX data-product are used here to show changes in extremes of temperature and precipitation from 1960 to 2003. In some cases the data end prior to 2003. Table 5 summarises the data used for each country. Below, we give a short summary of the methods used to create the HadEX dataset (for a full description see Alexander et al. 2006).

To account for the uneven spatial coverage when creating the HadEX dataset, the indices for each station were gridded, and a land-sea mask from the HadCM3 model applied. The interpolation method used in the gridding process uses a decorrelation length scale (DLS) to determine which stations can influence the value of a given grid box. This DLS is calculated from the e-folding distance of the individual station correlations. The DLS is calculated separately for five latitude bands, and then linearly interpolated between the bands. There is a noticeable difference in spatial coverage between the indices due to these differences in decorrelation length scales. This means that there will be some grid-box data where in fact there are no stations underlying it. Here we apply black borders to grid-boxes where at least 3 stations are present to denote greater confidence in representation of the wider grid-box area there. The land-sea mask enables the dataset to be used directly for model comparison with output from HadCM3. It does mean, however, that some coastal regions and islands over which one may expect to find a grid-box are in fact empty because they have been treated as sea

### **Data sources used for updates to the HadEX analysis of extremes**

We use a number of different data sources to provide sufficient coverage to update as many countries as possible to present day. These are summarised in Table 5. In building the new datasets we have tried to use exactly the same methodology as was used to create the original HadEX to retain consistency with a product that was created through substantial international effort and widely used, but there are some differences, which are described in the next section.

Wherever new data have been used, the geographical distributions of the trends were compared to those obtained from HadEX, using the same grid size, time span and fitting method. If the pattern of the trends in the temperature or precipitation indices did not match that from HadEX, we used the HadEX data despite its generally shorter time span. Differences in the patterns of the trends in the indices can arise because the individual stations used to create the gridded results are different from those in HadEX, and the quality

control procedures used are also very likely to be different. Countries where we decided to use HadEX data despite the existence of more recent data are Egypt and Turkey.

GHCND:

The Global Historical Climate Network Daily data has near-global coverage. However, to ensure consistency with the HadEX database, the GHCND stations were compared to those stations in HadEX. We selected those stations which are within 1500m of the stations used in the HadEX database and have a high correlation with the HadEX stations. We only took the precipitation data if its  $r > 0.9$  and the temperature data if one of its  $r$ -values  $> 0.9$ . In addition, we required at least 5 years of data beyond 2000. These daily data were then converted to the indices using the *fclimdex* software.

ECA&D and SACA&D:

The European Climate Assessment and Dataset and the Southeast Asian Climate Assessment and Dataset data are pre-calculated indices comprising the core 27 indices from the ETCCDI as well as some extra ones. We kindly acknowledge the help of Albert Klein Tank, the KNMI<sup>1</sup> and the BMKG<sup>2</sup> for their assistance in obtaining these data.

Mexico:

The station data from Mexico has been kindly supplied by the SMN<sup>3</sup> and Jorge Vazquez. These daily data were then converted to the required indices using the *Fclimdex* software. There are a total of 5298 Mexican stations in the database. In order to select those which have sufficiently long data records and are likely to be the most reliable ones we performed a cross correlation between all stations. We selected those which had at least 20 years of data post 1960 and have a correlation with at least one other station with an  $r$ -value  $> 0.95$ . This resulted in 237 stations being selected for further processing and analysis.

Indian Gridded:

The India Meteorological Department provided daily gridded data (precipitation 1951-2007, temperature 1969-2009) on a 1° x 1° grid. These are the only gridded daily data in our analysis. In order to process these in as similar a way as possible the values for each grid

---

<sup>1</sup> <http://www.knmi.nl>

<sup>2</sup> <http://www.bmkg.go.id>

<sup>3</sup> <http://www.smn.mx>

were assumed to be analogous to a station located at the centre of the grid. We keep these data separate from the rest of the study, which is particularly important when calculating the decorrelation length scale, which is on the whole larger for these gridded data.

Country	Region box (red dashed boxes in Fig. 1 and on each map at beginning of chapter)	Data source (T = temperature, P = precipitation)	Period of data coverage (T = temperature, P = precipitation)	Indices included (see Table 1 for details)	Temporal resolution available	Notes
Argentina	73.125 to 54.375 ° W, 21.25 to 56.25 ° S	Matlde Rusticucci (T,P)	1960-2010 (T,P)	TN10p, TN90p, TX10p, TX90p, PRCPTOT, CDD, CWD	annual	
Australia	114.375 to 155.625 °E, 11.25 to 43.75 °S	GHCND (T,P)	1960-2010 (T,P)	TN10p, TN90p, TX10p, TX90p, PRCPTOT, CDD, CWD	monthly, seasonal and annual	Land-sea mask has been adapted to include Tasmania and the area around Brisbane
Bangladesh	88.125 to 91.875 ° E, 21.25 to 26.25 ° N	Indian Gridded data (T,P)	1960-2007 (P), 1970-2009 (T)	TN10p, TN90p, TX10p, TX90p, PRCPTOT, CDD, CWD	monthly, seasonal and annual	Interpolated from Indian Gridded data
Brazil	73.125 to 31.875 ° W, 6.25 ° N to 33.75 ° S	HadEX (T,P)	1960-2000 (P) 2002 (T)	TN10p, TN90p, TX10p, TX90p, PRCPTOT, CDD, CWD	annual	Spatial coverage is poor
China	73.125 to 133.125 ° E, 21.25 to 53.75 ° N	GHCND (T,P)	1960-1997 (P) 1960-2003 (T <sub>min</sub> ) 1960-2010 (T <sub>max</sub> )	TN10p, TN90p, TX10p, TX90p, PRCPTOT, CDD, CWD	monthly, seasonal and annual	Precipitation has very poor coverage beyond 1997 except in 2003-04, and no data at all in 2000-02, 2005-11
Egypt	24.375 to 35.625 ° E, 21.25 to 31.25 ° N	HadEX (T,P)	No data	TN10p, TN90p, TX10p, TX90p, PRCPTOT,	annual	There are no data for Egypt so all grid-box values have been interpolated from stations in Jordan, Israel, Libya and Sudan
France	5.625 ° W to 9.375 °E, 41.25 to 51.25 °N	ECA&D (T,P)	1960-2010 (T,P)	TN10p, TN90p, TX10p, TX90p, PRCPTOT, CDD, CWD	monthly, seasonal and annual	



Germany	5.625 to 16.875 ° E, 46.25 to 56.25 ° N	ECA&D (T,P)	1960-2010 (T,P)	TN10p, TN90p, TX10p, TX90p, PRCPTOT, CDD, CWD	monthly, seasonal and annual	
India	69.375 to 99.375 ° E, 6.25 to 36.25 ° N	Indian Gridded data (T,P)	1960-2003 (P), 1970-2009 (T)	TN10p, TN90p, TX10p, TX90p, PRCPTOT, CDD, CWD	monthly, seasonal and annual	
Indonesia	95.625 to 140.625 ° E, 6.25 ° N to 11.25 ° S	HadEX (T,P)	1968-2003 (T,P)	TN10p, TN90p, TX10p, TX90p, PRCPTOT,	annual	Spatial coverage is poor
Italy	5.625 to 16.875 ° E, 36.25 to 46.25 ° N	ECA&D (T,P)	1960-2010 (T,P)	TN10p, TN90p, TX10p, TX90p, PRCPTOT, CDD, CWD	monthly, seasonal and annual	Land-sea mask has been adapted to improve coverage of Italy
Japan	129.375 to 144.375 ° E, 31.25 to 46.25 ° N	HadEX (P) GHCND (T)	1960-2003 (P) 1960-2000 (T <sub>min</sub> ) 1960-2010 (T <sub>max</sub> )	TN10p, TN90p, TX10p, TX90p, PRCPTOT,	monthly, seasonal and annual (T), annual (P)	
Kenya	31.875 to 43.125 ° E, 6.25 ° N to 6.25 ° S	HadEX (T,P)	1960-1999 (P)	TN10p, TN90p, TX10p, TX90p, PRCPTOT	annual	There are no temperature data for Kenya and so grid-box values have been interpolated from neighbouring Uganda and the United Republic of Tanzania. Regional averages include grid-boxes from outside Kenya that enable continuation to 2003
Mexico	118.125 to 88.125 ° W, 13.75 to 33.75 ° N	Raw station data from the Servicio Meteorológico Nacional (SMN) (T,P)	1960-2009 (T,P)	TN10p, TN90p, TX10p, TX90p, PRCPTOT, CDD, CWD	monthly, seasonal and annual	237/5298 stations selected. Non uniform spatial coverage. Drop in T and P coverage in 2009.
Peru	84.735 to 65.625 ° W, 1.25° N to 18.75 ° S	HadEX (T,P)	1960-2002 (T,P)	TN10p, TN90p, TX10p, TX90p, PRCPTOT, CDD, CWD	annual	Intermittent coverage in TX90p, CDD and CWD

Russia	West Russia 28.125 to 106.875 ° E, 43.75 to 78.75 ° N, East Russia 103.125 to 189.375 ° E, 43.75 to 78.75 ° N	ECA&D (T,P)	1960-2010 (T,P)	TN10p, TN90p, TX10p, TX90p, PRCPTOT, CDD, CWD	monthly, seasonal and annual	Country split for presentation purposes only.
Saudi Arabia	31.875 to 54.375 ° E, 16.25 to 33.75 ° N	HadEX (T,P)	1960-2000 (T,P)	TN10p, TN90p, TX10p, TX90p, PRCPTOT	annual	Spatial coverage is poor
South Africa	13.125 to 35.625 ° W, 21.25 to 36.25 ° S	HadEX (T,P)	1960-2000 (T,P)	TN10p, TN90p, TX10p, TX90p, PRCPTOT, CDD, CWD	annual	---
Republic of Korea	125.625 to 129.375 ° E, 33.75 to 38.75 ° N	HadEX (T,P)	1960-2003 (T,P)	TN10p, TN90p, TX10p, TX90p, PRCPTOT, CDD	annual	There are too few data points for CWD to calculate trends or regional timeseries
Spain	9.375 ° W to 1.875 ° E, 36.25 to 43.75 ° N	ECA&D (T,P)	1960-2010 (T,P)	TN10p, TN90p, TX10p, TX90p, PRCPTOT, CDD, CWD	monthly, seasonal and annual	
Turkey	24.375 to 46.875 ° E, 36.25 to 43.75 ° N	HadEX (T,P)	1960-2003 (T,P)	TN10p, TN90p, TX10p, TX90p, PRCPTOT, CDD, CWD	annual	Intermittent coverage in CWD and CDD with no regional average beyond 2000
United Kingdom	9.375 ° W to 1.875 ° E, 51.25 to 58.75 ° N	ECA&D (T,P)	1960-2010 (T,P)	TN10p, TN90p, TX10p, TX90p, PRCPTOT, CDD, CWD	monthly, seasonal and annual	
United States of America	125.625 to 65.625 ° W, 23.75 to 48.75 ° N	GHCND (T,P)	1960-2010 (T,P)	TN10p, TN90p, TX10p, TX90p, PRCPTOT, CDD, CWD	monthly, seasonal and annual	

**Table 5. Summary of data used for each country**

## **Quality control and gridding procedure used for updates to the HadEX analysis of extremes**

In order to perform some basic quality control checks on the index data, we used a two-step process on the indices. Firstly, internal checks were carried out, to remove cases where the 5 day rainfall value is less than the 1 day rainfall value, the minimum  $T_{\min}$  is greater than the minimum  $T_{\max}$  and the maximum  $T_{\min}$  is greater than the maximum  $T_{\max}$ .

Although these are physically impossible, they could arise from transcription errors when creating the daily dataset, for example, a misplaced minus sign, an extra digit appearing in the record or a column transposition during digitisation. During these tests we also require that there are at least 20 years of data in the period of record for the index for that station, and that some data is found in each decade between 1961 and 1990, to allow a reasonable estimation of the climatology over that period.

Weather conditions are often similar over many tens of kilometres and the indices calculated in this work are even more coherent. The correlation coefficient between each station-pair combination in all the data obtained is calculated for each index (and month where appropriate), and plotted as a function of the separation. An exponential decay curve is fitted to the data, and the distance at which this curve has fallen by a factor  $1/e$  is taken as the decorrelation length scale (DLS). A DLS is calculated for each dataset separately. For the GHCND, a separate DLS is calculated for each hemisphere. We do not force the fitted decay curve to show perfect correlation at zero distance, which is different to the method employed when creating HadEX. For some of the indices in some countries, no clear decay pattern was observed in some data sets or the decay was so slow that no value for the DLS could be determined. In these cases a default value of 200km was used.

We then perform external checks on the index data by comparing the value for each station with that of its neighbours. As the station values are correlated, it is therefore likely that if one station measures a high value for an index for a given month, its neighbours will also be measuring high. We exploit this coherence to find further bad values or stations as follows. Although raw precipitation data shows a high degree of localisation, using indices which have monthly or annual resolution improves the coherence across wider areas and so this neighbour checking technique is a valid method of finding anomalous stations.


We calculate a climatology for each station (and month if appropriate) using the mean value for each index over the period 1961-1990. The values for each station are then anomalised

using this climatology by subtracting this mean value from the true values, so that it is clear if the station values are higher or lower than normal. This means that we do not need to take differences in elevation or topography into account when comparing neighbours, as we are not comparing actual values, but rather deviations from the mean value.

All stations which are within the DLS distance are investigated and their anomalised values noted. We then calculate the weighted median value from these stations to take into account the decay in the correlation with increasing distance. We use the median to reduce the sensitivity to outliers.

If the station value is greater than 7.5 median-absolute-deviations away from the weighted median value (this corresponds to about 5 standard deviations if the distribution is Gaussian, but is a robust measure of the spread of the distribution), then there is low confidence in the veracity of this value and so it is removed from the data.

To present the data, the individual stations are gridded on a  $3.75^\circ \times 2.5^\circ$  grid, matching the output from HadCM3. To determine the value of each grid box, the DLS is used to calculate which stations can reasonably contribute to the value. The value of each station is then weighted using the DLS to obtain a final grid box value. At least three stations need to have valid data and be near enough (within 1 DLS of the gridbox centre) to contribute in order for a value to be calculated for the grid point. As for the original HadEX, the HadCM3 land-sea mask is used. However, in three cases the mask has been adjusted as there are data over Tasmania, eastern Australia and Italy that would not be included otherwise (Figure 6).



**Figure 6.** Land-sea mask used for gridding the station data and regional areas allocated to each country as described in Table 5.

## **Presentation of extremes of temperature and precipitation**



Indices are displayed as regional gridded maps of decadal trends and regional average time-series with decadal trends where appropriate. Trends are fitted using the median of pairwise slopes method (Sen 1968, Lanzante 1996). Trends are considered to be significantly different from a zero trend if the 5<sup>th</sup> to 95<sup>th</sup> percentiles of the pairwise slopes do not encompass zero. This is shown by a black dot in the centre of the grid-box or by a solid line on time-series plots. This infers that there is high confidence in the sign (positive or negative) of the sign. Confidence in the trend magnitude can be inferred by the spread of the 5<sup>th</sup> to 95<sup>th</sup> percentiles of the pairwise slopes which is given for the regional average decadal trends. Trends are only calculated when there are data present for at least 50% of years in the period of record and for the updated data (not HadEX) there must be at least one year in each decade.

Due to the practice of data-interpolation during the gridding stage (using the DLS) there are values for some grid boxes when no actual station lies within the grid box. There is more confidence in grid boxes for which there are underlying data. For this reason, we identify those grid boxes which contain at least 3 stations by a black contour line on the maps. The DLS differs with region, season and index which leads to large differences in the spatial coverage. The indices, by their nature of being largely threshold driven, can be intermittent over time which also effects spatial and temporal coverage (see Table 4).

Each index (and each month for the indices for which there is monthly data) has a different DLS, and so the coverage between different indices and datasets can be different. The restrictions on having at least 20 years of data present for each input station, at least 50% of years in the period of record and at least one year in each decade for the trending calculation, combined with the DLS, can restrict the coverage to only those regions with a dense station network reporting reliably.


Each country has a rectangular region assigned as shown by the red dashed box on the map in Figure 1 and listed in Table 2, which is used for the creation of the regional average. This is sometimes identical to the attribution region shown in grey on the map in Figure 1. This region is again shown on the maps accompanying the time series of the regional averages as a reminder of the region and grid boxes used in the calculation. Regional averages are created by weighting grid box values by the cosine of their grid box centre latitude. To ensure consistency over time a regional average is only calculated when there are a sufficient number of grid boxes present. The full-period median number of grid-boxes present is calculated. For regions with a median of more than six grid-boxes there must be at

least 80% of the median number of grid boxes present for any one year to calculate a regional average. For regions with six or fewer median grid boxes this is relaxed to 50%. These limitations ensure that a single station or grid box which has a longer period of record than its neighbours cannot skew the timeseries trend. So sometimes there may be grid-boxes present but no regional average time series. The trends for the regional averages are calculated in the same way as for the individual grid boxes, using the median of pairwise slopes method (Sen 1968, Lanzante 1996). Confidence in the trend is also determined if the 5<sup>th</sup> to 95<sup>th</sup> percentiles of the pairwise slopes are of the same sign and thus inconsistent with a zero trend. As well as the trend in quantity per decade, we also show the full change in the quantity from 1960 to 2010 that this fitted linear trend implies.



Monthly: 2.20% per decade (1.80 to 2.61)  
Total change of 11.02% from 1960 to 2011 (9.00% to 13.06%)

Annual: 2.28% per decade (1.69 to 2.86)  
Total change of 11.41% from 1960 to 2010 (8.43% to 14.28%)



**Figure 7.** Examples of the plots shown in the data section. Left: From ECA&D data between 1960-2010 for the number of warm nights, and Right: from HadEX data (1960-2003) for the total precipitation. A full explanation of the plots is given in the text below.

The results are presented in the form of a map and a time series for each country and index. The map shows the grid box decadal trend in the index over the period for which there are data. High confidence, as determined above, is shown by a black dot in the grid box centre. To show the variation over time, the values for each year (and month if available) are shown



in a time series for a regional average. The values of the indices have been normalised to a base period of 1961-1990 (except the Indian gridded data which use a 1971 to 1990 period), both in HadEX and in the new data acquired for this project. Therefore, for example, the percentage of nights exceeding the 90<sup>th</sup> percentile for a temperature is 10% for that period.

There are two influences on whether a grid box contains a value or not – the land-sea mask, and the decorrelation length scale. The land-sea mask is shown in Figure 6. There are grid boxes which contain some land but are mostly sea and so are not considered. The decorrelation length scale sets the maximum distance a grid box can be from stations before no value is assigned to it. Grid boxes containing three or more stations are highlighted by a thick border. This indicates regions where the value shown is likely to be more representative of the grid box area mean as opposed to a single station location.

On the maps for the new data there is a box indicating which grid boxes have been extracted to calculate the area average for the time series. This box is the same as shown in Figure 1 at the beginning of each country's document. These selected grid boxes are combined using area (cosine) weighting to calculate the regional average (both annual [thick lines] and monthly [thin lines] where available). Monthly (orange) and annual (blue) trends are fitted to these time series using the method described above. The decadal trend and total change over the period where there are data are shown with 5th to 95th percentile confidence intervals in parentheses. High confidence, as determined above, is shown by a solid line as opposed to a dotted one. The green vertical lines on the time series show the dates of some of the notable events outlined in each section.


## **Attribution**

Regional distributions of seasonal mean temperatures in the 2000s are computed with and without the effect of anthropogenic influences on the climate. The analysis considers temperatures averaged over the regions shown in Figure 8. These are also identified as grey boxes on the maps in Figure 1. The coordinates of the regions are given in Table 6. The methodology combines information from observations and model simulations using the approach originally introduced in Christidis et al., 2010 and later extended in Christidis et al., 2011, where more details can be found. The analysis requires spatial scales greater than about 2,500km and for that reason the selected regions (Fig.8 and Table 6) are often larger

than individual countries, or include several smaller countries in a single region (for example UK, Germany and France are grouped in one region).

Observations of land temperature come from the CRUTEM3 gridded dataset (Brohan et al., 2006) and model simulations from two coupled GCMs, namely the Hadley Centre HadGEM1 model (Martin et al., 2006) and version 3.2 of the MIROC model (K-1 Developers, 2004). The use of two GCMs helps investigate the sensitivity of the results to the model used in the analysis. Ensembles of model simulations from two types of experiments are used to partition the temperature response to external forcings between its anthropogenic and natural components. The first experiment (ALL) simulates the combined effect of natural and anthropogenic forcings on the climate system and the second (ANTHRO) includes anthropogenic forcings only. The difference of the two gives an estimate of the effect of the natural forcings (NAT). Estimates of the effect of internal climate variability are derived from long control simulations of the unforced climate. Distributions of the regional summer mean temperature are computed as follows:

- a) A global optimal fingerprinting analysis (Allen and Tett, 1999; Allen and Stott, 2003) is first carried out that scales the global simulated patterns (fingerprints) of climate change attributed to different combinations of external forcings to best match them to the observations. The uncertainty in the scaling that originates from internal variability leads to samples of the scaled fingerprints, i.e. several realisations that are plausibly consistent with the observations. The 2000-2009 decade is then extracted from the scaled patterns and two samples of the decadal mean temperature averaged over the reference region are then computed with and without human influences, which provide the Probability Density Functions (PDFs) of the decadal mean temperature attributable to ALL and NAT forcings.
- b) Model-derived estimates of noise are added to the distributions to take into account the uncertainty in the simulated fingerprints.
- c) In the same way, additional noise from control model simulations is introduced to the distributions to represent the effect of internal variability in the annual values of the seasonal mean temperatures. The result is a pair of estimated distributions of the annual values of the seasonal mean temperature in the region with and without the effect of human activity on the climate. The temperatures throughout the analysis are expressed as anomalies relative to period 1961-1990.



**Figure 8.** The regions used in the attribution analysis. Regions marked with dashed orange boundaries correspond to non-G20 countries that were also included in the analysis.

Region	Region Coordinates
Argentina	74-58W, 55-23S
Australia	110-160E, 47-10S
Bangladesh	80-100E, 10-35N
Brazil	73-35W, 30S-5N
Canada-Alaska	170-55W, 47-75N
China	75-133E, 18-50N
Egypt	18-40E, 15-35N
France-Germany-UK	10W-20E, 40-60N
India	64-93E, 7-40N
Indonesia	90-143E, 14S-13N
Italy-Spain	9W-20E, 35-50N
Japan-Republic of Korea	122-150E, 30-48N
Kenya	35-45E, 10S-10N
Mexico	120-85W, 15-35N
Peru	85-65W, 20-0S
Russia	30-185E, 45-78N
Saudi Arabia	35-55E, 15-31N
South Africa	10-40E, 35-20S
Turkey	18-46E, 32-45N

**Table 6.** The coordinates of the regions used in the attribution analysis.

## References

ALEXANDER, L.V. and D.E. PARKER. 2004. Europe in State of the Climate in 2003. *Bulletin of the American Meteorological Society* 85, S46.

ALEXANDER, L. V., ZHANG. X., PETERSON, T. C., CAESAR, J., GLEASON, B., KLEIN TANK, A. M. G., HAYLOCK, M., COLLINS, D., TREWIN, B., RAHIMZADEH, F., TAGIPOUR, A., RUPA KUMAR, K., REVADEKAR, J., GRIFFITHS, G., VINCENT, L., STEPHENSON, D. B., BURN, J., AGUILAR, E., BRUNET, M., TAYLOR, M., NEW, M., ZHAI, P., RUSTICUCCI, M. and VAZQUEZ-AGUIRRE, J. L. 2006. Global observed changes in daily climate extremes of temperature and precipitation. *J. Geophys. Res.* 111, D05109. doi:10.1029/2005JD006290.

ALLEN, M. R., TETT S. F. B. 1999. Checking for model consistency in optimal fingerprinting. *Clim Dyn* 15: 419-434.

ALLEN M. R., STOTT P. A. 2003. Estimating signal amplitudes in optimal fingerprinting, part I: theory. *Clim Dyn* 21: 477-491.

BBC NEWS. 2010. Deadly flash floods hit southern France. Available online: <http://www.bbc.co.uk/news/10327034> accessed on 14 October 2011.

BELL, G. & T. EICHLER. 2004. Summer Heat Wave in State of the Climate 2003. *Bulletin of the American Meteorological Society* 85, S46-48.

BROHAN, P., KENNEDY, J.J., HARRIS, I., TETT, S.F.B. and JONES, P.D. 2006. Uncertainty estimates in regional and global observed temperature changes: a new dataset from 1850. *J. Geophys. Res.* 111, D12106. doi:10.1029/2005JD006548.

CHRISTIDIS N., STOTT. P A., ZWIERS, F. W., SHIOGAMA, H., NOZAWA, T. 2010. Probabilistic estimates of recent changes in temperature: a multi-scale attribution analysis. *Clim Dyn* 34: 1139-1156.

CHRISTIDIS, N., STOTT, P. A., ZWIERS, F. W., SHIOGAMA, H., NOZAWA, T. 2011. The contribution of anthropogenic forcings to regional changes in temperature during the last decade. *Climate Dynamics* in press.

COMPO, G. P., J.S. WHITAKER, P.D. SARDESHMUKH, N. MATSUI, R.J. ALLAN, X. YIN, B.E. GLEASON, R.S. VOSE, G. RUTLEDGE, P. BESSEMOULIN, S. BRÖNNIMANN, M.

BRUNET, R.I. CROUTHAMEL, A.N. GRANT, P.Y. GROISMAN, P.D. JONES, M.C. KRUK, A.C. KRUGER, G.J. MARSHALL, M. MAUGERI, H.Y. MOK, Ø. NORDLI, T.F. ROSS, R.M. TRIGO, X.L. WANG, S.D. WOODRUFF and S.J. WORLEY. 2011. The Twentieth Century Reanalysis Project, *Q. J. R.Met.S.* 137, 1-28, doi: 10.1002/qj.776.

CORNES, R. C., and P. D. JONES. 2011. An examination of storm activity in the northeast Atlantic region over the 1851–2003 period using the EMULATE gridded MSLP data series. *J. Geophys. Res.* 116, D16110, doi:10.1029/2011JD016007.

K-1 MODEL DEVELOPERS (2004) K-1 coupled GCM (MIROC) description, K-1 Tech Rep, H Hasumi and S Emori (eds), Centre for Clim Sys Res, Univ of Tokyo.

KLEIN TANK, A.M.G. et al. 2002. Daily dataset of 20th-century surface air temperature and precipitation series for the European Climate Assessment. *Int. J. of Climatol.* 22, 1441-1453.

LANZANTE, J. R. 1996. Resistant, robust and non-parametric techniques for the analysis of climate data: theory and examples, including applications to historical radiosonde station data. *Int. J. Clim.* 16, 1197–226.

LEVINSON, D.H. and A. M. WAPLE, eds. 2004. State of the Climate in 2003. *Bulletin of the American Meteorological Society* 85, S1-72.

MAIER, F., A. OBREGÓN, P. BISSOLLI, J. J. KENNEDY, and D. E. PARKER. 2011. Central and Western Europe in State of the Climate in 2010. *Bulletin of the American Meteorological Society* 92, S203.

MARTIN G.M., RINGER. M. A., POPE V. D., JONES, A., DEARDEN, C., HINTON, T. 2006. The physical properties of the atmosphere in the new Hadley Centre Global Environmental Model (HadGEM1). Part I: Model description and global climatology. *J Clim* 19: 1274-1301.

METEOFRACTANCE. 2010. Catastrophe de Draguignan. Available online: <http://pluiesextremes.meteo.fr/2010-06-15/catastrophe-de-draguignan.html> accessed 4 August 2011.

OBREGÓN, A., P. BISSOLLI, and J. J. KENNEDY. 2007. WESTERN AND NORTHWESTERN EUROPE Europe in State of the Climate in 2006. *Bulletin of the American Meteorological Society* 88, S109.

- OBREGÓN, A., P. BISSOLLI, J. J. KENNEDY, and D.E. PARKER. 2010. Central and Western Europe in State of the Climate in 2009. *Bulletin of the American Meteorological Society* 91, S163.
- PETERSON, T.C., VAUTARD, R., McVICAR, T.R., THÉPAUT, J-N., and BERRISFORD, P. 2011. [Global Climate] Surface Winds over Land in State of the Climate 2010. *Bulletin of the American Meteorological Society* 92 (6), S57.
- SANCHEZ-LUGO, A., KENNEDY, J.J. and BERRISFORD, P. 2011. [Global Climate] Surface Temperatures in State of the Climate 2010. *Bulletin of the American Meteorological Society* 92 (6), S36-S37.
- SEN, P. K. 1968. Estimates of the regression coefficient based on Kendall's tau. *J. Am. Stat. Assoc.* 63, 1379–89.
- TRIGO, R., R. GARCIA-HERRERA and D. PAREDES. 2006. Iberia in State of the Climate in 2005. *Bulletin of the American Meteorological Society* 87, S84.
- USA TODAY. 2006. 64 people in France dead from heat wave. Available online: [http://www.usatoday.com/weather/news/2006-07-26-france-heat-wave\\_x.htm](http://www.usatoday.com/weather/news/2006-07-26-france-heat-wave_x.htm) accessed on 14 October 2011.
- WAPLE, A. M., J. H. LAWRIMORE, M. S. HALPERT, G. D. BELL, W. HIGGINS, B. LYON, M. J. MENNE, K. L. GLEASON, R. C. SCHNELL, J. R. CHRISTY, W. THIAW, W. J. WRIGHT, M. J. SALINGER, L. ALEXANDER, R. S. STONE, and S. J. CAMARGO. 2002. Europe in State of the Climate in 2001. *Bulletin of the American Meteorological Society* 83, S45-46.
- WMO WORLD METEOROLOGICAL ORGANIZATION. 2001. Statement on Status of the Global Climate in 2000, WMO-No. 920.  
[http://www.wmo.int/pages/prog/wcp/wcdmp/statement/wmostatement\\_en.html](http://www.wmo.int/pages/prog/wcp/wcdmp/statement/wmostatement_en.html)
- WMO WORLD METEOROLOGICAL ORGANIZATION. 2004. Statement on Status of the Global Climate in 2003, WMO-No. 966.  
[http://www.wmo.int/pages/prog/wcp/wcdmp/statement/wmostatement\\_en.html](http://www.wmo.int/pages/prog/wcp/wcdmp/statement/wmostatement_en.html)
- WMO WORLD METEOROLOGICAL ORGANIZATION. 2009. Statement on Status of the Global Climate in 2008, WMO-No. 1039.  
[http://www.wmo.int/pages/prog/wcp/wcdmp/statement/wmostatement\\_en.html](http://www.wmo.int/pages/prog/wcp/wcdmp/statement/wmostatement_en.html)

WMO WORLD METEOROLOGICAL ORGANIZATION. 2011. Statement on Status of the Global Climate in 2010, WMO-No. 1074.

[http://www.wmo.int/pages/prog/wcp/wcdmp/statement/wmostatement\\_en.html](http://www.wmo.int/pages/prog/wcp/wcdmp/statement/wmostatement_en.html)

# Acknowledgements

Data for this work were taken from the ECA&D dataset (Klein Tank et al. 2002). We thank Lisa Alexander and Markus Donat (University of New South Wales) for their help and advice. We also thank reviewers from France for their valuable input and advice.

②

②

②

②

②



?

?

?

## **Chapter 2 – Climate Change Projections**

?

## Introduction

Climate models are used to understand how the climate will evolve over time and typically represent the atmosphere, ocean, land surface, cryosphere, and biogeochemical processes, and solve the equations governing their evolution on a geographical grid covering the globe. Some processes are represented explicitly within climate models, large-scale circulations for instance, while others are represented by simplified parameterisations. The use of these parameterisations is sometimes due to processes taking place on scales smaller than the typical grid size of a climate model (a Global Climate Model (GCM) has a typical horizontal resolution of between 250 and 600km) or sometimes to the current limited understanding of these processes. Different climate modelling institutions use different plausible representations of the climate system, which is why climate projections for a single greenhouse gas emissions scenario differ between modelling institutes. This gives rise to “climate model structural uncertainty”.

In response to a proposed activity of the World Climate Research Programme's (WCRP's; <http://www.wcrp-climate.org/>) Working Group on Coupled Modelling (WGCM), the Program for Climate Model Diagnosis and Intercomparison (PCMDI; <http://www.pcmdi.llnl.gov/>) volunteered to collect model output contributed by leading climate modelling centres around the world. Climate model output from simulations of the past, present and future climate was collected by PCMDI mostly during the years 2005 and 2006, and this archived data constitutes phase 3 of the Coupled Model Intercomparison Project (CMIP3). In part, the WGCM organised this activity to enable those outside the major modelling centres to perform research of relevance to climate scientists preparing the IPCC Fourth Assessment Report (AR4). This unprecedented collection of recent model output is commonly known as the “CMIP3 multi-model dataset”. The GCMs included in this dataset are referred to regularly throughout this review, although not exclusively.

The CMIP3 multi-model ensemble has been widely used in studies of regional climate change and associated impacts. Each of the constituent models was subject to extensive testing by the contributing institute, and the ensemble has the advantage of having been constructed from a large pool of alternative model components, therefore sampling alternative structural assumptions in how best to represent the physical climate system. Being assembled on an opportunity basis, however, the CMIP3 ensemble was not designed to represent model uncertainties in a systematic manner, so it does not, in isolation, support robust estimates of the risk of different levels of future climate change, especially at a regional level.

Since CMIP3, a new (CMIP5) generation of coupled ocean-atmosphere models has been developed, which is only just beginning to be available and is being used for new projections for the IPCC Fifth Assessment Report (AR5).


These newer models typically feature higher spatial resolution than their CMIP3 counterparts, including in some models a more realistic representation of stratosphere-troposphere interactions. The CMIP5 models also benefit from several years of development in their parameterisations of small scale processes, which, together with resolution increases, are expected to result in a general improvement in the accuracy of their simulations of historical climate, and in the credibility of their projections of future changes. The CMIP5 programme also includes a number of comprehensive Earth System Models (ESMs) which explicitly simulate the earth's carbon cycle and key aspects of atmospheric chemistry, and also contain more sophisticated representations of aerosols compared to CMIP3 models.

The CMIP3 results should be interpreted as a useful interim set of plausible outcomes. However, their neglect of uncertainties, for instance in carbon cycle feedbacks, implies that higher levels of warming outside the CMIP3 envelope cannot be ruled out. In future, CMIP5 coupled model and ESM projections can be expected to produce improved advice on future regional changes. In particular, ensembles of ESM projections will be needed to provide a more comprehensive survey of possible future changes and their relative likelihoods of occurrence. This is likely to require analysis of the CMIP5 multi-model ESM projections, augmented by larger ensembles of ESM simulations in which uncertainties in physical and biogeochemical feedback processes can be explored more systematically, for example via ensembles of model runs in which key aspects of the climate model are slightly adjusted. Note that such an exercise might lead to the specification of wider rather than narrower uncertainties compared to CMIP3 results, if the effects of representing a wider range of earth system processes outweigh the effects of refinements in the simulation of physical atmosphere-ocean processes already included in the CMIP3 models.


# Climate projections

The Met Office Hadley Centre is currently producing perturbed parameter ensembles of a single model configuration known as HadCM3C, to explore uncertainties in physical and biogeochemical feedback processes. The results of this analysis will become available in the next year and will supplement the CMIP5 multi-model ESM projections, providing a more comprehensive set of data to help progress understanding of future climate change.

However, many of the studies covered in the chapter on climate impacts have used CMIP3 model output. For this reason, and because it is still the most widely used set of projections available, the CMIP3 ensemble output for temperature and precipitation, for the A1B emission scenario, for France and the surrounding region is shown below.



**Figure 1.** Percentage change in average annual temperature by 2100 from 1960-1990 baseline climate, averaged over 21 CMIP3 models. The size of each pixel represents the level of agreement between models on the magnitude of the change.



**Figure 2.** Percentage change in average annual precipitation by 2100 from 1960-1990 baseline climate, averaged over 21 CMIP3 models. The size of each pixel represents the level of agreement between models on the sign of the change.

## Summary of temperature change in France

Figure 1 shows the percentage change in average annual temperature by 2100 from 1960-1990 baseline climate, averaged over 21 CMIP3 models. All of the models in the CMIP3 ensemble project increased temperatures in the future, but the size of each pixel indicates how well the models agree over the magnitude of the increase.

Projected increases in temperature over France are generally between 2.5°C and 3.5°C. Increases are lower in the north, and higher in southern regions. The agreement between models is generally moderate over most of the country.

## Summary of precipitation change in France

Figure 2 shows the percentage change in average annual precipitation by 2100 from 1960-1990 baseline climate, averaged over 21 CMIP3 models. Unlike for temperature, the models sometimes disagree over whether precipitation is increasing or decreasing over a region, so

in this case the size of each pixel indicates the percentage of the models in the ensemble that agree on the sign of the change in precipitation.

Europe shows a strong contrast in projected precipitation changes, with large decreases in the south and large increases in the north. France falls towards the southern region with decreasing precipitation, with projected decreases of up to 20% in the southwest, but with smaller decreases of up to 5% further north. There is generally moderate agreement between ensemble members over central and southern France, and low agreement in the north, indicating uncertainty in the position of the transition zone between increasing and decreasing precipitation over Europe.

□

2

## **Chapter 3 – Climate Change Impact Projections**

# Introduction

## Aims and approach

This chapter looks at research on a range of projected climate change impacts, with focus on results for France. It includes projections taken from the AVOID programme, for some of the impact sectors.

The aim of this work is to take a 'top down' approach to assessing global impacts studies, both from the literature and from new research undertaken by the AVOID programme. This project covers 23 countries, with summaries from global studies provided for each of these. This global approach allows some level of comparison between countries, whilst presenting information on a scale most meaningful to inform international policy.

The literature covered in this chapter focuses on research published since the Fourth Assessment Report (AR4) of the Intergovernmental Panel on Climate Change (IPCC) and should be read in conjunction with IPCC AR4 WG1 and WG2 reports. For some sectors considered, an absence of research developments since the IPCC AR4, means earlier work is cited as this helps describe the current level of scientific understanding. This report focuses on assessing scientific research about climate change impacts within sectors; it does not present an integrated analysis of climate change adaptation policies.

Some national and sub-national scale literature is reported to a limited extent to provide some regional context.

## Impact sectors considered and methods

This report reviews the evidence for the impact of climate change on a number of sectors, for France. The following sectors are considered in turn in this report:

- Crop yields
- Food security
- Water stress and drought
- Pluvial flooding and rainfall
- Fluvial flooding



- Tropical cyclones (where applicable)
- Coastal regions

## **Supporting literature**

Literature searches were conducted for each sector with the Thomson Reuters Web of Science (WoS., 2011) and Google Scholar academic search engines respectively. Furthermore, climate change impact experts from each of the 23 countries reviewed were contacted. These experts were selected through a combination of government nomination and from experts known to the Met Office. They were asked to provide literature that they felt would be of relevance to this review. Where appropriate, such evidence has been included. A wide range of evidence was considered, including; research from international peer-reviewed journal papers; reports from governments, non-governmental organisations, and private businesses (e.g. reinsurance companies), and research papers published in national journals.

For each impact sector, results from assessments that include a global- or regional-scale perspective are considered separately from research that has been conducted at the national- or sub-national-scale. The consideration of global- and regional-scale studies facilitates a comparison of impacts across different countries, because such studies apply a consistent methodology for each country. While results from national- and sub-national-scale studies are not easily comparable between countries, they can provide a level of detail that is not always possible with larger-scale studies. However, the national- and sub-national scale literature included in this project does not represent a comprehensive coverage of regional-based research and cannot, and should not, replace individual, detailed impacts studies in countries. The review aims to present an up-to-date assessment of the impact of climate change on each of the sectors considered.

## **AVOID programme results**

Much of the work in this report is drawn from modelling results and analyses coming out of the AVOID programme. The AVOID programme is a research consortium funded by DECC and Defra and led by the UK Met Office and also comprises the Walker Institute at the University of Reading, the Tyndall Centre represented through the University of East Anglia,

and the Grantham Institute for Climate Change at Imperial College. The expertise in the AVOID programme includes climate change research and modelling, climate change impacts in natural and human systems, socio-economic sciences, mitigation and technology. The unique expertise of the programme is in bringing these research areas together to produce integrated and policy-relevant results. The experts who work within the programme were also well suited to review the literature assessment part of this report. In this report the modelling of sea level rise impacts was carried out for the AVOID programme by the University of Southampton.

The AVOID programme uses the same emissions scenarios across the different impact sectors studied. These are a business as usual (IPCC SRES A1B) and an aggressive mitigation (the AVOID A1B-2016-5-L) scenario. Model output for both scenarios was taken from more than 20 GCMs and averaged for use in the impact models. The impact models are sector specific, and frequently employ further analytical techniques such as pattern scaling and downscaling in the crop yield models.

Data and analysis from AVOID programme research is provided for the following impact sectors:

- Crop yields
- Water stress and drought
- Fluvial flooding
- Coastal regions

## **Uncertainty in climate change impact assessment**

There are many uncertainties in future projections of climate change and its impacts. Several of these are well-recognised, but some are not. One category of uncertainty arises because we don't yet know how mankind will alter the climate in the future. For instance, uncertainties in future greenhouse gas emissions depends on the future socio-economic pathway, which, in turn, depends on factors such as population, economic growth, technology development, energy demand and methods of supply, and land use. The usual approach to dealing with this is to consider a range of possible future scenarios.

Another category of uncertainties relate to our incomplete understanding of the climate system, or an inability to adequately model some aspects of the system. This includes:

- Uncertainties in translating emissions of greenhouse gases into atmospheric concentrations and radiative forcing. Atmospheric CO<sub>2</sub> concentrations are currently rising at approximately 50% of the rate of anthropogenic emissions, with the remaining 50% being offset by a net uptake of CO<sub>2</sub> into the oceans and land biosphere. However, this rate of uptake itself probably depends on climate, and evidence suggests it may weaken under a warming climate, causing the CO<sub>2</sub> rise to be larger proportion of emissions. The extent of this feedback is highly uncertain, but it not considered in most studies. The 3<sup>rd</sup> Coupled Model Intercomparison Project (CMIP3), which provided the future climate projections for the IPCC 4<sup>th</sup> Assessment Report, used a single estimate of CO<sub>2</sub> concentration rise for each emissions scenario, so the CMIP3 projections (which were used in most studies presented here, including AVOID) do not account for this uncertainty.
- Uncertainty in climate response to the forcing by greenhouse gases and aerosols. One aspect of this is the response of global mean temperature (“climate sensitivity”), but a more relevant aspect for impacts studies is the response of regional climates, including temperature, precipitation and other meteorological variables. Different climate models can give very different results in some regions, while giving similar results in other regions. Confidence in regional projections requires more than just agreement between models: physical understanding of the relevant atmospheric, ocean and land surface processes is also important, to establish whether the models are likely to be realistic.
- Additional forcings of regional climate. Greenhouse gas changes are not the only anthropogenic driver of climate change; atmospheric aerosols and land cover change are also important, and unlike greenhouse gases, the strength of their influence varies significantly from place to place. The CMIP3 models used in most impacts studies generally account for aerosols but not land cover change.
- Uncertainty in impacts processes. The consequences of a given changes in weather or climatic conditions for biophysical impacts such as river flows, drought, flooding, crop yield or ecosystem distribution and functioning depend on many other processes which are often poorly-understood, especially at large scales. In particular, the extent to which different biophysical impacts interact with each other has been hardly studied, but may be crucial; for example, impacts of climate change on crop yield may depend not only on local climate changes affecting rain-fed crops, but also remote climate changes affecting river flows providing water for irrigation.


- Uncertainties in non-climate effects of some greenhouse gases. As well as being a greenhouse gas, CO<sub>2</sub> exerts physiological influences on plants, affecting photosynthesis and transpiration. Under higher CO<sub>2</sub> concentrations, and with no other limiting factors, photosynthesis can increase, while the requirements of water for transpiration can decrease. However, while this has been extensively studied under experimental conditions, including in some cases in the free atmosphere, the extent to which the ongoing rise in ambient CO<sub>2</sub> affects crop yields and natural vegetation functioning remains uncertain and controversial. Many impacts projections assume CO<sub>2</sub> physiological effects to be significant, while others assume it to be non-existent. Studies of climate change impacts on crops and ecosystems should therefore be examined with care to establish which assumptions have been made.

In addition to these uncertainties, the climate varies significantly through natural processes from year-to-year and also decade-to-decade, and this variability can be significant in comparison to anthropogenic forcings on shorter timescales (the next few decades) particularly at regional scales. Whilst we can characterise the natural variability it will not be possible to give a precise forecast for a particular year decades into the future.

A further category of uncertainty in projections arises as a result of using different methods to correct for uncertainties and limitations in climate models. Despite being painstakingly developed in order to represent current climate as closely as possible, current climate models are nevertheless subject to systematic errors such as simulating too little or too much rainfall in some regions. In order to reduce the impact of these, '*bias correction*' techniques are often employed, in which the climate model as a source of information on the *change* in climate which is then applied to the observed present-day climate state (rather than using the model's own simulation of the present-day state). However, these bias-corrections typically introduce their own uncertainties and errors, and can lead to inconsistencies between the projected impacts and the driving climate change (such as river flows changing by an amount which is not matched by the original change in precipitation). Currently, this source of uncertainty is rarely considered

When climate change projections from climate models are applied to climate change impact models (e.g. a global hydrological model), the climate model structural uncertainty carries through to the impact estimates. Additional uncertainties include changes in future emissions and population, as well as parameterisations within the impact models (this is rarely considered). Figure 1 highlights the importance of considering climate model structural

uncertainty in climate change impacts assessment. Figure 1 shows that for 2°C prescribed global-mean warming, the magnitude of, and sign of change in average annual runoff from present, simulated by an impacts model, can differ depending upon the GCM that provides the climate change projections that drive the impact model. This example also shows that the choice of impact model, in this case a global hydrological model (GHM) or catchment-scale hydrological model (CHM), can affect the magnitude of impact and sign of change from present (e.g. see IPSL CM4 and MPI ECHAM5 simulations for the Xiangxi). To this end, throughout this review, the number of climate models applied in each study reviewed, and the other sources of uncertainty (e.g. emissions scenarios) are noted. Very few studies consider the application of multiple impacts models and it is recommended that future studies address this.



**Figure 1.** Change in average annual runoff relative to present (vertical axis; %), when a global hydrological model (GHM) and a catchment-scale hydrological model (CHM) are driven with climate change projections from 7 GCMs (horizontal axis), under a 2°C prescribed global-mean warming scenario, for six river catchments. The figure is from Gosling et al. (2011).

Uncertainties in the large scale climate relevant to France include the Atlantic Ocean has a Meridional Overturning Circulation (MOC) which transports large amounts of heat northwards in the Atlantic from the Equator. A key part of this is called the thermohaline circulation (THC). Disruption of the MOC could have a major impact on the Northern Hemisphere climate, including that of France, with likely detrimental impacts on human and animal systems. The IPCC AR4 concluded that "... it is very likely that the Atlantic Ocean Meridional Overturning Circulation could slow down during the course of the 21st century. A multi-model ensemble shows an average reduction of 25% with a broad range from virtually no change to a reduction of over 50% averaged over 2080 to 2099" (IPCC, 2007a).

Schneider et al. (2007) analysed simulations from several GCMs that were reviewed in the IPCC AR4 and found that projections of MOC change indicate it may weaken by 25-30% by the year 2100. Recent monitoring (Cunningham et al., 2007, Kanzow et al., 2007) has revealed large variability in the strength of the MOC on daily to seasonal timescales. This significant variability casts doubt on a previous report of decreases in MOC transport from several hydrographic sections (Bryden et al., 2005), although it does not explain the observed water mass changes below 3000m. Recent results based on radar altimeter and Argo data also suggest that there has been no slowdown, at least over the altimeter era (1993-present) (Willis, 2010). In contrast, two ocean state estimation studies (Balmaseda et al., 2007, Wunsch and Heimbach, 2006) indicated an MOC slow down. It has been suggested, based on model studies, that anthropogenic aerosols have slowed the weakening of the MOC and such weakening could only become significant several decades into the 21<sup>st</sup> century (Delworth and Dixon, 2006).

Regarding the possibility of MOC shutdown, a recent study presented by Swingedouw et al. (2007) with one climate model found that additional melt from Greenland could lead to complete AMOC shutdown in a CO<sub>2</sub> stabilisation experiment. However, a previous study with a different model (Ridley et al., 2005) found no effect from similar levels of meltwater input. Mikolajewicz et al. (2007) coupled an earth system model with atmospheric and ocean GCMs and observed a complete shutdown of the AMOC under a high emission scenario (SRES A2), but not before 2100. Moreover, Mikolajewicz et al. (2007) observed only a temporary weakening of the deep water formation in the North Atlantic by 2100 under a low emission scenario (B1).

Reversibility following AMOC shutdown is a key issue. Hofmann and Rahmstorf (2009) showed that hysteresis still occurs in a new low-diffusivity model. This is contrary to previous theoretical arguments that hysteresis is a product of diffusivity of the low-resolution simplified ocean models which are applied to perform the long-term simulations that are required to investigate this issue.

There is some new work on the impacts of AMOC weakening. Two studies (Kuhlbrodt et al., 2009, Vellinga and Wood, 2008) found SLR of several tens of cm along parts of the North Atlantic coast. They studies found that regional cooling could partially offset the greenhouse gas warming, and various other impacts may be substantial but hard to quantify such as change in tropical precipitation patterns and change in ocean currents leading to declining fish stocks and ecosystems (Schmittner, 2005).

In conclusion, large uncertainty remains in the probability of a complete MOC shutdown (Kriegler et al., 2009, Zickfeld et al., 2007). However, for the high temperature scenario considered by a recent expert elicitation exercise (centred on 4.5°C by 2100, 6.5°C by 2200) (Kriegler et al., 2009), the probability of complete shutdown was assessed to be at least 10% (according to several experts). Comparable results were found by the exercise reported by Zickfeld et al. (2007). To this end, it is thought unlikely that the AMOC could significantly weaken with 2°C global-mean warming.

□






# Summary of findings for each sector

## Crop yields

- Quantitative crop yield projections under climate change scenarios for France vary across studies due to the application of different models, assumptions and emissions scenarios.
- However, global- and regional-scale studies generally project yield gains for wheat and yield losses for maize, two of France's major crops, with climate change.
- National- and sub-national-scale studies included here also concur with the results from global-scale studies. In addition they note that repeated dry spells during the growing season may have a large impact on yields, although how the frequency of such events may be affected by climate change remains poorly understood.
- Important knowledge gaps and key uncertainties include the quantification of yield increases due to CO<sub>2</sub> fertilisation, quantification of yield reductions due to ozone damage and the extent to which crop diseases might affect crop yields with climate change.

## Food security

-  France is currently a country with extremely low levels of undernourishment. Global-scale studies included here generally project that France will not face serious food security issues over the next 40 years as a consequence of climate change.
- Recent research by the AVOID programme and others demonstrates that adaptive measures will be key to ensuring France's food security under climate change.
- One study concluded that the national economy of France presents a very low vulnerability to climate change impacts on fisheries by the 2050s.

## **Water stress and drought**

- The majority of global and regional assessments included here project that drought could become more frequent in France with climate change, and that water stress could increase.
- National-scale studies included here suggest that agricultural drought could increase in duration and intensity with climate change, to an extent never observed in the historical record. However there remains uncertainty in the magnitude to which such changes may be realised.
- Recent simulations by the AVOID programme project a median increase of 12% in the population may be exposed to increased water stress by 2100 under the A1B emissions scenario.

## **Pluvial flooding and rainfall**

- Recent modelling studies confirm conclusions from the IPCC AR4 that heavy winter precipitation and flooding could increase with climate change for continental France, although these projections are based upon large-scale climate modelling experiments across the European domain.
- However, recent regional climate model simulations for France suggest that extreme precipitation may not change under climate change.
- To this end, large uncertainty remains in quantifying the degree to which pluvial flooding could be affected by climate change.

## **Fluvial flooding**

- Projected changes in flood hazard in France are subject to large uncertainties owing to large natural variability and uncertainties in the simulated climate change signal from models.

- European-scale simulations have projected an increase in extreme flow levels in the main French rivers by the end of the century.
- Recent national-scale studies mostly found little change in flooding in France with climate change, however.
- Simulations by the AVOID programme support this, with the majority of the projections showing a greater tendency towards decreasing flood risk at first. Later in the century the models become more evenly divided between increases and decreases in flood risk, especially under the A1B scenario.
- Research should continue on the influence of the downscaling techniques used in hydrological impact studies on the simulation of flow extremes, and future assessments should focus on a reduction of the biases and on creating probabilistic scenarios based on larger ensembles of RCM simulations.

## **Tropical cyclones**

- Continental France is not impacted by tropical cyclones.

## **Coastal regions**

- Continental France's coasts and port cities could be affected by sea level rise (SLR) but to a lesser extent than many other countries across the globe.
- Results from the limited number of studies available indicate that by the 2080s under a high SLR scenario and without adaptation, the average annual number of people flooded in France could be around 463,000. This is greatly reduced with adaptation, to around 2,500. Under a low SLR scenario, 3,000 people could be flooded annually without adaptation and 1,800 could be flooded with adaptation.
- Some French districts in the West Indies, South America, Indian Ocean and Pacific Ocean could be affected by SLR, but these were not considered in this review.

# Crop yields

## Headline

Crop yield projections under climate change scenarios for France vary across studies due to the application of different models, assumptions and emissions scenarios. However, generally, projections imply yield gains for wheat and yield losses for maize, with climate change.

Results from the AVOID programme for France indicate that the balance is more towards declining suitability than improving suitability in the early part of the 21<sup>st</sup> century, and this increases further over time particularly in the A1B scenario.

## Supporting literature

### Introduction

The impacts of climate change on crop productivity are highly uncertain due to the complexity of the processes involved. Most current studies are limited in their ability to capture the uncertainty in regional climate projections, and often omit potentially important aspects such as extreme events and changes in pests and diseases. Importantly, there is a lack of clarity on how climate change impacts on drought are best quantified from an agricultural perspective, with different metrics giving very different impressions of future risk. The dependence of some regional agriculture on remote rainfall, snowmelt and glaciers adds to the complexity - these factors are rarely taken into account, and most studies focus solely on the impacts of local climate change on rain-fed agriculture. However, irrigated agricultural land produces approximately 40-45 % of the world's food (Doll and Siebert 2002), and the water for irrigation is often extracted from rivers which can depend on climatic conditions far from the point of extraction. Hence, impacts of climate change on crop productivity often need to take account of remote as well as local climate changes. Indirect impacts via sea-level rise, storms and diseases have also not been quantified. Perhaps most seriously, there is high uncertainty in the extent to which the direct effects of CO<sub>2</sub> rise on plant physiology will interact with climate change in affecting productivity. Therefore, at present, the aggregate impacts of climate change on large-scale agricultural productivity cannot be reliably

quantified (Gornall et al, 2010). This section summarises findings from a range of post IPCC AR4 assessments to inform and contextualise the analysis performed by AVOID programme for this project. The results from the AVOID work are discussed in the next section.

Wheat, maize and barley are important food crops in France (see Table 1). When it comes to value, grapes, sugar beets and rapeseed also play a vital role (FAO, (2008)).

□

Harvested area (ha)		Quantity (Metric ton)		Value (\$1000)	
Wheat	5490000	Wheat	39000000	Wheat	4380000
Barley	1790000	Sugar beet	30300000	Grapes	2620000
Maize	1700000	Maize	15800000	Sugar beet	1390000
Rapeseed	1420000	Barley	12100000	Rapeseed	1330000
Grapes	814000	Potatoes	6870000	Potatoes	921000
Sunflower seed	629000	Grapes	5670000	Maize	908000
Sugar beet	349000	Rapeseed	4720000	Barley	869000

**Table 1.** The top 7 crops by harvested area, quantity and value according to the FAO (2008) in France. Crops that feature in all lists are shaded green; crops that feature in two top 7 lists are shaded amber. Data is from FAO (2008) and has been rounded down to three significant figures.

A number of impact model studies looking at crop yield which include results for some of the main crops in France have been conducted. They apply a variety of methodological approaches, including using different climate model inputs and treatment of other factors that might affect yield, such as impact of increased CO<sub>2</sub> in the atmosphere on plant growth and adaption of agricultural practises to changing climate conditions. Some studies report projections for geographic or climatic areas larger than France alone and it is not always clear to what extent the crop yield projections are representative for France only in these cases. These different models, assumptions and emissions scenarios mean that there are a range of crop yield projections for France. However, the majority of studies explored in this report show yield declines with climate change for maize, but a gain in yield for wheat.

Important knowledge gaps and key uncertainties which are applicable to France as well as at the global-scale, include; the quantification of yield increases due to CO<sub>2</sub> fertilisation and yield reductions due to ozone damage (Ainsworth and McGrath, 2010, Iglesias et al., 2009), and the extent crop diseases could affect crop yields with climate change (Luck et al., 2011). Recent work has highlighted that repeated small water stresses during the growing season may have a strong impact on yields. However, the magnitude to which climate change might exacerbate this phenomenon remains poorly understood.

Most crop simulation models do not include the direct effect of extreme temperatures on crop development and growth, thus only changes in mean climate conditions are considered to affect crop yields for the studies included here.

## Assessments that include a global or regional perspective

### Recent Past

Crop yield changes could be due to a variety of factors, which might include, but not be confined to, a changing climate. In order to assess the impact of recent climate change (1980-2008) on wheat, maize, rice and soybean, Lobell et al. (2011) looked at how the overall yield trend in these crops changed in response to changes in climate over the period studied. The study was conducted at the global-scale but national estimates for France were also calculated. Lobell et al. (2011) divided the climate-induced yield trend by the overall yield trend for 1980–2008, to produce a simple metric of the importance of climate relative to all other factors. The ratio produced indicates the influence of climate on the productivity trend. So for example a value of  $-0.1$  represents a 10% reduction in yield gain due to climate change, compared to the increase that could have been achieved without climate change, but with technology and other gains. This can also be expressed as 10 years of climate trend being equivalent to the loss of roughly 1 year of technology gains. For France negative impacts were estimated to have occurred for maize and wheat, relative to what could have been achieved without the climate trends (see Table 2).

□

Crop	Trend
Maize	-0.1 to 0.0
Rice	0.1 to 0.2
Wheat	-0.2 to -0.1
Soybean	-0.2 to -0.1

**Table 2.** The estimated net impact of climate trends for 1980-2008 on crop yields in France. Climate-induced yield trend divided by overall yield trend. Data is from Lobell et al. (2011).

□

### Climate change studies

Several recent studies have applied climate projections from Global Climate Models (GCMs) to crop models to assess the global-scale impact of climate change on crop yields (Iglesias and Rosenzweig, 2009, Giannakopoulos et al., 2005, Moriondo et al., 2010, Olesen et al., 2007). Most of these studies include impact estimates at the national-scale for France which are presented in this section. The process of CO<sub>2</sub> fertilisation of some crops is usually included in most climate impact studies of yields. However, other gases can influence crop yield and are not always included in impacts models. An example of this is ozone (O<sub>3</sub>) and

so a study which attempts to quantify the potential impact on crop yield of changes in ozone in the atmosphere is also included (Avnery et al., 2011). In addition to these studies, the AVOID programme analysed the patterns of climate change for 21 GCMs, to establish an index of 'climate suitability' of agricultural land. Climate suitability is not directly equivalent to crop yields, but is a means of looking at a standard metric across all the countries including in this project, and of assessing the level of agreement on variables that affect crop production, between all 21 GCMs.

Iglesias and Rosenzweig (2009) repeated an earlier study presented by Parry et al. (2004) by applying climate projections from the HadCM3 GCM (instead of HadCM2, which was applied by Parry et al. (2004)), under seven SRES emissions scenarios and for three future time periods. This study used a globally consistent crop simulation methodologies and climate change scenarios, and weighted the model site results by their contribution to regional and national, and rain-fed and irrigated production. The study also applied a quantitative estimation of physiological CO<sub>2</sub> effects on crop yields and considered the affect of adaptation by assessing the country or regional potential for reaching optimal crop yield. The results from the study for France are presented in Table 3 and Table 4. Wheat yield was projected above baseline (1970-2000) levels for each future timeslice. Maize yields in 2020 and 2050 were slightly (<5%) lower than the baseline but by 2080 a small gain was projected with the A1FI and A2 scenarios, but not with the B1 and B2 scenarios.

□

Scenario	Year	Wheat	Maize
A1FI	2020	4.19	-1.68
	2050	9.28	-1.37
	2080	7.48	1.19
A2a	2020	5.67	-1.16
	2050	9.20	-1.32
	2080	13.14	0.45
A2b	2020	3.49	-0.72
	2050	8.92	-1.66
	2080	13.15	0.77
A2c	2020	3.34	-1.48
	2050	9.07	-1.81
	2080	13.51	0.44
B1a	2020	1.61	-1.71
	2050	5.28	-2.83
	2080	6.86	-3.21
B2a	2020	3.66	-2.60
	2050	5.13	-3.19
	2080	7.24	-1.42
B2b	2020	3.16	-2.55
	2050	5.50	-3.13
	2080	8.85	-1.35

**Table 3.** Wheat and maize yield changes (%) in France relative to baseline scenario (1970-2000) for different emission scenarios and future time periods. Some emissions scenarios were run in an ensemble simulation (e.g. A2a, A2b, A2c). Data is from Iglesias and Rosenzweig (2009).

□

	Wheat		Maize	
	Up	Down	Up	Down
<b>Baseline to 2020</b>	7	0	0	7
<b>Baseline to 2050</b>	7	0	0	7
<b>Baseline to 2080</b>	7	0	4	3
<b>2020 to 2050</b>	7	0	1	6
<b>2050 to 2080</b>	6	1	6	1


**Table 4.** The number of emission scenarios that predict yield gains (“Up”) or yield losses (“Down”) for wheat and maize in France between two points in time. Data is from Iglesias and Rosenzweig (2009).

Giannakopoulos et al. (2005, 2009) applied climate projections with the HadCM3 GCM under the SRES A2 and B2 emissions scenarios to assess climate change impacts for the Mediterranean basin for the period 2031-2060 under the A2 and B2 emissions scenarios. Climate data were used as input to the CROPSYST (Cropping Systems Simulation Model) (Stockle et al., 2003) crop model to project crop productivity changes (compared to 1961-1990) for a range of different crop types.

The crop types were divided into ‘C4’ summer crop, ‘C3’ summer crop, legumes, tuber crops and cereals, where ‘C4’ and ‘C3’ refer to two plant physiology types that affect the way



plants take up CO<sub>2</sub> from the atmosphere. 'C3' crops are able to benefit from CO<sub>2</sub> enrichment of the atmosphere, whereas 'C4' crops are not. This process is simulated by CROPSYST. The process is important because the benefit from CO<sub>2</sub> enrichment can potentially off-set some of the negative impacts of climate change for that crop. For France the 'C4' summer crop studied was irrigated maize, the 'C3' summer crop was rain-fed sunflowers, the legume was rain-fed soybeans, the tuber crop was irrigated potato and the cereal was rain-fed wheat. Yield increases were projected for all crops and in each emission scenario except for tuber crops (potato) with B2 emissions. These gains were smallest for cereals (wheat) and largest for C4 summer crops (maize), and intermediate for legumes (soybean) and C3 summer crops (sunflower).



**Figure 2.** Impact of climate change on crop productivity for different types of crops in France. The Y-axis is expressed as percentage difference between future (A2 and B2 scenarios respectively) and present yields. After Giannakopoulos et al. (2005, 2009).

Moriondo et al. (2010) simulated relative changes in crop yield for sunflower, soybean, spring wheat and durum wheat for a global mean warming of 2°C warmer than present climate change scenario with A2 socioeconomics. The study accounted for changes in extreme events such as droughts and the CO<sub>2</sub> fertilisation effect. Moriondo et al. (2010) compared the effectiveness of various adaptation options relative to no adaptation. No quantitative information on impacts is available from the study but estimates can be made whether, on average, a relative yield loss or a yield gain was projected for a given crop, adaptation method and country (see Table 5). The results indicate that for the 2030-2060 time horizon, yield gains were on average expected for sunflower and durum wheat on

application of certain adaptation methods. Whether longer cycle varieties have the potential to cause yield increases relative to the present period for soybean is unclear.

□

□

	No adaptation <sup>1</sup>	Advanced sowing	Delayed sowing	Shorter cycle varieties	Longer cycle varieties	Irrigation
<b>Sunflower</b>	+ -	+	-	-	+	+
<b>Soybean</b>	-	+	-	-	+ -	+
<b>Spring wheat</b>	-	+	-	-	+	+
<b>Durum wheat</b>	+	-	+	-	+	+

□

□

**Table 5.** Relative change in yield of four crops in a +2 °C world under SRES A2 socioeconomics for France. The relative change is calculated with respect to the same +2°C scenario without adaptation (left column). “+” = relative yield gain, “-” = relative yield loss, “+ -” = high spatial variability and uncertainty over sign of average yield change. After Moriondo et al. (2010).

□

Olesen et al (2007) addressed the issue of uncertainty in projecting impacts of climate change on agriculture. They projected rain-fed winter wheat yield across the European domain using nine different RCMs with HadAM3H as the bounding GCM, under SRES A2 emissions. For more than 50% of the cropping area, particularly in south-western France, RCMs conflicted in the direction of response. However, for northern France, the RCMs agreed on a yield increase.

Elsewhere, several recent studies have assessed the impact of climate change on a global-scale or regional-scale and include impact estimates for Western Europe or the Mediterranean as a whole (Ciscar et al., 2009, Ferrise et al., 2011, Iglesias et al., 2009, Tatsumi et al., 2011, Arnell et al., 2010a). Whilst these studies provide a useful indicator of crop yields under climate change for the larger region, it should be noted that the crop yields presented in such cases are not definitive national estimates. This is because the yields are averaged over the entire region, which includes other countries as well as France.

Tatsumi et al. (2011) applied an improved version of the GAEZ crop model (iGAEZ) to simulate crop yields on a global scale for wheat, potato, cassava, soybean, rice, sweet potato, maize, green beans. The impact of global warming on crop yields from the 1990s to 2090s was assessed by projecting five GCM outputs under the SRES A1B scenario and comparing the results for crop yields as calculated using the iGAEZ model for the period of 1990-1999. The results for Western Europe, the regional grouping which includes France,

are displayed in Table 6 and suggest increased yields for wheat, potato and green beans, but a decline for rice and maize.

□

□

Wheat	Potato	Cassava	Soybean	Rice	Sweet potato	Maize	Green beans
7.81	4.07	-	-0.03	-1.11	-	-5.17	7.34


**Table 6.** Average change in yield (%), during 1990s-2090s in Western Europe. Data is from Tatsumi et al. (2011).

□

Arnell et al. (2010a) applied five climate change projections from five GCMs to assess the effects of climate change on crop productivity and food security. Specifically, the crop simulation model GLAM-maize was used to simulate the effect of climate change on maize productivity. For Western Europe a yield loss of around 40% was projected, relative to the baseline (1961-1990) in the absence of adaptation and mitigation strategies, under an SRES A1B emissions scenario. Implementing a climate change mitigation scenario, which reduced emissions by 5%/year from 2016 onwards to a low emissions floor, mitigated the negative impact by approximately 20% and 30% in 2050 and 2100 respectively.

Ferrise et al. (2011) developed a probabilistic framework for evaluating the risk of durum wheat yield shortfall for the Mediterranean Basin. An artificial neural network, trained to emulate the outputs of a process-based crop growth model, was adopted to create yield response surfaces which were then overlaid with probabilistic projections of future temperature and precipitation changes in order to estimate probabilistic projections of future yields. To estimate the climatic risk of durum wheat shortfall in the next century, the future yield projections were compared with a critical threshold, calculated as the 30-year mean yield for the reference period (1961–1990). The climatic risk of durum wheat yield shortfall was then defined as the relative frequency of future yield projections below the threshold. Results were only presented as plotted maps of the spatial distribution of climatic risk of durum wheat shortfall (see Figure 3) but it is evident nonetheless that the projected probability of future yield being below the baseline is higher than 50% for most locations and time slices. Assuming crop yield probability distributions do not deviate much from normality it can be inferred that for most grid cells in France (and other Mediterranean countries) future durum crop yield is projected to decline.

□



**Figure 3.** Spatial distribution of risk of durum wheat yield shortfall by: (a) 2010–2030, (b) 2030–2050, (c) 2050–2070 and (d) 2070–2090. Risk is defined as the relative frequency of future projected yields that are lower than the selected threshold (30-year mean yield for 1961-1990). Figure is from Ferrise et al. (2011).

□


The PESETA project estimated the impacts of climate change on crop yields for different regions in the EU (Ciscar et al., 2009, Iglesias et al., 2009). Climate scenarios were created for the 2070-2100 time horizon using a combination of two GCMs and SRES emissions scenarios (A2 and B2). Crop yield simulations (winter wheat, spring wheat, rice, grassland, maize and soybeans) were then conducted using the DSSAT suite of crop models. The results for the “Central Europe South” region, which includes France and other countries, are displayed in Table 7. As mentioned previously, it should be noted that the projected yield changes may vary widely within a geographic region. The Central Europe South average is probably not representative (too optimistic) for France. Nevertheless, the PESETA project includes useful maps that show projected changes in crop yield for each emissions scenario, from which impacts for France can be inferred (see Figure 4). By the end of the century the results generally indicate a general decline in crop yields, especially in the western and southwestern parts of the country.

□

2011-2040		2071-2100		
A2 ECHAM4	A2 HadAM3h	B2 HadAM3h	A2 ECHAM4	B2 ECHAM4
7	5	5	-3	3

**Table 7.** Projected crop yield changes (%), compared to 1961-1990 period for the “Central Europe South” region, which includes France. Data is from Ciscar et al. (2009).

□



**Figure 4.** Crop yield changes under the HadCM3/HIRHAM A2 and B2 scenarios for the period 2071 - 2100 and for the ECHAM4/RCA3 A2 and B2 scenarios for the period 2011 – 2040 compared to baseline. The figure is from (Iglesias et al., 2009), p.31.

In addition to the studies looking at the effect of changes in climate and CO<sub>2</sub> concentrations on crop yield, Avnery et al. (2011) investigated the effects of ozone surface exposure on crop yield losses for soybeans, maize and wheat under the SRES A2 and B1 scenarios respectively. Two metrics of ozone exposure were investigated; seasonal daytime (08:00-19:59) mean O<sub>3</sub> (“M12”) and accumulated O<sub>3</sub> above a threshold of 40 ppbv (“AOT40”). The effect of the ozone exposure was considered in isolation from climate and other changes. The results for France are presented in Table 8.

□

	A2		B1	
	M12	AOT40	M12	AOT40
<b>Soybeans</b>	-	-	-	-
<b>Maize</b>	8-10	4-6	6-8	2-4
<b>Wheat</b>	4-6	15-20	4-6	10-15

**Table 8.** National relative crop yield losses (%) for 2030 under A2 and B1 emission scenarios according to the M12 (seasonal daytime (08:00–19:59) mean) and AOT40 (accumulated O<sub>3</sub> above a threshold of 40 ppbv) metrics of O<sub>3</sub> exposure. Data is from Avnery et al. (2011).

## **National-scale or sub-national scale assessments**

In this section we present results from recent studies that have looked at ongoing trends in crop yields, or have produced national or sub-national scale projections of future crop yields in France.

### **Recent past**

Brisson et al. (2010) analysed yield trends in France in recent decades using national and regional statistics, scattered trials and results of agro-climatic models using climatic data. Effects in genetic changes through breeding, agronomy and climate were investigated as possible causes for the stagnation of wheat yield. The results showed that genetic progress did not decline but was partly counteracted, from 1990 onwards, by increased frequencies of heat stress during grain filling and drought during stem elongation. However, the authors could not unequivocally single out climate trends to be responsible for the yield stagnation, as the decline of legumes in the cereal rotations (replaced by oilseed rape) and to a lesser extent the decrease in nitrogen fertilization are also likely to have contributed.

### **Climate change studies**

Brisson & Levraut (2010) report of a large nationwide study into the projected effects of future climate on agriculture and forestry in France. For a range of crops, yield projections and feasibility of cropping area was estimated under A1B emissions. Absolute yield changes, if significant, were calculated for 12 reference sites and are qualitatively summarised in Table 9. The main conclusion is that on average maize yields were projected to decrease but yields for wheat (soft and durum) and other crops were projected to increase. Yield and quality of grapes for wine was extremely variable depending on the characteristics of the site and growing system considered.

		Wheat	Maize	Sorghum	Oilseed rape	Sunflower
<b>Near future</b>	Increase	6	4	4	5	5
	Not-significant	6	1	5	7	7
	Decrease	0	7	3	0	0
<b>Distant future</b>	Increase	8	4	7	6	6
	Not-significant	4	1	4	6	6
	Decrease	0	7	1	0	0

**Table 9.** Frequency of significant and non-significant yield changes relative to the reference period (1970-1999) for the near (2020-2049) and distant future (2070-2099) for five crops at 12 reference sites in France, under A1B emissions. Data from Brisson & Levrault (2010).

□

Recent work has highlighted that repeated small water stresses during the growing season may have a strong impact on yields (Gate et al., 2010). However, the magnitude to which climate change might exacerbate this phenomenon remains poorly understood.

A recent national-scale assessment quantified the impacts of climate change on the agricultural sector in France (MEEDDM, 2008b, Galliot et al., 2009). Growth models showed an increase in yield in response to climate change for France, notably for wheat up to 2100. This increase did not take into account inter-annual variability and declines in water availability, however. The inclusion of these variability factors, which are still poorly integrated into the growth models, could enable the results to be refined and the anticipated increase in yield to be moderated. For example, increased events like the 2003 heatwave could, in 2100, represent a cost of up to more than €300 million per year for a crop such as wheat in the absence of any adaptation measure (MEEDDM, 2008b, Galliot et al., 2009). The assessment also showed that French viticulture could be affected by climate change, with high territorial differences and effects on the quality of the wines.

## **AVOID programme results**

To further quantify the impact of climate change on crops, the AVOID programme simulated the effect of climate change on the suitability of land for crop cultivation for all countries reviewed in this literature assessment based upon the patterns of climate change from 21 GCMs (Warren et al., 2010). This ensures a consistent methodological approach across all countries and takes consideration of climate modelling uncertainties.

## Methodology

The effect of climate change on the suitability of land for crop cultivation is characterised here by an index which defines the percentage of cropland in a region with 1) a decrease in suitability or 2) an increase in suitability. A threshold change of 5% is applied here to characterise decrease or increase in suitability. The crop suitability index is calculated at a spatial resolution of  $0.5^{\circ} \times 0.5^{\circ}$ , and is based on climate and soil properties (Ramankutty et al., 2002). The baseline crop suitability index, against which the future changes are measured, is representative of conditions circa 2000. The key features of the climate for the crop suitability index are temperature and the availability of water for plants. Changes in these were derived from climate model projections of future changes in temperature and precipitation, with some further calculations then being used to estimate actual and potential evapotranspiration as an indicator of water availability. It should be noted that changes in atmospheric  $\text{CO}_2$  concentrations can decrease evapotranspiration by increasing the efficiency of water use by plants (Ramankutty et al., 2002), but that aspect of the index was not included in the analysis here. Increased  $\text{CO}_2$  can also increase photosynthesis and improve yield to a small extent, but again these effects are not included. Exclusion of these effects may lead to an overestimate of decreases in suitability.

The index here is calculated only for grid cells which contain cropland circa 2000, as defined in the global crop extent data set described by Ramankutty et al. (2008) which was derived from satellite measurements. It is assumed that crop extent does not change over time. The crop suitability index varies significantly for current croplands across the world (Ramankutty et al., 2002), with the suitability being low in some current cropland areas according to this index. Therefore, while climate change clearly has the potential to decrease suitability for cultivation if temperature and precipitation regimes become less favourable, there is also scope for climate change to increase suitability in some existing cropland areas if conditions become more favourable in areas where the suitability index is not at its maximum value of 1. It should be noted that some areas which are not currently croplands may already be suitable for cultivation or may become suitable as a result of future climate change, and may become used as croplands in the future either as part of climate change adaptation or changes in land use arising for other reasons. Such areas are not included in this analysis.




## Results

Crop suitability was estimated under the pattern of climate change from 21 GCMs with two emissions scenarios; 1) SRES A1B and 2) an aggressive mitigation scenario where emissions follow A1B up to 2016 but then decline at a rate of 5% per year thereafter to a low emissions floor (denoted A1B-2016-5-L). The application of 21 GCMs is an attempt to quantify the uncertainty due to climate modelling, although it is acknowledged that only one crop suitability impacts model is applied. Simulations were performed for the years 2030, 2050, 2080 and 2100. The results for France are presented in Figure 5.

By 2030 in both emissions scenarios, all models projected 4% of current French cropland areas to undergo an improvement of suitability of cultivation. Over the 21<sup>st</sup> century the difference between models becomes slightly larger over time for both scenarios but slightly more so for A1B, with the range of croplands showing improving suitability expanding to approximately 1%-7% for A1B and 4%-6% for the mitigation scenario by 2100. For both scenarios, between 0% and 32% of current French croplands are projected to undergo declining suitability by 2030. By 2100 this rises to 1%-62% under the mitigation scenario and 5%-92% under A1B.

So, for France, the balance is more towards declining suitability than improving suitability in the early part of the 21<sup>st</sup> century, and this increases further over time particularly in the A1B scenario.

2



**Figure 5.** Box and whisker plots for the impact of climate change on increased crop suitability (top panel) and decreased crop suitability (bottom panel) for France, from 21 GCMs under two emissions scenarios (A1B and A1B-2016-5-L), for four time horizons. The plots show the 25th, 50th, and 75th percentiles (represented by the boxes), and the maximum and minimum values (shown by the extent of the whiskers).

2

27

# Food security

## Headline

Several studies show that France could remain food-secure with climate change. Recent research by the AVOID programme and others demonstrates that adaptive measures could be key to ensuring France's food security under climate change.

## Supporting literature

### Introduction

Food security is a concept that encompasses more than just crop production, but is a complex interaction between food availability and socio-economic, policy and health factors that influence access to food, utilisation and stability of food supplies. In 1996 the World Food Summit defined food security as existing 'when all people, at all times, have physical and economic access to sufficient, safe and nutritious food to meet their dietary needs, and their food preferences are met for an active and healthy life'. As such this section cannot be a comprehensive analysis of all the factors that are important in determining food security, but does attempt to assess a selection of the available literature on how climate change, combined with projections of global and regional population and policy responses, may influence food security.

### Assessments that include a global or regional perspective

France is a country of very low concern for food security, relative to other countries across the globe. According to FAO statistics (FAO, 2010) France has an extremely low level of undernourishment (less than 5% of the population). Moreover, a number of global studies point towards a generally optimistic and positive outlook for the impact of climate change on food security in France, largely as a result of its high adaptive capacity.

Several recent studies have analysed food security under climate change across the globe. For example, Wu et al. (2011) simulated crop yields with the GIS-based Environmental Policy Integrated Climate (EPIC) model. This was combined with crop areas simulated by a crop choice decision model to calculate total food production and per capita food availability

across the globe, which was used to represent the status of food availability and stability. The study focussed on the SRES A1 scenario and applied climate change simulations for the 2000s (1991–2000) and 2020s (2011–2020). The climate simulations were performed by MIROC (Model for Interdisciplinary Research on Climate) version 3.2., which means the effects of climate model uncertainty were not considered. Downscaled population and GDP data from the International Institute for Applied Systems Analysis (IIASA) were applied in the simulations. Whilst France appeared as a hot spot for food security vulnerability in 2020, Wu et al. (2011) note that France's population could still be food-secure as their population is less reliant on subsistence agriculture and because France possess a high capability of importing food due to strong purchasing power and financial support. Also, France presents substantial adaptive capacity and possesses proactive food management systems.

Falkenmark et al. (2009) present a global analysis of food security under climate change scenarios for the 2050s that considers the importance of water availability for ensuring global food security. The study presents an analysis of water constraints and opportunities for global food production on current croplands and assesses five main factors:

- 1) how far improved land and water management might go towards achieving global food security,
- 2) the water deficits that would remain in regions currently experiencing water scarcity and which are aiming at food self-sufficiency,
- 3) how the water deficits above may be met by importing food,
- 4) the cropland expansion required in low income countries without the needed purchasing power for such imports, and
- 5) the proportion of that expansion pressure which will remain unresolved due to potential lack of accessible land.

Similar to the study presented by Wu et al. (2011), there is no major treatment of modelling uncertainty; simulations were generated by only the LPJml dynamic global vegetation and water balance model (Gerten et al. 2004) with population growth and climate change under the SRES A2 emission scenario. Falkenmark et al. (2009) summarise the impacts of future improvements (or lack thereof) in water productivity for each country across the globe and show that this generates either a deficit or a surplus of water in relation to food water requirements in each country. These can be met either by trade or by horizontal expansion (by converting other terrestrial ecosystems to crop land). The study estimated that in 2050

around one third of the world's population will live in each of three regions: those that export food, those that import food, and those that have to expand their croplands at the expense of other ecosystems because they do not have enough purchasing power to import their food. The simulations demonstrated that France was a food exporting country in 2050.

The International Food Policy Research Institute (IFPRI) have produced a report and online tool that describes the possible impact of climate change on two major indicators of food security; 1) the number of children aged 0-5 malnourished, and 2) the average daily kilocalorie availability (Nelson et al., 2010, IFPRI, 2010). The study considered three broad socio-economic scenarios; 1) a 'pessimistic' scenario, which is representative of the lowest of the four GDP growth rate scenarios from the Millennium Ecosystem Assessment GDP scenarios and equivalent to the UN high variant of future population change, 2) a 'baseline' scenario, which is based on future GDP rates estimated by the World Bank and a population change scenario equivalent to the UN medium variant, and 3) an 'optimistic' scenario that is representative of the highest of the four GDP growth rate scenarios from the Millennium Ecosystem Assessment GDP scenarios and equivalent to the UN low variant of future population change. Nelson et al. (2010) also considered climate modelling and emission uncertainty and included a factor to account for CO<sub>2</sub> fertilisation in their work. The study applied two GCMs, the CSIRO GCM and the MIROC GCM, and forced each GCM with two SRES emissions scenarios (A1B and B1). They also considered a no climate change emissions scenario, which they called 'perfect mitigation' (note that in most other climate change impact studies that this is referred to as the baseline). The perfect mitigation scenario is useful to compare the effect of climate change against what might have happened without, but is not a realistic scenario itself. IFPRI have not published projections for child malnourishment in France but information on average daily kilocalorie availability has been made available. Table 10 displays the average daily kilocalorie availability simulated under different climate and socioeconomic scenarios for France and Figure 6 displays the effect of climate change, calculated by comparing the 'perfect mitigation' scenario with each baseline, optimistic and pessimistic scenario. Whilst by 2050 climate change is attributable for up to a 6% decline in kilocalorie availability, the absolute value of available kilocalories remains high (above 3,000) under all scenarios, which suggests France may not face food security issues in 2050. Figure 6 shows how the changes projected for France compare with the projections for the rest of the globe (IFPRI, 2010).

□

?


Scenario	2010	2050
Baseline CSI A1B	3452	3461
Baseline CSI B1	3457	3483
Baseline MIR A1B	3437	3392
Baseline MIR B1	3447	3443
Baseline Perfect Mitigation	3486	3622
Pessimistic CSI A1B	3488	3134
Pessimistic CSI B1	3492	3152
Pessimistic MIR A1B	3472	3074
Pessimistic MIR B1	3479	3103
Pessimistic Perfect Mitigation	3521	3269
Optimistic CSI A1B	3450	3589
Optimistic CSI B1	3454	3605
Optimistic MIR A1B	3434	3510
Optimistic MIR B1	3441	3542
Optimistic Perfect Mitigation	3483	3749

**Table 10.** Average daily kilocalorie availability simulated under different climate and socioeconomic scenarios, for France (IFPRI, 2010).

?


?

?



**Figure 6.** The impact of climate change on average daily kilocalorie availability (IFPRI, 2010).

?




**Figure 7.** Average daily kilocalorie availability simulated by the CSIRO GCM (CSI) under an A1B emissions scenario and the baseline socioeconomic scenario, for 2010 (top panel), 2030 (middle panel) and 2050 (bottom panel). The figure is from IFPRI (IFPRI, 2010). The changes show the combination of both climate change and socio-economic changes.

Arnell et al. (2010b) considered the impacts of global climate change and mitigation policy on food security for eleven countries. The study applied climate change patterns from the HadCM3 GCM and explored food security under two emissions scenarios; a business as usual scenario (SRES A1B) and four mitigations scenarios where emissions peak in 2030 and subsequently reduce at 2% per year to a high emissions floor (referred to as 2030-2-H) or 5% per year to a low emissions floor (2030-5-L), or where they peak in 2016 and subsequently reduce at 2% per year to a high emissions floor (referred to as 2016-2-H) or 5% per year to a low emissions floor (2016-5-L). The study also considered a series of structural adjustments that could be made in the future to adapt to food security issues, including that 1) if there is a shortfall of any per-capita food availability due to crop yield and/or population changes, then original (baseline) food amounts are made up by reducing or removing export amounts; and 2) if, after the above adjustments, there is still a shortfall, then the amount of crops going to animal feed is reduced or removed to try to make up to the original (baseline) food amounts. The model simulations presented by Arnell et al. (2010b) characterise the numbers of people *exposed to undernourishment* in the absence of increased crop production and imports, not actual numbers of undernourished people. The results are presented in Figure 8. Arnell et al. (2010b) showed that a 14% increase in population in France combined with 16-23% decreases in crop production by 2050 leads to a substantial increase in exposure to undernourishment. Without mitigation or structural adjustment, 52% of the population were projected to be exposed to undernourishment by 2050. 2030-5-L and 2030-2-H policies were associated with 42% and 50% of the population undernourished respectively by 2100, whereas this was substantially decreased by mitigation scenarios 2016-2-H and 2016-5-L which projected 37% and 32% undernourished by 2100 respectively. However, when structural adjustments were incorporated into the simulations, climate change ceased to have a discernible impact on food security. This is largely supportive of the findings presented by Wu et al. (2011), discussed previously.

□





**Figure 8.** Total projected population exposed to undernourishment in France. The left panel shows total exposure under the A1B emissions scenario (“A1b REF”), plus the A1B scenario with exports reduced or removed (“A1b-EXP”) and the A1B scenario with exports removed and allocation to feed reduced or removed (“A1b-EXP-FEED”). The right panel shows the total exposure under the A1b-EXP-FEED and three mitigation scenarios. The figure is from Arnell et al. (2010b).

□


□

It is important to note that up until recently, projections of climate change impacts on global food supply have tended to focus solely on production from terrestrial biomes, with the large contribution of animal protein from marine capture fisheries often ignored. However, recent studies have addressed this knowledge gap (e.g., Allison et al., 2009). In addition to the direct affects of climate change, changes in the acidity of the oceans, due to increases in CO<sub>2</sub> levels, could also have an impact of marine ecosystems, which could also affect fish stocks. However, this relationship is complex and not well understood, and studies today have not been able to begin to quantify the impact of ocean acidification on fish stocks.

Allison et al. (2009) present a global analysis that compares the vulnerability of 132 national economies to potential climate change impacts on their capture fisheries. The study considered a country’s vulnerability to be a function of the combined effect of projected climate change, the relative importance of fisheries to national economies and diets, and the national societal capacity to adapt to potential impacts and opportunities. Climate change projections from a single GCM under two emissions scenarios (SRES A1FI and B2) were used in the analysis. Allison et al. (2009) concluded that the national economy of France presented a very low vulnerability to climate change impacts on fisheries. In contrast, countries in Central and Western Africa (e.g. Malawi, Guinea, Senegal, and Uganda), Peru and Colombia in north-western South America, and four tropical Asian countries (Bangladesh, Cambodia, Pakistan, and Yemen) were identified as most vulnerable (see Figure 9). It should be noted, however, that results from studies that have applied only a

single climate model or climate change scenario should be interpreted with caution. This is because they do not consider other possible climate change scenarios which could result in a different impact outcome, in terms of magnitude and in some cases sign of change.

2



2

**Figure 9.** Vulnerability of national economies to potential climate change impacts on fisheries under SRES B2 (Allison et al., 2009). Colours represent quartiles with dark brown for the upper quartile (highest index value), yellow for the lowest quartile, and grey where no data were available.

2

### National-scale or sub-national scale assessments

Literature searches yielded no results for national-scale or sub-national scale studies for this impact sector.

2

# Water stress and drought

## Headline

Most studies suggest that droughts could become more frequent in France with climate change and that water stress might increase to a moderate level, although there remains a degree of uncertainty in the magnitude to which these changes could be realised. Recent simulations by the AVOID programme support this.

## Supporting literature

### Introduction

For the purposes of this report droughts are considered to be extreme events at the lower bound of climate variability; episodes of prolonged absence or marked deficiency of precipitation. Water stress is considered as the situation where water stores and fluxes (e.g. groundwater and river discharge) are not replenished at a sufficient rate to adequately meet water demand and consumption.

A number of impact model studies looking at water stress and drought for the present (recent past) and future (climate change scenario) have been conducted. These studies are conducted at global or national scale and include the application of global water 'availability' or 'stress' models driven by one or more climate change scenario from one or more GCM. The approaches variously include other factors and assumptions that might affect water availability, such as the impact of changing demographics and infrastructure investment, etc. These different models (hydrological and climate), assumptions and emissions scenarios mean that there are a range of water stress projections for France. This section summarises findings from these studies to inform and contextualise the analysis performed by the AVOID programme for this project. The results from the AVOID work and discussed in the next section.

Important knowledge gaps and key uncertainties which are applicable to France as well as at the global-scale, include; the appropriate coupling of surface water and groundwater in hydrological models, including the recharge process, improved soil moisture and evaporation

dynamics, inclusion of water quality, inclusion of water management (Wood et al. 2011) and further refinement of the down-scaling methodologies used for the climate driving variables (Harding et al. 2011).

## **Assessments that include a global or regional perspective**

### **Recent Past**

Recent research presented by Vörösmarty et al. (2010) describes the calculation of an 'Adjusted Human Water Security Threat' (HWS) indicator. The indicator is a function of the cumulative impacts of 23 biophysical and chemical drivers simulated globally across 46,517 grid cells representing 99.2 million km<sup>2</sup>. With a digital terrain model at its base, the calculations in each of the grid boxes of this model take account of the multiple pressures on the environment, and the way these combine with each other, as water flows in river basins. The level of investment in water infrastructure is also considered. This infrastructure measure (the *investment benefits factor*) is based on actual existing built infrastructure, rather than on the financial value of investments made in the water sector, which is a very unreliable and incomplete dataset. The analysis described by Vörösmarty et al. (2010) represents the current state-of-the-art in applied policy-focussed water resource assessment. In this measure of water security, the method reveals those areas where this is lacking, which is a representation of human water stress. One drawback of this method is that no analysis is provided in places where there is 'no appreciable flow', where rivers do not flow, or only do so for such short periods that they cannot be reliably measured. This method also does not address places where water supplies depend wholly on groundwater or desalination, being piped in, or based on wastewater reuse. It is based on what is known from all verified peer reviewed sources about surface water resources as generated by natural ecosystem processes and modified by river and other hydraulic infrastructure (Vörösmarty et al. (2010).


Here, the present day HWS is mapped for France. The model applied operates at 50km resolution, so, larger countries appear to have smoother coverage than smaller countries, but all are mapped and calculated on the same scale, with the same data and model, and thus comparisons between places are legitimate. It is important to note that this analysis is a comparative one, where each place is assessed *relative* to the rest of the globe. In this way, this presents a realistic comparison of conditions across the globe. As a result of this, however, some places may seem to be less stressed than may be originally considered. One example is Australia, which is noted for its droughts and long dry spells, and while there are some densely populated cities in that country where water stress is a real issue, for most

of the country, *relative to the rest of the world*, the measure suggests water stress (as measured by HWS defined by Vörösmarty et al. (2010)), is not a serious problem.

Figure 10 presents the results of this analysis for France, showing moderate degrees of water security threat, particularly in the north, and in Mediterranean areas. Most of the country however is not currently exposed to a water security threat.

□

□



□

**Figure 10.** Present Adjusted Human Water Security Threat (HWS) for France, calculated following the method described by Vörösmarty et al. (2010).


□

Smakhtin et al. (2004) present a first attempt to estimate the volume of water required for the maintenance of freshwater-dependent ecosystems at the global scale. This total environmental water requirement (EWR) consists of ecologically relevant low-flow and high-flow components. The authors argue that the relationship between water availability, total use and the EWR may be described by the water stress indicator (WSI). If WSI exceeds 1.0, the basin is classified as “environmentally water scarce”. In such a basin, the discharge has already been reduced by total withdrawals to such levels that the amount of water left in the basin is less than EWR. Smaller index values indicate progressively lower water resources exploitation and lower risk of “environmental water scarcity.” Basins where WSI is greater

than 0.6 but less than 1.0 are arbitrarily defined as heavily exploited or “environmentally water stressed” and basins where WSI is greater than 0.3 but less than 0.6 are defined as moderately exploited. In these basins, 0-40% and 40-70% of the utilizable water respectively is still available before water withdrawals come in conflict with the EWR. Environmentally “safe” basins are defined as those where WSI is less than 0.3. The global distribution of WSI for the 1961-1990 time horizon is shown in Figure 11. The results show that for the basins considered, France exhibits moderate water stress in the north of the country.

□

□



**Figure 11.** A map of the major river basins across the globe and the water stress indicator (WSI) for the 1961-1990 time horizon. The figure is from Smakhtin et al.

□

□

### Climate change studies


The IPCC AR4 (2007a) noted that annual precipitation changes by the end of the 21<sup>st</sup> Century under the A1B emissions scenario range between 0% and 16% in Northern Europe. The sign of changes in summer vary between models, but most models simulate decreased precipitation over this region. In Northern Europe, the CMIP3 multi-model dataset disagrees on whether summer soil moisture could increase or decrease, due to the competition between increased precipitation on one hand, and earlier snowmelt and increased evaporation on the other. The IPCC AR4 (2007a) also notes that a number of studies have suggested no major change in dry-spell length in northern Europe.

Rockstrom et al. (2009) applied the LPJml vegetation and water balance model (Gerten et al 2004) to assess green-blue water (irrigation and infiltrated water) availability and requirements. The authors applied observed climate data from the CRU TS2.1 gridded dataset for a present-day simulation, and climate change projections from the HadCM2 GCM under the SRES A2 scenario to represent the climate change scenario for the year 2050. The study assumed that if water availability was less than 1,300m<sup>3</sup>/capita/year, then the country was considered to present insufficient water for food self-sufficiency. The simulations presented by Rockstrom et al. (2009) should not be considered as definitive, however, because the study only applied one climate model, which means climate modelling uncertainty was overlooked. The results from the two simulations are presented in Figure 12. Rockstrom et al. (2009) found that globally in 2050 and under the SRES A2 scenario, around 59% of the world's population could be exposed to "blue water shortage" (i.e. irrigation water shortage), and 36% exposed to "green water shortages" (i.e. infiltrated rain shortage). For France, Rockstrom et al. (2009) found that blue-green water availability was well above the 1,300m<sup>3</sup>/capita/year threshold in present conditions and under climate change. This indicates that at a national level, France's water resource requirements should be met by 2050.

□

□






**Figure 12.** Simulated blue-green water availability ( $m^3/capita/year$ ) for present climate (top panel) and including both demographic and climate change under the SRES A2 scenario in 2050 (bottom panel). The study assumed that if water availability was less than  $1,300m^3/capita/year$ , then the country was considered to present insufficient water for food self-sufficiency. The figure is from Rockstrom et al. (2009).

Doll (2009) presents updated estimates of the impact of climate change on groundwater resources by applying a new version of the WaterGAP hydrological model. The study accounted for the number of people affected by changes in groundwater resources under climate change relative to present (1961-1990). To this end, the study provides an assessment of the vulnerability of humans to decreases in available groundwater resources (GWR). This indicator was termed the “Vulnerability Index” (VI), defined as;  $VI = -\% \text{ change GWR} * \text{Sensitivity Index (SI)}$ . The SI component was a function of three more specific sensitivity indicators that include an indicator of water scarcity (calculated from the ratio between consumptive water use to low flows), an indicator for the dependence upon groundwater supplies, and an indicator for the adaptive capacity of the human system. Doll



(2009) applied climate projections from two GCMs (ECHAM4 and HadCM3) to WaterGAP, for two scenarios (SRES A2 and B2), for the 2050s. Figure 13 presents each of these four simulations respectively. There is variation across scenarios and GCMs. With HadCM3, the simulated VI for France is negligible, because the simulated GWR decreases by less than 10% with climate change. However, this is not the case with the ECHAM4 GCM under SRES A2 emissions; here much of France exhibits a low to moderate VI, with VI higher in the south of the country.






**Figure 13.** Vulnerability index (VI) showing human vulnerability to climate change induced decreases of renewable groundwater resources (GWR) by the 2050s under two emissions scenarios for two GCMs. VI is only defined for areas with a GWR decrease of at least 10% relative to present (1961-1990). Also shown is VI for the Mediterranean region with ECHAM4 under A2 emissions. The figure is from Doll (2009).

□

□

Lehner et al. (2006) assessed the impact of climate change on European drought risk. The authors accounted for future human water use and assessed future flood and drought frequencies by applying the WaterGAP hydrological model, driven by climate projections from the HadCM3 and ECHAM4 GCMs, under a 1%/year CO<sub>2</sub> increase emissions scenario. The simulations are presented in Figure 14 and Figure 15. The results reflect the general consensus from other studies that southern and south-eastern Europe could experience


increased drought frequencies, leading to water stress. This in part due to increased water use but the impacts are much more pronounced and wide spread when climate change is factored in (Lehner et al., 2006). Long term projections indicate those drought events expected to occur once every 100 years could become much more frequent, to around every 40 years in the most extreme areas, including much of the Mediterranean. For France, both GCMs simulated that 100-year droughts could become slightly more likely in the 2020s, but could increase significantly by the 2070s. Large areas of south and southwest France could experience more frequent 100-year droughts. Results from the HadCM3 model also simulate that these droughts could become more intense by this time horizon. Similarly, Menzel and Matovelle (2010) showed that out of six simulations with climate and socio-economic changes, all of them broadly agreed that areas of northeast France could be under severe water stress by 2050 as expressed by the withdrawals to abstractions ratio threshold being greater than 0.4.



**Figure 14.** Change in recurrence of 100-year droughts, based on comparisons between today's climate and water use (1961–1990) and simulations for the 2020s and 2070s (ECHAM4 and HadCM3 GCMs), under a 1%/year CO<sub>2</sub> increase emissions scenario. The figure is from Lehner et al. (2006).

?

???



?

**Figure 15.** Change in intensity of 100-year droughts, based on comparison between today's climate and water use (1961–1990) and simulations for the 2070s (left map: HadCM3 GCM; right map: only water use scenario, no climate change), under a 1%/year CO<sub>2</sub> increase emissions scenario.

### National-scale or sub-national scale assessments

Boe and Terray (2008) applied an analysis of observed weather-types over France and found an increase of precipitation in northern France, and a decrease in southern France during the 20<sup>th</sup> Century. They suggest that the role of greenhouse gas and sulphate aerosol emissions in the changing weather types was limited, at least in the historic record. For the Pyrenees, Lopez-Moreno and Beniston (2009) found a tendency towards increasing drought periods from RCM simulations for the 21<sup>st</sup> Century. Likewise, Blenkinsop and Fowler (2007) showed that RCM projections favour a reduction in summer precipitation over most of Europe by the end of the 21<sup>st</sup> Century. However, in some parts of France (e.g. the Meuse basin) there was uncertainty as to the future change in direction of both long and short duration events.

A recent assessment presented by Soubeyrou et al. (2011) describes the possible impact of climate change on agricultural drought for France, under three emissions scenarios (A2, A1B and B1). The authors found that drought could become more frequent in the 2050s and that by the 2080s, droughts could be substantially more intense and longer than any in the observed record for France, especially in mountainous regions. While there was uncertainty,

all climate simulations were in agreement that drought could increase in duration and intensity with climate change.

## **AVOID Programme Results**

To further quantify the impact of climate change on water stress and the inherent uncertainties, the AVOID programme calculated water stress indices for all countries reviewed in this literature assessment based upon the patterns of climate change from 21 GCMs, following the method described by Gosling et al. (2010) and Arnell (2004). This ensures a consistent methodological approach across all countries and takes consideration of climate modelling uncertainties.

### **Methodology**

The indicator of the effect of climate change on exposure to water resources stress has two components. The first is the number of people within a region with an *increase in exposure to stress*, calculated as the sum of 1) people living in water-stressed watersheds with a significant reduction in runoff due to climate change and 2) people living in watersheds which become water-stressed due to a reduction in runoff. The second is the number of people within a region with a *decrease in exposure to stress*, calculated as the sum of 1) people living in water-stressed watersheds with a significant increase in runoff due to climate change and 2) people living in watersheds which cease to be water-stressed due to an increase in runoff. It is not appropriate to calculate the net effect of “increase in exposure” and “decrease in exposure”, because the consequences of the two are not equivalent. A water-stressed watershed has an average annual runoff less than 1000m<sup>3</sup>/capita/year, a widely used indicator of water scarcity. This indicator may underestimate water stress in watersheds where per capita withdrawals are high, such as in watersheds with large withdrawals for irrigation.

Average annual runoff (30-year mean) is simulated at a spatial resolution of 0.5°x0.5° using a global hydrological model, MacPDM (Gosling and Arnell, 2011), and summed to the watershed scale. Climate change has a “significant” effect on average annual runoff when the change from the baseline is greater than the estimated standard deviation of 30-year mean annual runoff: this varies between 5 and 10%, with higher values in drier areas.


The pattern of climate change from 21 GCMs was applied to MacPDM, under two emissions scenarios; 1) SRES A1B and 2) an aggressive mitigation scenario where emissions follow A1B up to 2016 but then decline at a rate of 5% per year thereafter to a low emissions floor (denoted A1B-2016-5-L). Both scenarios assume that population changes through the 21st century following the SRES A1 scenario as implemented in IMAGE 2.3 (van Vuuren et al., 2007). The application of 21 GCMs is an attempt to quantify the uncertainty due to climate modelling, although it is acknowledged that only one impacts model is applied (MacPDM). Simulations were performed for the years 2030, 2050, 2080 and 2100. Following Warren et al. (2010), changes in the population affected by increasing or decreasing water stress represent the additional percentage of population affected due to climate change, not the absolute change in the percentage of the affected population relative to present day.

## **Results**

The results for France are presented in Figure 16. They show that by 2100 under A1B, the median percentage of the population exposed to increased water stress is around 12% but uncertainty is high. Conversely, there is higher certainty that none of France's population might see a decrease in water stress with climate change.



2



**Figure 16.** Box and whisker plots for the impact of climate change on increased water stress (top panel) and decreased water stress (bottom panel) in France, from 21 GCMs under two emissions scenarios (A1B and A1B-2016-5-L), for four time horizons. The plots show the 25th, 50th, and 75th percentiles (represented by the boxes), and the maximum and minimum values (shown by the extent of the whiskers).

# Pluvial flooding and rainfall

## Headline

Recent modelling studies confirm conclusions from the IPCC AR4 that heavy winter precipitation and flooding could increase with climate change for continental France (French districts in West Indies, South America, Indian Ocean and Pacific Ocean are not considered in this review), although these projections are based upon large-scale climate modelling experiments across the European domain. France covers the transition zone between Northern Europe and Mediterranean Europe, and therefore extreme precipitation events over the north and south of the country may show differing responses to climate change.

## Supporting literature

### Introduction

Pluvial flooding can be defined as flooding derived directly from heavy rainfall, which results in overland flow if it is either not able to soak into the ground or exceeds the capacity of artificial drainage systems. This is in contrast to fluvial flooding, which involves flow in rivers either exceeding the capacity of the river channel or breaking through the river banks, and so inundating the floodplain. Pluvial flooding can occur far from river channels, and is usually caused by high intensity, short-duration rainfall events, although it can be caused by lower intensity, longer-duration events, or sometimes by snowmelt. Changes in mean annual or seasonal rainfall are unlikely to be good indicators of change in pluvial flooding; changes in extreme rainfall are of much greater significance. However, even increases in daily rainfall extremes will not necessarily result in increases in pluvial flooding, as this is likely to be dependent on the sub-daily distribution of the rainfall as well as local factors such as soil type, antecedent soil moisture, land cover (especially urbanisation), capacity and maintenance of artificial drainage systems etc. It should be noted that both pluvial and fluvial flooding can potentially result from the same rainfall event.



## **Assessments that include a global or regional perspective**

### **Climate change studies**

The IPCC AR4 (2007a) noted that annual precipitation is very likely to increase in most of northern Europe, and daily precipitation extremes are likely to increase under the A1B emissions scenario during the 21<sup>st</sup> Century. However, although the simulated responses between models are qualitatively consistent, significant uncertainties remain particularly regarding the magnitude and geographical details of precipitation change (IPCC, 2007a). The substantial natural variability of European climate is also a major uncertainty, particularly with respect to near-term climate projections. Annual precipitation changes by the end of the 21<sup>st</sup> Century under the A1B emissions scenario range between 0% and 16% in Northern Europe (IPCC, 2007a). The largest increases are simulated during winter. The sign of changes in summer vary between models, but most models simulate decreased precipitation over this region. Precipitation extremes during winter are also very likely to increase in both magnitude and frequency (IPCC, 2007a). Since southern France lies in the Mediterranean region, the IPCC AR4 (2007a) notes here that annual precipitation is very likely to decrease, the annual number of precipitation days is very likely to decrease, and the risk of summer drought is likely to increase. Extreme short-term precipitation may either increase (due to the increased water vapour content of a warmer atmosphere) or decrease (due to a decreased number of precipitation days, which if acting alone could also make heavy precipitation less common).

More recently, Bates et al. (2008) note that for Europe, based on climate projections from the ECHAM4 and HadCM3 GCMs, in the 2020s there is a projected increased risk of winter floods in northern Europe, and increased risk of flash floods over the whole of Europe. Sillmann and Roeckner (2008) find that over a broad region they classify as central Europe (12W–40E 45N–55N) there is an increase of both maximum 5-day rainfall (RX5day) and 95<sup>th</sup> percentile rainfall (R95p) with climate change under the A1B emissions scenario. In general, the differences between A1B and B1 were not pronounced until the decade approaching 2100.

Similarly, Beniston et al. (2007) found that heavy winter precipitation could increase in central and northern Europe with climate change. These changes were weaker for the B2 emissions scenarios than for the A2 scenario. However, the authors found that model choices can have greater effects on the magnitude (RCM) and pattern (GCM) of response than the choice of scenario. Analysing projections under the A2 and B2 scenarios from an

ensemble of nine RCMs, they found that changes in maximum 5-day rainfall simulated under the B2 scenario were smaller than those simulated under the A2 scenario in two cases, and similar in the other two cases. However, there were no systematic differences in projected increases in maximum 1-day rainfall between the scenarios, though the increases were positive and up to about 40%.

Goubanova and Li (2007) used a variable grid atmospheric GCM with a zoom over the Mediterranean region run with the A2 emissions scenario. They found that projections for the 21st Century showed an increase in precipitation extremes and variability over the Mediterranean region in winter, spring and autumn seasons. This is despite an overall decrease in mean precipitation.

## **National-scale or sub-national scale assessments**

### **Climate change studies**

López-Moreno and Beniston (2009) investigated a set of climate parameters (including mean precipitation, number of wet days, daily intensity, and number of days with more than 50 mm rainfall) over the Pyrenees for the 21<sup>st</sup> Century using a set of six RCMs under the SRES A2 emissions scenario. The climate of this region is very complex, but the projections indicated an intensification of extremes, increasing daily rainfall intensity, and increasing contribution of intense events to total precipitation. Quintana-Seguí et al. (2011) explored the impact of climate change on flash-floods in Cévennes region of the Massif Central. The authors found that an increase in the 10-year flood level of sometimes up to 100% was possible by the 2050s, and that the expected return period of what was a 10-year flood at the end of the 20<sup>th</sup> century could reduce to 2 years by the 2050s. Furthermore, Quintana Seguí et al. (2010) showed that by the 2050s, there might be significant increases of winter precipitation in this region and significant decreases of summer precipitation in most of the region.

Deque (2007) found that the index for number of winter days above 10mm of precipitation increases in the northern half of France by the late 2100s using the Météo-France atmospheric model ARPEGE/Climate. The number increases by 24% over France as a whole on average. They note that this is related to a change in the tails of the distribution not simply a shift in mean precipitation.

In contrast, for the French Alps, a study presented by Jomelli et al. (2009), which applied the ARPEGE model with the SRES A2 emissions scenario showed that the most significant climatic trends for the end of the 21<sup>st</sup> century could be a decrease in intense rainfall events, particularly during the autumn and winter. Similar results were simulated by two French

regional climate models; ARPEGE and LMDz, in a study presented by Peings et al. (2011). The simulated number of days with over 20 mm of precipitation, was observed to change little from present, throughout the 21<sup>st</sup> century (Figure 17).

?

?

?


?

?

?

?

?



**Figure 17.** The number of days with over 20mm of precipitation for the present (1990), and the differences between future time horizons and present, under the SRES B2 and A2 scenarios. Units are days. The figure is from Peings et al. (2011).

# Fluvial flooding

## Headline

Projections of changes in flood hazard in France with climate change are subject to large uncertainties due to large natural variability and large uncertainties in the simulated climate signal from climate models. European-scale simulations have projected an increase in extreme flow levels in the main French rivers by the end of the century. Recent national-scale assessments mostly found little change in flooding in France with climate change, however. This is confirmed by simulations by the AVOID programme, in which the majority of the projections show a greater tendency towards decreasing flood risk at first. Later in the century the models become more evenly divided between increases and decreases in flood risk, especially under the A1B scenario. Research should continue to focus on the influence of the downscaling techniques used in hydrological impact studies on the simulation of flow extremes. Future assessments should focus on a reduction of the biases and on creating probabilistic scenarios based on larger ensembles of RCM simulations.

## Supporting literature

### Introduction

This section summarises findings from a number of post IPCC AR4 assessments on river flooding in France to inform and contextualise the analysis performed by the AVOID programme for this project. The results from the AVOID work are discussed in the next section.

Fluvial flooding involves flow in rivers either exceeding the capacity of the river channel or breaking through the river banks, and so inundating the floodplain. A complex set of processes is involved in the translation of precipitation into runoff and subsequently river flow (routing of runoff along river channels). Some of the factors involved are; the partitioning of precipitation into rainfall and snowfall, soil type, antecedent soil moisture, infiltration, land cover, evaporation and plant transpiration, topography, groundwater storage. Determining whether a given river flow exceeds the channel capacity, and where any excess flow will go, is also not straightforward, and is complicated by the presence of artificial river

embankments and other man-made structures for example. Hydrological models attempt to simplify and conceptualise these factors and processes, to allow the simulation of runoff and/or river flow under different conditions. However, the results from global-scale hydrological modelling need to be interpreted with caution, especially for smaller regions, due to the necessarily coarse resolution of such modelling and the assumptions and simplifications this entails (e.g. a 0.5° grid corresponds to landscape features spatially averaged to around 50-55km for mid- to low-latitudes). Such results provide a consistent, high-level picture, but will not show any finer resolution detail or variability. Smaller-scale or catchment-scale hydrological modelling can allow for more local factors affecting the hydrology, but will also involve further sources of uncertainty, such as in the downscaling of global climate model data to the necessary scale for the hydrological models. Furthermore, the application of different hydrological models and analysis techniques often makes it difficult to compare results for different catchments.

In a comprehensive analysis of streamflow records in France, Renard et al. (2008) found no general trends in hydrological extremes, with three exceptions: in the northeast of the country, flood peaks were found to have increased; in the Pyrenees high and low flows showed decreasing trends, and in the Alps, earlier snowmelt-related floods were detected, along with less severe drought and increasing runoff due to glacier melting (Renard et al. 2008). Several other studies have looked at potential changes in flood hazard in France under future climate change, which are discussed below.

### **Assessments that include a global or regional perspective**

A European-scale study presented by Dankers and Feyen (2008) applied very high resolution (~12 km) RCM simulations to drive a flood forecasting model to make projections of future changes in flood hazard. The results showed an increase in the 100-year flood level of most major rivers in France by the end of the century (2071-2100) under the A2 emissions scenario. For some rivers, this increase reached up to 40%, for example in parts of the Loire and Garonne catchments. For the Loire river, the increase in flood levels was more prominent at shorter return periods. In all of the four main rivers (Rhône, Garonne, Loire and Seine) the return period of the current (1961-1990) 100-year flood level was projected to reduce to 50 years or less. On a seasonal basis, the 100-year flood level generally increased during winter and spring, but decreased in summer and autumn.

In a follow-up study using an ensemble of two RCMs with two different emission scenarios (A2 and B2), Dankers and Feyen (2009) found the projected increase in extreme flood levels in most of the French rivers to be relatively robust; although not all model experiments

presented an increase, very few projected a decrease in the 100-year flood level by the end of the century. In the Loire, for example, the projected changes ranged from slightly negative in 2 model experiments to +30%, with one experiment projecting an increase in the 100-year flow level of more than 50%. Generally speaking the changes were, however, less strong than projected by the high resolution simulations described by Dankers and Feyen (2008), averaging to increases of +5% to +10%. Also, there was little difference between the two emission scenarios (A2 and B2) with regards to the direction and magnitude of change in extreme flow levels. Some of these changes in simulated flood hazard may partly be attributed to large, decadal-scale variability in the simulated climate, although this effect seemed smaller than in other major European river basins.

As highlighted by Dankers and Feyen (2009), projections of changes in flood hazard with climate change are subject to large uncertainties due to large natural variability and large uncertainties in the simulated climate signal. Research should continue to focus on the influence of the downscaling techniques used in hydrological impact studies on the simulation of flow extremes. Future assessments should focus on a reduction of the biases and on creating probabilistic scenarios based on larger ensembles of RCM simulations.

### **National-scale or sub-national scale assessments**

Several local studies have been undertaken in the Seine and Loire basins. Applying different models and downscaling techniques, these studies do not confirm the projected general rise in flood hazard found by Dankers and Feyen (2008, 2009). In the Seine River, Ducharme et al. (2011) found a slight decrease in high flow levels that was less robust than the general decrease in low flows. In their simulations the 10-year return level did not change significantly during the 21<sup>st</sup> century and also the 100-year return level remained of the same order of magnitude. Similar results were found for the Loire river, where Moatar et al. (2010) reported little significant change in the 10-year flood level by the middle of the century, and slightly negative trends towards the end, albeit with a large spread highlighting important uncertainties. Boé et al. (2009) showed that despite large uncertainties linked to climate models, some robust signals appear by the 2050s in simulations of French river discharge. In particular, a decrease in mean discharges in summer and autumn and more frequent low flows. However, changes in the intensity of high flows were more uncertain (Boé et al., 2009). Studies like these may use models that are better optimised to local conditions than global-scale studies, however the results may also to a large extent be influenced by the choice of downscaling method, as demonstrated by Quintana-Seguí et al. (2011). In the area of the Cévennes, already seriously affected by flash-floods, the authors found an increase in the

10-year flood level of sometimes up to 100%, and the expected return period of what was a 10-year flood at the end of the 20th century was projected to reduce to 2 years by the middle of this century.

## **AVOID programme results**

To quantify the impact of climate change on fluvial flooding and the inherent uncertainties, the AVOID programme calculated an indicator of flood risk for all countries reviewed in this literature assessment based upon the patterns of climate change from 21 GCMs (Warren et al., 2010). This ensures a consistent methodological approach across all countries and takes consideration of climate modelling uncertainties.

### **Methodology**

The effect of climate change on fluvial flooding is shown here using an indicator representing the percentage change in average annual flood risk within a country, calculated by assuming a standardised relationship between flood magnitude and loss. The indicator is based on the estimated present-day (1961-1990) and future flood frequency curve, derived from the time series of runoff simulated at a spatial resolution of  $0.5^{\circ} \times 0.5^{\circ}$  using a global hydrological model, MacPDM (Gosling and Arnell, 2011). The flood frequency curve was combined with a generic flood magnitude–damage curve to estimate the average annual flood damage in each grid cell. This was then multiplied by grid cell population and summed across a region, producing in effect a population-weighted average annual damage. Flood damage is thus assumed to be proportional to population in each grid cell, not the value of exposed assets, and the proportion of people exposed to flood is assumed to be constant across each grid cell (Warren et al., 2010).

The national values are calculated across major floodplains, based on the UN PREVIEW Global Risk Data Platform ([preview.grid.unep.ch](http://preview.grid.unep.ch)). This database contains gridded estimates, at a spatial resolution of 30 arc-seconds ( $0.00833^{\circ} \times 0.00833^{\circ}$ ), of the estimated frequency of flooding. From this database the proportion of each  $0.5^{\circ} \times 0.5^{\circ}$  grid cell defined as floodplain was determined, along with the numbers of people living in each  $0.5^{\circ} \times 0.5^{\circ}$  grid cell in flood-prone areas. The floodplain data set does not include “small” floodplains, so underestimates actual exposure to flooding. The pattern of climate change from 21 GCMs was applied to MacPDM, under two emissions scenarios; 1) SRES A1B and 2) an aggressive mitigation scenario where emissions follow A1B up to 2016 but then decline at a rate of 5% per year



thereafter to a low emissions floor (denoted A1B-2016-5-L). Both scenarios assume that population changes through the 21<sup>st</sup> century following the SRES A1 scenario as implemented in IMAGE 2.3 (van Vuuren et al., 2007). The application of 21 GCMs is an attempt to quantify the uncertainty due to climate modelling, although it is acknowledged that only one impacts model is applied (MacPDM). Simulations were performed for the years 2030, 2050, 2080 and 2100. The result represents the change in flood risk due to climate change, not the change in flood risk relative to present day (Warren et al., 2010).

## Results


The results for France are presented in Figure 18. By the 2030s, the models project a range of changes in mean fluvial flooding risk over France in both scenarios, with some models projecting decreases and others increases. However, the balance is more towards lower flood risk, with over 75% of the models projecting a decrease in the average annual flood risk. The largest decline projected for the 2030s is around -40%, and the largest increase around +40%. The mean across all projections is a decrease of approximately 18%.

By 2100 the model projections become more balanced between increased and decreased flood risk in both scenarios, and the difference in projections from the different models also becomes greater. Both these aspects of the results are more pronounced for the A1B scenario than the mitigation scenario. Under the mitigation scenario, a majority models still project a lower flood risk (with a maximum decrease of 50%), but several models project an increase. The mean projection over all models is a decrease of 8%, and the largest increase is around +70%. Under the A1B scenario, less than half the models project a lower flood risk (with a maximum decrease of -60%). The largest projected increase in average annual flood risk is approximately +200%, with a mean projected increase of 10%.

So for France, the models show a greater tendency towards decreasing flood risk at first, but later in the century the models become more evenly divided between increases and decreases, especially so under the A1B scenario. The differences between the model projections are greater later in the Century and particularly for A1B.

□

□



**Figure 18.** Box and whisker plots for the percentage change in average annual flood risk within France, from 21 GCMs under two emissions scenarios (A1B and A1B-2016-5-L), for four time horizons. The plots show the 25th, 50th, and 75th percentiles (represented by the boxes), and the maximum and minimum values (shown by the extent of the whiskers).

## Tropical cyclones

Continental France is not impacted by tropical cyclones (French districts in West Indies, South America, Indian Ocean and Pacific Ocean are not considered in this review).

☐

# Coastal regions

## Headline

France's coasts and port cities could be affected by sea level rise (SLR) but to a lesser extent than many other countries across the globe. Adaptation (e.g. coastal defences) is important in mediating the impacts. Results show that by the 2080s under a high SLR scenario and without adaptation, the average annual number of people flooded in France could be around 463,000. This is greatly reduced with adaptation, to around 2,500. Under a low SLR scenario, 3,000 people could be flooded annually without adaptation and 1,800 could be flooded with adaptation. However, these results are based upon a limited number of studies and the impact of SLR for France requires further quantification. Also, some French districts in the West Indies, South America, Indian Ocean and Pacific Ocean could be affected by SLR, but these were not considered in this review.

## Supporting literature

### Assessments that include a global or regional perspective

#### Climate change studies

The IPCC AR4 concluded that at the time, understanding was too limited to provide a best estimate or an upper bound for global SLR in the twenty-first century (IPCC, 2007b). However, a range of SLR, excluding accelerated ice loss effects was published, ranging from 0.19m to 0.59m by the 2090s (relative to 1980-2000), for a range of scenarios (SRES A1FI to B1). The IPCC AR4 also provided an illustrative estimate of an additional SLR term of up to 17cm from acceleration of ice sheet outlet glaciers and ice streams, but did not suggest this is the upper value that could occur. Although there are published projections of SLR in excess of IPCC AR4 values (Nicholls et al., 2011), many of these typically use semi-empirical methods that suffer from limited physical validity and further research is required to produce a more robust estimate. Linking sea level rise projections to temperature must also be done with caution because of the different response times of these two climate variables to a given radiative forcing change.

Nicholls and Lowe (2004) previously showed that mitigation alone would not avoid all of the impacts due to rising sea levels, adaptation would likely be needed too. Recent work by van Vuuren et al. (2011) estimated that, for a world where global mean near surface temperatures reach around 2°C by 2100, global mean SLR could be 0.49m above present levels by the end of the century. Their sea level rise estimate for a world with global mean temperatures reaching 4°C by 2100 was 0.71m, suggesting around 40% of the future increase in sea level to the end of the 21<sup>st</sup> century could be avoided by mitigation. A qualitatively similar conclusion was reached in a study by Pardaens et al. (2011), which examined climate change projections from two GCMs. They found that around a third of global-mean SLR over the 21st century could potentially be avoided by a mitigation scenario under which global-mean surface air temperature is near-stabilised at around 2°C relative to pre-industrial times. Under their baseline business-as-usual scenario the projected increase in temperature over the 21st century is around 4°C, and the sea level rise range is 0.29-0.51m (by 2090-2099 relative to 1980-1999; 5% to 95% uncertainties arising from treatment of land-based ice melt and following the methodology used by the IPCC AR4). Under the mitigation scenario, global mean SLR in this study is projected to be 0.17-0.34m.

The IPCC 4th assessment (IPCCa) followed Nicholls and Lowe (2004) for estimates of the numbers of people affected by coastal flooding due to sea level rise. Nicholls and Lowe (2004) projected for the North and West Europe region that an additional 100 thousand people per year could be flooded due to sea level rise by the 2080s relative to the 1990s for the SRES A2 Scenario (note this region also includes other countries, such as UK and Norway). For the North Mediterranean region, this figure is less than 200 thousand people per year. However, it is important to note that this calculation assumed that protection standards increased as GDP increased, although there is no additional adaptation for sea level rise. More recently, Nicholls et al. (2011) also examined the potential impacts of sea level rise in a scenario that gave around 4°C of warming by 2100. Readings from Figure 3 from Nicholls *et al.* (2011) for the North and West Europe region suggest that less than an approximate 1 million additional people could be flooded for a 0.5 m SLR (assuming no additional protection), with less than 2 million additional people flooded in the north Mediterranean region. Nicholls et al. (2011) also looked at the consequence of a 2m SLR by 2100, however as we consider this rate of SLR to have a low probability we don't report these figures here.

The European Commission (2009) assessed the vulnerability of several European countries to SLR. The study showed 4.7% of France's coastline is comprised of 10km long stretches that are below 5m elevation and that 2055km is subject to erosion. The study also calculated

that 17% of GDP is located within 50km of the coast and that 26% of the country's population live within this zone.

Recent results from the PESETA (Projection of Economic impacts of climate change in Sectors of the European Union based on bottom-up Analysis) project have afforded consistent quantitative projections of the impact of SLR for several European countries (Richards and Nicholls, 2009). These are advantageous because previous European assessments have tended to be more qualitative in nature (Nicholls, 2000). The results show that while Europe is potentially highly threatened by SLR, adaptation (in the form of the two protection options considered) can greatly reduce these impacts to levels which appear manageable. The adaptation methods and costs assessed were the raising of flood dykes and the application of beach nourishment. Richards and Nicholls (2009) show that there are almost immediate benefits of adaptation, and the analysis suggests that widespread adaptation to SLR across Europe could be prudent. The assessment considered SLR projections from two GCMs, ECHAM4 and HadCM3. For each of these, SLR estimates for low, medium and high climate sensitivities were applied, and under the A2 and B2 emissions scenarios. To further quantify uncertainty, the upper and lower estimates of global SLR from the IPCC TAR (IPCC, 2001) were also applied. The estimates of global SLR considered by Richards and Nicholls (2009) are summarised in Table 11. Given that the IPCC TAR estimates of SLR encompass the full range of uncertainty that Richards and Nicholls (2009) considered, impacts for the IPCC TAR low and high scenarios are presented in Table 12. The results show that by the 2080s under the high SLR scenario and without adaptation, the average annual number of people flooded is around 463,000. This is greatly reduced with adaptation, to around 2,500. Under the low SLR scenario, 3,000 people are flooded annually without adaptation and 1,800 are flooded with adaptation. The results highlight the importance of climate sensitivity in determining the impacts as well as demonstrating clear potential benefits of adaptive measures, which by the 2080s can almost completely remove any incremental climate change effect. Adaptation could also effectively reduce the financial impacts, although it should be noted that the baseline of the damage costs for coastal flooding in France (€253.2 million), is greater than the total average cost to the whole French insurance system of all floods occurring in France. To this end, it is possible that Richards and Nicholls (2009) over-estimated the financial cost of coastal flooding for France due to cost modelling biases.

?

?

?

?

<b>GCM</b>	<b>ECHAM4</b>		<b>HadCM3</b>		<b>IPCC TAR</b>
<b>SRES scenario</b>	<b>A2</b>	<b>B2</b>	<b>A2</b>	<b>B2</b>	<b>A2/B2</b>
<b>Climate sensitivity</b>					
<i>Low</i>	29.2	22.6	25.3	19.4	9
<i>Medium</i>	43.8	36.7	40.8	34.1	
<i>High</i>	58.5	50.8	56.4	48.8	88

**Table 11.** Global SLR (cm) for low, medium and high climate sensitivities at 2100, for the A2 and B2 SRES scenarios, that were applied by Richards and Nicholls (2009).

Country	Indicator	Baseline (1995)	IPCC A2 2020s Low SLR		IPCC A2 2020s High SLR		IPCC A2 2080s Low SLR		IPCC A2 2080s High SLR	
			No adaptation	With adaptation	No adaptation	With adaptation	No adaptation	With adaptation	No adaptation	With adaptation
France	Total damage costs (millions €/year)	253.2	362.7	228.8	434.9	291.7	673.3	220.9	3359.8	456.6
	Land loss (submergence) (km <sup>2</sup> /year)	0	0.2	0.2	0.6	0.6	0.2	0.2	130.8	1.6
	Migration due to land loss (1000s of people/year)	0	0	0	0	0	0	0	19.5	0
	Net loss of wetland area (km <sup>2</sup> /year)	12.4	300.7	300.7	688.7	688.7	950.9	950.9	1963.8	1963.8
	People actually flooded (1000s/year)	1.96	2.4	2.0	2.9	2.1	3.0	1.8	463.7	2.5
	Salinity intrusion costs (millions €/year)	135.85	145.3	145.3	153.5	153.5	199.9	199.9	268.7	268.7
	Total damage costs (millions €/year)	428.1	473.7	398.9	590.3	448.3	782.2	344.6	2607.6	686.9
Germany	Land loss (submergence) (km <sup>2</sup> /year)	0	0	0	0	0	0	0	178.1	0
	Migration due to land loss (1000s of people/year)	0	0	0	0	0	0	0	14.3	0.1
	Net loss of wetland area (km <sup>2</sup> /year)	0.3	146.4	146.4	418.3	418.3	673.2	673.2	1938.5	1938.5
	People actually flooded (1000s/year)	2.23	2.6	2.3	3.7	2.5	4.4	2.1	293.3	2.9
	Salinity intrusion costs (millions €/year)	274.78	240.8	240.8	251.2	251.2	343.3	343.3	431.4	431.4
	Total damage costs (millions €/year)	8.5	18.5	6.6	271.9	7.1	444.2	9.6	8044.2	17.0
	Land loss (submergence) (km <sup>2</sup> /year)	0	0	0	0	0	0	0	188.5	0.1
Italy	Migration due to land loss (1000s of people/year)	0	0	0	0	0	0	0	55.2	0
	Net loss of wetland area (km <sup>2</sup> /year)	9.6	78.2	78.2	155.5	155.5	221.8	221.8	393.7	393.7
	People actually flooded (1000s/year)	1.49	1.9	1.5	3.1	1.6	3.3	1.5	512.9	2.3
	Salinity intrusion costs (millions €/year)	5.84	6.4	6.4	6.7	6.7	9.2	9.2	11.9	11.9
	Total damage costs (millions €/year)	8.5	18.5	6.6	271.9	7.1	444.2	9.6	8044.2	17.0



<b>Spain</b>	Total damage costs (millions €/year)	15.3	23.6	9.4	55.6	14.5	75.6	3.9	749.7	32.0	
	Land loss (submergence) (km <sup>2</sup> /year)	0	0	0	0	0	0	0	38.3	0.1	
	Migration due to land loss (1000s of people/year)	0	0	0	0	0	0	0	7.0	0	
	Net loss of wetland area (km <sup>2</sup> /year)	0.2	64.4	64.4	174.3	174.3	174.3	242.5	242.5	659.8	659.8
	People actually flooded (1000s/year)	0.73	0.9	0.7	1.3	0.8	1.1	1.1	0.7	321.8	1.1
	Salinity intrusion costs (millions €/year)	0.92	1.0	1.0	1.0	1.0	1.4	1.4	1.4	1.8	1.8
	Total damage costs (millions €/year)	189.5	414.2	50.1	743.5	136.4	1087.0	6.3	10062.6	155.1	155.1
	Land loss (submergence) (km <sup>2</sup> /year)	0	0	0	0	0	0	0	0	239.3	0
	Migration due to land loss (1000s of people/year)	0	0	0	0	0	0	0	0	73.6	0
<b>UK</b>	Net loss of wetland area (km <sup>2</sup> /year)	8.5	175.1	175.1	522.6	522.6	789.4	789.4	1690.2	1690.2	1690.2
	People actually flooded (1000s/year)	4.85	5.7	4.4	6.1	4.9	6.8	4.3	986.3	5.6	5.6
	Salinity intrusion costs (millions €/year)	2.32	2.5	2.5	2.8	2.8	3.6	3.6	5.7	5.7	5.7

**Table 12.** Estimates of the impact of SLR with the low and high estimates of SLR from the IPCC TAR in the 2020s and 2080s, and assuming no adaptation or adaptation. Baseline impacts (1995) are also presented. Data is from Richards and Nicholls (2009).

Hanson et al. (2010) investigated population exposure to global SLR, natural and human subsidence/uplift, and more intense storms and higher storm surges, for 136 port cities across the globe. Future city populations were calculated using global population and economic projections, based on the SRES A1 scenario up to 2030. The study accounted for uncertainty on future urbanization rates, but estimates of population exposure were only presented for a rapid urbanisation scenario, which involved the direct extrapolation of population from 2030 to 2080. All scenarios assumed that new inhabitants of cities in the future will have the same relative exposure to flood risk as current inhabitants. The study is similar to a later study presented by Hanson et al. (2011) except here, different climate change scenarios were considered, and published estimates of exposure are available for more countries, including France. Future water levels were generated from temperature and thermal expansion data related to greenhouse gas emissions with SRES A1B (un-mitigated climate change) and under a mitigation scenario where emissions peak in 2016 and decrease subsequently at 5% per year to a low emissions floor (2016-5-L). Table 13 shows the aspects of SLR that were considered for various scenarios and Table 14 displays port city population exposure for each scenario in the 2030s, 2050s and 2070s. The result show that France is one of the countries where port cities are likely to impacted least by SLR (e.g. compare the projections in Table 14 with the estimates for exposure in the absence of climate change that are presented in Table 15). For example, at present, 13,000 people in port cities are exposed to SLR in France. By the 2070s in the absence of climate change 18,000 are exposed. With climate change in the 2070s, 23,000 people are exposed. Hanson et al. (2010) also demonstrated that aggressive mitigation could avoid an exposure of around 2,000 people in France, relative to un-mitigated climate change (see Table 15).

Scenario		Water				
		Climate			Subsidence	
Code	Description	More intense storms	Sea-level change	Higher storm surges	Natural	Anthropogenic
FNC	Future city	V	x	x	X	x
FRSLC	Future City Sea-Level Change	V	V	x	V	x
FCC	Future City Climate Change	V	V	V	V	x
FAC	Future City All Changes	V	V	V	V	V

**Table 13.** Summary of the aspects of SLR considered by Hanson et al. (2010). 'V' denotes that the aspect was considered in the scenario and 'x' that it was not.

Rapid urbanisation projection																	
2030						2050						2070					
Country	Ports	Water level projection			Country	Ports	Water level projection			Country	Ports	Water level projection					
		FAC	FCC	FRSL C			FNC	FAC	FCC			FRSLC	FNC	FAC	FCC	FRSL C	FNC
CHINA	15	17,100	15,500	15,400	14,600	CHINA	15	23,000	19,700	18,700	17,400	CHINA	15	27,700	22,600	20,800	18,600
INDIA	6	11,600	10,800	10,300	9,970	INDIA	6	16,400	14,600	13,600	12,500	INDIA	6	20,600	17,900	15,600	13,900
US	17	8,990	8,960	8,830	8,460	US	17	11,300	11,200	10,800	9,970	US	17	12,800	12,700	12,100	10,700
JAPAN	6	5,260	4,610	4,430	4,390	JAPAN	6	6,440	5,280	5,000	4,760	JAPAN	6	7,800	5,970	5,580	5,070
INDONESIA	4	1,420	1,200	1,200	1,170	INDONESIA	4	2,110	1,610	1,610	1,500	INDONESIA	4	2,680	1,830	1,830	1,530
BRAZIL	10	833	833	833	802	BRAZIL	10	929	929	929	879	BRAZIL	10	940	940	940	864
UK	2	497	497	478	459	UK	2	609	609	564	521	UK	2	716	716	640	569
CANADA	2	459	433	422	405	CANADA	2	549	512	486	457	CANADA	2	614	585	545	489
REP. OF KOREA	3	344	344	331	441	REP. OF KOREA	3	361	361	341	318	REP. OF KOREA	3	377	377	325	303
GERMANY	1	257	257	253	248	GERMANY	1	287	287	273	269	GERMANY	1	309	309	290	280
RUSSIA	1	177	177	177	177	RUSSIA	1	202	202	173	173	RUSSIA	1	226	226	197	169
AUSTRALIA	5	162	162	157	157	AUSTRALIA	5	197	197	191	181	AUSTRALIA	5	196	196	186	175
SAUDI ARABIA	1	24	24	24	22	SAUDI ARABIA	1	33	33	33	27	SAUDI ARABIA	1	38	38	38	29
SOUTH AFRICA	2	30	30	30	29	SOUTH AFRICA	2	28	28	28	27	SOUTH AFRICA	2	30	30	30	27
FRANCE	1	15	15	15	15	FRANCE	1	19	19	19	17	FRANCE	1	23	23	23	18
ITALY	1	2	2	2	2	ITALY	1	4	4	4	3	ITALY	1	6	6	6	4
MEXICO	0	0	0	0	0	MEXICO	0	0	0	0	0	MEXICO	0	0	0	0	0

**Table 14.** Estimates of port city population exposure (1,000s) for each water level projection (ranked according to exposure with the FAC (Future City All Changes) scenario) under a rapid urbanisation projection for the 2030s, 2050s and 2070s. Estimates for present day exposure and in the absence of climate change (for 2070 only) for comparison are presented in Table 15. Data is from Hanson et al. (2010) and has been rounded down to three significant figures.

Country	Ports	Population exposure				Exposure avoided
		Current	2070. Rapid urbanisation, FAC water level scenario			
			No climate change	A1B un-mitigated	Mitigated (2016-5-L)	
CHINA	15	8,740	18,600	27,700	26,500	1,140
UNITED STATES	17	6,680	10,700	12,800	12,300	505
RUSSIA	1	189	169	226	197	28
JAPAN	6	3,680	5,070	7,800	7,290	515
SOUTH AFRICA	2	24	27	30	29	0
INDIA	6	5,540	13,900	20,600	18,900	1,670
BRAZIL	10	555	864	940	926	14
MEXICO	0	0	0	0	0	0
CANADA	2	308	489	614	599	15
AUSTRALIA	5	99	175	196	190	6
INDONESIA	4	602	1,530	2,680	2,520	156
REP. OF KOREA	3	294	303	377	343	34
UK	2	414	569	716	665	51
FRANCE	1	13	18	23	20	2
ITALY	1	2	4	6	6	0
GERMANY	1	261	280	309	295	15
SAUDI ARABIA	1	15	29	38	35	3

**Table 15.** Exposed port-city population (1,000s) in present (current), and in the 2070s in the absence of climate change (no climate change), with unmitigated climate change (A1B un-mitigated), and mitigated climate change (mitigated 2016-5-L), under the rapid urbanisation and (Future City All Changes) water level scenarios. The final column shows the potential avoided exposure, as a result of mitigation. Data is from Hanson et al. (2010) and has been rounded down to three significant figures.

To further quantify the impact of SLR and some of the inherent uncertainties, the DIVA model was used to calculate the number of people flooded per year for global mean sea level increases (Brown et al., 2011). The DIVA model (DINAS-COAST, 2006) is an integrated model of coastal systems that combines scenarios of water level changes with socio-economic information, such as increases in population. The study uses two climate scenarios; 1) the SRES A1B scenario and 2) a mitigation scenario, RCP2.6. In both cases an SRES A1B population scenario was used. The results are shown in Table 16.

?

?	A1B?		RCP?	?
?	Low?	High?	Low?	High?
Additional people flooded (1000s)?	7.00	124.01	4.62?	37.25
Loss of wetlands area (% of country's total wetland)?	33.45%?	48.46%?	29.28%?	47.23%?

**Table 16.** Number of additional people flooded (1000s), and percentage of total wetlands lost by the 2080s under the high and low SRES A1B and mitigation (RCP 2.6) scenarios (Brown et al., 2011).

## National-scale or sub-national scale assessments

### Climate change studies

Poumadère (2008) conducted an expert and stakeholder elicitation exercise to investigate the national response to a 5-6m SLR induced by the collapse of the West Antarctic Ice Sheet (WAIS) for the period 2030–2130, for the Rhone delta. Group process analyses demonstrate that there was a high level of trust and cooperation between stakeholders to face extreme SLR issue, despite potentially diverging interests. A conservative “wait and see” option was decided when the risk of the WAIS collapse was announced in 2030 but when the WAIS collapse generates an effective 1m SLR by 2050, decisions are taken for total retreat and rendering of the Rhone delta to its hydrological function.

The estimates presented by the European-scale assessment of Richards and Nicholls (2009) are consistent with the coastal impacts from SLR in France that are presented by the Ministère de l'Écologie, du Développement durable, des transports et du Logement (MEEDDM, 2008b).

Vinchon et al. (2009) demonstrated that the Aquitaine and Languedoc Roussillon regions of France are particularly vulnerable to SLR. Moreover, the authors argue that there is a need for better knowledge on physical resilience capacity of the coast, assets values and vulnerability, and the urge of an integrated management approach of coastal risk in France. Vinchon et al. (2011) also note the vulnerability of Languedoc Roussillon to SLR. The authors demonstrated that a 1m SLR by 2100 could result in 4000 ha of coastal land in Languedoc Roussillon being permanently lost, including existing beaches and natural lands, as well as 1000 ha of urbanised areas. However, it should be noted that dynamics of simulated SLR in the Mediterranean basin are highly uncertain (Tsimplis et al., 2008).

□

□

## References

- AINSWORTH, E. A. & MCGRATH, J. M. 2010. Direct Effects of Rising Atmospheric Carbon Dioxide and Ozone on Crop Yields. *In: LOBELL, D. & BURKE, M. (eds.) Climate Change and Food Security*. Springer Netherlands.
- ALLISON, E. H., PERRY, A. L., BADJECK, M.-C., NEIL ADGER, W., BROWN, K., CONWAY, D., HALLS, A. S., PILLING, G. M., REYNOLDS, J. D., ANDREW, N. L. & DULVY, N. K. 2009. Vulnerability of national economies to the impacts of climate change on fisheries. *Fish and Fisheries*, 10, 173-196.
- ARNELL, N., OSBORNE, T., HOOKER, J., DAWSON, T., PERRYMAN, A. & WHEELER, T. 2010a. Simulation of AVOIDed impacts on crop productivity and food security. *Work stream 2, Report 17 of the AVOID programme (AV/WS2/D1/R17)*.
- ARNELL, N., WHEELER, T., OSBORNE, T., ROSE, G., GOSLING, S., DAWSON, T., PENN, A. & PERRYMAN, A. 2010b. The implications of climate policy for avoided impacts on water and food security. *Work stream 2, Report 6 of the AVOID programme (AV/WS2/D1/R06)*. London: Department for Energy and Climate Change (DECC).
- ARNELL, N. W. 2004. Climate change and global water resources: SRES emissions and socio-economic scenarios. *Global Environmental Change*, 14, 31-52.
- AVNER, S., MAUZERALL, D. L., LIU, J. F. & HOROWITZ, L. W. 2011. Global crop yield reductions due to surface ozone exposure: 2. Year 2030 potential crop production losses and economic damage under two scenarios of O<sub>3</sub> pollution. *Atmospheric Environment*, 45, 2297-2309.
- BALMASEDA, M. A., SMITH, G. C., HAINES, K., ANDERSON, D., PALMER, T. N. & VIDARD, A. 2007. Historical reconstruction of the Atlantic Meridional Overturning Circulation from the ECMWF operational ocean reanalysis. *Geophysical Research Letters*, 34, -.
- BATES, B. C., KUNDZEWICZ, Z. W., WU, S. & PALUTIKOF, J. 2008. Climate Change and Water. *Technical Paper of the Intergovernmental Panel on Climate Change, IPCC Secretariat, Geneva*, 210pp.
- BENESTAD, R. 2008. Heating degree days, Cooling degree days, and precipitation in Europe: Analysis for the CELECT-project.

BENISTON, M. 2004. The 2003 heat wave in Europe: A shape of things to come? An analysis based on Swiss climatological data and model simulations. *Geophysical Research Letters*, 31, L02202.

BENISTON, M., STEPHENSON, D. B., CHRISTENSEN, O. B., FERRO, C. A. T., FREI, C., GOYETTE, S., HALSNAES, K., HOLT, T., JYLHA, K., KOFFI, B., PALUTIKOF, J., SCHOLL, R., SEMMLER, T. & WOTH, K. 2007. Future extreme events in European climate: an exploration of regional climate model projections. *Climatic Change*, 81, 71-95.

BLENKINSOP, S. & FOWLER, H. J. 2007. Changes in European drought characteristics projected by the PRUDENCE regional climate models. *International Journal of Climatology*, 27, 1595-1610.

BOE, J. & TERRAY, L. 2008. A weather-type approach to analyzing winter precipitation in France: Twentieth-century trends and the role of anthropogenic forcing. *Journal of Climate*, 21, 3118-3133.

BOÉ, J., TERRAY, L., MARTIN, E. & HABETS, F. 2009. Projected changes in components of the hydrological cycle in French river basins during the 21st century. *Water Resources Research*, 45, W08426.

BONAZZA, A., MESSINA, P., SABBIONI, C., GROSSI, C. M. & BRIMBLECOMBE, P. 2009a. Mapping the impact of climate change on surface recession of carbonate buildings in Europe. *Science of The Total Environment*, 407, 2039-2050.

BONAZZA, A., SABBIONI, C., MESSINA, P., GUARALDI, C. & DE NUNTIIS, P. 2009b. Climate change impact: Mapping thermal stress on Carrara marble in Europe. *Science of The Total Environment*, 407, 4506-4512.

BOTZEN, W. J. W. & VAN DEN BERGH, J. C. J. M. 2008. Insurance against climate change and flooding in the Netherlands: Present, future, and comparison with other countries. *Risk Analysis*, 28, 413-426.

BRISSON, N., GATE, P., GOUACHE, D., CHARMET, G., OURY, F. X. & HUARD, F. 2010. Why are wheat yields stagnating in Europe? A comprehensive data analysis for France. *Field Crops Research*, 119, 201-212.

BRISSON, N. & LEVRAULT, F. 2010. *Climate change, agriculture and forests in France: simulations of the impacts on the main species. The Green Book of the CLIMATOR project (2007-2010)*. ADEME.



BROWN, S., NICHOLLS, R., LOWE, J.A. and PARDAENS, A. (2011), Sea level rise impacts in 24 countries. Faculty of Engineering and the Environment and Tyndall Centre for Climate Change Research, University of Southampton.

BRYDEN, H. L., LONGWORTH, H. R. & CUNNINGHAM, S. A. 2005. Slowing of the Atlantic meridional overturning circulation at 25 degrees N. *Nature*, 438, 655-657.

CCR 2007. Natural disasters in France. Paris: Caisse Centrale de Reassurance.

CHAKRABORTY, S. & NEWTON, A. C. 2011. Climate change, plant diseases and food security: an overview. *Plant Pathology*, 60, 2-14.

CHATRY, C., LE QUENTREC, M., LAURENS, D., LE GALLOU, J.-Y., LAFITTE, J.-J. & CREUCHET, B. 2010. Changement climatique et extension des zones sensibles aux feux de forêt. Ministère De L'Alimentation, De L' Agriculture Et De La Peche.

CIA 2011. World Factbook. US Central Intelligence Agency.

CISCAR, J. C., IGLESIAS, A., FEYEN, L., GOODESS, C. M., SZABÓ, L., CHRISTENSEN, O. B., NICHOLLS, R., AMELUNG, B., WATKISS, P., BOSELLO, F., DANKERS, R., GARROTE, L., HUNT, A., HORROCKS, L., MONEO, M., MORENO, M., PYE, S., QUIROGA, A., VAN REGEMORTER, D., RICHARDS, J., ROSON, R. & SORIA, A. 2009. Climate change impacts in Europe. Final report of the PESETA research project. *In: CISCAR, J. C. (ed.) EUR 24093 EN, JRC Scientific and Technical Reports.*

CORTI, T., MUCCIONE, V., KÖLLNER-HECK, P., BRESCH, D. & SENEVIRATNE, S. I. 2009. Simulating past droughts and associated building damages in France. *Hydrol. Earth Syst. Sci.*, 13, 1739-1747.

CUNNINGHAM, S. A., KANZOW, T., RAYNER, D., BARINGER, M. O., JOHNS, W. E., MAROTZKE, J., LONGWORTH, H. R., GRANT, E. M., HIRSCHI, J. J. M., BEAL, L. M., MEINEN, C. S. & BRYDEN, H. L. 2007. Temporal variability of the Atlantic meridional overturning circulation at 26.5 degrees N. *Science*, 317, 935-938.

DANKERS, R. & FEYEN, L. 2008. Climate change impact on flood hazard in Europe: An assessment based on high-resolution climate simulations. *Journal of Geophysical Research-Atmospheres*, 113.

DANKERS, R. & FEYEN, L. 2009. Flood hazard in Europe in an ensemble of regional climate scenarios. *Journal of Geophysical Research-Atmospheres*, 114.

- DELWORTH, T. L. & DIXON, K. W. 2006. Have anthropogenic aerosols delayed a greenhouse gas-induced weakening of the North Atlantic thermohaline circulation? *Geophysical Research Letters*, 33, -.
- DEQUE, M. 2007. Frequency of precipitation and temperature extremes over France in an anthropogenic scenario: Model results and statistical correction according to observed values *Global and Planetary Change*, Vol. 57, No. 1-2. pp. 16-26.
- DINAS-COAST Consortium. 2006 DIVA 1.5.5. Potsdam, Germany: Potsdam Institute for Climate Impact Research (on CD-ROM).
- DOLL, P. 2009. Vulnerability to the impact of climate change on renewable groundwater resources: a global-scale assessment. *Environmental Research Letters*, 4.
- DOLL, P. & SIEBERT, S. 2002. Global modeling of irrigation water requirements. *Water Resources Research*. Vol: 38 Issue: 4. Doi: 10.1029/2001WR000355 .
- DUCHARNE, A., SAUQUET, E., HABETS, F., DÉQUÉ, M., GASCOIN, S., HACHOUR, A., MARTIN, E., OUDIN, L., PAGÉ, C., TERRAY, L., THIÉRY, D. & VIENNOT, P. 2011. Évolution potentielle du régime des crues de la Seine sous changement climatique (Potential evolution of the Seine River flood regime under climate change). *La Houille Blanche* 1, 51–57.
- ESKELAND, G. A. & MIDEKSA, T. K. 2009. Climate Change and Residential Electricity Demand in Europe. *Center for International Climate and Environmental Research (CICERO)*.
- EUROPEAN COMMISSION, E. C. 2009. The economics of climate change adaptation in EU coastal areas. Brussels, Belgium: European Commission.
- FALKENMARK, M., ROCKSTRÖM, J. & KARLBERG, L. 2009. Present and future water requirements for feeding humanity. *Food Security*, 1, 59-69.
- FAO. 2008. *Food and Agricultural commodities production* [Online]. Available: <http://faostat.fao.org/site/339/default.aspx> [Accessed 1 June 2011].
- FAO. 2010. *Food Security country profiles* [Online]. Available: <http://www.fao.org/economic/ess/ess-fs/ess-fs-country/en/> [Accessed 1 Sept 2011].

FERRISE, R., MORIONDO, M. & BINDI, M. 2011. Probabilistic assessments of climate change impacts on durum wheat in the Mediterranean region. *Nat. Hazards Earth Syst. Sci.*, 11, 1293-1302.

FILLEUL, L., CASSADOU, S., MÉDINA, S., FABRES, P., LEFRANC, A., EILSTEIN, D., LE TERTRE, A., PASCAL, L., CHARDON, B. & BLANCHARD, M. 2006. The relation between temperature, ozone, and mortality in nine French cities during the heat wave of 2003. *Environmental health perspectives*, 114, 1344.

FOUILLET, A., REY, G., WAGNER, V., LAAIDI, K., EMPEREUR-BISSONNET, P., LE TERTRE, A., FRAYSSINET, P., BESSEMOULIN, P., LAURENT, F. & DE CROUY-CHANEL, P. 2008. Has the impact of heat waves on mortality changed in France since the European heat wave of summer 2003? A study of the 2006 heat wave. *International journal of epidemiology*, 37, 309.

FRONZEK, S. & CARTER, T. 2007. Assessing uncertainties in climate change impacts on resource potential for Europe based on projections from RCMs and GCMs. *Climatic Change*, 81, 357-371.

FSF 2010. Financial Standards Report France: Insurance Core Principles. Financial Standards Foundation.

GALLIOT, M., MONLÉON, M., REYSSET, B., RUSSO, L. & VERGÈS, P. 2009. Climate change: costs of impacts and lines of adaptation. Paris, France: National Observatory for the Impacts of Global Warming.

GATE, P., BRISSON, N. & GOUACHE, D. 2010. Les Causes du plafonnement du rendement du ble en France: D'abord une origine climatique. *Comptes rendus de l'Académie d'agriculture de France*, 96, 17-23.

GERTEN D., SCHAPHOFF S., HABERLANDT U., LUCHT W., SITCH S. 2004 . Terrestrial vegetation and water balance: hydrological evaluation of a dynamic global vegetation model. *International Journal Water Resource Development*, 286,249–270.

GIANNAKOPOULOS, C., BINDI, M., MORIONDO, M., LESAGER, P. & TIN, T. 2005. Climate change impacts in the Mediterranean resulting from a 2C global temperature rise. *WWF Report*. Gland, Switzerland: WWF.

GIANNAKOPOULOS, C., LE SAGER, P., BINDI, M., MORIONDO, M., KOSTOPOULOU, E. & GOODESS, C. M. 2009. Climatic changes and associated impacts in the Mediterranean resulting from a 2 °C global warming. *Global and Planetary Change*, 68, 209-224.

GORNALL, J., BETTS, R., BURKE, E., CLARK, R., CAMP, J., WILLETT, K., WILTSHIRE, A. 2010. Implications of climate change for agricultural productivity in the early twenty-first century. *Phil. Trans. R. Soc. B*, DOI: 10.1098/rstb.2010.0158.

GOSLING, S., TAYLOR, R., ARNELL, N. & TODD, M. 2011. A comparative analysis of projected impacts of climate change on river runoff from global and catchment-scale hydrological models. *Hydrology and Earth System Sciences*, 15, 279–294.

GOSLING, S. N. & ARNELL, N. W. 2011. Simulating current global river runoff with a global hydrological model: model revisions, validation, and sensitivity analysis. *Hydrological Processes*, 25, 1129-1145.

GOSLING, S. N., BRETHERTON, D., HAINES, K. & ARNELL, N. W. 2010. Global hydrology modelling and uncertainty: running multiple ensembles with a campus grid. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 368, 4005-4021.

GOUBANOVA, K. & Li, L., 2007. Extremes in temperature and precipitation around the Mediterranean basin in an ensemble of future climate scenario simulations. *Global & Planetary Change*, 57, 27-42.

HANSON, S., NICHOLLS, R., RANGER, N., HALLEGATTE, S., CORFEE-MORLOT, J., HERWEIJER, C. & CHATEAU, J. 2011. A global ranking of port cities with high exposure to climate extremes. *Climatic Change*, 104, 89-111.

HANSON, S., NICHOLLS, R., S, H. & CORFEE-MORLOT, J. 2010. The effects of climate mitigation on the exposure of worlds large port cities to extreme coastal water levels. London, UK.

HARDING, R., BEST, M., BLYTH, E., HAGEMANN, D., KABAT, P., TALLAKSEN, L.M., WARNAARS, T., WIBERG, D., WEEDON, G.P., van LANEN, H., LUDWIG, F., HADDELAND, I. 2011. Preface to the “Water and Global Change (WATCH)” special collection: Current knowledge of the terrestrial global water cycle. *Journal of Hydrometeorology*, DOI: 10.1175/JHM-D-11-024.1

- HOFMANN, M. & RAHMSTORF, S. 2009. On the stability of the Atlantic meridional overturning circulation. *Proceedings of the National Academy of Sciences of the United States of America*, 106, 20584-20589.
- HOUGHTON, J. T., DING, Y., GRIGGS, D. J., NOGUER, M., VAN DER LINDEN, P. J., DAI, X., MASKELL, K. & JOHNSON, C.A. (Eds.). IPCC (2001) *Climate Change 2001: The Scientific Basis*. Cambridge: Cambridge University.
- IFPRI. 2010. *International Food Policy Research Institute (IFPRI) Food Security CASE maps. Generated by IFPRI in collaboration with StatPlanet*. [Online]. Available: [www.ifpri.org/climatechange/casemaps.html](http://www.ifpri.org/climatechange/casemaps.html) [Accessed 21 June 2010].
- IGLESIAS, A., GARROTE, L., QUIROGA, S. & MONEO, M. 2009. Impacts of climate change in agriculture in Europe. PESETA-Agriculture study. *JRC Scientific and Technical Reports*.
- IGLESIAS, A. & ROSENZWEIG, C. 2009. Effects of Climate Change on Global Food Production under Special Report on Emissions Scenarios (SRES) Emissions and Socioeconomic Scenarios: Data from a Crop Modeling Study. Palisades, NY: Socioeconomic Data and Applications Center (SEDAC), Columbia University.
- IPCC 2007a. *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* In: SOLOMON, S., QIN, D., MANNING, M., CHEN, Z., MARQUIS, M., AVERYT, K. B., TIGNOR, M. & MILLER, H. L. (eds.). Cambridge, United Kingdom and New York, NY, USA.
- IPCC 2007b. Summary for Policymakers. In: PARRY, M. L., CANZIANI, O. F., PALUTIKOF, J. P., VAN DER LINDEN, P. J. & HANSON, C. E. (eds.) *Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge: Cambridge University Press.
- JOMELLI, V., BRUNSTEIN, D., DEQUE, M., VRAC, M. & GRANCHER, D. 2009. Impacts of future climatic change (2070-2099) on the potential occurrence of debris flows: a case study in the Massif des Ecrins (French Alps). *Climatic Change*, 97, 171-191.
- KANZOW, T., CUNNINGHAM, S. A., RAYNER, D., HIRSCHI, J. J. M., JOHNS, W. E., BARINGER, M. O., BRYDEN, H. L., BEAL, L. M., MEINEN, C. S. & MAROTZKE, J. 2007.

Observed flow compensation associated with the MOC at 26.5 degrees N in the Atlantic. *Science*, 317, 938-941.

KRIEGLER, E., HALL, J. W., HELD, H., DAWSON, R. & SCHELLNHUBER, H. J. 2009. Imprecise probability assessment of tipping points in the climate system. *Proceedings of the National Academy of Sciences of the United States of America*, 106, 5041-5046.

KUHLBRODT, T., RAHMSTORF, S., ZICKFELD, K., VIKEBO, F. B., SUNDBY, S., HOFMANN, M., LINK, P. M., BONDEAU, A., CRAMER, W. & JAEGER, C. 2009. An Integrated Assessment of changes in the thermohaline circulation. *Climatic Change*, 96, 489-537.

LAGADEC, P. 2004. Understanding the French 2003 heat wave experience: beyond the heat, a multi-layered challenge. *Journal of Contingencies and Crisis Management*, 12, 160-169.

LE TERTRE, A., LEFRANC, A., EILSTEIN, D., DECLERCQ, C., MEDINA, S., BLANCHARD, M., CHARDON, B. Œ., FABRE, P., FILLEUL, L. & JUSOT, J. F. 2006. Impact of the 2003 heatwave on all-cause mortality in 9 French cities. *Epidemiology*, 17, 75.

LEHNER, B., CZISCH, G. & VASSOLO, S. 2005. The impact of global change on the hydropower potential of Europe: a model-based analysis. *Energy Policy*, 33, 839-855.

LEHNER, B., DOLL, P., ALCAMO, J., HENRICHS, T. & KASPAR, F. 2006. Estimating the impact of global change on flood and drought risks in europe: A continental, integrated analysis. *Climatic Change*, 75, 273-299.

LÉTARD, V., FLANDRE, H. & LEPELTIER, S. 2004. La France et les Français face à la canicule: les leçons d'une crise. Report No. 195 (2003-2004) to the Sénat.

LINDGREN, E. & JAENSON, T. G. T. 2006. Lyme borreliosis in Europe: influences of climate and climate change, epidemiology, ecology and adaptation measures. Denmark: WHO Regional Office for Europe Copenhagen.

LOBELL, D. B., SCHLENKER, W. & COSTA-ROBERTS, J. 2011. Climate Trends and Global Crop Production Since 1980. *Science*.

LÓPEZ-MORENO, J. & BENISTON, M. 2009. Daily precipitation intensity projected for the 21st century: seasonal changes over the Pyrenees. *Theoretical and Applied Climatology*, 95, 375-384.

LUCK, J., SPACKMAN, M., FREEMAN, A., TREBICKI, P., GRIFFITHS, W., FINLAY, K. & CHAKRABORTY, S. 2011. Climate change and diseases of food crops. *Plant Pathology*, 60, 113-121.

MEEDDM 2008a. Evaluation du cout des impacts du changement climatique de de l'adaptation en France. Rapport de la deuxième phase Partie II – Rapports des groupes transversaux. . Paris: Ministère de l'Ecologie, du Développement durable, des transports et du Logement.

MEEDDM 2008b. Evaluation du coût des impacts du changement climatique et de l'adaptation en France. Rapport de la seconde phase Partie III – Rapports des groupes sectoriels Ministère de l'Ecologie, du Développement durable, des transports et du Logement.

MEEHL, G. A. & TEBALDI, C. 2004. More intense, more frequent, and longer lasting heat waves in the 21st century. *Science*, 305, 994.

MENZEL, L. & MATOVELLE, A. 2010. Current state and future development of blue water availability and blue water demand: A view at seven case studies. *Journal of Hydrology*, 384, 245-263.

MIKOLAJEWICZ, U., GROGER, M., MAIER-REIMER, E., SCHURGERS, G., VIZCAINO, M. & WINGUTH, A. M. E. 2007. Long-term effects of anthropogenic CO<sub>2</sub> emissions simulated with a complex earth system model. *Climate Dynamics*, 28, 599-631.

MOATAR, F., DUCHARNE, A., THIÉRY, D., BUSTILLO, V., SAUQUET, E. & VIDAL, J.-P. 2010. La Loire à l'épreuve du changement climatique. *Géosciences* 12, 78-87.

MORIONDO, M., BINDI, M., KUNDZEWICZ, Z. W., SZWED, M., CHORYNSKI, A., MATCZAK, P., RADZIEJEWSKI, M., MCEVOY, D. & WREFORD, A. 2010. Impact and adaptation opportunities for European agriculture in response to climatic change and variability. (Special Issue: Assessing adaptation to extreme weather events in Europe.). *Mitigation and Adaptation Strategies for Global Change*, 15, 657-679.

NELSON, G. C., ROSEGRANT, M. W., PALAZZO, A., GRAY, I., INGERSOLL, C., ROBERTSON, R., TOKGOZ, S., ZHU, T., SULSER, T. & RINGLER, C. 2010. Food Security, Farming and Climate Change to 2050. *Research Monograph, International Food Policy Research Institute*. Washington, DC.

NICHOLLS, R. 2000. Coastal zones. In: PARRY, M. (ed.) *Assessment of potential effects and adaptations for climate change in Europe: The Europe ACACIA Project*. Norwich, UK: University of East Anglia.

NICHOLLS, R. J. and LOWE, J. A. (2004). "Benefits of mitigation of climate change for coastal areas." *Global Environmental Change* 14(3): 229-244.

NICHOLLS, R. J., MARINOVA, N., LOWE, J. A., BROWN, S., VELLINGA, P., DE GUSMÃO, G., HINKEL, J. and TOL, R. S. J. (2011). "Sea-level rise and its possible impacts given a 'beyond 4°C world' in the twenty-first century." *Philosophical Transactions of the Royal Society A* 369: 1-21.

OLESEN, J. E., CARTER, T. R., DIAZ-AMBRONA, C. H., FRONZEK, S., HEIDMANN, T., HICKLER, T., HOLT, T., MINGUEZ, M. I., MORALES, P., PALUTIKOF, J. P., QUEMADA, M., RUIZ-RAMOS, M., RUBAEK, G. H., SAU, F., SMITH, B. & SYKES, M. T. 2007.

Uncertainties in projected impacts of climate change on European agriculture and terrestrial ecosystems based on scenarios from regional climate models. (Special Issue: Prediction of regional scenarios and uncertainties for defining European climate change risks and effects: the PRUDENCE project.). *Climatic Change*, 81, 123-143.

PARDAENS, A. K., LOWE, J., S, B., NICHOLLS, R. & DE GUSMÃO, D. 2011. Sea-level rise and impacts projections under a future scenario with large greenhouse gas emission reductions. *Geophysical Research Letters*, 38, L12604.

PARRY, M. L., ROSENZWEIG, C., IGLESIAS, A., LIVERMORE, M. & FISCHER, G. 2004. Effects of climate change on global food production under SRES emissions and socio-economic scenarios. *Global Environmental Change-Human and Policy Dimensions*, 14, 53-67.

PEINGS, Y., JAMOUS, M., PLANTON, S. & LE TREUT, H. 2011. Scénarios climatiques : indices sur la France métropolitaine pour les modèles français ARPEGE-Climat et LMDz et quelques projections pour les DOM-COM. Toulouse: Météo France.

POUMADÈRE, M., MAYS, C., PFEIFLE, G. & VAFEIDIS, A. 2008. Worst case scenario as stakeholder decision support: a 5- to 6-m sea level rise in the Rhone delta, France. *Climatic Change*, 91, 123-143.



QUINTANA-SEGUI, P., HABETS, F. & MARTIN, E. 2011. Comparison of past and future Mediterranean high and low extremes of precipitation and river flow projected using different statistical downscaling methods. *Natural Hazards and Earth System Sciences*, 11, 1411-1432.

QUINTANA SEGUÍ, P., RIBES, A., MARTIN, E., HABETS, F. & BOÉ, J. 2010. Comparison of three downscaling methods in simulating the impact of climate change on the hydrology of Mediterranean basins. *Journal of Hydrology*, 383, 111-124.

RAHMSTORF, S. 2010. A new view on sea level rise. *Nature Reports Climate Change*, 4, 44-45.

RAMANKUTTY, N., EVAN, A. T., MONFREDA, C. & FOLEY, J. A. 2008. Farming the planet: 1. Geographic distribution of global agricultural lands in the year 2000. *Global Biogeochemical Cycles*, 22, GB1003.

RAMANKUTTY, N., FOLEY, J. A., NORMAN, J. & MCSWEENEY, K. 2002. The global distribution of cultivable lands: current patterns and sensitivity to possible climate change. *Global Ecology and Biogeography*, 11, 377-392.

RICHARDS, J. A. & NICHOLLS, R. J. 2009. Impacts of climate change in coastal systems in Europe. PESETA-Coastal Systems study. Luxembourg.

RIDLEY, J. K., HUYBRECHTS, P., GREGORY, J. M. & LOWE, J. A. 2005. Elimination of the Greenland ice sheet in a high CO<sub>2</sub> climate. *Journal of Climate*, 18, 3409-3427.

ROCKSTROM, J., FALKENMARK, M., KARLBERG, L., HOFF, H., ROST, S. & GERTEN, D. 2009. Future water availability for global food production: The potential of green water for increasing resilience to global change. *Water Resources Research*, 45.

SCHMITTNER, A. 2005. Decline of the marine ecosystem caused by a reduction in the Atlantic overturning circulation. *Nature*, 434, 628-633.

SCHNEIDER, B., LATIF, M. & SCHMITTNER, A. 2007. Evaluation of different methods to assess model projections of the future evolution of the Atlantic meridional overturning circulation. *Journal of Climate*, 20, 2121-2132.

SCHWIERZ, C., KÖLLNER-HECK, P., ZENKLUSEN MUTTER, E., BRESCH, D., VIDALE, P.-L., WILD, M. & SCHÄR, C. 2010. Modelling European winter wind storm losses in current and future climate. *Climatic Change*, 101, 485-514.

- SILLMANN, J. & ROECKNER, E. 2008. Indices for extreme events in projections of anthropogenic climate change. *Climatic Change*, 86, 83-104.
- SMAKHTIN, V., REVENGA, C. & DOLL, P. 2004. A pilot global assessment of environmental water requirements and scarcity. *Water International*, 29, 307-317.
- SOUBEYROUX, J.-M., VIDAL, J.-P., NAJAC, J., KITOVA, N., BLANCHARD, M., DANDIN, P., MARTIN, E., PAGÉ, C. & HABETS, F. 2011. Impact du changement climatique en France sur la sécheresse et l'eau du sol *Projet ClimSec*. Toulouse: Meteo France.
- STOCKLE, C.O., DONATELLI, M., NELSON, R., 2003 CropSyst, a cropping systems simulation model. *European Journal of Agronomy* 18, 289-307.
- SWINGEDOUW, D., BRACONNOT, P., DELECLUSE, P., GUILYARDI, E. & MARTI, O. 2007. Quantifying the AMOC feedbacks during a 2xCO(2) stabilization experiment with land-ice melting. *Climate Dynamics*, 29, 521-534.
- TATSUMI, K., YAMASHIKI, Y., VALMIR DA SILVA, R., TAKARA, K., MATSUOKA, Y., TAKAHASHI, K., MARUYAMA, K. & KAWAHARA, N. 2011. Estimation of potential changes in cereals production under climate change scenarios. *Hydrological Processes*, Special Issue: Japan Society of Hydrology and water resources, 25 (17), 2715-2725.
- TSIMPLIS, M. N., MARCOS, M. & SOMOT, S. 2008. 21st century Mediterranean sea level rise: Steric and atmospheric pressure contributions from a regional model. *Global and Planetary Change*, 63, 105-111.
- VAN VUUREN, D., DEN ELZEN, M., LUCAS, P., EICKHOUT, B., STRENGERS, B., VAN RUIJVEN, B., WONINK, S. & VAN HOUDT, R. 2007. Stabilizing greenhouse gas concentrations at low levels: an assessment of reduction strategies and costs. *Climatic Change*, 81, 119-159.
- VAN VUUREN, D. P., ISAAC, M., KUNDZEWICZ, Z. W., ARNELL, N., BARKER, T., CRIQUI, P., BERKHOUT, F., HILDERINK, H., HINKEL, J., HOF, A., KITOUS, A., KRAM, T., MECHLER, R. & SCRIECIU, S. 2011. The use of scenarios as the basis for combined assessment of climate change mitigation and adaptation. *Global Environmental Change*, 21, 575-591.
- VELLINGA, M. & WOOD, R. A. 2008. Impacts of thermohaline circulation shutdown in the twenty-first century. *Climatic Change*, 91, 43-63.

VINCHON, C., AUBIE, S., BALOUIN, Y., CLOSSET, L., GARCIN, M., IDIER, D. & MALLET, C. 2009. Anticipate response of climate change on coastal risks at regional scale in Aquitaine and Languedoc Roussillon (France). *Ocean & Coastal Management*, 52, 47-56.

VINCHON, C., BARON-YELLES, N., BERTHELIER, E., HÉRIVAUX, C., LECACHEUX, S., MEUR-FEREC, C., PEDREROS, R., REY-VALETTE, H. & RULLEAU, B. 2011. MISEEVA : Set up of a transdisciplinary approach to assess vulnerability of the coastal zone to marine inundation at regional and local scale, within a global change context. *Littoral*, 11003.

VOROSMARTY, C. J., MCINTYRE, P. B., GESSNER, M. O., DUDGEON, D., PRUSEVICH, A., GREEN, P., GLIDDEN, S., BUNN, S. E., SULLIVAN, C. A., LIERMANN, C. R. & DAVIES, P. M. 2010. Global threats to human water security and river biodiversity. *Nature*, 467, 555-561.

WARREN, R., ARNELL, N., BERRY, P., BROWN, S., DICKS, L., GOSLING, S., HANKIN, R., HOPE, C., LOWE, J., MATSUMOTO, K., MASUI, T., NICHOLLS, R., O'HANLEY, J., OSBORN, T., SCRIECRU, S. (2010) The Economics and Climate Change Impacts of Various Greenhouse Gas Emissions Pathways: A comparison between baseline and policy emissions scenarios, AVOID Report, AV/WS1/D3/R01.  
[http://www.metoffice.gov.uk/avoid/files/resources-researchers/AVOID\\_WS1\\_D3\\_01\\_20100122.pdf](http://www.metoffice.gov.uk/avoid/files/resources-researchers/AVOID_WS1_D3_01_20100122.pdf).

WATKISS, P., HORROCKS, L., PYE, S., SEARL, A. & HUNT, A. 2009. Impacts of climate change in human health in Europe. PESETA-Human health study. Luxembourg.

WILLIS, J. K. 2010. Can in situ floats and satellite altimeters detect long-term changes in Atlantic Ocean overturning? *Geophysical Research Letters*, 37, -.

WOOD, E.F., ROUNDY, J.K., TROY, T.J., van BEEK, L.P.H., BIERKENS, M.F.P., BLYTH, E., de ROO, A., DOLL, P., EK, M., FAMIGLIETTI, J., GOCHIS, D., van de GIESEN, N., HOUSER, P., JAFFE, P.R., KOLLET, S., LEHNER, B., LETTENMAIER, D.P., PETERS-LIDARD, C., SIVAPALAN, M., SHEFFIELD, J., WADE, A. & WHITEHEAD, P. 2011. Hyperresolution global land surface modelling: Meeting a grand challenge for monitoring Earth's terrestrial water. *Water Resources Research*, 47, W05301.

WOS. 2011. *Web of Science* [Online]. Available:  
[http://thomsonreuters.com/products\\_services/science/science\\_products/a-z/web\\_of\\_science](http://thomsonreuters.com/products_services/science/science_products/a-z/web_of_science) [Accessed August 2011].

WU, W., TANG, H., YANG, P., YOU, L., ZHOU, Q., CHEN, Z. & SHIBASAKI, R. 2011. Scenario-based assessment of future food security. *Journal of Geographical Sciences*, 21, 3-17.

WUNSCH, C. & HEIMBACH, P. 2006. Estimated decadal changes in the North Atlantic meridional overturning circulation and heat flux 1993-2004. *Journal of Physical Oceanography*, 36, 2012-2024.

ZICKFELD, K., LEVERMANN, A., MORGAN, M. G., KUHLBRODT, T., RAHMSTORF, S. & KEITH, D. W. 2007. Expert judgements on the response of the Atlantic meridional overturning circulation to climate change. *Climatic Change*, 82, 235-265.

☐

## **Acknowledgements**

Funding for this work was provided by the UK Government Department of Energy and Climate Change, along with information on the policy relevance of the results.

The research was led by the UK Met Office in collaboration with experts from the University of Nottingham, Walker Institute at the University of Reading, Centre for Ecology and Hydrology, University of Leeds, Tyndall Centre – University of East Anglia, and Tyndall Centre – University of Southampton.

Some of the results described in this report are from work done in the AVOID programme by the UK Met Office, Walker Institute at the University of Reading, Tyndall Centre – University of East Anglia, and Tyndall Centre – University of Southampton.

The AVOID results are built on a wider body of research conducted by experts in climate and impact models at these institutions, and in supporting techniques such as statistical downscaling and pattern scaling.

The help provided by experts in each country is gratefully acknowledged – for the climate information they suggested and the reviews they provided, which enhanced the content and scientific integrity of the reports.

The work of the independent expert reviewers at the Centre for Ecology and Hydrology, University of Oxford, and Fiona's Red Kite Climate Consultancy is gratefully acknowledged.

Finally, thanks go to the designers, copy editors and project managers who worked on the reports.

Met Office  
FitzRoy Road, Exeter  
Devon, EX1 3PB  
United Kingdom

Tel: 0870 900 0100  
Fax: 0870 900 5050  
[enquiries@metoffice.gov.uk](mailto:enquiries@metoffice.gov.uk)  
[www.metoffice.gov.uk](http://www.metoffice.gov.uk)

Produced by the Met Office.  
© Crown copyright 2011 11/0209g  
Met Office and the Met Office logo  
are registered trademarks