



Cine MRI assessment of motility in the unprepared small bowel in the fasting and fed state: Beyond the breath-hold

Journal:	<i>Neurogastroenterology and Motility</i>
Manuscript ID	NMO-00087-2018.R2
Manuscript Type:	Original Article
Date Submitted by the Author:	06-Aug-2018
Complete List of Authors:	<p>Khalaf, Asseel; University of Nottingham School of Medicine, Nottingham Digestive Diseases Centre; Nottingham University Hospitals NHS Trust, NIHR Nottingham Biomedical Research Centre (BRC)</p> <p>Nowak, Adam; Nottingham University Hospitals NHS Trust, NIHR Nottingham Biomedical Research Centre (BRC)</p> <p>Menys, Alex; University College London Medical School, Centre for Medical Imaging</p> <p>Marciani, Luca; University of Nottingham School of Medicine, Nottingham Digestive Diseases Centre; Nottingham University Hospitals NHS Trust, NIHR Nottingham Biomedical Research Centre (BRC)</p> <p>Taylor, Stuart; University College London Medical School, Centre for Medical Imaging</p> <p>Spiller, Robin; University of Nottingham School of Medicine, Nottingham Digestive Diseases Centre; Nottingham University Hospitals NHS Trust, NIHR Nottingham Biomedical Research Centre (BRC)</p> <p>Gowland, Penny; University of Nottingham School of Physics and Astronomy, Sir Peter Mansfield Imaging Centre</p> <p>Moran, Gordon; University of Nottingham School of Medicine, Nottingham Digestive Diseases Centre; Nottingham University Hospitals NHS Trust, NIHR Nottingham Biomedical Research Centre (BRC)</p> <p>Hoad, Caroline; Nottingham University Hospitals NHS Trust, NIHR Nottingham Biomedical Research Centre (BRC); University of Nottingham School of Physics and Astronomy, Sir Peter Mansfield Imaging Centre</p>
Key Words:	Gastrointestinal motility, MRI, Cine MRI, Fasting, Fed state

Khalaf 1

**Cine MRI assessment of motility in the unprepared small bowel in the fasting
and fed state: Beyond the breath-hold**

**Running head: Optimised analysis technique for the assessment of small
bowel motility**

Asseel Khalaf^{1,2}, Adam Nowak², Alex Menys³, Luca Marciani^{1,2}, Stuart A. Taylor³,
Robin C. Spiller^{1,2}, Penny A. Gowland⁴, Gordon W. Moran^{1,2}, and Caroline L. Hoad^{2,4}

¹Nottingham Digestive Diseases Centre, School of Medicine, University of Nottingham UK

²NIHR Nottingham Biomedical Research Centre (BRC), Nottingham University Hospitals NHS Trust and the University of Nottingham, Nottingham, UK.

³ Centre for Medical Imaging, Division of Medicine, UCL, London, UK

⁴ Sir Peter Mansfield Imaging Centre, School of Physics and Astronomy, University of Nottingham, Nottingham UK

Corresponding author:

Dr Caroline Hoad

Senior Research Fellow

Sir Peter Mansfield Imaging Centre, School of Physics and Astronomy, University of Nottingham

NIHR Nottingham Biomedical Research Centre (BRC), Nottingham University Hospitals NHS Trust and the University of Nottingham, Nottingham, UK

Nottingham NG7 2UH, U.K.

E-mail: caroline.l.hoad@nottingham.ac.uk

Abstract

Background:

The symptoms of functional bowel disorders are common in postprandial but investigations are generally undertaken in the fasted state using invasive procedures. MRI provides a non-invasive tool to study the gastro-intestinal tract in an unperturbed, fed state. The aim of this study was to develop a technique to assess small bowel motility from cine MRI data in the unprepared bowel in fasting and fed states.

Methods:

Fifteen healthy volunteers underwent a baseline MRI scan after which they consumed a 400g soup. Subjects then underwent a postprandial scan followed by further scans at regular intervals. Small bowel motility was assessed using single-slice bTFE cine MRI.

An optimised processing technique was used to generate motility data based on power spectrum analysis of voxel-signal changes with time. Inter-observer variability (n=15) and intra-observer (n=6) variability were assessed. Changes in the motility index were compared between fasted and immediate postprandial state.

Results:

Excellent agreement between observers was seen across the range of motility measurements acquired, with intraclass correlation coefficient (ICC) of 0.979 (p<0.0001) and Bland-Altman limits of agreement 95% CI: -28.9 to 45.9a.u. Intra-observer variability was low with ICC of 0.992 and 0.960 (2 observers, P<0.0001).

Changes from the fasted to immediately postprandial state showed an average increase of 122.4±98.7% (n=15)

1
2
3
4 Conclusion:

5
6 This optimised technique showed excellent inter and intra observer agreement. It was
7
8 sensitive to changes in motility induced feeding. This technique will be useful to study
9
10 contractile activity and regional patterns along the gastrointestinal tract under
11
12 physiological conditions.
13
14
15
16

17 **KEY WORDS:** Gastrointestinal motility; MRI; cine MRI; fasting; fed state
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Khalaf 4

Key points:

- Many patients' symptoms are more prevalent after eating a meal. However most investigative MR paradigms are undertaken in the fasting patient using bowel preparation, which does not represent normal physiological motion.
- This study developed an optimised technique for the analysis of gut motility from cine MRI images, specifically addressing the differences in appearance and motility of the small bowel wall in unprepared cine MR images in both fasting and fed states.
- With the clinical availability of MRI scanners, this technique has the potential to improve our knowledge of the pathophysiology of gastrointestinal tract under normal physiological conditions.

For Peer Review

Introduction

Conventional manometry of small bowel motility has provided valuable insights into motor function pathophysiology of the gastrointestinal tract^{1,2}. The technique has limitations, with naso-duodenal or oro-duodenal intubation being a difficult, uncomfortable and invasive procedure for patients. Moreover, manometry techniques are not generally used in the lower sections of the small bowel due to difficulties with access and the invasiveness of the technique^{3,4}. Furthermore, misinterpretation of the manometry recordings can occur if non-occlusive contractions occur and large spacing between ports mean that motor patterns can be mis-defined⁵. The tube may also interfere with normal feeding making it particularly difficult to study physiological changes from the fasting state and the effect of nutrient intake.

Over the last 10 years, MRI has proven to be a useful tool to probe the unprepared physiology of the gastro-intestinal tract⁶⁻⁹. It is particularly suitable for longitudinal or repeated studies, and its versatility allows for multiple physiological parameters to be monitored in a single scanning session. Magnetic resonance enterography (MRE) is used to evaluate the small bowel after the ingestion of an oral contrast agent. It involves distending the bowel artificially to produce detailed images of the bowel wall¹⁰ and induces bowel wall motion to move the large amount of oral contrast agent through the GI tract, which can then be studied using cine MRI.

Motility measurements following oral contrast preparation using a cine MRI acquisition have made significant advances in recent years^{8,11-17}, but to date quantification of wall motion has either involved looking for contractions across the lumen^{13,14,16} or using registration methods^{15,17}, which work well in the deliberately distended bowel where the walls are clearly visualised. However, bowel distension with a hypo-osmotic solution may not be truly physiological and so cannot study true

Khalaf 6

1
2
3
4 fasting motility patterns and may not represent the full range of motility patterns in the
5
6 postprandial state. The ability to study motility in the postprandial state, or the
7
8 transition between the two states, has many potential advantages in furthering our
9
10 understanding of physiology and the origin of symptoms which many patients
11
12 experience after feeding. Moreover, it is particularly important for the pharmaceutical
13
14 sciences because the rate and extent of drug dissolution and absorption from solid
15
16 oral dosage forms is highly dependent upon gastrointestinal motility^{18,19}. Furthermore,
17
18 the use of bowel distension limits the use of MRE in paediatric and elderly
19
20 populations.
21
22

23
24 The unprepared small bowel can be imaged using the same high spatial and
25
26 temporal resolution cine acquisitions as for the prepared bowel^{11,12,14}. However, the
27
28 post processing techniques used to parameterise the motility may need to be refined
29
30 for images, which do not delineate the small bowel wall clearly and show different
31
32 patterns of motility.
33

34
35 The aim of this study was to develop an analysis technique to assess motility from
36
37 cine MRI data acquired in the fasted and fed, unprepared small bowel. Inter- and
38
39 intra-observer variability, and the sensitivity to changes in motility caused by feeding
40
41 were investigated.
42
43
44

45 **Materials and methods**

46
47 This study was approved by the local Ethics Committee of the University of
48
49 Nottingham (H19062014). This study is registered on www.clinicaltrials.gov with
50
51 identifier NCT02717117. All subjects gave informed written consent. The study
52
53 design, subjects, and data sets used have been reported previously²⁰.
54
55
56
57
58
59
60

Unprepared bowel data acquisition

Fifteen healthy volunteers (age 29 ± 10 years, BMI 24 ± 5 kg/m²) were recruited from the local campus population. Subjects with any disease or taking medication (e.g. loperamide, codeine, metoclopramide, hyoscine butylbromide, mebeverine, ondansetron) that affects gastric emptying or small bowel transit were excluded. Standard MRI exclusion criteria were applied.

This study was open label. Subjects were scanned using a 1.5T Philips Achieva MRI scanner (Philips Healthcare, Best, the Netherlands) using the 16 element torso (XL-TORSO, Philips Healthcare) coil at the Sir Peter Mansfield Imaging Centre, University of Nottingham. They underwent a baseline fasting scan defined as $t = -20$ min time point. They were asked to consume a soup meal (cream of chicken soup (400g) (or mushroom for vegetarians) (Heinz, Wigan, UK))²¹ within 20 min then the subjects underwent a first immediate postprandial scan (defined as $t = 0$ min). MRI data collection was subsequently repeated every 15 minutes for the first 60 minutes where the subjects remained in the scanner and then every 30 minutes up to 270 minutes where subjects were allowed to leave the scanner between scans if they chose to²⁰. The subjects were scanned using a range of sequences. At each time point scans were acquired to assess small bowel motility⁸ using a single slice bTFE cine MRI acquisition (with reconstructed in-plane resolution 1.49×1.7 mm², slice thickness 10 mm, echo time (TE)=1.52 ms, repetition time (TR)=3.0 ms, flip angle 80°, SENSE 2.0), of 1 minute free-breathing, temporal resolution of 1s, this was repeated at six contiguous parallel coronal planes through the small bowel as previously described²⁰. The total scan time for motility was 6 min. The subjects were instructed to take shallow gentle breaths for the duration of the motility acquisition.

Data analysis

Motility assessment

Free breathing data were processed using GIQuant™ (Motilent, Ford, UK). The algorithm corrects respiratory motion²² before applying the non-linear optic flow registration as described previously¹⁵ to correct local deformation caused by bowel wall motion and model intensity changes caused by luminal flow. The data output from the image registration were further analysed using a customised graphical user interface written in MATLAB® (MathWorks, Natick, MA, USA).

On MRI the unprepared small bowel has a very different appearance to the prepared bowel required for MRE (Figure 1). In the prepared bowel there is clear definition of the bowel walls and obvious peristaltic motion is visible through the time series across most of the small bowel. In the unprepared bowel, the bowel wall is not always visible and bolus movement of the chyme between segments is common post-prandially. Therefore a different approach for quantifying the motility of the unprepared small bowel was developed, based on the registration parameter C ¹⁵ which represents the change in signal intensity between time points, within a defined region of interest (ROI) placed over the small bowel loops. This parameter is modelled simultaneously with the deformation during the registration process and is intended to capture any signal intensity changes not occurring from in-plane motion (i.e. through plane motion and flow)¹⁵.

To allow sensitivity to both oscillatory events such as mixing of contents during peristalsis and forward propulsion of boluses of chyme, the power spectrum analysis of the image registration parameter C was developed similar to that proposed by van der Paardt et al²³ and Sprengers et al²⁴. Initially, to remove zero frequency data, the

1
2
3
4 mean of C through time for each pixel was calculated and subtracted from each pixel
5 value (Figure 2A). Then for each pixel in the data, the power spectrum (Fourier
6 transform of time course of mean-subtracted C, smoothed to reduce noise and with
7 the first time point removed to eliminate data not in the steady state, and then
8 multiplied by its complex conjugate) was calculated (Figure 2B), generating data with
9 frequency information up to 0.48 Hz. The area under the power spectrum was
10 calculated as a summary metric ($AUC_{\text{power spectrum}}$), and maps created to visualise the
11 regions of higher motility. This metric was intended to reflect both segmental
12 oscillations and bolus movement of contents, typically seen post-prandially²⁵. When
13 regions of interest were defined, average data for the ROI was calculated from the
14 pixel by pixel measurements within the ROI of the $AUC_{\text{power spectrum}}$ maps.

31 *Observer variability*

32
33 To determine the variability in the results due to observer definition of the region of
34 the small bowel loops, the following analyses were carried out.

- 35
36
37
38 1. Inter-observer variability: Two observers, 1 experienced (CH over 10 years) in
39 viewing small bowel MRI data and 1 inexperienced (AK less than 2 years)
40 drew regions around all the visible small bowel segments across the 6 coronal
41 slices acquired. This was repeated for all 15 subjects scanned at all time
42 points pre and post-test meal. The number of regions depended on the spatial
43 separation of the different bowel loops. If there was large amount of visceral
44 fat separating the loops, more regions were drawn to encompass all the small
45 bowel loops.
46
47
48
49
50
51
52
53
54
55 2. Intra-observer variability: The same two observers drew regions around the
56
57
58
59
60

Khalaf 10

1
2
3
4 visible small bowel segments from only six subjects (chosen to have different
5
6 body composition: 3 normal BMI (22.3, 22.9 22.9 kg/m²) and 3 high BMI (30.6,
7
8 27.1, 29.1 kg/m²). The changing body composition resulted in images with
9
10 very different contrast of the edges of the small bowel loops and therefore
11
12 represented the maximum range of tissue contrast that would be seen across
13
14 all subjects. Regions were defined by drawing the ROIs to encompass only the
15
16 visible small bowel loops ignoring intra-abdominal visceral fat and other
17
18 tissues; the regions were defined twice on the images with at least one month
19
20 interval between observations.
21
22
23
24
25
26

Changing motility in response to feeding using two motility metrics

27
28
29 We examined the strength of correlation between two motility analysis techniques
30
31 using the total power (AUC_{power spectrum}) and the standard deviation of the Jacobian
32
33 (SD_{JAC}, a previous published metric for motility⁸). SD_{JAC} looks at the geometric
34
35 changes from image registration and is currently used for small bowel motility in the
36
37 prepared bowel. In addition, we showed how both metrics change with feeding by
38
39 looking at the mean change in the metric between the fasted (t= -20min) and the
40
41 immediately postprandial data (t= 0 min).
42
43
44
45
46

Statistical analyses

47
48 All statistical analysis was carried out using Graph Pad Prism 7.0 (La Jolla, USA). All
49
50 data were tested for normality using the D'Agostino and Pearson normality test.
51
52 Inter-observer variability was investigated by using a Bland-Altman plot to determine
53
54 the 95% confidence limits of agreement. Correlation between observers was
55
56
57
58
59
60

measured using the Intra-class correlation coefficient (ICC) using a 2-way random effects model, with a single rater and absolute agreement²⁶. Intra-observer variability was also investigated with Intra-class correlation coefficients using a 2-way mixed effects model and single rater and absolute agreement. The 95% confidence limits of agreement were also calculated. Pearson correlation coefficient was used to measure the strength of correlation between the measurement of motility using $AUC_{\text{power spectrum}}$ and SD_{JAC} for the fasting and immediately postprandial data sets.

Results

Visualisation of high motility regions

Example maps of $AUC_{\text{power spectrum}}$ of fasting and postprandial data are shown in Figure 3 for one subject. Example maps of SD_{JAC} are also shown for comparison. The $AUC_{\text{power spectrum}}$ maps show a lower values in regions of known low motility (e.g. liver) compared to the SD_{JAC} maps. There also appears to be a larger change ($AUC_{\text{power spectrum}}$ 122.4±98.7%, SD_{JAC} 31.8±20.7%) (calculated by dividing the fed-fasted data by the fasted data across all the 15 subjects) across the small bowel between the fasted and immediately postprandial states which suggests the $AUC_{\text{power spectrum}}$ maps may have better range to define the differences between sporadic movements occurring during fasting and large scale movements following ingestion of the meal.

Inter-observer variability of small bowel motility

The variation in both $AUC_{\text{power spectrum}}$ and SD_{JAC} across different time points, averaged over all healthy volunteers at each time point, and covering all regions of

Khalaf 12

1
2
3
4 small bowel from the 6 slices, measured by each observer is shown in figure 4A
5 (error bars shown are SEM). This graph shows low measured motility at baseline in
6 the fasting state followed by a significant increase post-prandially which then persists
7 for the majority of the imaging period. The degree of correlation between two
8 observers was assessed using the ICC to be 0.979 and $p < 0.0001$, $n = 195$ (Figure
9 4B). Inter-observer variability was assessed using the Bland Altman plot ²⁷ (Figure
10 4C) which showed a mean difference of 8.5 a.u. between small bowel motility
11 measurements, with a 95% confidence interval of -28.9 to 45.9 a.u. as indicated by
12 the upper and lower dotted lines (figure 4C).
13
14
15
16
17
18
19
20
21
22
23
24
25
26

Intra-observer variability of small bowel motility

27
28
29 The correlation between the two analyses for $AUC_{\text{power spectrum}}$ performed by each
30 observer was also assessed using the ICC and Bland-Altman limits of agreement
31 (Table 1), showing good agreement between the analyses (ICC > 0.9 for all data).
32 Figure 5 plots $AUC_{\text{power spectrum}}$ measurements against time for the six subjects with
33 normal and high BMI, showing good agreement between analyses at most time
34 points for both body compositions.
35
36
37
38
39
40
41
42
43
44

Changing motility in response to feeding

45
46
47 The subjects showed an increase between their fasting and initial postprandial
48 motility measurement for both analysis methods ($AUC_{\text{power spectrum}} 122.4 \pm 98.7\%$,
49 $SD_{JAC} 31.8 \pm 20.7\%$), although there was considerable spread within the data across
50 the subjects (Figure 6). The correlation between the techniques was significant
51 ($r = 0.9$, $p < 0.0001$, $n = 30$).
52
53
54
55
56
57
58
59
60

Discussion

This study has described and evaluated an optimised technique for the analysis of gut motility from MRI images, specifically addressing the differences in appearance and motility of the small bowel wall in unprepared bowel MR images, compared to images obtained following bowel preparation with oral luminal contrast agent. The images in fasting and fed conditions showed poor small bowel wall definition in places. Coupled with the different motility patterns of bolus propulsion as well as peristalsis meant previously published analysis techniques, which are based on geometric changes over the measurement period to generate the motility metric, were not as appropriate to use in fasting and fed studies. Maps of $AUC_{\text{power spectrum}}$ indicated a better discrimination of the bowel tissue compared to the more widely published SD_{JAC} ^{8,15,28} with lower noise in the regions where non/low-motility is expected, and a bigger range of motility indices. The proposed method utilises changes in signal intensities that occur when the small bowel contents move between segments in regions showing bolus movement of contents (duodenum and jejunum) as well as those exhibiting more oscillatory motion (ileum), rather than looking for continuous motion throughout the time series.

The appearance of the unprepared bowel, which may contain multiple collapsed loops which are not always easily identified, means the definition of the ROIs is more subjective than for the prepared bowel with distended lumen. However our inter-observer variability data suggested that ROI definition had a small effect on the results compared to the changes in motility seen following eating the meal. Excellent correlation between observers was seen across the whole range of $AUC_{\text{power spectrum}}$ measurement acquired and the Bland-Altman limits of agreement were low compared

Khalaf 14

1
2
3
4 to the range of values measured. The postprandial changes over time show a rapid
5
6 increase in motility following ingestion of the soup meal with levels returning towards
7
8 baseline much later after the meal had been consumed ²⁰.
9

10
11 Intra-observer variability was also low with a small range of Bland-Altman limits of
12
13 agreements for both observers across two very different body shapes. The ICC was
14
15 excellent with all data greater than 0.9.
16

17
18 This study presents an optimised analysis of MRI data to assess the motility of the
19
20 unprepared fasting and fed small bowel. The MRI method proposed removes the
21
22 need for intubating subjects, which can be a stressful procedure, often requiring
23
24 fluoroscopy to place the catheter and rarely covering the entire length of the small
25
26 bowel. Orocecal transit times can be measured using breath tests ²⁹ but these
27
28 include changes due to gastric emptying and do not give information about the specific
29
30 motor function of the small bowel but are an indirect measure of small bowel motility.
31
32 Scintigraphy transit studies ³⁰ involving ionising radiation do not provide information
33
34 about the motor patterns seen in the small bowel. The proposed techniques
35
36 (including further frequency analysis of the power spectra) will be useful to study the
37
38 time scales of contractile activity and regional patterns along the gastrointestinal tract
39
40 in health and disease. Information on small bowel motility can be obtained in
41
42 conjunction with mapping the bowel liquid pockets using MRI ³¹ furthering
43
44 understanding of the effects of motility on the fluid environment of the bowel. These
45
46 combined insights could help with advancing in vitro/in vivo predictive dissolution
47
48 studies of oral dosage forms under similar, undisturbed conditions.
49
50

51
52 The data from the normal and high BMI subjects would indicate that there is a
53
54 potential for over estimating motility in the higher BMI subjects. These subjects all
55
56 presented with high motility indices postprandially, however only one of the 3 showed
57
58
59
60

1
2
3
4 high baseline data. These larger motility indices may have been measured as
5 regions of high signal intensity in the fat as it moves into and out of the imaging plane
6 during respiratory motion. As this may not be fully corrected by the registration
7 algorithm these movements will be interpreted as bowel motility. Further studies of
8 higher BMI subjects is needed to understand the factors contributing to the larger
9 motility metric measured and whether poor registration is a factor.
10
11
12
13
14
15
16

17 Other factors, which could also influence the signal intensity, are field
18 inhomogeneities and metallic artefacts. To some extent, overall changes in image
19 intensity across the image due to these factors are removed from the $AUC_{\text{power spectrum}}$
20 analysis by using the registration parameter C which models the signal changes and
21 not the absolute values. An empty bowel has a different intensity to a filled bowel,
22 however movement of the contents either between loops or from one section to
23 another show similar changes in intensity levels. Other meal contents should have
24 similar motility patterns to the meal in this study, but would need investigating to
25 determine whether the sensitivity is the same as the soup meal, particularly for a
26 more solid meal, which may have a much lower signal intensity in the small bowel.
27
28
29
30
31
32
33
34
35
36
37

38 There were limitations to our study. Due to the time consuming nature of drawing all
39 the individual ROIs on the motility data the intra-observer repeated measurements
40 were confined to just 6 subjects, not the full 15 available (used for the inter-observer
41 data). Drawing of ROIs for a single time point took around 5-10 minutes depending
42 on the anatomy including the loading of each data set into the software. However, the
43 intra-observer data were chosen from subjects who had very different small bowel
44 anatomical appearances due to their differing BMIs, providing the observers with
45 contrasting data for drawing the regions. Smoothing of the data before calculating the
46 power spectrum reduces the effects of isolated poor mis-registration of the data.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Khalaf 16

1
2
3
4 However it will not eliminate the effects completely and these datasets will slightly
5
6 overestimate the small bowel motility present.
7

8
9 In conclusion, this study describes an optimised analysis technique to evaluate small
10
11 bowel motility in the physiological fasting and postprandial states using registered
12
13 cine MRI datasets. This method showed excellent agreement between
14
15 measurements of intra and inter observers as well as showing the sensitivity of the
16
17 technique to changes in motility induced by ingestion of a meal. Cine MRI scanning is
18
19 available on most clinical scanners worldwide therefore future studies have real
20
21 potential to translate and improve our knowledge of the small bowel environment in
22
23 health and disease.
24
25
26
27

28 **Acknowledgements**

29
30 This study was supported by the National Institute for Health Research Nottingham
31
32 Biomedical Research Centre. The views expressed are those of the authors and not
33
34 necessarily those of the National Health Service (NHS, U.K.), the National Institute
35
36 for Health Research (NIHR, U.K.) or the Department of Health. This study was
37
38 funded by a scholarship from the Kuwait University and from a research grant from
39
40 the Nottingham University Hospitals' Charity (PP-Gordon Moran-Nov14). The views
41
42 expressed are those of the author(s) and not necessarily those of the NHS, the NIHR
43
44 or the Department of Health.
45
46

47
48 LM is supported in part by Award # HHSF223201510157C: "In Vivo Predictive
49
50 Dissolution (IPD) to Advance Oral Product Bioequivalence (BE) Regulation" by the
51
52 U.S. Food and Drug Administration (FDA); this manuscript represents the position of
53
54 the authors and not necessarily that of the FDA.
55

56 AM is the CEO of Motilent Limited, a medical imaging analysis company.
57
58
59
60

Competing Interests: the remaining authors have no competing interests.

AK, CLH, LM and GWM designed the research. AK, AN recruited the patients. AK collected the data. AK, AM, and CLH, analysed the data. AK, CLH, LM and GWM wrote the manuscript draft. All authors revised the final manuscript.

References

1. Hansen MB. Small intestinal manometry. *Physiol Res*. 2002;51(6):541-556.
2. Malagelada JR, Stanghellini V. Manometric Evaluation of Functional Upper Gut Symptoms. *Gastroenterology*. 1985;88(5):1223-1231.
3. Malagelada C, Malagelada JR. Small Bowel Motility. *Curr Gastroenterol Rep*. Jun 2017;19(6):26.
4. Seidl H, Gundling F, Pfeiffer A, Pehl C, Schepp W, Schmidt T. Comparison of small-bowel motility of the human jejunum and ileum. *Neurogastroenterol Motil*. Aug 2012;24(8):e373-380.
5. Arkwright JW, Dickson A, Maunder SA, et al. The effect of luminal content and rate of occlusion on the interpretation of colonic manometry. *Neurogastroenterol Motil*. Jan 2013;25(1):E52-E59.
6. Schwizer W, Steingoetter A, Fox M. Magnetic resonance imaging for the assessment of gastrointestinal function. *Scand J Gastroenterol*. Nov 2006;41(11):1245-1260.
7. Fruehauf H, Menne D, Kwiatek MA, et al. Inter-observer reproducibility and analysis of gastric volume measurements and gastric emptying assessed with magnetic resonance imaging. *Neurogastroenterol Motil*. Sep 2011;23(9):854-861.
8. Menys A, Taylor SA, Emmanuel A, et al. Global Small Bowel Motility: Assessment with Dynamic MR Imaging. *Radiology*. Nov 2013;269(2):442-449.
9. Marciani L, Garsed KC, Hoad CL, et al. Stimulation of colonic motility by oral PEG electrolyte bowel preparation assessed by MRI: comparison of split vs single dose. *Neurogastroenterol Motil*. Oct 2014;26(10):1426-1436.
10. Griffin N, Grant LA, Anderson S, Irving P, Sanderson J. Small bowel MR enterography: problem solving in Crohn's disease. *Insights Imaging*. Jun 2012;3(3):251-263.
11. Wakamiya M, Furukawa A, Kanasaki S, Murata K. Assessment of Small Bowel Motility Function With Cine-MRI Using Balanced Steady-State Free Precession Sequence. *J Magn Reson Imaging*. May 2011;33(5):1235-1240.
12. Ohkubo H, Kessoku T, Fuyuki A, et al. Assessment of Small Bowel Motility in Patients With Chronic Intestinal Pseudo-Obstruction Using Cine-MRI. *Am J Gastroenterol*. Jul 2013;108(7):1130-1139.

13. Menys A, Atkinson D, Odille F, et al. Quantified terminal ileal motility during MR enterography as a potential biomarker of Crohn's disease activity: a preliminary study. *Eur Radiol*. Nov 2012;22(11):2494-2501.
14. Froehlich JM, Waldherr C, Stoupis C, Erturk SM, Patak MA. MR motility imaging in Crohn's disease improves lesion detection compared with standard MR imaging. *Eur Radiol*. Aug 2010;20(8):1945-1951.
15. Odille F, Menys A, Ahmed A, Punwani S, Taylor SA, Atkinson D. Quantitative assessment of small bowel motility by nonrigid registration of dynamic MR images. *Magn Reson Med*. Sep 2012;68(3):783-793.
16. Bickelhaupt S, Froehlich JM, Cattin R, et al. Differentiation between active and chronic Crohn's disease using MRI small-bowel motility examinations - Initial experience. *Clin Radiol*. Dec 2013;68(12):1247-1253.
17. Hahnemann ML, Nensa F, Kinner S, Gerken G, Lauenstein TC. Motility mapping as evaluation tool for bowel motility: initial results on the development of an automated color-coding algorithm in cine MRI. *J Magn Reson Imaging*. Feb 2015;41(2):354-360.
18. Oberle RL, Amidon GL. The influence of variable gastric emptying and intestinal transit rates on the plasma level curve of cimetidine: an explanation for the double peak phenomenon. *J Pharmacokinet Biopharm*. Oct 1987;15(5):529-544.
19. Oberle RL, Chen TS, Lloyd C, et al. The influence of the interdigestive migrating myoelectric complex on the gastric emptying of liquids. *Gastroenterology*. Nov 1990;99(5):1275-1282.
20. Khalaf A, Hoad CL, Menys A, et al. MRI assessment of the postprandial gastrointestinal motility and peptide response in healthy humans. *Neurogastroenterol Motil*. Aug 31 2017.
21. Moran GW, Leslie FC, McLaughlin JT. Crohn's disease affecting the small bowel is associated with reduced appetite and elevated levels of circulating gut peptides. *Clin Nutr*. Jun 2013;32(3):404-411.
22. Hamy V, Dikaïos N, Punwani S, et al. Respiratory motion correction in dynamic MRI using robust data decomposition registration - Application to DCE-MRI. *Med Image Anal*. Feb 2014;18(2):301-313.
23. van der Paardt MP, Sprengers AMJ, Zijta FM, Lamerichs R, Nederveen AJ, Stoker J. Noninvasive Automated Motion Assessment of Intestinal Motility by Continuously Tagged MR Imaging. *J Magn Reson Imaging*. Jan 2014;39(1):9-16.
24. Sprengers AMJ, van der Paardt MP, Zijta FM, et al. Use of continuously MR tagged imaging for automated motion assessment in the abdomen: A feasibility study. *J Magn Reson Imaging*. Aug 2012;36(2):492-497.
25. Silverthorn DU. *Human Physiology: an integrated Approach*. London: Pearson; 2012.
26. Koo TK, Li MY. A Guideline of Selecting and Reporting Intraclass Correlation Coefficients for Reliability Research. *J Chiropr Med*. Jun 2016;15(2):155-163.
27. Bland JM, Altman DG. Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet*. Feb 08 1986;1(8476):307-310.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
28. Plumb AA, Menys A, Russo E, et al. Magnetic resonance imaging-quantified small bowel motility is a sensitive marker of response to medical therapy in Crohn's disease. *Aliment Pharmacol Ther.* Aug 2015;42(3):343-355.
 29. Geypens B, Bennink R, Peeters M, et al. Validation of the lactose-[C-13]ureide breath test for determination of orocecal transit time by scintigraphy. *J Nucl Med.* Sep 1999;40(9):1451-1455.
 30. Maurer AH. Gastrointestinal Motility, Part 2: Small-Bowel and Colon Transit. *J Nucl Med Technol.* Mar 2016;44(1):12-18.
 31. Mudie DM, Murray K, Hoad CL, et al. Quantification of gastrointestinal liquid volumes and distribution following a 240 mL dose of water in the fasted state. *Mol Pharm.* Sep 2014;11(9):3039-3047.
- For Peer Review

Tables

Table 1

	ICC	Bland-Altman
Observer 1	0.992 P<0.0001 n=78	MD 6.1 a.u. 95% CI -7.2 to 19.4 a.u. Measurement Range (27.2 – 254.5 a.u.)
Observer 2	0.960 P<0.0001 n=78	MD 1.0 a.u. 95% CI -41.2 to 43.19 a.u. Measurement Range (28.4 – 281.0 a.u.)

Table legends

Table 1: Table summarising the AUC_{power spectrum} ICC and Bland-Altman from observer 1 and observer 2. MD – Mean Difference; CI – Confidence Interval

Figure legends

Figure 1. Figure illustrating the difference between (A) the prepared bowel with clear definition of the bowel wall and bright luminal contents (data not from the current study, bowel preparation of 2% mannitol with 0.2% locust bean gum) and (B) the unprepared bowel with less visible bowel wall and brighter contents only when chyme moves into the segment.

Figure 2 A. Graph showing the variation in intensity of the C parameter with time for 3 abdominal regions. Solid black line is a small ROI from the upper small bowel

(jejunum), dashed black line is a small ROI from the lower small bowel (ileum), solid grey line is a small ROI from the ascending colon representing a known low motility region of the GI tract. B. Corresponding power spectrum of the data in A.

Figure 3. Example of motility maps generated by the software for a single volunteer across the 6 slices acquired, visualising the areas of high motility. A and B illustrate the fasting and the fed state motility maps. C represents the different motility maps generated by the $AUC_{\text{power spectrum}}$ and SD_{JAC} motility parameters. S: slice number. Regions of small bowel have been highlighted on the images.

Figure 4. A. Graph showing the mean small bowel motility assessed with $AUC_{\text{power spectrum}}$ and SD_{JAC} across time points, measured by two observers, error bars are SEM. B. Graph showing correlation of Inter-observer data for $AUC_{\text{power spectrum}}$ data. C. Bland–Altman plot showing the 95% limits of agreement in $AUC_{\text{power spectrum}}$ results. (Mean difference: thick solid line, mean ± 2 standard deviations: dotted lines).

Figure 5. Graphs showing the repeated analysis of $AUC_{\text{power spectrum}}$ for the normal (A,B,C) and high (D,E,F) BMI subjects measured by the two observers.

Figure 6: Graph illustrating the difference in small bowel motility during fasting and immediately after feeding using the two different motility parameters across all the 15 subjects. A. The fasting and fed small bowel motility measured by $AUC_{\text{power spectrum}}$. B. The fasting and fed small bowel motility measured by SD_{JAC} . C. Correlation between the two parameters of the fasting and fed data. Individual subjects have the same colour coding across A and B.

List of abbreviations:

AUC _{power spectrum}	Area under the power spectrum
ICC	Intra-class correlation coefficient
MRE	Magnetic resonance enterography
ROI	Region of interest
SD _{JAC}	Standard deviation of the Jacobian

For Peer Review




Figure 1. Figure illustrating the difference between (A) the prepared bowel with clear definition of the bowel wall and bright luminal contents (data not from the current study, bowel preparation of 2% mannitol with 0.2% locust bean gum) and (B) the unprepared bowel with less visible bowel wall and brighter contents only when chyme moves into the segment.

Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2 A. Graph showing the variation in intensity of the C parameter with time for 3 abdominal regions. Solid black line is a small ROI from the upper small bowel (jejunum), dashed black line is a small ROI from the lower small bowel (ileum), solid grey line is a small ROI from the ascending colon representing a known low motility region of the GI tract. B. Corresponding power spectrum of the data in A.




Figure 3. Example of motility maps generated by the software for a single volunteer across the 6 slices acquired, visualising the areas of high motility. A and B illustrate the fasting and the fed state motility maps. C represents the different motility maps generated by the $AUC_{\text{power spectrum}}$ and SD_{JAC} motility parameters. S: slice number. Regions of small bowel have been highlighted on the images.




Figure 4. A. Graph showing the mean small bowel motility assessed with AUCpower spectrum and SDJAC across time points, measured by two observers, error bars are SEM. B. Graph showing correlation of Inter-observer data for AUCpower spectrum data. C. Bland-Altman plot showing the 95% limits of agreement in AUCpower spectrum results. (Mean difference: thick solid line, mean ± 2 standard deviations: dotted lines).




Figure 5. Graphs showing the repeated analysis of AUCpower spectrum for the normal (A,B,C) and high (D,E,F) BMI subjects measured by the two observers.

Review




Figure 6: Graph illustrating the difference in small bowel motility during fasting and immediately after feeding using the two different motility parameters across all the 15 subjects. A. The fasting and fed small bowel motility measured by AUC_{power spectrum}. B. The fasting and fed small bowel motility measured by SD_{JAC}. C. Correlation between the two parameters of the fasting and fed data. Individual subjects have the same colour coding across A and B.