

1 **Conservation on the blink: deficient technical reports threaten conservation in the**
2 **Natura 2000 network**

3
4 Fulgencio Lisón^{1*}, Adison Altamirano¹, Richard Field² and Gareth Jones³

5
6 ¹Laboratorio de Ecología del Paisaje Forestal, Departamento de Ciencias Forestales, Universidad de La
7 Frontera, P.D. Box 45-D, Temuco, Chile.

8 ²School of Geography, University of Nottingham, Nottingham NG7 2RD, United Kingdom.

9 ³School of Biological Sciences, University of Bristol, 24 Tyndall Avenue, BS8 1TQ Bristol, United
10 Kingdom.

11
12 *Corresponding author: e-mail: fulgencio.lison@ufrontera.cl; Phone: +56 45 2591043; Fax: +56 45
13 2325634

14
15 **Running title:** Technical deficiencies in Natura 2000 network

16
17 **Keywords:** Chiroptera, Conservation, Habitats Directive 92/43/EEC, Management policies, Spain,
18 Special Areas of Conservation.

19
20 Type of article: Research papers

21 Words in abstract: 254

22 Words, including abstract, main text, acknowledgments, references, figures and tables (with each
23 display piece counted as 300 words): 6519

24
25 Number of references: 59

26 Number of figures: 2

27 Number of tables: 1

28 Supplementary material: SM1 – data sources; SM2 – official and known presences of each bat species
29 across each Autonomous Community

30 **Abstract**

31 Globally, laws define both where protected areas are and their level of protection. Usually, legal
32 protection is not absolute and alternative land-uses can be implemented if perceived gains outweigh
33 conservation losses. Technical reports, describing the importance of each protected area, are therefore
34 crucial for decision-making, impact assessments, mitigation policies and management plans, and thus
35 effective conservation. While much research has focused on protected areas themselves, including the
36 biodiversity they contain and the impact of illegal activities, almost no research has evaluated the
37 adequacy of the technical reports. Given high levels of data availability, the European Natura 2000
38 network (N2000) might be expected to represent best practice. Here we compare known bat presences
39 with records from Standard Data Forms (SDFs) of Spanish N2000 Special Areas of Conservation
40 (SACs); the Habitats Directive protects all European bat species. Across 1206 SACs, we found far fewer
41 bat species listed in the SDFs than are known to occur in the SACs they represent, for both Annex II
42 and particularly Annex IV bat species. These findings have serious conservation implications, including
43 that decisions are systematically biased against conservation outcomes: if SDFs greatly underestimate
44 the conservation value of their SACs, development of the land (or sea) is much more likely to be
45 permitted. Incorporating known species presences into the SDFs of SACs is low in cost and
46 straightforward, and can potentially achieve tremendous conservation benefits for minimal outlay; it
47 should therefore be a top conservation priority globally, and conservation scientists should urgently
48 engage with government agencies, accordingly.

49

50 **Keywords:** Chiroptera, Habitats Directive 92/43/EEC, Management policies, Special Areas of
51 Conservation, Spain.

52

53 **1. Introduction**

54 Despite protected areas now covering 15.4% of the world's land surface (UNEP-WCMC, 2016), global
55 biodiversity continues to decline more than 20 years after the United Nations Convention of Biological
56 Diversity (Butchart et al., 2010). The extent to which reserves fulfil their role of protecting biodiversity
57 depends on how well they meet two objectives (Margules and Pressey, 2000). The first is
58 representativeness, a long-established goal referring to the need for reserves to represent, or sample, the
59 full variety of biodiversity. This is addressed mainly in the original designation of areas as protected,
60 and has been much researched (for example, using gap analyses and metrics of irreplaceability; Kukkala
61 and Moilanen, 2013; Le Saout et al., 2013; Kukkala et al., 2016). The second objective is persistence:
62 reserves, once established, should promote the long-term survival of the species and other elements of
63 biodiversity they contain by maintaining natural processes and viable populations (Margules and
64 Pressey, 2000) – increasingly important because of ongoing climate change that could undermine
65 conservation efforts (Araújo et al., 2011; D'Amen et al., 2011; de Koning et al., 2014). An integral part
66 of biodiversity conservation is ensuring the ongoing protection of reserves in the face of competing
67 demands for the use of that land (or sea), such as development for housing, agriculture or extraction
68 industries. In most cases, the legislation protecting any given reserve allows alternative uses of the area
69 if the benefits are considered to outweigh the costs to a sufficient degree (Lee et al., 2007). Evaluating
70 the conservation costs usually involves considering the likely impacts on the species known to be present
71 in the reserve – as listed in official documentation associated with each reserve. However, this official
72 documentation may not include all species known to occur within the reserve, and no research that we
73 are aware of has examined this. Here we evaluate the extent to which official documentation reflects the
74 bat species known to be present in Natura 2000 protected areas throughout Spain.

75 In Europe, the Natura 2000 network of protected areas (N2000) has become the mainstay of
76 current conservation policies (Gaston et al., 2008). N2000 is based on two European Directives: (i) the
77 Birds Directive (Official Journal of the European Union, 2009), which provides a list of birds for which
78 the member states are required to designate Special Protection Areas (SPAs); and (ii) the Habitats
79 Directive (Council Directive, 1992), which aims to protect specific animals (other than birds), plants

80 and habitats, for which each member state is required to designate Special Areas of Conservation
81 (SACs). Together, SPAs and SACs form the N2000 network.

82 In the case of SACs, the presence of a species in a particular annex of the Habitats Directive can
83 be used as a proxy for conservation interest. The species of highest conservation interest tend to be those
84 appearing in Annex II and Annex IV of the Habitats Directive. Annex II species are defined as “*animal*
85 *or plant species of community interest whose conservation requires the designation of SAC sites*”
86 (Council Directive, 1992). For Annex IV species “*a strict protection regime must be applied across*
87 *their entire natural range within the EU, both within and outside Natura 2000 sites*” (Council Directive,
88 1992). The Habitats Directive recognizes caves as Priority Habitat (Code 8310) and many SAC sites
89 were designated specifically for bats, and to include caves. For each SAC site, there is a Standard Data
90 Form (SDF; Official Journal of the European Union, 2011), which defines its characteristics, location,
91 size and the species present, for which it was designated. These SDFs are the basis for developing future
92 management and recovery plans for these species. They are also crucial for decision-making because
93 they are consulted to make Environmental Impact Assessments, for decisions on the development of
94 infrastructures and urban areas, and to prioritise allocation of public funds for agriculture and forestry.

95 A number of studies have evaluated the effectiveness of the N2000 network in representing
96 particular taxonomic groups, including plants (Chiarucci et al., 2008; Kallimanis et al., 2014),
97 invertebrates (Abellán et al., 2007; Sánchez-Fernández et al., 2008) and various vertebrate groups
98 (Abellán and Sánchez-Fernández, 2015; Albuquerque et al., 2013; Lisón et al., 2015b; Maiorano et al.,
99 2015). Some have examined species of conservation concern in different taxonomic groups, for example
100 species included in Annex II of the Habitats Directive (Gruber et al., 2012; Lisón et al., 2013) or
101 threatened species (Trochet and Schmeller, 2013). However, there are no studies, to our knowledge, that
102 investigate whether the SDFs are accurate in their documentation of the biodiversity present in the SAC
103 sites they support, or whether this accuracy differs between threat categories for the species according
104 to their status in the Habitats Directive’s annexes.

105 Bats are an appropriate group to explore these questions because all European bat species are
106 included in either Annex II or Annex IV of the Habitats Directive. European bats form a relatively
107 homogeneous group of mammals, with similar ecological requirements (Dietz et al., 2009; Jones et al.,

108 2009). Bats are also threatened (Jones et al., 2009), with most species' populations decreasing in recent
109 decades (Dietz et al., 2009). In some European countries, however, an increase in the range and size of
110 bat populations has been observed and attributed to application of specific management measures
111 (Haysom et al., 2013). Because of their position in food webs, bats are very important species for the
112 maintenance of ecosystem functions such as suppression of insect pests, pollination, seed dispersal, and
113 nutrient cycling through their guano in terrestrial, aquatic and cave ecosystems (Civantos et al., 2012;
114 Jones et al., 2009; Kunz et al., 2011). As well as providing such valuable ecosystem services for humans,
115 bats serve as bioindicator and sentinel species (Boyles et al., 2011; Jones et al., 2009; Lisón et al., 2015a).
116 Nevertheless, rather few studies have specifically examined the relationship between bat distributions
117 and the N2000 network (Lisón et al., 2013, 2015b; Maiorano et al., 2015; Zehetmair et al., 2015).

118 Spain represents approximately 18% of the surface of N2000 within the EU, with more protected
119 land surface area in N2000 than any other country (European Commission, 2010). With 31 bat species,
120 the Iberian Peninsula contains a high proportion of European bat biodiversity (approximately 80% of
121 European bat species; Dietz et al., 2009; Palomo et al., 2007). Using data on bat species' distributions
122 from mainland Spain and the Balearic Islands, we test the accuracy of data documented in SDFs for
123 each SAC site, with respect to the bat biodiversity contained within these protected areas. We do this by
124 comparing the species listed in the SDFs with the known distributions of European bat species. We also
125 test for differences in this accuracy of the SDFs between Annex II and Annex IV species. Finally, we
126 discuss the implications of the deficiencies we find for achieving the N2000 aims, and for conservation
127 more generally.

128

129 **2. Materials and methods**

130 Our dataset for bat occurrences in mainland Spain and the Balearic Islands was based initially on
131 distribution maps published in the Atlas and Red Book of Spanish terrestrial mammals (Palomo et al.,
132 2007), with a resolution of 10 x 10 km UTM cells. Given the large home ranges of most bats and the
133 fact that SACs frequently contain bat roosts, it is reasonable to attribute species' presences to SACs
134 based on this distribution information. We added to the dataset 3708 new records at the same resolution
135 from our own field sampling, and from a comprehensive compilation of recent literature (see Appendix

136 A). This information was generated by bat experts using different methodologies (roost searching,
137 acoustic surveys and trapping), minimizing the risk of bias in species detection (Flaquer et al., 2007),
138 and providing important information even for rare species. The size of the spatial units used is
139 appropriate given that the home-range of bats usually spans several kilometres (Dietz et al., 2009;
140 Rainho and Palmeirim, 2011) and the roosts of most of the bats are in SACs (Lisón et al., 2013; Rainho
141 and Palmeirim, 2013). We used this dataset to determine which bat species are present in each Spanish
142 terrestrial SAC, according to the best available information. We refer to this as the ‘known presences’
143 of bat species in SACs.

144 We used data for 29 of the 31 species present in the study area. We did not include *Myotis*
145 *nattereri/escalerai*, because these cryptic species have only recently been separated and their
146 distributions are not yet well known (Palomo et al., 2007). For the sibling species *Eptesicus serotinus*
147 and *E. isabellinus*, we considered the distribution of the latter to be confined to Andalusia and Murcia
148 in south-southeast Iberia (Lisón et al., 2015b), while the former is distributed throughout the rest of the
149 Iberian Peninsula, although there may be a contact zone (Santos et al., 2014). Eleven out of the 29
150 species in our dataset are listed in Annex II of the Habitats Directive and the other 18 are listed in Annex
151 IV (see Appendices B and C).

152 In Spain, Autonomous Communities (AACCs or ‘regions’) are responsible for designating the
153 SACs, so we aggregated data at the AACC level for analysis and display purposes. After determining
154 which bat species are present in which SACs, according to our distribution dataset (known presences),
155 we calculated the number of SACs in each region that contain at least one Annex II bat species, and the
156 number that contain at least one Annex IV bat species (Appendix C).

157 In parallel, we analysed the Standard Data Form (SDF) for each SAC (Spanish Government,
158 2013), which were updated in 2013. We recorded which bat species are officially documented as being
159 present in each Spanish terrestrial SAC, according to its SDF. We refer to this as the ‘official presences’
160 of bat species in SACs. Using this parallel dataset, we again calculated the number of SACs in each
161 region that contain at least one Annex II bat species, and the number that contain at least one Annex IV
162 bat species (Appendix B).

163 For each region, we divided the number of SACs that have official presences of Annex II or
164 Annex IV bat species by the number of SACs in the region, to determine the percentage of SACs
165 containing these target species. We refer to this as the ‘official percentage of SACs’ with Annex II or
166 with Annex IV species. Then we repeated the process using known presences instead of official
167 presences, to produce the ‘known percentage of SACs’ with Annex II or with Annex IV bat species. For
168 each Annex type separately, we used a Wilcoxon test to determine whether there were significant
169 differences between the official and actual percentages of SACs with species in that Annex. All analyses
170 were performed using R software (R Core Team 2014), including the *Rcmdr* package.

171 Some SACs are underground roosts and their designation as SACs was exclusively due to the
172 presence of bats. We refer to these SACs as ‘exclusive’, to indicate the centrality of bats to their
173 protected status. These sites are usually very small because only the entrance to the roost (cave, mine or
174 tunnel) has been protected, but we highlight their importance since they have been protected specifically
175 for their bat communities. Usually, the information contained in the SDFs is based on a regional atlas
176 compiled by regional experts. However, it is difficult to know when surveys were done, what the survey
177 effort was (and thus how complete the species list is likely to be for any particular SAC) or the aims of
178 the surveys, because this information is normally not published and not available.

179

180 **3. Results**

181 We analysed a total of 1206 SDFs for SAC sites in Spain (excluding the Canary Islands). According to
182 this official documentation, 481 SACs (39.9%) have Annex II species and 123 (10.2%) have Annex IV
183 species (Table 1). These numbers are much lower than when we recalculate them using known
184 presences: 944 SACs (78.3 %) contain Annex II species and 953 (79.0 %) contain Annex IV species.
185 Eighty-one SACs (6.7 %) were designated exclusively for bats (Table 1); five regions had no SACs of
186 this type, while the highest amount was 17 of the 84 (20.2%) SACs in Comunidad Valenciana being of
187 this type. All regions (AACCs) had Annex II species in the technical reports for at least some of their
188 SACs; proportions ranged from 16.9% (30 out of 178) of the SACs in Andalucía containing at least one
189 Annex II species to 100% for La Rioja (all six) and Madrid (all seven). Again, these numbers are much
190 higher when using known presences instead of official presences (Table 1). For Annex IV bat species,

191 six regions did not report the presence of any of these species in their SDFs for any of their SACs, while
192 the highest percentage was Madrid, in which six of the seven (86%) of the SACs contained Annex IV
193 species. Once again, the numbers were much higher for known presences than for official presences
194 (Table 1).

195 Considering each species separately, the number and percentage of SACs in each region that
196 contain individual Annex II and Annex IV bat species are shown in Appendix B (official presences) and
197 Appendix C (known presences). Across all species and regions, we found that the percentage of SACs
198 with Annex II bat species officially present was significantly lower than for known presences (Wilcoxon
199 test, $w = 215$, $P < 0.001$; Figure 1A). We found the same for Annex IV bat species (Wilcoxon test, $w =$
200 242 , $P < 0.001$; Figure 1B). Also, the percentage of SACs with Annex II bat species officially present
201 was significantly higher than that for Annex IV bat species (Wilcoxon test, $w = 233$, $P < 0.001$; Figure
202 1), but the percentages of known presences did not differ significantly between the two annexes
203 (Wilcoxon test, $w = 170$, $P = 0.119$).

204 Similarly, when we analysed regions separately we found significant underrepresentation of
205 Annex II bat species in the SDFs (Figure 2A): there were always higher levels of known presence than
206 official presence for these species. Importantly, there was much more marked underrepresentation of
207 the presence of Annex IV species in the SDFs (Figure 2B).

208

209 **4. Discussion**

210 We have shown that the Standard Data Forms for Special Areas of Conservation in Spain seriously
211 under-represent the occurrences and diversity of protected bat species known to be present in those
212 SACs. This deficiency is particularly large for the Annex IV species. While the scientific community
213 has carefully studied the effectiveness of the Natura 2000 network in terms of whether the SACs and
214 SPAs contain various taxonomic groups and their habitats (e.g. Abellán and Sánchez-Fernández, 2015;
215 Lisón et al., 2013, 2015b; Maiorano et al., 2015; Sánchez-Fernández et al., 2008), very little attention
216 has been paid to whether the SDFs supporting these protected areas accurately represent the biodiversity
217 known to be within them. This issue is very important because the SDFs underpin management and
218 conservation plans, and are central to the practical implementation of sustainable development (Popescu

219 et al., 2014): conservation planning should be established in accordance with actual biodiversity patterns
220 (Pressey et al., 2003; Jeanmougin et al., 2016). Further, the SDFs are the defining documents used in
221 decision-making by governments regarding territorial planning, land use, future infrastructure
222 development, Environmental Impact Assessment, N2000 connectivity, public funds for agroforestry and
223 mitigating policies (Mazaris et al., 2013; Romano and Zullo, 2015; Stone et al., 2013). The SDFs can
224 be used to indicate conservation needs in N2000 sites, and they are keystone documents for achieving
225 the N2000 conservation aims (Hochkirch et al., 2013; Kati et al., 2015) and Aichi Targets (Convention
226 on Biological Diversity, 2010).

227 Some studies have questioned the effectiveness of the N2000 network in protecting bat species
228 (Lisón et al., 2013; Zehetmair et al., 2015) and whether the Annex II bat species act as ‘umbrella species’
229 representing the remaining bat species (Lisón et al., 2015b), although some Annex IV species are
230 threatened or rare (Palomo et al., 2007). Our results add a new dimension to a growing body of research
231 that identifies shortcomings in conservation policies with respect to biological knowledge (Rodhouse et
232 al., 2016). Also, they highlight the importance of improving communication between ecologists and
233 managers, as well as the need to implement open communication channels that will help make protected-
234 area management more effective. Part of a new emphasis on better data in official documentation for
235 protected areas can be the involvement of the public in gathering such data (citizen science; see e.g.
236 Bonney et al., 2009; Silvertown, 2009; Barlow et al., 2015; Newson et al., 2015).

237 Our quantification of the very large under-representation of bats in the SDFs of Special Areas
238 of Conservation is partly based on the assumption that distribution data at 10 km x 10 km resolution
239 indicate the presence or absence of species in SACs within those grid squares. This is potentially
240 problematic for two main reasons. First, some of the distribution data may reflect historic presences but
241 the species may now be absent from the grid square. Second, species do not completely fill the
242 landscape, so presence within a grid square does not necessarily mean presence in a SAC within the grid
243 square. However, given what we know about bat ecology and movement capacity (Dietz et al., 2009;
244 Rainho and Palmeirim, 2013) and the habitats contained within SACs, we consider it likely that most
245 bat species appearing in areas surrounding SACs use those sites for roosting or feeding, or that the SACs
246 contain important habitat for them in some other respect (Lisón et al., 2013, 2015b; Rainho and

247 Palmeirim, 2013). Further, part of our dataset was from recent field surveys. Certainly it is inconceivable
248 that the massive under-representation of known occurrences that we found in the SDFs is entirely due
249 to artefacts of the methods. Instead, the great differences found in almost all the bat species between the
250 official and known occurrences show that there are serious deficiencies in the application of Natura
251 2000, especially for Annex IV bat species, even though most of these species are easily detected by
252 ultrasound surveys (Flaquer et al., 2007) and many have quite a wide distribution (Appendix C; Palomo
253 et al., 2007).

254 The under-reporting of species in the SDFs could also be due in part to those reports being
255 written several years ago (Lisón et al., 2015b; Palomo et al., 2007), when there was less information
256 about bat distributions in some regions. However, SDFs were updated in 2013 after the last report made
257 during the period between 2007 and 2012 under art. 17 of the Habitats Directive. At this time, the Atlas
258 of Mammals of Spain was revised and contained considerable information on bat distributions.
259 Therefore, the necessary information was available and this problem could be solved simply by updating
260 the dataset (see Appendices). There may also be a lack of political will to ensure that the reports are up
261 to date, and greater willingness would strengthen the protection of species offered by N2000 (Rojas-
262 Briales, 2000; Orlikowska et al., 2016).

263 We cannot reject the possibility that our results represent an issue restricted to Spain, though we
264 consider it highly unlikely. Given the seriousness of the implications of our findings, similar studies in
265 other countries and with different taxa should be undertaken as a matter of urgency. Further, the Spanish
266 situation in itself is of international interest because of the high importance of Spain within the N2000
267 network, and its high biodiversity within the European context.

268 Member States are not specifically required to list the Annex IV species present in the Natura
269 2000 sites (Official Journal of the European Union, 2011). This does not explain under-representation
270 as a general phenomenon because Annex II species are also significantly under-represented in the
271 official documentation. However, it is almost certainly one reason why the under-representation is much
272 more extreme for Annex IV than Annex II species. Given the need to conserve Annex IV species, many
273 of which are threatened, our results thus highlight the lack of requirement to list Annex IV species in

274 SDFs as a major problem with the Habitats Directive; it would be interesting to explore why this
275 requirement is missing from the legislation.

276 Certainly, the under-representation of protected species in the SDFs biases the outcomes and
277 associated decision-making of N2000 against conservation objectives, and favours developers and other
278 parties that may oppose conservation goals (Apostolopoulou and Pantis, 2009; Margules and Pressey,
279 2000). Thus it is very likely to negatively affect conservation and the aims of N2000. But these aims are
280 very important. For one thing, some Annex IV bat species are rare, and often endemic and cryptic, and
281 therefore could face high levels of threat (see national Red List in Palomo et al., 2007 and regional Red
282 List in Lisón et al., 2011). Further, bats have important roles in ecosystem structure and function (Boyles
283 et al., 2011; Jones et al., 2009; Kunz et al., 2011) to ignore their presence in the SDFs will hinder the
284 development of strong conservation plans to guarantee the integrity and coherence of the N2000 network
285 (Orlikowska et al., 2016).

286

287 **5. Conclusions**

288 The considerable deficit of occurrence information for species of conservation importance in the SDFs
289 of protected areas is a serious concern, undermining the effectiveness of conservation networks such as
290 Natura 2000. The technical reports are the keystone to management and conservation plans and decision-
291 making within protected areas. There is an urgent need to update them with much more complete
292 information about which species are in which protected conservation sites, and also to open up
293 communication channels between conservation scientists and the managers of protected areas:
294 incorporating citizen science would enhance this. Better knowledge of the biodiversity of protected
295 areas, and their conservation status, will aid decision-making at all levels.

296

297 **6. Acknowledgements**

298 We are grateful for the important contribution of bat workers and SECEMU members to the
299 development of the distribution of bat fauna of Spain. We thank Ángeles Haz for correcting the English
300 and José F. Calvo and David Sánchez-Fernández for his support and help. FL was supported by a
301 fellowship (Programa MECE Educación Superior) from the Chilean Ministry of Education and

302 postdoctoral fellowship (Programa de Formación de Investigadores Postdoctorales) from Universidad
303 de La Frontera, Chile.

304

305 7. References

306 Abellán, P., Sánchez-Fernández, D., 2015. A gap analysis comparing the effectiveness of Natura 2000
307 and national protected area networks in representing European amphibians and reptiles.
308 *Biodivers. Conserv.* 24, 1377–1390. doi:10.1007/s10531-015-0862-3

309 Abellán, P., Sánchez-Fernández, D., Velasco, J., Millán, A., 2007. Effectiveness of protected area
310 networks in representing freshwater biodiversity: the case of a Mediterranean river basin (south-
311 eastern Spain). *Aquat. Conserv. Mar. Freshw. Ecosyst.* 17, 361–364. doi:10.1002/aqc

312 Albuquerque, F.S., Assunção-Albuquerque, M.J.T., Cayuela, L., Zamora, R., Benito, B.M., 2013.
313 European Bird distribution is “well” represented by Special Protected Areas: mission
314 accomplished? *Biol. Conserv.* 159, 45–50. doi:10.1016/j.biocon.2012.10.012

315 Apostolopoulou, E., Pantis, J.D., 2009. Conceptual gaps in the national strategy for the implementation
316 of the European Natura 2000 conservation policy in Greece. *Biol. Conserv.* 142, 221–237.
317 doi:10.1016/j.biocon.2008.10.021

318 Araújo, M.B., Alagador, D., Cabeza, M., Nogués-Bravo, D., Thuiller, W., 2011. Climate change
319 threatens European conservation areas. *Ecol. Lett.* 14, 484–492. doi:10.1111/j.1461-
320 0248.2011.01610.x

321 Barlow, K.E., Briggs, P.A., Haysom, K.A., Hutson, A.M., Lechiara, N.L., Racey, P.A., Walsh, A.L.,
322 Langton, S.D., 2015. Citizen science reveals trends in bat populations: The National Bat
323 Monitoring Programme in Great Britain. *Biol. Conserv.* 182, 14–26.
324 doi:10.1016/j.biocon.2014.11.022

325 Bonney, R., Cooper, C.B., Dickinson, J., Kelling, S., Phillips, T., Rosenberg, K. V., Shirk, J., 2009.
326 Citizen science: a developing tool for expanding science knowledge and scientific literacy.
327 *Bioscience* 59, 977–984. doi:10.1525/bio.2009.59.11.9

328 Boyles, J.G., Cryan, P.M., McCracken, G.F., Kunz, T.K., 2011. Economic importance of bats in
329 agriculture. *Science* 332, 41–42. doi:10.1126/science.1201366

330 Butchart, S.H.M., Walpole, M., Collen, B., van Strien, A., Scharlemann, J.P.W., Almond, R.E. a, Baillie,
331 J.E.M., Bomhard, B., Brown, C., Bruno, J., Carpenter, K.E., Carr, G.M., Chanson, J., Chenery,
332 A.M., Csirke, J., Davidson, N.C., Dentener, F., Foster, M., Galli, A., Galloway, J.N., Genovesi,
333 P., Gregory, R.D., Hockings, M., Kapos, V., Lamarque, J.-F., Leverington, F., Loh, J.,
334 McGeoch, M. a, McRae, L., Minasyan, A., Hernández Morcillo, M., Oldfield, T.E.E., Pauly,
335 D., Quader, S., Revenga, C., Sauer, J.R., Skolnik, B., Spear, D., Stanwell-Smith, D., Stuart,

336 S.N., Symes, A., Tierney, M., Tyrrell, T.D., Vié, J.-C., Watson, R., 2010. Global biodiversity:
337 indicators of recent declines. *Science* 328, 1164–1168. doi:10.1126/science.1187512

338 Chiarucci, A., Bacaro, G., Rocchini, D., 2008. Quantifying plant species diversity in a Natura 2000
339 network: old ideas and new proposals. *Biol. Conserv.* 141, 2608–2618.
340 doi:10.1016/j.biocon.2008.07.024

341 Civantos, E., Thuiller, W., Maiorano, L., Guisan, A., Araújo, M.B., 2012. Potential impacts of climate
342 change on ecosystem services in Europe: the case of pest control by vertebrates. *Bioscience* 62,
343 658–666. doi:10.1525/bio.2012.62.7.8

344 Convention on Biological Diversity, 2010. Aichi Biodiversity Targets 9–10.

345 Council Directive, 1992. Council Directive 92/43/EEC of 21 May 1992 on the conservation of natural
346 habitats and of wild fauna and flora. Available at: [http://eur-lex.europa.eu/legal-](http://eur-lex.europa.eu/legal-content/EN/ALL/?uri=CELEX:01992L0043-20070101)
347 [content/EN/ALL/?uri=CELEX:01992L0043-20070101](http://eur-lex.europa.eu/legal-content/EN/ALL/?uri=CELEX:01992L0043-20070101)

348 D’Amen, M., Bombi, P., Pearman, P.B., Schmatz, D.R., Zimmermann, N.E., Bologna, M.A., 2011. Will
349 climate change reduce the efficacy of protected areas for amphibian conservation in Italy? *Biol.*
350 *Conserv.* 144, 989–997. doi:<http://dx.doi.org/10.1016/j.biocon.2010.11.004>

351 de Koning, J., Winkel, G., Sotirov, M., Blondet, M., Borrás, L., Ferranti, F., Geitzner, M., 2014.
352 Natura 2000 and climate change—Polarisation, uncertainty, and pragmatism in discourses on
353 forest conservation and management in Europe. *Environ. Sci. Policy* 39, 129–138.
354 doi:<http://dx.doi.org/10.1016/j.envsci.2013.08.010>

355 Dietz, C., von Helversen, O., Nill, D., 2009. *Bats of Britain, Europe and northwest Africa*. A&C Black
356 Publishers Ltd., London.

357 European Commission, 2010. Natura 2000 database and GIS. Available at:
358 http://ec.europa.eu/environment/nature/natura2000/db_gis/index_en.htm#area_calc

359 Flaquer, C., Torre, I., Arrizabalaga, A., Torre, I., 2007. Comparison of sampling methods for inventory
360 of bat communities. *J. Mammal.* 88, 526–533. doi:10.1644/06-MAMM-A-135R1.1

361 Gaston, K.J., Jackson, S.F., Nagy, A., Cantú-Salazar, L., Johnson, M., 2008. Protected areas in Europe:
362 principle and practice. *Ann. N. Y. Acad. Sci.* 1134, 97–119. doi:10.1196/annals.1439.006

363 Gruber, B., Evans, D., Henle, K., Bauch, B., Schmeller, D.S., Dziock, F., Henry, P.-Y., Lengyel, S.,
364 Margules, C., Dormann, C.F., 2012. “Mind the gap!” – How well does Natura 2000 cover
365 species of European interest? *Nat. Conserv.* 3, 45–63. doi:10.3897/natureconservation.3.3732

366 Haysom, K., Dekker, J., Russ, J., van der Meij, T., van Strien, A., 2013. European bat population trends.
367 A prototype biodiversity indicator, EEA Technical report No19. doi:10.2800/10311

368 Hochkirch, A., Schmitt, T., Beninde, J., Hiery, M., Kinitz, T., Kirschey, J., Matenaar, D., Rohde, K.,
369 Stoefen, A., Wagner, N., Zink, A., Lötters, S., Veith, M., Proelss, A., 2013. Europe needs a new
370 vision for a Natura 2020 Network. *Conserv. Lett.* 6, 462–467. doi:10.1111/conl.12006

371 Jeanmougin, M., Dehais, C., Meinard, Y., 2016. Mismatch between habitat science and habitat directive
372 – lessons from the French (counter) example. *Conserv. Lett.*, doi: 10.1111/conl.12330

373 Jones, G., Jacobs, D.S., Kunz, T.H., Wilig, M.R., Racey, P.A., 2009. Carpe noctem: The importance of
374 bats as bioindicators. *Endanger. Species Res.* 8, 93–115. doi:10.3354/esr00182

375 Kallimanis, A.S., Touloumis, K., Tzanopoulos, J., Mazaris, A.D., Apostolopoulou, E., Stefanidou, S.,
376 Scott, A. V., Potts, S.G., Pantis, J.D., 2014. Vegetation coverage change in the EU: patterns
377 inside and outside Natura 2000 protected areas. *Biodivers. Conserv.* 24, 579–591.
378 doi:10.1007/s10531-014-0837-9

379 Kati, V., Hovardas, T., Dieterich, M., Ibisch, P.L., Mihok, B., Selva, N., 2015. The challenge of
380 implementing the European network of protected areas Natura 2000. *Conserv. Biol.* 29, 260–
381 270. doi:10.1111/cobi.12366

382 Kukkala, A.S., Arponen, A., Maiorano, L., Moilanen, A., Thuiller, W., Toivonen, T., Zupan, L.,
383 Brotons, L., Cabeza, M., 2016. Matches and mismatches between national and EU-wide
384 priorities: examining the Natura 2000 network in vertebrate species conservation. *Biol. Conserv.*
385 198, 193–201. doi:10.1016/j.biocon.2016.04.016

386 Kukkala, A.S., Moilanen, A., 2013. Core concepts of spatial prioritisation in systematic conservation
387 planning. *Biol. Rev.* 88, 443–464. doi:10.1111/brv.12008

388 Kunz, T.H., de Torrez, E.B., Bauer, D., Lobova, T., Fleming, T.H., 2011. Ecosystem services provided
389 by bats. *Ann. N. Y. Acad. Sci.* 1223, 1–38. doi:10.1111/j.1749-6632.2011.06004.x

390 Lee, T.M., Sodhi, N.S., Prawiradilaga, D.M., 2007. The importance of protected areas for the forest and
391 endemic avifauna of Sulawesi (Indonesia). *Ecol. Applic.* 17: 1272–1741. doi: 10.1890/06-
392 1256.1

393 Lisón, F., Aledo, E., Calvo, J.F., 2011. Los murciélagos (Mammalia: Chiroptera) de la Región de Murcia
394 (SE España): distribución y estado de conservación. *An. Biol.* 33, 79–92.

395 Lisón, F., López-Espinosa, J.A., Calvo, J.F., Jones, G., 2015a. Diet of the meridional serotine *Eptesicus*
396 *isabellinus* in an urban semiarid Mediterranean landscape. *Acta Chiropterol.* 17, 371–378.
397 doi:10.3161/15081109ACC2015.17.2.013

398 Lisón, F., Palazón, J.A., Calvo, J.F., 2013. Effectiveness of the Natura 2000 Network for the
399 conservation of cave-dwelling bats in a Mediterranean region. *Anim. Conserv.* 16, 528–537.
400 doi:10.1111/acv.12025

401 Lisón, F., Sánchez-Fernández, D., Calvo, J.F., 2015b. Are species listed in the Annex II of the Habitats
402 Directive better represented in Natura 2000 network than the remaining species? A test using
403 Spanish bats. *Biodivers. Conserv.* 24, 2459–2473.

404 Maiorano, L., Amori, G., Montemaggiori, A., Rondinini, C., Santini, L., Saura, S., Boitani, L., 2015.
405 On how much biodiversity is covered in Europe by national protected areas and by the Natura
406 2000 network: insights from terrestrial vertebrates. *Conserv. Biol.* 29, 956–995.
407 doi:10.1111/cobi.12535

408 Margules, C.R., Pressey, R.L., 2000. Systematic conservation planning. *Nature* 405, 243–53.
409 doi:10.1038/35012251

410 Mazaris, A.D., Papanikolaou, A.D., Barbet-Massin, M., Kallimanis, A.S., Jiguet, F., Schmeller, D.S.,
411 Pantis, J.D., 2013. Evaluating the connectivity of a protected areas' network under the prism of
412 global change: the efficiency of the European Natura 2000 Network for four birds of prey. *PLoS*
413 *One* 8, e59640. doi:10.1371/journal.pone.0059640

414 Newson, S.E., Evans, H.E., Gillings, S., 2015. A novel citizen science approach for large-scale
415 standardised monitoring of bat activity and distribution, evaluated in eastern England. *Biol.*
416 *Conserv.* 191, 38–49. doi:10.1016/j.biocon.2015.06.009

417 Official Journal of the European Union, 2009. Directive 2009/147/EC of the European Parliament and
418 of the Council of 30 November 2009 on the conservation of wild birds. Available at: [http://eur-](http://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX:32009L0147)
419 [lex.europa.eu/legal-content/EN/TXT/?uri=CELEX:32009L0147](http://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX:32009L0147)

420 Official Journal of the European Union, 2011. Commission implementing decision of 11 July 2011
421 concerning a site information format for Natura 2000 sites. Available at: [http://eur-](http://eur-lex.europa.eu/legal-content/EN/ALL/?uri=CELEX%3A32011D0484)
422 [lex.europa.eu/legal-content/EN/ALL/?uri=CELEX%3A32011D0484](http://eur-lex.europa.eu/legal-content/EN/ALL/?uri=CELEX%3A32011D0484)

423 Orlikowska, E.H., Roberge, J.M., Blicharska, M., Mikusiński, G., 2016. Gaps in ecological research on
424 the world's largest internationally coordinated network of protected areas: A review of Natura
425 2000. *Biol. Conserv.* 200, 216-227. doi: 10.1016/j.biocon.2016.06.015

426 Palomo, L.J., Gisbert, J., Blanco, J.C., 2007. *Atlas y Libro Rojo de los mamíferos terrestres de España.*
427 Dirección General de la Biodiversidad-SECEM-SECEMU, Madrid.

428 Popescu, V.D., Rozyłowicz, L., Niculae, I.M., Cucu, A.L., Hartel, T., 2014. Species, habitats, society:
429 an evaluation of research supporting EU's Natura 2000 network. *PLoS One* 9, e113648.
430 doi:10.1371/journal.pone.0113648

431 Pressey, R.L., Cowling, R.M., Rouget, M., 2003. Formulating conservation targets for biodiversity
432 pattern and process in the Cape Floristic Region, South Africa. *Biol. Conserv.* 112, 99–127.
433 doi:10.1016/S0006-3207(02)00424-X

434 R Core Team, 2014. *R: A Language and Environment for Statistical Computing.* R Foundation for
435 Statistical Computing, Vienna, Austria.

436 Rainho, A., Palmeirim, J.M., 2013. Prioritizing conservation areas around multispecies bat colonies
437 using spatial modeling. *Anim. Conserv.* 16, 438–448. doi:10.1111/acv.12013

438 Rainho, A., Palmeirim, J.M., 2011. The importance of distance to resources in the spatial modelling of
439 bat foraging habitat. *PLoS One* 6. doi:10.1371/journal.pone.0019227

440 Rodhouse, T.J., Philippi, T.E., Monahan, W.B., Castle, K.T., 2016. A macroecological perspective on
441 strategic bat conservation in the U.S. National Park Service. *Ecosphere*, 7, e01576. doi:
442 10.1002/ecs2.1576.

443 Rojas-Briales, E., 2000. Socio-economics of nature protection policies in the perspective of the
444 implementation of Natura 2000 Network: the Spanish case. *Forestry* 73, 199–207.
445 doi:10.1093/forestry/73.2.199

446 Romano, B., Zullo, F., 2015. Protected areas, Natura 2000 Sites and landscape: divergent policies on
447 converging values, in: Gambino, R., Peano, A. (Eds.), *Nature Policies and Landscape Policies*.
448 Springer, pp. 127–136.

449 Sánchez-Fernández, D., Bilton, D.T., Abellán, P., Ribera, I., Velasco, J., Millán, A., 2008. Are the
450 endemic water beetles of the Iberian Peninsula and the Balearic Islands effectively protected?
451 *Biol. Conserv.* 141, 1612–1627. doi:10.1016/j.biocon.2008.04.005

452 Santos, H., Juste, J., Ibáñez, C., Palmeirim, J.M., Godinho, R., Amorim, F., Alves, P., Costa, H., de Paz,
453 O., Pérez-Suarez, G., Martínez-Alos, S., Jones, G., Rebelo, H., 2014. Influences of ecology and
454 biogeography on shaping the distributions of cryptic species: three bat tales in Iberia. *Biol. J.*
455 *Linn. Soc.* 112, 150–162. doi:10.1111/bij.12247

456 Le Saout, S., Hoffmann, M., Shi, Y., Hughes, A., 2013. Protected areas and effective biodiversity
457 conservation. *Science* 342, 803–805. doi:10.1126/science.1239268

458 Silvertown, J., 2009. A new dawn for citizen science. *Trends Ecol. Evol.* 24, 467–71.
459 doi:10.1016/j.tree.2009.03.017

460 Spanish Government, 2013. Lugares de Importancia Comunitaria (LIC). Available at:
461 [http://www.magrama.gob.es/es/biodiversidad/temas/espacios-protegidos/red-natura-](http://www.magrama.gob.es/es/biodiversidad/temas/espacios-protegidos/red-natura-2000/lic.aspx)
462 [2000/lic.aspx](http://www.magrama.gob.es/es/biodiversidad/temas/espacios-protegidos/red-natura-2000/lic.aspx)

463 Stone, E.L., Jones, G., Harris, S., 2013. Mitigating the effect of development on bats in England with
464 derogation licensing. *Conserv. Biol.* 27, 1324–1334. doi:10.1111/cobi.12154

465 Trochet, A., Schmeller, D.S., 2013. Effectiveness of the Natura 2000 network to cover threatened
466 species. *Nat. Conserv.* 4, 35–53. doi:10.3897/natureconservation.4.3626

467 UNEP-WCMC, 2016. Mapping the world's special places. Opening access to data on global protected
468 areas. Available at: [http://www.unep-wcmc.org/featured-projects/mapping-the-worlds-special-](http://www.unep-wcmc.org/featured-projects/mapping-the-worlds-special-places)
469 [places](http://www.unep-wcmc.org/featured-projects/mapping-the-worlds-special-places)


470 Zehetmair, T., Müller, J., Runkel, V., Stahlschmidt, P., Winter, S., Zharov, A., Gruppe, A., 2015. Poor
471 effectiveness of Natura 2000 beech forests in protecting forest-dwelling bats. *J. Nat. Conserv.*
472 23, 53–60. doi:10.1016/j.jnc.2014.07.003
473

474 Table 1: Number of Special Areas of Conservation (SACs) containing Annex II and Annex IV bat
 475 species in each Autonomous Community (AACC) of Spain. ‘Abbrev.’ is the abbreviation used for each
 476 AACC in Figure 2 and in the supplementary online material. ‘No. of SACs’ is the number of SACs in
 477 each AACC, and ‘Exclusive’ is the number of these that were designated exclusively for bats. The other
 478 columns contain the number of SACs in each region that have Annex II or Annex IV bat species
 479 according to the Standard Data Forms (official presences) or known presences. Numbers in parentheses
 480 are percentages of the total number of SACs in the AACC.
 481

AACC	Abbrev.	No. of SACs	Official presences		Known presences		Exclusive
			Annex II	Annex IV	Annex II	Annex IV	
Andalucía	And	178	30 (16.9%)	3 (1.7%)	118 (66.3%)	87 (48.9%)	3 (8.4%)
Aragón	Ara	156	69 (44.2%)	29 (18.6%)	106 (67.9%)	119 (76.3%)	12 (7.7%)
Asturias	Ast	49	19 (38.8%)	0 (0.0%)	44 (89.6%)	44 (89.6%)	1 (2.0%)
Baleares	Bal	111	29 (26.1%)	5 (4.5%)	68 (61.3%)	79 (71.2%)	8 (7.2%)
Cantabria	Can	21	10 (47.6%)	0 (0.0%)	20 (95.2%)	19 (90.5%)	2 (9.5%)
Castilla León	Cle	122	70 (57.4%)	0 (0.0%)	110 (90.2%)	118 (96.7%)	3 (2.5%)
Castilla La Mancha	Cma	74	19 (25.7%)	0 (0.0%)	55 (74.3%)	56 (75.7%)	5 (6.8%)
Cataluña	Cat	111	81 (73.0%)	0 (0.0%)	105 (94.6%)	99 (89.2%)	2 (1.8%)
Extremadura	Ext	88	34 (38.6%)	17 (19.3%)	69 (78.4%)	87 (98.9%)	14 (15.9%)
Galicia	Gal	58	11 (19.0%)	0 (0.0%)	48 (82.8%)	44 (75.9%)	0 (0.0%)
Madrid	Mad	7	7 (100.0%)	6 (85.7%)	7 (100.0%)	7 (100.0%)	0 (0.0%)
Región de Murcia	Mur	47	28 (59.6%)	31 (66.0%)	38 (80.9%)	38 (80.9%)	2 (4.3%)
Navarra	Nav	42	16 (38.1%)	9 (21.4%)	35 (83.3%)	42 (100.0%)	0 (0.0%)
La Rioja	Rio	6	6 (100.0%)	4 (66.7%)	6 (100.0%)	6 (100.0%)	0 (0.0%)
Comunidad Valenciana	Val	84	37 (44.0%)	9 (10.7%)	63 (75.0%)	56 (66.7%)	17 (20.2%)
País Vasco	Vas	52	15 (28.8%)	10 (19.2%)	52 (100.0%)	52 (100.0%)	0 (0.0%)
Total		1206	481 (39.9%)	123 (10.2%)	944 (78.3%)	953 (79.0%)	81 (6.7%)


482

483




484
 485 Figure 1: Boxplots representing the variation across regions (Autonomous Communities) in the percentage of Special Areas of Conservation that contain at least
 486 one Annex II (A) or Annex IV (B) bat species. Each percentage value (vertical axis, labeled “% SACs containing bats”) refers to one region.
 487

488 (A) Annex II



489

490 (B) Annex IV



491

492 Figure 2: Under-representation of bat species in the Standard Data Forms (SDFs) supporting Special
493 Areas of Conservation (SACs; 'Official' – white bars), as judged by known occurrences ('Known' –
494 black bars). For each region (Autonomous Community; abbreviations explained in Table 1), the mean
495 percentage across all the species is shown (error bars are one standard error of the mean), where each
496 percentage is the number of SACs in the region that contain the species, expressed as a percentage of
497 the total number of SACs in the region. Annex II (A) and Annex IV (B) bat species.

498

499

500 Appendix A: List of bibliographic references used to complete the database of bat distributions and known
501 occurrences in Special Areas of Conservation in Spain.
502

503 Alberdi, A., Aihartza, J.R., Albero, J.C., Aizpurua, O., López-Baucells, A., Freixas, L., Puig, X., Flaquer, C.,
504 Garín, I., 2012. First records of the parti-coloured bat *Vespertilio murinus* (Chiroptera: Vespertilionidae)
505 in the Pyrenees. *Mammalia* 76, 109-111.

506 Alcalde, J.T., Trujillo, D., Artázcoz, A., Agirre-Mendi, P.T., 2008. Distribución y estado de conservación de los
507 quirópteros en Aragón. *Graellsia* 64, 3-16.

508 Alcalde, J.T., 2009. *Myotis alcathoe* Helversen & Heller, 2001 y *Pipistrellus pygmaeus* (Leach, 1825), nuevas
509 especies de quirópteros para Navarra. *Munibe* 57, 225-236.

510 Duarte, J., Farfán, M.A., 2009. Nuevos datos sobre factores de riesgo y presencia del murciélago rabudo *Tadarida*
511 *teniotis* (Rafinesque, 1814) en Andalucía. *Galemys* 21, 76-77.

512 Encabo, I., Barba, E., Belda, E.J., Monrós, J.S., 2007. Área de campeo de quirópteros en el término municipal de
513 Carcaixent (Valencia): Nuevas citas para el Atlas de los Mamíferos Terrestres. *Galemys* 19, 37-44.

514 Flaquer, C., Puig, X., Fàbregas, E., Guixé, D., Torre, I., Ràfols, R.G., Páramo, F., Camprodon, J., Cumplido, J.M.,
515 Ruíz-Jarillo, R., Baucells, A.L., Freixas, L., Arrizabalaga, A., 2010. Revisión y aportación de datos sobre
516 quirópteros de Catalunya: Propuesta de Lista Roja. *Galemys* 22, 29-61.

517 García, D., Arbona, P., 2009. Presencia del murciélago ratonero pardo *Myotis emarginatus* (Geoffroy, 1806)
518 (Chiroptera: Vespertilionidae) en Mallorca (Islas Baleares). *Endins* 33, 121-124.

519 Hermida, R.J., Lamas, F.J., Graña, D.A., Rial, S., Cerqueira, F., Arzúa, M., Seage, R., 2012. Contribución al
520 conocimiento de la distribución de los Murciélagos (O. Chiroptera) en Galicia. *Galemys* 24, 13-23.

521 Lisón, F., Aledo, E., Calvo, J.F., 2011. Los murciélagos (Mammalia: Chiroptera) de la Región de Murcia (SE
522 España): distribución y estado de conservación. *An. Biol.* 33, 79-92.

523 Lisón, F., Yelo, N.D., Haz, Á., Calvo, J.F., 2010. Contribución al conocimiento de la distribución de la fauna
524 quiropterológica de la Región de Murcia. *Galemys* 22, 11-28.

525 Lisón, F., 2012. Murciélago de cueva – *Miniopterus schreibersii*. In: Salvador, A., Cassinello, J. (Eds).
526 Enciclopedia Virtual de los Vertebrados Españoles. Museo Nacional de Ciencias Naturales, Madrid.
527 Available from <http://www.vertebradosibericos.org/>

528 Lisón, F., Picazo, J., López, M., 2012. Primera cita del murciélago ratonero patudo *Myotis capaccinii* (Bonaparte,
529 1837) en el Parque Natural Lagunas de Ruidera (Castilla-La Mancha). *Galemys* 24, 65-66.

- 530 Marco, O., Castaño, J., Carpena, F.J., Ortuño, A., Rico, F., Sánchez, I., Lisón, F., 2015. Los murciélagos
531 (Mammalia: Chiroptera) del término municipal de Yecla (Región de Murcia, SE España): distribución y
532 estado de conservación. *An. Biol.* 37, 133-141.
- 533 Picazo, F., Lisón, F., 2013. Fauna quiropterológica del término municipal de Villagordo del Júcar y sus
534 alrededores (Castilla-La Mancha). *An. Biol.* 35, 1-8.
- 535 Serra-Cobo, J., Amengual, B., López-Roig, M., Márquez, J., Bayer, X., Guasch, C., Sánchez, A., Oliver, J.A.,
536 2007. Quinze anys d'estudis quiropterològics a les Illes Balears (1993-2007). *Endins* 31, 125-140.
- 537 de Paz, O., de Lucas, J., Martínez-Alós, S., Pérez-Suárez, G., 2015. Distribución de Quirópteros (Mammalia,
538 Chiroptera) en Madrid y Castilla La Mancha, España Central. *Boletín de la Real Sociedad Española de*
539 *Historia Natural, Sección Biología* 109, 21-34.
- 540 Trujillo, D., García, D., 2009. Primera cita del murciélago de Nathusius *Pipistrellus nathusii* (Keyserling y Blasius,
541 1839) para las Islas Baleares. *Galemys* 21, 39-46.
- 542

543 Appendix B: Official presences of each bat species in Special Areas of Conservation (SACs) in each Autonomous Community (AACC) in Spain. The values shown are
 544 percentages: in each case, the percentage of all the SACs in the AACC that are occupied by the species, according to the Standard Data Forms of the SACs. Abbreviations of
 545 AACCs are explained in Table 1; those of species are explained in the footnote below.

	Sp	And	Ara	Ast	Bal	Can	Cle	Cma	Cat	Ext	Gal	Mad	Mur	Nav	Rio	Val	Vas
Annex II	Bbar	0.6	19.9	18.4	0.9	14.3	9.0	2.7	9.9	4.5	6.9	42.9	0.0	16.7	33.3	6.0	7.7
	Mbec	1.1	0.0	0.0	0.0	14.3	3.3	4.1	1.8	9.1	1.7	0.0	0.0	4.8	33.3	3.6	3.8
	Mbly	7.9	6.4	0.0	0.0	23.8	8.2	4.1	24.3	14.8	1.7	71.4	19.1	7.1	33.3	31.0	1.9
	Mcap	0.6	1.9	0.0	10.8	0.0	0.0	0.0	20.7	0.0	0.0	0.0	27.7	0.0	0.0	23.8	0.0
	Mema	2.8	16.0	10.2	0.9	14.3	9.8	9.5	29.7	13.6	3.4	42.9	8.5	14.3	66.7	11.9	15.4
	Mmyo	11.2	18.6	6.1	18.0	42.9	32.8	10.8	17.1	31.8	69.0	85.7	34.0	14.3	33.3	28.6	15.4
	Msch	14.0	18.6	22.4	17.1	38.1	29.	20.3	41.4	31.8	5.2	71.4	40.4	19.0	50.0	33.3	23.1
	Reur	10.7	19.9	22.4	0.0	47.6	9.8	13.5	38.7	21.6	5.2	85.7	14.9	4.8	83.3	32.1	11.5
	Rfer	12.9	28.8	32.7	7.2	47.6	41.0	21.6	55.9	35.2	98.3	100.0	44.7	23.8	66.7	36.9	26.9
	Rhip	7.9	34.0	30.6	5.4	19.0	36.9	14.9	44.1	20.5	100.0	57.1	25.5	19.0	100.0	19.0	23.1
Rmeh	7.9	0.0	0.0	4.5	4.8	0.0	10.8	2.7	31.8	1.7	57.1	10.6	0.0	0.0	16.7	0.0	
Annex IV	Eser	0.0	0.0	0.0	0.9	0.0	0.0	0.0	0.0	1.1	3.4	0.0	0.0	0.0	66.7	0.0	9.6
	Eisa	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	51.1	0.0	0.0	0.0	0.0
	Hsav	0.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	5.7	0.0	0.0	2.1	0.0	66.7	0.0	0.0
	Malc	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.4	0.0	0.0	0.0

Mdau	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	6.8	0.0	0.0	8.5	0.0	66.7	1.2	7.7
Mmys	0.0	12.2	0.0	0.0	0.0	0.0	0.0	0.0	2.3	0.0	0.0	0.0	0.0	33.3	0.0	1.9
Nlas	0.0	9.0	0.0	0.0	0.0	0.0	0.0	0.0	1.1	0.0	0.0	2.1	7.1	33.3	0.0	0.0
Nlei	0.0	11.5	0.0	0.0	0.0	0.0	0.0	0.0	5.7	0.0	0.0	0.0	16.7	50.0	0.0	13.5
Nnoc	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	50.0	0.0	1.9
Paur	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.3	12.1	0.0	0.0	7.1	50.0	1.2	15.4
Paus	1.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	5.7	1.7	42.9	25.5	2.4	66.7	2.4	13.5
Pmac	0.0	3.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.4	0.0	0.0	0.0
Pnat	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.1	0.0	0.0	0.0	0.0
Pkuh	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.3	0.0	14.3	36.2	0.0	50.0	0.0	7.7
Ppip	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3.4	5.2	71.4	57.4	0.0	50.0	0.0	11.5
Ppyg	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.3	0.0	0.0	53.2	0.0	0.0	0.0	0.0
Tten	0.0	0.0	0.0	1.8	0.0	0.0	0.0	0.0	1.1	0.0	14.3	42.6	2.4	66.7	0.0	5.8
Vmur	0.0	5.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

546 Species abbreviations: Bbar: *Barbastella barbastellus*; Mbec: *Myotis bechsteinii*; Mbly: *Myotis blythii*; Mcap: *Myotis capaccinii*; Mema: *Myotis emarginatus*; Mmyo: *Myotis*
547 *myotis*; Msch: *Miniopterus schreibersii*; Reur: *Rhinolophus euryale*; Rfer: *Rhinolophus ferrumequinum*; Rhip: *Rhinolophus hipposideros*; Rmeh: *Rhinolophus mehelyi*; Eser:
548 *Eptesicus serotinus*; Eisa: *Eptesicus isabellinus*; Hsav: *Hypsugo savii*; Malc: *Myotis alcaethoe*; Mdau: *Myotis daubentonii*; Mmys: *Myotis mystacinus*; Nlas: *Nyctalus lasiopterus*;
549 *Nyctalus leisleri*; Nnoc: *Nyctalus noctula*; Paur: *Plecotus auritus*; Paus: *Plecotus austriacus*; Pmac: *Plecotus macrobullaris*; Pnat: *Pipistrellus nathusii*; Pkuh: *Pipistrellus*
550 *kuhlii*; Ppip: *Pipistrellus pipistrellus*; Ppyg: *Pipistrellus pygmaeus*; Tten: *Tadarida teniotis*; Vmur: *Vespertilio murinus*.

551

552 Appendix C: Known presences of each bat species in Special Areas of Conservation (SACs) in each Autonomous Community (AACC) in Spain. The values shown are
 553 percentages: in each case, the percentage of all the SACs in the AACC that are occupied by the species, according to the best available information on bat distributions.
 554 Abbreviations of AACCs are explained in Table 1; those of species are explained in the footnote below.

	Sp	And	Ara	Ast	Bal	Can	Cle	Cma	Cat	Ext	Gal	Mad	Mur	Nav	Rio	Val	Vas
Annex II	Bbar	3.9	26.3	34.7	1.8	38.1	50.0	6.8	42.3	8.0	29.3	42.9	0.0	40.5	50.0	6.0	38.5
	Mbec	6.7	1.3	0.0	0.0	19.0	10.7	10.8	13.5	21.6	10.3	57.1	0.0	4.8	50.0	0.0	15.4
	Mbly	25.8	7.7	6.1	0.0	42.9	26.2	17.6	42.3	33.0	1.7	85.7	31.9	16.7	66.7	22.6	11.5
	Mcap	2.8	3.8	0.0	15.3	0.0	0.0	2.7	39.6	0.0	0.0	0.0	34.0	0.0	0.0	29.8	0.0
	Mema	17.4	29.5	18.4	0.0	33.3	27.9	32.4	58.6	22.7	22.4	71.4	29.8	42.9	66.7	22.6	44.2
	Mmyo	40.4	28.2	18.4	27.9	90.5	70.5	43.2	44.1	62.5	24.1	100.0	36.2	40.5	16.7	45.2	50.0
	Msch	47.8	24.4	53.1	24.3	85.7	64.8	45.9	64.0	62.5	12.1	100.0	57.4	33.3	100.0	58.3	75.0
	Reur	34.8	22.4	42.9	0.0	66.7	39.3	37.8	52.3	39.8	15.5	85.7	17.0	23.8	83.3	50.0	44.2
	Rfer	55.1	44.2	71.4	28.8	95.2	75.4	66.2	82.9	70.5	58.6	100.0	66.0	69.0	100.0	70.2	78.8
	Rhip	37.1	50.0	83.7	45.0	76.2	79.5	55.4	76.6	36.4	77.6	85.7	48.9	71.4	100.0	34.5	76.9
Rmeh	29.8	0.0	2.0	11.7	47.6	6.6	20.3	5.4	52.3	0.0	71.4	21.3	0.0	0.0	26.2	0.0	
Annex IV	Eser	0.0	42.3	51.0	18.9	42.9	68.9	27.0	57.7	51.1	50.0	85.7	0.0	76.2	83.3	29.8	98.1
	Eisa	23.0	0.0	0.0	0.0	0.0	0.0	6.8	0.0	5.7	0.0	0.0	63.8	0.0	0.0	2.4	0.0
	Hsav	12.4	48.7	16.3	28.8	9.5	47.5	36.5	62.2	13.6	10.3	100.0	8.5	45.2	66.7	25.0	19.2
	Malc	0.0	0.0	0.0	0.0	0.0	0.8	0.0	7.2	0.0	3.4	0.0	0.0	14.3	16.7	0.0	0.0

Mdau	18.5	32.7	59.2	0.0	38.1	80.3	41.9	51.4	29.5	41.0	85.7	17.0	78.6	50.0	10.7	76.9
Mmys	1.1	9.6	12.2	0.0	19.0	22.1	6.8	9.0	6.8	10.3	42.9	0.0	23.8	16.7	0.0	30.8
Nlas	9.0	12.2	6.1	0.0	0.0	23.0	10.8	2.7	6.8	5.2	57.1	0.0	14.3	16.7	0.0	0.0
Nlei	11.2	26.9	12.2	34.2	23.8	44.3	27.0	45.0	28.4	25.9	57.1	0.0	76.2	66.7	2.4	86.5
Nnoc	0.6	0.0	2.0	0.0	9.5	6.6	1.4	0.0	2.3	0.0	14.3	0.0	4.8	16.7	0.0	3.8
Paur	0.0	25.6	26.5	0.0	38.1	47.5	12.2	34.2	8.0	37.9	57.1	0.0	31.0	16.7	1.2	57.7
Paus	13.5	40.4	10.2	22.5	19.0	74.6	44.6	55.0	33.0	46.6	85.7	46.8	47.6	83.3	22.6	42.3
Pmacc	0.0	1.9	0.0	0.0	0.0	0.0	0.0	1.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Pnat	0.0	0.0	0.0	1.8	0.0	0.8	50.0	17.1	1.1	0.0	14.3	2.1	14.3	0.0	1.2	15.4
Pkuh	15.2	52.6	4.1	46.8	14.3	52.5	0.0	72.1	43.2	1.7	85.7	48.9	81.0	83.3	26.2	96.2
Ppip	22.5	64.7	73.5	56.8	81.0	95.9	54.1	69.4	97.7	63.8	85.7	59.6	100.0	100.0	48.8	100.0
Ppyg	23.0	10.9	71.4	22.5	81.0	94.3	47.3	72.1	97.7	15.5	85.7	53.2	40.5	100.0	47.6	100.0
Tten	15.2	46.2	55.1	46.8	33.3	72.1	48.6	67.6	88.6	24.1	85.7	51.1	78.6	83.3	39.3	42.3
Vmur	0.0	5.8	0.0	0.0	0.0	0.0	0.0	0.9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

555 Species abbreviations: Bbar: *Barbastella barbastellus*; Mbec: *Myotis bechsteinii*; Mbly: *Myotis blythii*; Mcap: *Myotis capaccinii*; Mema: *Myotis emarginatus*; Mmyo: *Myotis*
556 *myotis*; Msch: *Miniopterus schreibersii*; Reur: *Rhinolophus euryale*; Rfer: *Rhinolophus ferrumequinum*; Rhip: *Rhinolophus hipposideros*; Rmeh: *Rhinolophus mehelyi*; Eser:
557 *Eptesicus serotinus*; Eisa: *Eptesicus isabellinus*; Hsav: *Hypsugo savii*; Malc: *Myotis alcaethoe*; Mdau: *Myotis daubentonii*; Mmys: *Myotis mystacinus*; Nlas: *Nysctalus lasiopterus*;
558 *Nyctalus leisleri*; Nnoc: *Nyctalus noctula*; Paur: *Plecotus auritus*; Paus: *Plecotus austriacus*; Pmac: *Plecotus macrobullaris*; Pnat: *Pipistrellus nathusii*; Pkuh: *Pipistrellus*
559 *kuhlii*; Ppip: *Pipistrellus pipistrellus*; Ppyg: *Pipistrellus pygmaeus*; Tten: *Tadarida teniotis*; Vmur: *Vespertilio murinus*.

560