

RESEARCH

Open Access



Factors influencing COVID-19 health protective behaviours in Zambian university students with symptoms of low mood

Joyce M. Ncheke¹, J. Anitha Menon^{2,3}, E Bethan Davies^{4,5}, Ravi Paul¹, Sydney O. C. Mwaba², John Mudenda⁶, Heather Wharrad⁷, Parisa Toutouchi⁵ and Cris Glazebrook^{4,5*}

Abstract

Background Health protective behaviours are crucial in the prevention of the spread of COVID-19, particularly in university students who typically live and study in large groups. Depression and anxiety are common in students and can impact young people's motivations to follow health advice. The study aims to assess the relationship between mental health and COVID-19 health-protective behaviours in Zambian university students with symptoms of low mood.

Methods The study was a cross-sectional, online survey of Zambian university students. Participants were also invited to take part in a semi-structured interview to explore views about COVID-19 vaccination. Invitation emails were sent explaining the study aims and directed students who self-identified as having low mood in the past two weeks to an online survey. Measures included COVID-19 preventive behaviours, COVID-19-related self-efficacy, and Hospital and Anxiety Depression scale.

Results A total of 620 students ($n=308$ female, $n=306$ male) participated in the study, with a mean participant age of 22.47 ± 3.29 years (range 18–51). Students reported a mean protective behaviour score of 74.09/105 and 74% scored above the threshold for possible anxiety disorder. Three-way ANOVA showed lower COVID-19 protective behaviours in students with possible anxiety disorder ($p=.024$) and those with low self-efficacy ($p<0.001$). Only 168 (27%) said they would accept vaccination against COVID-19, with male students being twice as likely to be willing to accept COVID-19 vaccination ($p<0.001$). Of 50 students interviewed, 30 (60%) expressed fears about the vaccination and 16 (32%) were concerned about a lack of information. Only 8 (16%) participants expressed doubts about effectiveness.

Conclusion Students who self-identify as having symptoms of depression have high levels of anxiety. The results suggest that interventions to reduce anxiety and promote self-efficacy might enhance students' COVID-19 protective behaviours. Qualitative data provided insight into the high rates of vaccine hesitancy in this population.

Keywords Mental health, COVID-19 protective behaviours, Self-efficacy, Gender

*Correspondence:

Cris Glazebrook
cris.glazebrook@nottingham.ac.uk

¹ Department of Psychiatry, School of Medicine, University of Zambia, Lusaka, Zambia

² Department of Psychology, School of Humanities and Social Sciences, University of Zambia, Lusaka, Zambia

³ Department of International Studies, American University of Sharjah, Sharjah, UAE

⁴ Institute of Mental Health, NIHR MindTech MedTech Co-Operative, The University of Nottingham, Nottingham, UK

⁵ Clinical Neurosciences and Mental Health, School of Medicine, The University of Nottingham, Nottingham, UK

⁶ Lusaka Apex Medical University, Lusaka, Zambia

⁷ School of Health Sciences, University of Nottingham, Nottingham, UK



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

The first two cases of SARS-CoV-2 (COVID-19) in Zambia were recorded on 18 March 2020 [1]. Since then, despite a relatively low reported death toll, the secondary effects of the pandemic have stretched social and economic sectors, and rural healthcare systems have been pushed beyond capacity [2]. A systematic review of excess deaths globally suggests that the impact of the COVID-19 pandemic on mortality rates is likely to have been more devastating in sub-Saharan Africa than that indicated by the official statistics for COVID-19 deaths. [3]. For university students, additional disruptions to academic life such as lack of access to Information and Communications Technology (ICT) off-campus prevent participation in e-learning classes, thus causing a widening socio-economic divide in higher education and jeopardising students' academic achievements and future careers [4].

Although seen as disruptive by students, health-protective behaviours such as mask-wearing, social distancing, and hand washing are crucial to preventing the spread of COVID-19: particularly in university settings where young people live and study in large groups. However, COVID-19 restrictions may be challenging for university students at a time when they are forging new social relationships and subject to intense peer pressure. A recent cross-sectional survey of Ethiopian students found that, although 70% of students were willing to wear a mask, the proportion of students with high levels of preparedness was low (19%), with male students being less prepared than female students. Furthermore, only half of the sample scored highly for self-efficacy, the belief in one's ability to implement a particular behaviour, which is a strong determinant of health-protective behaviours [5].

To implement effective health-protective strategies, it is important to understand barriers and facilitators for behaviour change and pandemics can provide an important context for such research. A review of 26 studies of health-protective behaviours associated with previous pandemics, including SARS and swine flu, explored social cognition and demographic factors influencing compliance with recommended behavioural adaptations [6]. Factors that emerged consistently across some studies as increasing rates of health-protective behaviours or intentions to adhere to prevention strategies included female gender, older age, higher educational level, increased perceived susceptibility to the pandemic disease, and higher levels of self-efficacy.

Although less physically vulnerable to COVID-19, arguably the impact of the pandemic on student mental health may be significant, due to disruption of social support and additional educational pressures. A meta-analysis of studies evaluating the mental health of

college students conducted between December 2019 and October 2020 concluded that rates of depression had increased substantially from 21% (95% CI: 16–25%) before March 2020 to 54% (95% CI: 40–67%) between December 2019 and October 2020 [7]. Rates of anxiety had also increased substantially after the start of pandemic restrictions. A longitudinal US study found that rates of moderate to severe depression in students increased from 10% to 31.7% in the four months following the start of the pandemic with women being more affected than men [8]. Students in low-income countries may be particularly affected by the constraints of the pandemic due to limited access to remote learning and greater economic vulnerability. A survey of South African undergraduate students conducted during the pandemic reported high levels of psychological distress and found that higher levels of hopelessness and anxiety were associated with lower levels of health-protective behaviours [9]. This may reflect the strong association between poor mental health and low self-efficacy [10].

The COVID-19 pandemic offers an opportunity to explore the impact of poor mental health on facilitators of healthy behaviour such as self-efficacy, and also directly on health-protective behaviours themselves. These relationships are likely to be particularly significant for less resilient students. Therefore, this study recruited a large sample of students with self-reported low mood during the peak of the COVID-19 outbreak in Zambia. The findings will inform our understanding of the health promotion strategies needed to protect vulnerable students during a pandemic. The study is underpinned by social cognition theories (Theory of Planned Behaviour and the Revised Health Model) which argue that health behaviour is driven by beliefs about outcomes of the behaviour and motivation and feelings of capability regarding the ability to carry out the behaviour. Objectives are to determine rates of health-protective behaviours in students with symptoms of low mood and to explore the association between mental health and self-efficacy and health-protective behaviours in vulnerable students. A further objective is to explore rates of vaccine hesitancy and to understand factors influencing up take of COVID-19 vaccination. We hypothesize that students with low self-efficacy will report fewer health protective behaviours.

Methods

Design

This was a cross-sectional online survey, and the participants were Zambian university students at risk for depression. Data were collected between February–July 2021 as baseline assessment for a longitudinal, mixed-methods feasibility study of an online cognitive

behavioural therapy (CBT) based online intervention to promote emotional and behavioural resilience during the COVID-19 pandemic. Participants were also given the opportunity to take part in a semi-structured interview and the responses to a question about COVID-19 vaccination were analysed as part of this study.

Participants and data collection procedures

Non-probability sampling was used to recruit students at risk of depression from the University of Zambia (UNZA) and Lusaka Apex Medical University (LAMU). The pragmatic target sample size was 1000 students. To be eligible to participate, university students had to be aged ≥ 18 years, in their second year of study or beyond, and self-identify as having experienced at least one of two key symptoms of depression in the previous two weeks [11]. Due to the longitudinal nature of the intervention, students in their final year of study were not eligible to participate.

To avoid potential bias invitation, emails were sent to all registered students meeting the inclusion criteria for the year of study ($n=21,450$) via UNZA Centre for Information Communication Technologies (CICT) and LAMU information technology department. The invitation email explained the study aims and directed students who self-identified as having low mood in the past two weeks (little interest or pleasure in doing things and/or, feeling down, depressed, or hopeless) to an online survey hosted on JISC Online Surveys [12]. Invitation reminders were emailed two and four weeks after the initial invitation. The anonymous survey included a detailed participant information sheet and a tick-box consent form and took approximately 20 min to complete. It was written in English and comprised questions on socio-economic factors, health-protective behaviours, beliefs and attitudes towards COVID-19 and academic performance (not reported in this study). Fifty students who agreed to a recorded telephone/ online interview were contacted by email to arrange a suitable time.

Measures

Participants reported socio-demographic details including age, gender, year of study, university major and parental education. An adapted version of the Material Affluence Scale (MAS) [13] was used to assess family economic status. This 7-item scale was devised specifically for use in low-income countries and probes economic indicators associated with the family home such as car ownership, electricity and type of construction of the home. Total MAS scores range from 0 to 7, with a higher MAS score indicating higher family economic status. Total MAS scores are a valid and easily collected measure of family economic status, correlating with parental

social economic status, parental education and parental occupation. Cronbach's alpha was acceptable (0.62) for the 9-item version which included 2 items with very low item total correlations which were not included in the present study [13].

Preventative behaviours

The Preventive Behaviours Scale (PBS) was developed for the study based on WHO guidelines [14] and adapted from a questionnaire developed for the influenza H1N1 pandemic [15]. The 16 items rated the frequency of recommended COVID-19 preventive behaviours in the past seven days such as hand washing, isolation, vaccination and wearing face masks, on a 1 to 7 scale from 'not at all' to 'very much so'. Total scores (excluding the item rating self-isolation if tested positive) ranged from 15–105, with higher scores indicating stronger compliance with COVID-19 prevention behaviours.

Self-efficacy

To assess perceived self-efficacy to protect oneself against COVID-19, a valid two-item scale [16] again adapted from WHO guidelines [14] was used to assess confidence in the ability to protect oneself against COVID-19. The items are measured on a scale from 1 to 7, with total scores ranging from 2 to 14, and higher scores indicating a higher level of self-efficacy and preparedness to carry out COVID-19 protective behaviours. Participants scoring below the median (2 to 10) were allocated to the low self-efficacy group and participants scoring above the median (≥ 11) were allocated to the high self-efficacy group.

Vaccination uptake

A single item rated 'Yes', 'No' or 'Unsure' was used to assess the willingness to take up a potential offer of a COVID-19 vaccination. To create a binary variable, 'No' or 'Unsure' responses were recoded as 0 (not accept), and 'Yes' responses recoded as 1 (accept).

Mental health

The Hospital Anxiety and Depression Scale (HADS) assessed students' depressive and anxiety symptomology [17]. It consists of two seven-item subscales reflecting depressive and anxiety symptomology respectively, with higher scores indicating greater symptomology. A threshold score of ≥ 8 was used on each subscale to classify participants as experiencing possible clinically significant anxiety or depression [18]. The HADS has been used previously with South African students to determine rates of depression and anxiety among medical students and to examine the associations with various socio-demographic variables [19]. One review found that across 747 papers

that used HADS, the mean Cronbach's alpha for the anxiety subscale was $\alpha=0.83$ (range 0.68 to 0.93), and $\alpha=0.82$ for the depression scale (range of 0.67 to 0.90) [18].

The interview proforma included an open question probing views about vaccination (How do you feel about vaccination against COVID-19?).

Ethical approval

Ethical approval was obtained from UNZA Biomedical Research ethics committee (UNZABREC), the National Health Research Authority (NHRA) and permissions were sought from the management of the University of Zambia and Lusaka Apex Medical University. Participants were required to complete an online, tick box informed consent form to participate in the anonymous survey and the follow-up interview.

Data analysis

Data were exported and analysed using SPSS version 27 [20]. Pearson correlations were conducted to explore relationships between PBS scores and demographic and mental health. A three-way Analysis of Variance (ANOVA) was conducted with mental health (depression/no-depression), self-efficacy (high/low), and gender as the independent variables and the total Protective Behaviour Scale score as the dependent variable. Chi-square was used to examine the relationship between vaccine willingness and depression and anxiety groups, and independent t-tests to examine group differences. Participants with missing values for the HADS were excluded but no other adjustments for missing values were required. Content analysis was used to analyse responses to the question about views on the COVID-19 vaccination.

Results

A total of 633 participants completed the online survey. One participant was excluded from the database for not meeting the inclusion criteria, and a further 12 participants (4 females, 8 male) were removed from the data set due to missing data values for the HADS. The remaining 620 participants (308 females, 306 males, 6 unspecified) had a mean age of 22.47 years ($SD=3.29$; range 18–51). Table 1 shows the full demographic information for the sample.

Internal consistency was acceptable for the HADS depression subscale ($\alpha=0.68$) and good for the HADS anxiety subscale ($\alpha=0.81$) and the Protective Behaviours Scale ($\alpha=0.89$).

The two-item self-efficacy scale had only moderate internal consistency ($\alpha=0.51$). The study found that 335 students (54%) scored above the threshold for possible depressive disorder and 460 students (74.2%)

Table 1 Socio-demographic characteristics of the sample

Gender	
Male	306 (49.40%)
Female	308 (49.70%)
Other	1 (0.20%)
Prefer not to say	5 (0.80%)
Age	
18 – 19	39 (6.30%)
20 – 29	511 (82.40%)
30 +	17 (3.00%)
Prefer not to say	53 (8.50%)
Mean age (SD)	22.47 (SD 29.00)
Marital status	
Single	596 (96.10%)
Married	17 (2.70%)
Divorced	1 (0.20%)
Prefer not to say	6 (1.00%)
Year of study	
Second year	256 (41.30%)
Third year	217 (35.00%)
Fourth year	83 (13.40%)
Sixth year	29 (4.70%)
Prefer not to say	35 (5.60%)
Department of study	
Agriculture UNZA	19 (3.10%)
Engineering UNZA	50 (8.10%)
Education UNZA	103 (16.60%)
Humanities and social sciences UNZA	90 (14.50%)
Law UNZA	55 (8.90%)
Mines UNZA	5 (0.80%)
Medicine UNZA	129 (20.80%)
Natural sciences UNZA	64 (10.30%)
Veterinary medicine UNZA	9 (1.50%)
Medicine LAMU	81 (13.10%)
Prefer not to say	15 (2.40%)
Mean Material Affluence Scale score (SD)	9.0 (SD 1.79)

scored above the threshold for possible anxiety disorder. Females had significantly higher scores for depression ($t=2.01$, $df=612$, $p=0.045$) and anxiety ($t=-3.27$, $df=612$, $p<0.001$). Compared to males, a higher proportion of females scored above the cutoff for anxiety ($X^2=11.27$, $df=1$, $p<0.001$) and above the cutoff for depression ($X^2=5.48$, $df=1$, $p=0.019$). There was no difference in self-efficacy between males and females ($p=0.05$) (Table 2).

Factors influencing health-protective behaviours

Students reported a mean protective behaviour score (PBS) of 74.09 (out of 105). Just over a quarter ($n=168$, 27.00%) said they would accept vaccination against

Table 2 Mental health outcomes and self-efficacy by gender

	Male (n = 306)	Female (n = 308)	Total (n = 620)	P value
Mean HADS-D (SD)	7.6 (SD 3.64)	8.26 (3.71)	7.99 (3.67)	0.045
Case depression (%)	150 (49.00%)	180 (58.40%)	335 (54%)	0.019
Mean HADS-A (SD)	10.18 (SD 4.60)	11.36 (SD 4.31)	10.81 (4.52)	<0.001
Case anxiety (%)	208 (68.00%)	246 (79.90%)	460 (74.2%)	<0.001
Mean self-efficacy (SD)	11.00 (2.45)	11.02 (2.32)	11.00 (2.34)	ns

ns not significant

COVID-19, $n=333$ (54.20%) were unsure and $n=113$ (18.40%) would refuse a vaccination if they were offered one. There was no significant difference in total PBS scores between males and females ($p>0.05$) but over a third of males ($n=105$, 34.30%) reported they would accept a vaccination, compared to only a fifth ($n=63$, 20.50%) of female students ($X^2=14.83$, $df=1$, $P<0.001$; OR 2.03, CIs 1.41- 2.92). Thus, males were twice as likely to agree that they would accept a COVID-19 vaccination if they were offered one.

Higher reported total PBS scores were associated with higher self-efficacy ($r=0.24$, $n=620$, $p<0.001$) and lower levels of anxiety ($r=-0.103$, $n=620$, $p=0.01$). Age, family affluence and depression were not significantly related to PBS (all $p>0.05$). Both depression ($r=-0.106$, $p<0.001$) and anxiety ($r=-0.1$, $p=0.013$) were negatively associated with perceptions of self-efficacy for protecting against COVID-19. Participants scoring above the cut-off for possible clinically significant anxiety had lower PBS scores ($t=2.4$, $df=618$, $p=0.017$) and lower self-efficacy scores ($t=2.25$, $df=218$, $p=0.025$). Participants scoring above the cut-off for possible clinically significant depression disorder had lower self-efficacy scores ($t=2.29$, $df=218$, $p=0.02$) but not lower PBS scores (Table 3).

A three-way ANOVA found main effects for anxiety group ($F=5.00$, $df=1,618$, $p=0.024$, $\eta^2=0.008$) and self-efficacy group ($F=13.1$, $df=1,618$, $p<0.001$, $\eta^2=0.021$). Students scoring above the cut-off for possible clinically significant anxiety and students with low-self efficacy had lower self-reported total COVID-19 protective behaviour scores (Table 3). There were no significant interactions between the variables. There was no relationship between willingness to be vaccinated and

either anxiety or depression. Mean scores for individual COVID-19 protective behaviours are shown in Table 4. The anxiety group had lower scores for mask-wearing in class, mask-wearing outside when socially distancing was not possible, and mask-wearing when visiting family and friends. They also had lower scores for keeping social distance in class and keeping social distance outside but higher scores for isolating if they suspected they had COVID-19 symptoms.

Content analysis of views about vaccination

Fifty participants were interviewed (26 males and 24 females). Thirty (60%) participants (15 males and 15 females) expressed fears about the vaccination and its side effects. A common anxiety was that the vaccination programme was part of a plot to reduce the population.

"Some are saying they are injecting chips. I don't know, to control us here in Africa, to kill us" (P 38).

Linked to this were concerns that the vaccination had magnetic properties.

"I just saw on social media, the injection site leaves a magnetic effect. Now the site can switch on a bulb without any connection" (P 30).

These views seemed to be reinforced by the perception that Africa had been sent vaccinations not used in higher income countries.

"Why is it labelled for export only? Is it just for us... the Africans? We need answers to these questions?" (P25).

Table 3 PBS scores and self-efficacy by anxiety caseness

	Possible case of clinically significant anxiety (n = 460)	Non-case of clinically significant anxiety (n = 160)	P value
Mean PBS (SD)	72.95 (19.98)	77.33 (19.65)	0.017
Mean self-efficacy (SD)	10.87 (2.32)	11.36 (2.37)	0.025

Table 4 Health protective behaviours ($n=620$) and differences in mental health status

	Mean behaviour (SD)	Never carried out behaviour (%)	Case anxiety vs Non-anxiety Case
Frequently washed my hands with soap and water	5.43 (1.76)	31 (5.00%)	ns
Avoided touching my eyes, nose and mouth with unwashed hands	4.45 (1.99)	73 (11.80%)	ns
Used disinfectants to clean hands when soap and water were not available	2.104 (5.19)	76 (12.30%)	ns
Avoided mixing with large groups of friends	4.94 (2.10)	72 (11.60%)	ns
Wore a mask during classes	5.27 (2.39)	126 (20.30%)	$t=2.11, p=0.03$
Wore a mask on the bus	5.77 (2.0)	67 (10.80%)	ns
Wore a mask in shops	5.96 (1.83)	49 (7.90%)	ns
Wore a mask outside when I couldn't socially distance	5.30 (2.12)	80 (12.90%)	$t=2.42, p=0.016$
Wore a mask to visit family or friends	4.39 (2.19)	114 (18.40%)	$t=3.23, p=0.001$
Kept my distance with a group of friends outside	4.33 (2.14)	100 (16.10%)	$t=2.91, p=0.004$
Kept my distance with a group of friends inside	3.90 (2.17)	135 (21.80%)	ns
Isolated myself when I thought I might have COVID-19 symptoms	3.99 (2.58)	226 (36.50%)	$t=-2.57, p=0.01$ (Anx > Non-Anx)
Wore a mask outside in a large group	5.53 (2.10)	79 (12.70%)	ns
Kept my distance in shops	5.39 (2.00)	67 (10.800%)	ns
Kept my distance in class	4.25 (2.43)	167 (26.90%)	$t=2.04, p=0.041$
Isolated myself when I had a positive COVID-19 test ^a	4.73 (2.56)	51 (26.40%)	ns

ns not significant

^a $n = 193$ (Not included in PBS total)

Nearly a third (32%) expressed a need for more information about the vaccine (9 males, 7 females).

..people are hearing a lot on social media. If the people giving can say something... we will listen to then, we have no other choice. The people giving vaccinations are not giving out any information, they need to say how the vaccine will help" (P20).

But only 8 (16%) participants voiced doubts about the effectiveness of the vaccination.

Discussion

In this group of students who identified as having at least one symptom of low mood during the COVID-19 pandemic, over half (54.00%) scored above the cut-off for possible depression. There were very high rates of anxiety (74.00%) within this vulnerable sample, with females having higher scores for both anxiety and depression. Levels of self-reported protective behaviours were generally high but a fifth of students would not wear a mask in class and a quarter did not socially distance themselves in class. There was evidence of high levels of vaccine hesitancy with only 27.00% of students reporting they would accept vaccination; with the majority reporting, they were unsure.

Higher levels of depression and anxiety were associated with lower self-efficacy for protecting against COVID-19 but only anxiety was associated with poorer COVID-19

protective behaviours. ANOVA revealed independent main effects for low self-efficacy and possible anxiety disorder on lower PBS scores but no effect of gender and no significant interactions.

A systematic review of 36 studies evaluating the prevalence of anxiety disorders in university students during the COVID-19 pandemic reported rates ranging between 11.00% and 89.00%, with an average prevalence of 41.00% (95% CI=0.34–0.49) [21]. The majority of studies were carried out in Asia, with only one from Africa (Egypt). The results of our study suggest that Zambian students who identify as having symptoms of low mood have relatively high rates of possible anxiety disorder. However, a study of Bangladeshi students also found high rates of anxiety with 87.70% of the 392 students recruited above the threshold for mild to severe anxiety [22].

The high rate of vaccine hesitancy in this student sample (73.00%) is in contrast to studies of student populations in Lebanon (13.00%) [23], Italy (13.90%) [24], the US (46.00%) [25] and Ethiopia (41.20%) [26]. Concerns about safety and lack of awareness of effectiveness appear to be driving the high rates of vaccine hesitancy in Zambia [27, 28]. Although official figures appear to suggest a low death rate in Zambia from COVID-19, addressing vaccine hesitancy is vital as morgue data and excess death figures paint a much bleaker picture [3, 29]. Our qualitative data confirmed high levels of fear associated with the COVID-19 vaccine, even in this educated sample,

and suggested that social media had fueled those fears in the absence of information from reliable sources. Only a small proportion of our sub-sample expressed concerns about vaccine effectiveness suggesting that information should also focus on safety and health care providers' motivation for providing treatment. A recent study with Nigerian victims of insurgency demonstrated that pictorial information about COVID-19 vaccination was associated with greater self-efficacy and stronger intentions to take up vaccination [30].

In our study, female students had greater vaccine hesitancy which has also been observed in other studies with student populations [25]. A systematic review to explore gender differences in COVID-19 vaccination intentions found males were more likely to be intending to accept a COVID-19 vaccine in 35/60 studies (58.00%). Meta-analysis of data from 46 of the included studies found the odds of men accepting vaccination were 1.4 times higher than women. (95% CI 1.28–1.55) [31]. In our study men had twice the acceptance rate compared to women. Our qualitative data failed to illuminate the reasons for gender differences in vaccine hesitancy, suggesting a common pattern of fears and concerns. However, since women have been shown to be less individualistic than men in terms of their decision making and more risk-averse, the same fears and uncertainties may have a differential impact for women when they come to make a judgement about whether to accept COVID-19 vaccination [32, 33].

We found no significant difference in total protective behaviour scores between males and females and overall scores suggested reasonable compliance to recommended protective behaviours. However, some behaviours had relatively high rates of non-compliance with over a fifth of students reporting that they never wore masks in class or socially distanced from friends indoors and over a third reported that they did not socially isolate if they had COVID-19 symptoms. Anxiety was negatively associated with protective behaviours and students scoring above the threshold for possible anxiety disorder had lower rates of total protective behaviours. Students in the high anxiety group were significantly less likely to wear a mask in class, when visiting family or outside when they couldn't socially distance themselves.

They were also less likely to socially distance themselves in class or outside but were more likely to socially isolate themselves if they suspected symptoms. Relatively few studies have explored the impact of student mental health on COVID-19 protective behaviours but some studies have provided support for the relationship between high anxiety and poorer adherence to health advice. For example, a study in Western India reported an inverse correlation between anxiety and health-protective behaviours in nursing students [34]. However, analysis of data from

the US and Australia suggested that anxiety may have functional benefits by mediating the association between information consumption and improved COVID-19 protective behaviours [35]. This supports the role of effective health education to help channel students' anxiety and build self-efficacy.

We found high self-efficacy was the strongest independent predictor of better health-protective behaviours and both depression and anxiety were inversely related to students' perceptions of their ability to manage their COVID-19 risk. A study in China found that students with high self-efficacy used more active coping strategies and had better mental health [36], and a study of pregnant women in China also found self-efficacy to be protective against anxiety and depression during the COVID-19 pandemic [37]. Since self-efficacy has been found to mediate the relationship between knowledge of dengue fever and dengue protective behaviour during a dengue fever outbreak in Malaysia [38], it is possible that promoting self-efficacy through effective education could improve health protective behaviours through its impact on self-efficacy and mental health [39].

In recognition of the need for appropriate, culturally tailored Covid-19 education for students delivered through accessible channels such as mobile technology [40], the research team has co-created with students from Africa and other stakeholders, a freely available, multimedia educational package titled 'Covid-19 Education for African Students (COVEDAS)' to educate and reassure students about how to keep them and their communities safe from Covid-19 [41]. Sections on COVID-19 vaccination have been informed by the qualitative data from this study and aim to address common myths and concerns. Reducing anxiety may help to increase responsiveness to effective and culturally sensitive health education, thus building self-efficacy for health-protective behaviours, with consequent public health benefits.

The strengths of the current study are that it had a large sample of students ($N=620$) from a low-income country and one which is underrepresented in COVID-19 research. An additional strength is the high proportion of males recruited since they are also under-represented in this type of study. We excluded only 12 students with missing values for the HADS and the remaining sample had complete data for health protective behaviours. A limitation of the study is that we cannot estimate the response rate as only students who self-identified as having symptoms of depression were eligible to participate. Furthermore, recruitment was disrupted by COVID-19 lockdowns with many students returning to their family homes, often with limited internet access. The cross-sectional nature of this study limited our ability to identify causal relationships. However, to our knowledge, this is

the first research to explore the potential of mental health to impact COVID-19 prevention behaviours in an African student sample and the first to use a mixed-methods approach to understanding vaccine hesitancy. We demonstrated that participants with poorer mental health perceived themselves as less able to protect themselves from COVID-19 and that more anxious students practised fewer protective health behaviours. Future research should consider replicating this research to a wider African population, including students outside of Zambia to see if the results are replicable to the general African student population. Our findings suggest that interventions are needed to support anxious students during times of health threat.

Conclusion

Students from a low-income country who self-identified as having symptoms of depression had high levels of anxiety during the COVID-19 pandemic. The results of the study suggest that interventions to reduce anxiety and promote self-efficacy might enhance students' COVID-19 protective behaviours. It is possible that promoting self-efficacy through effective education could improve health-protective behaviours through its impact on self-efficacy and mental health, but longitudinal research is needed to explore causal relationships in more detail. Qualitative research suggested that high levels of vaccine hesitancy may reflect the influence of poor and misleading information, but more research is needed to understand gender differences in rates of vaccine hesitancy in this population.

Abbreviations

CICT	Centre for Information Communication Technologies
PBS	Preventive Behaviours Score
WHO	World Health Organisation
UNZA	University of Zambia
LAMU	Lusaka Apex Medical University
UNZABREC	University of Zambia Biomedical Research ethics committee
NHRA	National Health Research Authority
OR	Odds Ratio
CI	Confidence intervals
MAS	Material Affluence Scale
HADS	Hospital Anxiety and Depression Scale

Acknowledgements

The project received support from UK Research Innovation (UKRI) Global Challenges Research Fund through the University of Nottingham to conduct the research. EBD is funded by the National Institute for Health (NIHR) MindTech Medtech Cooperative and the NIHR Nottingham Biomedical Centre. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care. The project also received support from UNZA CICT and UNZA Projects office.

Authors' contributions

CG and JAM conceived the study idea. JMN and EBD participated in recruitment and data collection. CG and JAM led the analysis, wrote the manuscript and prepared the analysis plan, designed the study, critically reviewed the manuscript and finalised the write-up of the manuscript. RP, SM and JMN

oversaw the recruitment of participants. HW contributed to study design. PT participated in writing the introduction and data analysis. All authors read and approved the final manuscript. All authors contributed to the development of a freely available multimedia educational package titled 'Covid-19 Education for African Students (COVEDAS).

Funding

The project was funded by UK Research Innovation (UKRI) Engineering and Physical Sciences Research Council grant number EP/V034529/1) and the Global Challenges Research Fund (GCRF)/Newton Fund, UK.

Availability of data and materials

Data are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

Ethical approval was obtained from the University of Zambia Biomedical Research ethics committee [UNZABREC] under the ethical approval reference number 1296–2020, National Health Research Authority (NHRA) and permission was sought from the management of UNZA and LAMU. Participants were required to complete an online, tick box informed consent form to participate in the anonymous survey. All methods were carried out in accordance with relevant guidelines and regulations.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Received: 9 July 2022 Accepted: 17 January 2023

Published online: 15 February 2023

References

1. Simulundu E, Mupeta F, Chanda-Kapata P, Saasa N, Changula K, Muleya W. First COVID-19 case in Zambia-Comparative Phylogenetic analyses of SARS-COV-2 detected in African countries. *Int J Infect Dis.* 2021;102(1):455–9.
2. Mathew N, Deborah I, Karonga T, Rumbidzai C. The impact of COVID-19 lockdown in a developing country: Narratives of self-employed women in Ndola Zambia. *Health Care Women Int.* 2020;41(11–12):1370–83. <https://doi.org/10.1080/07399332.2020.1823983>.
3. Excess Mortality Collaborators Estimating excess mortality due to the COVID-19 pandemic: a systematic analysis of COVID-19-related mortality, 2020–21 COVID-19 Published Online March 10, 2022, [https://doi.org/10.1016/S0140-6736\(21\)02796-3](https://doi.org/10.1016/S0140-6736(21)02796-3).
4. Kayombo KM, Mwiinga B. Acceptability and challenges of online higher education in the covid-19 era in Zambia Academia. *Int Multidiscip Res J.* 2021;11(2):1628–45. <https://doi.org/10.5958/2249-7137.2021.00568.1>.
5. Tadesse M, Mihretie A. Attitude, preparedness, and perceived self-efficacy in controlling COVID-19 pandemics and associated factors among university students during school reopening. *PLoS ONE.* 2021;16(9):e0255121. <https://doi.org/10.1371/journal.pone.0255121>.
6. Bish A, Michie S. Demographic and attitudinal determinants of protective behaviours during a pandemic: A review. *Br J Health Psychol.* 2010;15(4):797–824. <https://doi.org/10.1348/135910710X485826>.
7. Li Y, Wang A, Wu Y, Han N, Huang H. Impact of the COVID-19 Pandemic on the Mental Health of College Students: a systematic review and meta-analysis. *Front Psychol.* 2021;12:669119. <https://doi.org/10.3389/fpsyg.2021.669119>.
8. Fruehwirth JC, Biswas S, Perreira KM. The Covid-19 pandemic and mental health of first-year college students: Examining the effect of Covid-19 stressors using longitudinal data. *PLoS ONE.* 2021;16(3):e0247999. <https://doi.org/10.1371/journal.pone.0247999>.
9. Padmanabhanunni A, Pretorius S. Behaviour is the Key in a Pandemic: The Direct and Indirect Effects of COVID-19-Related Variables on

- Psychological Wellbeing; *Psychological Reports*;20210(0):1–14. <https://doi.org/10.1177/00332941211025269>.
10. Bavojean MR, Towhidi A, Rahmati A. The Relationship between Mental Health and General Self-Efficacy Beliefs, Coping Strategies and Locus of Control in Male Drug Abusers. *Addict & Health*. 2011;3(3–4):111–8.
 11. Kroenke K, Spitzer R, Williams J, Löwe B. An ultra-brief screening scale for anxiety and depression: the PHQ-4. *Psychosomatics*. 2009;50:613–21.
 12. Jisc. (2021). Online surveys. <https://www.onlinesurveys.ac.uk/>.
 13. Doku D, Koivusilta L, Rimpelä A. Indicators for measuring material affluence of adolescents in health inequality research in developing countries. *Child Ind Res*. 2010;3:243–26014.
 14. World Health Organisation. Monitoring knowledge, risk perceptions, preventive behaviours and trust to inform pandemic outbreak response. 2020. https://www.euro.who.int/__data/assets/pdf_file/0007/436705/COVID-19-survey-tool-and-guidance.pdf.
 15. SteelFisher GK, Blendon RJ, Ward JR, Rapoport R, Kahn EB, Kohl KS. Public response to the 2009 influenza A H1N1 pandemic: a polling study in five countries. *Lancet Infect Dis*. 2012;12(11):845–50. [https://doi.org/10.1016/S1473-3099\(12\)70206-2](https://doi.org/10.1016/S1473-3099(12)70206-2).
 16. Bandura A. Guide for constructing self-efficacy scales. Self-efficacy beliefs of adolescents. 2006;5(1):307–337.
 17. Zigmond AS, Snaith RP. The Hospital Anxiety and Depression Scale. *Acta Psychiatr Scand*. 1983;67(6):361–70. <https://doi.org/10.1111/j.1600-0447.1983.tb09716.x>
 18. Bjelland I, Dahl AA, Haug TT, Neckelmann D. The validity of the Hospital Anxiety and Depression Scale: An updated literature review. *J Psychosom Res*. 2002;52(2):69–77. [https://doi.org/10.1016/S0022-3999\(01\)00296-3](https://doi.org/10.1016/S0022-3999(01)00296-3).
 19. Sweetland AC, Belkin GS, Verdelli H. Measuring depression and anxiety in sub-Saharan Africa. *Depress Anxiety*. 2014;31(3):223–32. <https://doi.org/10.1002/da.22142>.
 20. IBM Corp. (2020). IBM SPSS Statistics for Windows, version 27.0 Armonk, NY: IBM Corp.
 21. Liyanage S, Saqib K, Khan AF, Thobani TR, Tang WC, Chiarot CB. A Prevalence of Anxiety in University Students during the COVID-19 Pandemic: A Systematic Review. *Int J Environ Res Public Health*. 2022;19:62. <https://doi.org/10.3390/ijerph19010062>.
 22. Islam A, Barna SD, Raihan H, Khan NA, Hossain T. Depression and anxiety among university students during the COVID-19 pandemic in Bangladesh: A web-based Cross-sectional Survey. *PLoS ONE*. 2020. <https://doi.org/10.1371/journal.pone.02238162>.
 23. Bou Hamdan M, Singh S, Polavarapu M, Jordan TR, Melhem NM. COVID-19 vaccine hesitancy among university students in Lebanon. *Epidemiol Infect*. 2021;149(e242):1–11. <https://doi.org/10.1017/S0950268821002314>.
 24. Barello S, Nania T, Dellafiore F, Graffigna G, Caruso R. "Vaccine hesitancy" among university students in Italy during the COVID-19 pandemic. *Eur J Epidemiol*. 2020;35(8):781–3. <https://doi.org/10.1007/s10654-020-00670-z>.
 25. Elliott L, Yang K. Vaccine acceptance and hesitancy among college students in Nevada: a state-wide cross-sectional study. *Vaccines*. 2022;10:105. <https://doi.org/10.3390/vaccines10010105>.
 26. Mose A, Haile K, Timerga A. COVID-19 vaccine hesitancy among medical and health science students attending Wolkite University in Ethiopia. *PLoS One*. 2022;17(1):e0263081. <https://doi.org/10.1371/journal.pone.0263081>.
 27. GTCG. See the difference. Misinformation still hindering COVID-19 uptake in Zambia Xinhua News Agency, 01:57, 11-Oct-2021. http://www.news.cn/english/africa/2021-10/10/c_1310236285.htm.
 28. Mudenda S, Hikaambo CN, Daka V, Chileshe M, Mfuno RL, Kampamba M, Kasanga M, Phiri M, Mufwambi W, Banda M, Phiri MN, Mukosha M. Prevalence and factors associated with COVID-19 vaccine acceptance in Zambia: a web-based cross-sectional study. *Pan Afr Med J*. 2022;41:112. <https://doi.org/10.11604/pamj.2022.41.112.31219>.
 29. Kreier F. Morgue Data Hint at COVID's True Toll in Africa. *Nature News focus*. 2022;603:778–9.
 30. Ugwuoke JC, Talabi FO, Adelabu O, Sanusi B, Gever VC, Onuora C. Expanding the boundaries of vaccine discourse: impact of visual illustrations communication intervention on intention towards COVID-19 vaccination among victims of insecurity in Nigeria. *Human Vaccin Immunother*. 2021;17(10):3450–6. <https://doi.org/10.1080/21645515.2021.1886558>.
 31. Zintel S, Flock C, Arbogast AL, Forster A, Wagner V, Sieverding M. Gender differences in the intention to get vaccinated against COVID-19: a systematic review and meta-analysis. *J Public Health*. <https://doi.org/10.1007/s10389-021-01677-w>.
 32. Delaney R, Strough J, Parker AM, de Bruin WB. Variations in Decision-Making Profiles by Age and Gender: A Cluster-Analytic Approach. *Pers Individ Dif*. 2015;1(85):19–24. <https://doi.org/10.1016/j.paid.2015.04.034>.
 33. Harris CR, Jenkins M. Gender Differences in Risk Assessment: Why do Women Take Fewer Risks than Men? *Judgm Decis Mak*. 2006;1(1):48–63.
 34. Parihar A, Kumar A, Gomathi A, Nebhinani N, Rani R. COVID 19-induced anxiety and protective behaviour among nursing students: A survey from western India. *J Family Med Prim Care*. 2021;10:4483–8.
 35. Antwerpen N, Turnbull D & Searston RA. The role of anxiety in mediating the relationship between information consumption and COVID-19 protective behaviours, *Psychology, Health & Medicine*. 2021. <https://doi.org/10.1080/13548506.2021.2008994>.
 36. Zhou C, Dong Yue X, Zhang X, Shangguan F, Zhang XY. Self-efficacy and mental health problems during COVID-19 pandemic: a multiple mediation model based on the Health Belief Model. *Pers Individ Dif*. 2021;179:110893. <https://doi.org/10.1016/j.paid.2021.110893>.
 37. Cui C, Zhai L, Sznajder KK, Wang J, Sun X, Zhang W. Factors among Chinese pregnant women during the COVID-19 pandemic- A smart phone questionnaire survey study. *BMC Psychiatry*. 2021;21:619. <https://doi.org/10.1186/s12888-021-03624-1>.
 38. Isa A, Loke YK, Smith JR, Papageorgiou A, Hunter PR. Mediation Effects of Self-Efficacy Dimensions in the Relationship between Knowledge of Dengue and Dengue Preventive Behaviour with Respect to Control of Dengue Outbreaks: a Structural Equation Model of a Cross-Sectional Survey. *PLoS Negl Trop Dis*. 2013;7(9):e2401. <https://doi.org/10.1371/journal.pntd.0002401>.
 39. Mashudi DN, Ahmad N, Mohd Said S. Level of dengue preventive practices and associated factors in a Malaysian residential area during the COVID-19 pandemic: a cross-sectional study. *PLoS ONE*. 2022;17(4):e0267899. <https://doi.org/10.1371/journal.pone.0267899>.
 40. Ale V. A library-based model for explaining information exchange on Coronavirus disease in Nigeria. *Ianna Jo Interdiscip Stud*. 2021;2(1):1–11.
 41. Ncheka JM, Menon J, Davies B, Paul R, Mwaba SOC, Mudenda J, Wharrad H, Taylor M, Maboshe S, Glazebrook C. COVID-19 Education for African Students (COVEDAS). 2021 <https://www.nottingham.ac.uk/helmopen/rls/practice-learning/public-health/covedas/>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

