## **Tomorrow's doctors conference 2023**

https://www.rsm.ac.uk/events/students/2022-23/str02/

Sat 10 Jun 2023, Royal Society of Medicine.

## Attitudes and current practice in alcohol screening, brief intervention, and referral for treatment among staff working in urgent and emergency settings: a survey.

Holly Blake\*1,2, Mehmet Yildirim1, Vinuja Premakumar3, Lucy Morris4, Philip Miller5, Frank Coffey4.

## **Abstract**

**Background**: The aim of the study was to ascertain the views and experiences of those working in urgent and emergency care (UEC) settings towards screening, brief intervention, and referral to treatment (SBIRT) for alcohol, to inform future practice.

**Objectives**: To explore i) views towards health promotion, ii) views towards SBIRT, iii) experience of SBIRT in practice, iv) facilitators and barriers to delivering SBIRT in UEC settings and v) training needs to support future SBIRT practice.

**Methods**: Open cross-sectional survey, using an online self-administered questionnaire with closed and open-ended responses. Participants were ≥18 years of age, from any occupational group, working in urgent and emergency care (UEC) settings in any country or region.

**Results**: 362 respondents (aged 21-65 years, 87.8% shift workers) from 7 occupational groups including physicians (48.6%), nurses (22.4%) and advanced clinical practitioners (18.5%). Most believed that health promotion is part of their role, and that SBIRT for alcohol prevention is needed and appropriate in UEC settings. SBIRT was seen to be acceptable to patients. 66% currently provide brief alcohol advice, but fewer screen for alcohol problems or make alcohol-related referrals. The most common barriers were lack of knowledge and training on SBIRT, lack of access to high-quality resources, lack of timely referral pathways, and concerns about patient resistance to advice.

**Discussion**: UEC workers are willing to engage in SBIRT for alcohol prevention but there are challenges to implementation in UEC environments and concerns about workload impacts on already-burdened staff, particularly in the context of global workforce shortages. UEC workers advocate for clear guidelines and policies, increased staff capacity and/or dedicated health promotion teams onsite, SBIRT education/training/resources, appropriate physical spaces for SBIRT conversations and improved alcohol referral pathways to better funded services.

**Conclusions**: This is the first study to describe the views, experiences, barriers, and enablers of SBIRT for alcohol prevention, and health promotion more broadly, as reported by UEC workers across a range of occupational roles and settings. Implementation of SBIRT could contribute to improving population health and reducing service demand, but it requires significant and sustained commitment of time and resources for prevention across healthcare organisations.

**Keywords:** Emergency care, emergency department, urgent care, health promotion, health screening, brief intervention, alcohol, prevention.

<sup>&</sup>lt;sup>1</sup>School of Health Sciences, University of Nottingham, Nottingham, UK.

<sup>&</sup>lt;sup>2</sup>NIHR Nottingham Biomedical Research Centre, Nottingham, UK.

<sup>&</sup>lt;sup>3</sup>School of Medicine, University of Nottingham, Nottingham, UK.

<sup>&</sup>lt;sup>4</sup>Emergency Department, Nottingham University Hospitals NHS Trust, Nottingham, UK.

<sup>&</sup>lt;sup>5</sup>East Midlands Academic Health Sciences Network, Nottingham, UK.