

Table S1: Univariable and multivariable models for correlates among smokers of having a discussion with a health professional about stopping smoking

Baseline variables	DV: Discussion with a health professional about quitting (n=482)					
	Univariate model			Multivariable model		
	OR	p	95% CI	OR	p	95% CI
Smoked in a prior pregnancy	0.61	0.010	0.42,0.89			
Heaviness of Smoking Index	0.93	0.278	0.81,1.06			
Urges to smoke	0.99	0.785	0.90,1.08			
Has tried to quit (y/n)	3.69	<0.001	2.51,5.43	3.04	<0.001	2.00,4.61
Number of 24 hour quit attempts	0.99	0.455	0.96,1.02			
Seriously planning to quit	1.80	<0.001	1.53,2.11			
Determination to stop till baby born	1.81	<0.001	1.54,2.13			
Confidence to stop till baby born	1.28	0.001	1.11,1.48			
Confidence to stop alone	1.08	0.275	0.94,1.25			
Confidence to stop with health professional help	1.58	<0.001	1.35,1.85			
Beliefs on harms of smoking to baby	1.95	0.012	1.16,3.28			
Knows others who smoked in pregnancy	1.03	0.916	0.58,1.84			
People important to me think I should stop	2.45	<0.001	1.60,3.74			
Has support to stop	1.70	0.008	1.15,2.52			
Partner smokes	1.17	0.426	0.80,1.72			
General health	1.19	0.504	0.72,1.97			
Depression	1.11	0.658	0.71,1.73			
Perceived Stress Scale	1.02	0.529	0.96,1.09			
Gestation (weeks)	0.98	0.398	0.94,1.03			
Ethnic minority	1.17	0.701	0.52,2.64			
Age	1.02	0.324	0.98,1.05			
Deprivation (Index of Multiple Deprivation; IMD)	0.99	0.191	0.98,1.00			
Interest in support at baseline	6.17	<0.001	4.12,9.24	5.42	<0.001	3.57,8.23