



Gender congruence and body satisfaction in non-binary transgender people: A case control study

Journal:	<i>International Journal of Transgenderism</i>
Manuscript ID	Draft
Manuscript Type:	Original Research
Keywords:	Non-binary, gender congruence, body satisfaction, gender affirming medical treatments, transgender health services

SCHOLARONE™
Manuscripts

Gender congruence and body satisfaction in non-binary transgender people: A case control study

ABSTRACT

Background: Binary transgender people access gender affirming medical interventions to alleviate gender incongruence and increase body satisfaction. Despite the increase in non-binary transgender people, this population are less likely to access transgender health services compared to binary transgender people. No research has yet explored *why* by exploring levels of gender congruence and body satisfaction in non-binary transgender people.

Objective: The aim of this study was to compare levels of gender congruence and body satisfaction in non-binary transgender people to controls (binary transgender people and cisgender (non-trans) people).

Method: In total, 526 people from a community sample in the United Kingdom took part in the study (97 non-binary, 91 binary and 338 cisgender identifying people). Participants were asked to complete an online survey about gender congruence and body satisfaction.

Results: There were differences in gender congruence and body satisfaction between non-binary and binary transgender people. On sex specific parts of the body (i.e., chest, genitalia and secondary sex characteristics), non-binary transgender people reported significantly higher levels of gender congruence and body satisfaction compared to binary transgender people. However, there was no difference in congruence and satisfaction with social gender role between the two transgender groups (non-binary and binary). Cisgender people reported significantly higher levels of gender congruence and body satisfaction compared to transgender people (non-binary and binary).

Conclusion: There are differences in gender congruence and body satisfaction between non-binary and binary transgender people. Non-binary individuals may be less likely to access transgender health services due to experiencing less gender incongruence and more body satisfaction compared to binary transgender people. Transgender health services need to be more inclusive of non-binary transgender people and their support and treatment needs, which may differ from those who identify within the binary gender system.

1
2
3 **KEYWORDS:** Non-binary; gender congruence; body satisfaction; gender affirming medical
4 treatments; transgender health services
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

Introduction

Transgender health research, which is a relatively new field, has demonstrated that many of the people accessing transgender health services experience very low levels of gender congruence (i.e., they do not feel that their body matches their gender identity) and body satisfaction (e.g., Bandini et al., 2013; Becker et al., 2016, 2018; de Vries et al., 2014; van de Grift et al., 2016, 2017; de Vries, Steensma, Doreleijers, & Cohen-Kettenis, 2011; Röder et al., 2018). Body satisfaction has been found to be much lower than what is found within the cisgender population (i.e., people who do not experience incongruence between their sex assigned at birth and gender identity) (Witcomb et al., 2015). However, gender congruence and body satisfaction of treatment-seeking transgender people have been found to increase following gender affirming medical treatments (GAMT), such as cross-sex hormones and genital affirming surgery (e.g., Bandini et al., 2013; Becker et al., 2018; de Vries et al., 2011, 2014; Jones, Haycraft, Murjan, & Arcelus, 2016). Research has shown that GAMT also increases mental well-being (e.g., Bouman et al., 2016, 2017; Dhejne et al., 2016; Davis & Meier, 2015) and quality of life (e.g., Gorin-Lazard et al., 2012; 2013; Nobili et al., 2018) in binary (treatment-seeking) transgender people. In this paper, when referring to *binary* transgender identity, we are describing a spectrum of transgender people, who seek cross-gender identification and a complete change in social gender role status (i.e. from one gender to the other; Richards & Barker, 2013).

As societies' understanding of gender expression and diversity has evolved, people have had the freedom to express their gender and to be able to identify outside and other than the binary gender concept of exclusively male or female. We use 'non-binary' as an adjective and umbrella term for people who self-identify as such and include all other identities than male or female, such as gender neutral, gender fluid, agender, non-gender, or gender queer, for example (Arcelus & Bouman, 2017; Bouman et al., 2017; Richards et al., 2016; Richards, Bouman, & Barker 2018). In the Western world, the number and visibility of people identifying as non-binary and binary transgender has increased considerably over the past few years (Richards et al., 2016, 2018). Kuyper & Wijzen (2014) examined self-reported gender identity and dysphoria in a large Dutch population sample (N.8064, aged 15–70 years old), and found that 4.6% of people assigned male at birth and 3.2% of people assigned female at birth reported an 'ambivalent gender identity' (defined as equal identification with other sex as with sex assigned at birth) and 1.1% of people assigned male at birth and 0.8% of people assigned female at birth reported an 'incongruent gender identity' (defined as stronger

1
2
3 identification with other sex as with sex assigned at birth). Similarly, van Caenegem et al.
4 (2015) reported results based on two population-based surveys, one of 1,832 Flemish persons
5 and one of 2,472 sexual minority individuals in Flanders, Belgium examining the prevalence
6 of 'gender ambivalence' and 'gender incongruence'. In the general population, gender
7 ambivalence was present in 2.2% of male and 1.9% of female participants, whereas gender
8 incongruence was found in 0.7% of men and 0.6% of women. In sexual minority individuals,
9 the prevalence of gender ambivalence and gender incongruence was 1.8% and 0.9% in men
10 and 4.1% and 2.1% in women, respectively (Bouman, de Vries, & T'Sjoen, 2016).

11 Historically, the medical establishment prevented non-binary people from GAMT, because
12 they were not considered good candidates for medical transition as they did not fit the
13 stereotypical binary concept. A lack of specific clinical guidelines and treatment protocols
14 for non-binary transgender people wanting to access clinical services, including GAMT,
15 further compounded the issue. Emphasis on a binary gender concept within treatment
16 guidelines may also deter patients from being open about their non-binary gender identity to
17 enable them to access GAMT and therefore patients may feel forced to present themselves as
18 a binary transgender person.
19
20
21
22
23
24
25
26
27
28
29

30 The amount of non-binary transgender people accessing transgender health services and
31 undergoing GAMT remains low, even though transgender health clinics have recently began
32 to provide more services for people not seeking a binary transition, and tailoring person-
33 centered approaches to gender transition (e.g., Beek et al., 2015; Clark, Veale, Townsend,
34 Frohard-Dourlent & Saewyc, 2018; Doan, 2016; Government Equalities Office, 2018; Jones
35 et al., 2017; Thorne, Witcomb, Nieder, Nixon & Arcelus, 2018). A recent survey from the
36 United Kingdom (UK) found that only 7% of non-binary transgender people accessed
37 transgender health services in comparison to 50% of transgender men and 43% of
38 transgender women (Government Equalities Office, 2018), which is in keeping with findings
39 in other European countries (Motmans & Burgwal, 2018). Additionally, in Canada fewer
40 non-binary transgender youth (13%) have been found to access cross-sex hormone treatment
41 compared to binary transgender youth (52%) (Clark et al., 2018). This study also found that
42 non-binary transgender youth had more difficulties accessing cross-sex hormones compared
43 to binary transgender youth (Clark et al., 2018). Similarly, Taylor, Zalewska, Gates and
44 Millon (2018) found that non-binary transgender patients, who would like to undergo some
45 GAMT were rejected by transgender health services due to their gender identity. The
46 National LGBT survey in the UK found that 76% of non-binary transgender people hide their
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 gender identity from others due to fear of negative evaluation (Government Equalities Office,
4 2018). People with non-binary gender identities have also reported feeling socially invisible
5 as they do not meet gender norms in Western society (Motmans & Burgwal, 2018; Scottish
6 Trans Alliance, 2015; Taylor et al., 2018). Therefore, a larger amount of non-binary
7 identifying transgender people than expected may be accessing clinical services, including
8 GAMT and withholding their non-binary gender identity. Alternatively, non-binary
9 transgender people may not be attending transgender health services as they experience
10 relatively adequate levels of gender congruence and body satisfaction and consequently do
11 not wish to undergo GAMT.
12
13
14
15
16
17
18

19 To date there has been no empirical research that has investigated why fewer non-binary
20 transgender people are accessing transgender health services by exploring levels of gender
21 congruence and body satisfaction in this population. Consequently, there is a lack of
22 knowledge about interventions that non-binary transgender people may wish to undergo to
23 increase gender congruence and body satisfaction. Previously, research with binary
24 transgender people has focused mainly on gender congruence and body satisfaction with sex-
25 specific characteristics (e.g., Jones, Haycraft, Murjan, & Arcelus, 2016, van de Grift et al.,
26 2016, 2017, 2018). However, one study involving binary transgender people found that body
27 features associated with social gender role recognition (e.g., hairstyle, clothes, mannerism)
28 which cannot be altered through GAMT were the strongest predictors of overall body
29 satisfaction (van de Grift et al., 2016). In addition to this, age has been found to have a
30 positive relationship with body satisfaction in both cisgender men and women (e.g., Peat,
31 Peyerl, Ferraro & Butler, 2011; Tiggemann & McCourt, 2013). Consequently, sex-specific
32 characteristics as well as social gender role recognition should be explored to obtain a
33 broader understanding of gender congruence and body satisfaction in non-binary transgender
34 people, whilst controlling for age.
35
36
37
38
39
40
41
42
43
44
45

46 As most transgender health research has recruited those attending clinical services and
47 accessing GAMT, the existing knowledge regarding the treatment wishes and needs of this
48 population are largely unknown and appear to be largely extrapolated from the wishes and
49 needs of binary (treatment-seeking) transgender people. Information regarding non-binary
50 transgender people accessing clinical services is limited to a few recent studies which have
51 found that this population appears to present to such services with higher levels of mental
52 health problems than binary transgender people (e.g., Rimes et al., 2018; Thorne et al., 2018).
53
54
55
56
57
58
59

Objective

In light of this gap within the literature, the aim of the current study was to explore levels of gender congruence and body satisfaction among non-binary transgender people and to compare these levels to controls (binary transgender people and cisgender people). Only transgender people who had *not* undergone GAMT were selected for this study as GAMT has been found to increase gender congruence and body satisfaction (e.g., Bandini et al., 2013; de Vries et al., 2011, 2014). Age was also controlled for as it has been shown to have a positive relationship with body satisfaction in both cisgender men and women (e.g., Peat, Peyerl, Ferraro & Butler, 2011; Tiggemann & McCourt, 2013). It was hypothesised that there would be a difference in gender congruence and body satisfaction between non-binary and binary transgender people. It was also hypothesised that cisgender people would report higher levels of gender congruence and body satisfaction compared to transgender people (non-binary and binary).

Method

Participants and recruitment

Transgender (non-binary and binary) and cisgender participants aged 18 and over were invited to take part through snowball sampling. The invitation to participate was initially distributed to LGBTQ organisations within the UK, via email and social media sites. The content of this advertisement was the same for transgender and cisgender people. All participants were asked to pass on the survey link to potentially interested parties.

Procedures

Participants were asked to read through the information sheet. Individuals who decided to take part were then asked to complete a survey online which took approximately 20 minutes to complete. The study was approved by Loughborough University Research Ethics Committee at Loughborough University, UK.

Measures

Socio-demographics

Participants were asked to provide information about their age, sex assigned at birth and gender identity. For this study, people who selected their gender identity as transgender male or transgender female were classified as binary transgender people. Participants who selected

1
2
3 androgynous, gender neutral, non-binary, pangender, bigender, gender queer, gender fluid or
4 other were classified as non-binary transgender people. Participants were also asked to
5 provide information about cross-sex hormone use and gender affirming surgeries.
6
7

8 9 *Gender congruence and body satisfaction*

10 To assess gender congruence and body satisfaction in transgender people the Transgender
11 Congruence Scale (Kozee, Tylka, & Bauerband, 2012) and Hamburg Body Drawing Scale
12 (Becker et al., 2016) have been employed (e.g., Jones et al., 2016, 2018). In addition to this, a
13 newly developed tool (Gender Congruence and Life Satisfaction Scale; Jones, Bouman,
14 Haycraft, & Arcelus, 2018) was employed. This measure was developed including non-
15 binary and binary transgender people and provides an understanding of gender congruence
16 and body satisfaction simultaneously by exploring specific body features (Jones et al., 2018).
17 The three measures employed in the study are discussed in more detail below.
18
19
20
21
22
23

24 25 *Transgender Congruence Scale (TCS; Kozee, Tylka, & Bauerband, 2012)*

26 This measure has 12 questions which assess gender congruence. Nine items correspond with
27 the 'appearance congruence' subscale (e.g., 'My outward appearance represents my gender
28 identity'; 'I am generally comfortable with how others perceive my gender identity when
29 they look at me') which was used in the current study. Responses are rated on a 5-point scale
30 from 1 = strongly disagree to 5 = strongly agree. A higher score indicates a higher level of
31 'appearance congruence'. This measure has been found to have good reliability ($\alpha=.93$;
32 Kozee et al., 2012) and the appearance congruence subscale had excellent reliability in the
33 current sample ($\alpha=.96$).
34
35
36
37
38
39
40

41 42 *Hamburg Body Drawing Scale (HBDS; originally developed by Appelt & Strauss, 1988;* 43 *revised version by Becker et al., 2016)*

44 This scale assesses body satisfaction and has been validated with the binary transgender
45 population (Becker et al., 2016). There is an item that assesses overall satisfaction with
46 appearance, which was the only item used in the current study. The subscales were not used
47 as they are different for people assigned male and female at birth and therefore subscales are
48 not comparable. Participants are asked to rate their responses on a 5-point Likert scale (1 =
49 very dissatisfied; 5 = very satisfied) and therefore a higher score indicates a higher level of
50 body satisfaction. Becker et al. (2016) found the HBDS subscales to have good reliability in a
51
52
53
54
55
56
57
58
59
60

1
2
3 transgender population ($\alpha=.62-.91$). As it is not possible to conduct reliability analysis with
4 just one item, this was not calculated in the current study.
5

6
7
8 *Gender Congruence and Life Satisfaction Scale* (GCLS; Jones, Bouman, Haycraft, &
9 Arcelus, 2018)

10
11 The GCLS aims to measure improvements in gender congruence, body satisfaction, related
12 psychological well-being, and life satisfaction during a gender transition. In addition to the
13 TCS and HBDS, this tool was chosen as it was specifically developed to be inclusive of non-
14 binary and binary transgender people (Jones et al., 2018). The items in the GCLS are gender
15 neutral and the same subscales can be administered to everyone regardless of sex assigned at
16 birth or gender identity. Participants are asked to rate their responses on a 5-point Likert scale
17 (always=1; never=5). A higher score is associated with a positive outcome (i.e., greater
18 gender congruence, greater body satisfaction, greater gender-related well-being, and greater
19 life satisfaction). Mean scores are also calculated for each of the seven subscales, four of
20 which (relating to body congruence) were used in the current study: genitalia, chest, other
21 secondary sex characteristics, and, social gender role recognition. Within the current sample,
22 the Cronbach's alphas for the subscales used within the current study was very good ($\alpha=.77-$
23 $.95$).
24
25
26
27
28
29
30
31
32

33 *Analysis*

34
35 The data were analysed using SPSS 23 (IBM, 2016). The data were not normally distributed
36 and as there were no non-parametric alternatives, robust parametric tests were selected (Field,
37 2009). To explore differences in gender congruence and body satisfaction between non-
38 binary transgender people and controls (binary transgender people and cisgender people), a
39 series of one-way ANCOVAs were conducted, controlling for age. The sample was divided
40 by sex assigned at birth (i.e., male or female) and these groups were analysed in relation to
41 gender identity (i.e., non-binary transgender, binary transgender or cisgender). This analysis
42 was then followed up with post-hoc tests (Sidak) to determine where any significant
43 difference lay. The significance level was set at $p<0.05$.
44
45
46
47
48
49
50

51 **Results**

52 *Descriptive analysis*

53
54 In total, 833 people were recruited in to the study. Seven people were removed as they
55 provided no information about their gender (sex assigned at birth or gender identity) and a
56
57
58

1
2
3 further 37 people were removed as they identified as cisgender, but reported that their sex
4 assigned at birth was different to their gender identity. For the purpose of this study, only
5 people who were yet to undergo GAMT were included within the analysis. Therefore, a
6 further 263 people were removed. The final sample consisted of 526 people. Of this sample,
7 97 were non-binary transgender people, 91 were binary transgender people and 338 were
8 cisgender people. The distribution of sex assigned at birth is displayed in Table 1.
9
10
11
12
13

14 ***Insert Table 1. about here***
15

16
17 There was a significant effect for sex assigned at birth between non-binary and binary
18 transgender people ($\chi^2 = 5.97$; $p < .015$; Cramer's $V = .18$; $p < .015$). The standardised
19 residuals demonstrated that there were significantly more people assigned male at birth in the
20 binary transgender group ($z = 1.4$) compared to the non-binary transgender group ($z = -1.3$). In
21 addition, there were significantly more people assigned female at birth in the non-binary
22 transgender group ($z = 1.1$) compared to the binary transgender group ($z = -1.1$).
23
24
25
26
27
28

29 The mean age of the whole sample was 35.70 years ($SD = 13.16$). For the mean age of each
30 group see Table 1. There was a significant group effect for age ($F(2, 519) = 3.14$, $p = .044$).
31 Post-hoc tests revealed that cisgender people were significantly older than non-binary
32 identifying transgender people (Mean difference = 3.80; $p = .037$). There was no significant
33 difference in age between cisgender people and binary identifying transgender people (Mean
34 difference = .69; $p = .960$).
35
36
37
38
39

40 *Body congruence in individuals assigned male at birth: comparing cisgender, non-binary and*
41 *binary transgender people*
42

43 A total of 31 non-binary people (assigned male at birth) were compared on measures of body
44 congruence to 45 transgender females and 84 cisgender males. There was a significant
45 difference between these three groups on the appearance congruence subscale from the TCS
46 after age was controlled for (see Table 2). Post-hoc tests revealed that on the TCS appearance
47 congruence subscale, cisgender males scored significantly higher (more congruence)
48 compared to both non-binary (Mean difference = 2.24; $p < .001$) and binary transgender
49 people (transgender females) (Mean difference = .251; $p < .001$). There was no significant
50 difference between non-binary and binary transgender people assigned male at birth (Mean
51
52
53
54
55
56
57
58
59
60

1
2
3 difference = .27; $p=.166$). This indicates that cisgender males experience more gender
4 congruence with their appearance compared to transgender people assigned male at birth
5 (non-binary and binary) and that there is no difference in the levels of appearance congruence
6 between non-binary and binary people assigned male at birth.
7
8

9
10
11 There was also a significant main effect for body satisfaction, as measured by the HBDS,
12 after controlling for age (see Table 2). As expected, cisgender people scored higher (most
13 body satisfaction) compared to non-binary transgender people (Mean difference= -.95;
14 $p<.001$) and binary transgender people (Mean difference = 1.79; $<.001$; See Table 2). Body
15 satisfaction scores on the HBDS were also significantly higher for the non-binary transgender
16 people when compared to the binary transgender people (Mean difference = .84; $p<.001$).
17 This indicates that while body satisfaction is highest in cisgender people, non-binary
18 transgender people have higher levels of body satisfaction (on the HBDS) than binary
19 transgender people assigned male at birth (transgender females), after controlling for age.
20
21
22
23
24
25

26
27 To obtain a more in-depth understanding regarding gender congruence and body satisfaction
28 in these three groups, differences were explored using the GCLS. For people that were
29 assigned male at birth there was a significant difference between non-binary transgender
30 people, binary transgender people (transgender females) and cisgender males on the chest,
31 genitalia, secondary sex characteristics and social gender role recognition subscales of the
32 GCLS (see Table 2). Post-hoc tests showed that cisgender people scored significantly higher
33 on the chest (Mean difference = .73; $p<.001$), genitalia (Mean difference = .60; $p<.001$) and
34 secondary sex characteristics (Mean difference = 2.05; $p<.001$) subscales of the GCLS
35 compared to non-binary transgender people. Cisgender people also scored higher compared
36 to binary transgender people on the chest (Mean difference = 2.26; $p<.001$), genitalia (Mean
37 difference = 1.81; $p<.002$), and secondary sex characteristics (Mean difference = 3.22;
38 $p<.001$) subscales of the GCLS. Non-binary transgender people scored significantly higher
39 on the chest (Mean difference = 1.54; $p<.001$), genitalia (Mean difference = 1.21; $p<.001$)
40 and secondary sex characteristics (Mean difference = 1.15; $p<.001$) subscales of the GCLS
41 compared to binary transgender people (transgender females). These findings indicate that for
42 those assigned male at birth, non-binary transgender people have higher levels of gender
43 congruence and body satisfaction regarding their chest, genitals and secondary sex
44 characteristics compared to binary transgender people (transgender females), but lower than
45 cisgender males after controlling for age.
46
47
48
49
50
51
52
53
54
55
56
57

1
2
3
4 On the social gender role recognition subscale of the GCLS, cisgender males scored
5 significantly higher compared to both non-binary (Mean difference = .84; $p < .001$) and binary
6 transgender people (transgender females) (Mean difference = 1.18; $p < .001$). There was no
7 significant difference in congruence between non-binary and binary transgender people on
8 the social gender role recognition subscale (Mean difference = .34; $p = .343$). This indicates
9 that transgender people (both binary and non-binary) assigned male at birth experience less
10 congruence and satisfaction with their social gender role compared to cisgender people but
11 there is no difference in social gender role congruence between both groups of transgender
12 people after controlling for age.
13
14
15
16
17
18
19
20

21 *Insert Table 2. about here*
22
23

24 *Body congruence in individuals assigned female at birth: comparing cisgender, non-binary*
25 *and binary transgender people*
26

27 In total, 66 non-binary transgender people were compared to 46 binary transgender people
28 (transgender males) and 254 cisgender females. It was found that there was a significant main
29 effect between these groups on the appearance congruence subscale from the TCS (see Table
30 3). Post-hoc tests showed that cisgender people scored significantly higher on the TCS
31 appearance congruence subscale compared to non-binary (Mean difference = 2.24; $p < .001$)
32 and binary transgender people (transgender males) (Mean difference = 2.43; $p < .001$). There
33 was no significant difference between the non-binary and binary transgender people on this
34 subscale (Mean difference = .19; $p = .356$). These findings demonstrate that cisgender females
35 experience more gender congruence with their appearance compared to both non-binary and
36 binary transgender people (transgender males) assigned female at birth after controlling for
37 age.
38
39
40
41
42
43
44
45

46 There was also a significant main effect of body satisfaction, as measured by the HBDS,
47 between the three groups (see Table 3). As expected, cisgender people scored significantly
48 higher compared to non-binary transgender people (Mean difference = .945 $p < .001$; see Table
49 3) and binary transgender people (Mean difference = 1.67; $p < .001$). It was also found that
50 non-binary transgender people scored significantly higher (more body satisfaction) compared
51 to binary transgender people (transgender males; Mean difference = .71; $p < .001$). Findings
52 from the HBDS therefore suggest that, after controlling for age, cisgender people are the most
53
54
55
56
57
58
59
60

1
2
3 satisfied with their body, followed by non-binary transgender people. Binary transgender
4 people (transgender males) are the least satisfied with their bodies.
5
6

7
8 Gender congruence and body satisfaction between these groups was explored in more detail
9 using the GCLS. There was a significant main effect for the chest, genitalia, secondary sex
10 characteristics and social gender role recognition subscales of the GCLS (see Table 3). Post-
11 hoc tests showed that cisgender females scored higher on the chest (Mean difference = 1.67;
12 $p < .001$), genitalia (Mean difference = .73; $p < .001$) and secondary sex characteristics (Mean
13 difference = 1.21; $p < .001$) subscales compared to non-binary transgender people. Cisgender
14 people also scored higher than binary transgender people on the chest (Mean difference = -
15 3.03; $p < .001$), genitalia (Mean difference = 2.02; $p < .001$) and secondary sex characteristics
16 (Mean difference = 2.81; $p < .001$) subscales of the GCLS. Non-binary transgender people
17 scored significantly higher (more congruence) on the chest (Mean difference = 1.36; $p < .001$),
18 genitalia (Mean difference = 1.28; $p < .001$) and secondary sex characteristics (Mean
19 difference = 1.60; $p < .001$) subscales of the GCLS compared to binary transgender people
20 (transgender males). These findings demonstrate that when age is controlled for, non-binary
21 transgender people experience more gender congruence and body satisfaction with their sex
22 characteristics compared to binary transgender people (transgender males). However,
23 cisgender females experience more gender congruence and body satisfaction with sex
24 characteristics when compared to transgender people (non-binary and binary).
25
26
27
28
29
30
31
32
33
34
35
36

37 On the social gender role recognition subscale of the GCLS, cisgender females scored
38 significantly higher (more congruence) compared to both non-binary transgender people
39 (Mean difference = 1.49; $p < .001$) and transgender males (Mean difference = 1.50; $p < .001$).
40 There was no significant difference in congruence between non-binary and binary
41 transgender people on the social gender role recognition subscale (Mean difference = .02;
42 $p = .994$). Cisgender people, as expected, therefore report experiencing more gender
43 congruence and body satisfaction with their social gender role compared to transgender
44 people assigned female at birth (non-binary and binary).
45
46
47
48
49
50

51
52 *Insert Table 3. about here*
53
54
55
56
57
58
59
60

Discussion

This study explored levels of gender congruence and body satisfaction among non-binary transgender people and compared these levels to two control groups, which included binary transgender people yet to undergo GAMT, and cisgender people. As hypothesised, there were differences in gender congruence and body satisfaction between non-binary and binary transgender people. On sex specific parts of the body (i.e., chest, genitalia and secondary sex characteristics), non-binary transgender people reported significantly higher levels of gender congruence and body satisfaction compared to binary transgender people. There was no difference in congruence and satisfaction with social gender role between the two transgender identifying groups (non-binary and binary). As expected, cisgender people reported significantly higher levels of gender congruence and body satisfaction compared to transgender people (non-binary and binary). The findings from this study may indicate that non-binary individuals may be less likely to access transgender health services due to experiencing less gender incongruence and more body satisfaction compared to binary transgender people (e.g., Beek et al., 2015; Clark et al., 2018; Government Equalities Office, 2018; Thorne et al., 2018). However, gender congruence and body satisfaction with the chest, genitalia and secondary sex characteristics is still much lower among non-binary transgender people compared to cisgender people. Consequently, some non-binary transgender people may wish to undergo GAMT. Clinicians working at transgender health services should be mindful of this and create an inclusive and supportive environment to enable transgender people to disclose their gender identity without this having any bearing on their desired gender affirming treatment, whether medical or other (Rider et al., 2018). Recent research in a UK transgender health service showed that clinicians need to adopt an affirmative approach to encourage non-binary transgender patients to articulate their gender identity and treatment requests (Taylor et al., 2018). Transgender health services should also review their treatment pathway and protocols to be inclusive of non-binary transgender people, and refrain from insisting on specific, fixed trajectories (i.e., hormone treatment before chest reconstructive surgery), as there is little evidence to underpin the benefit of these. Further research that explores the long-term effects of various different treatment trajectories (i.e., gender affirming hormone treatment and/or surgery before or without social gender role transition; chest reconstructive surgery before or without testosterone treatment, etcetera) among non-binary transgender people would also facilitate the development of evidenced-based treatment protocols for this population.

1
2
3 In this study both transgender groups (non-binary and binary) reported less gender
4 congruence and body satisfaction with their social gender role compared to cisgender people.
5 There was no significant difference in congruence and satisfaction with social gender role
6 between non-binary and binary transgender people. Non-binary transgender people have
7 reported feeling socially invisible in social settings in particular, and society in general,
8 which tend to adhere to and advocate a binary gender system, including a binaried language
9 system (see for instance, Bradford et al., 2018; Nicholas, 2018). This tends to leave non-
10 binary transgender people not feeling accepted or validated by society (Saltzburg & Davis,
11 2010). Without a societal structure and a language reflecting the existence and experiences of
12 non-binary identifying people underpinned by legislation and supported by one's country or
13 State non-binary people remain ignored and invisible. Invisibility and victimisation in non-
14 binary transgender youth has been found to have an association with poor mental well-being
15 (Rimes, Goodship, Ussher, Baker & West, 2018; Thorne et al., 2018). This is likely to further
16 contribute to low levels of congruence and satisfaction with social gender role – i.e., if non-
17 binary people feel that their social gender role is not recognised and widely accepted by those
18 around them. In contrast, binary transgender people have been found to be most dissatisfied
19 with unwanted body features that are difficult to hide in everyday social situations such as the
20 jaw, facial hair, hands and hips (van de Grift et al., 2016). This may increase anxiety and
21 distress relating to “passing” as their gender identity (i.e., male or female) (McGuire, Doty,
22 Catalpa & Ola, 2016). Therefore, it appears that non-binary and binary transgender people
23 experience (in)congruence and (dis)satisfaction with their social gender role in different
24 ways. This may explain why no significant difference in gender congruence and body
25 satisfaction with social gender role was found between the transgender groups in this study.
26 Societal awareness of transgender people, especially non-binary transgender people, should
27 continue to be increased. This can be achieved through mass media awareness campaigns on
28 television, by providing educational resources to schools, universities and workplaces, and in
29 many other ways (e.g., Nicholas, 2018).

30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48 The findings of this study are strengthened by the fact that participants were recruited from
49 the community as opposed to from a transgender health service. The number of non-binary
50 and binary transgender people recruited within this study were similar in size, which is
51 another strength of this study. In contrast, research conducted in transgender health services
52 has typically recruited much smaller samples of non-binary transgender people (Thorne et al.,
53 2018; Taylor et al., 2018). Participants in the current study may have felt empowered to be
54
55
56
57
58

1
2
3 open and honest about their gender identity rather than feeling a need to withhold their non-
4 binary identity to satisfy clinical expectations and medical gatekeeping in order to obtain
5 GAMT. Much of the current literature in transgender healthcare is often criticised for social
6 desirability bias. When recruiting from clinical transgender health services, participants may
7 over-report their distress and dissatisfaction and follow a specific binary transgender
8 narrative to access GAMT in a timely manner. The current community study supports
9 previous clinical literature that has found transgender people to report less body satisfaction
10 compared to cisgender people (e.g., Witcomb et al., 2015).
11
12
13
14
15

16
17 It is noteworthy that 'non-binary' is a very broad term which captures a wide variety of
18 different gender identities that fall between and outside the gender binary (Richards et al.,
19 2016; 2017; 2018). This study therefore failed to capture nuances between people with
20 different gender identities who fall under the umbrella term of 'non-binary'. Future research
21 therefore may wish to refine this group further to explore differences in specific gender
22 identities (e.g., gender fluid, gender queer, gender neutral). The findings of the study can also
23 only be generalised to transgender people within the UK. Countries differ in their tolerance of
24 transgender people which will affect how comfortable people feel identifying as transgender
25 (both binary and non-binary) (e.g., Ahmadzad-Asl et al., 2010; Dhejne, Öberg, Arver, &
26 Landén, 2014). Future research could consider exploring cross-cultural differences in non-
27 binary transgender individuals. The current study was also cross-sectional and future research
28 would benefit from a longitudinal research design so that potential changes in gender
29 congruence and body satisfaction in non-binary and binary transgender people can be
30 followed overtime.
31
32
33
34
35
36
37
38
39
40
41

42 **Conclusion**

43 Transgender health research has typically neglected the inclusion of non-binary transgender
44 people. This is an important omission as the current study has shown that there are nuances in
45 gender congruence and body satisfaction between non-binary and binary transgender people.
46 Consequently, the GAMT that non-binary transgender people wish to access in order to
47 increase their gender congruence and body satisfaction may be different from that desired by
48 binary transgender people. The implications of this research are that transgender health
49 services need to be more inclusive of non-binary transgender people and their treatment
50 needs and adjust their treatment guidelines accordingly. Future research should explore
51 gender congruence and body satisfaction longitudinally in non-binary transgender people.
52
53
54
55
56
57
58
59
60

References

- Ahmadzad-Asl, M., Jalali, A. H., Alavi, K., Naserbakht, M., Taban, M., Mohseninia-Omrani, K., & Eftekhar, M. (2010). The epidemiology of transsexualism in Iran. *Journal of Gay & Lesbian Mental Health*, 15(1), 83-93.
- Appelt, H., & Strauss, B. (1988). *Psychoendokrinologische Gynäkologie*. Stuttgart: Enke.
- Arcelus, J., & Bouman, W.P. (2017). *Language and Terminology*. In: W.P. Bouman & J. Arcelus (Eds.). *The Transgender Handbook – A Guide for Transgender People, their Families and Professionals* (pp. 1-12). New York: Nova Publishers.
- Bandini, E., Fisher, A. D., Castellini, G., Lo Sauro, C., Lelli, L., Meriggiola, M. C., ... Ricca, V. (2013). Gender identity disorder and eating disorders: Similarities and differences in terms of body uneasiness. *Journal of Sexual Medicine*, 10(4), 1012–1023.
- Becker, I., Nieder, T. O., Cerwenka, S., Briken, P., Kreukels, B. P., Cohen-Kettenis, P. T., ... Richter-Appelt, H. (2016). Body image in young gender dysphoric adults: A European multi-center study. *Archives of Sexual Behavior*, 45(3), 559-574.
- Becker, I., Auer, M., Barkmann, C., Fuss, J., Möller, B., Nieder, T. O., ... & Richter-Appelt, H. (2018). A Cross-Sectional Multicenter Study of Multidimensional Body Image in Adolescents and Adults with Gender Dysphoria Before and After Transition-Related Medical Interventions. *Archives of Sexual Behavior*. Advance online publication. DOI: 10.1007/s10508-018-1278-4
- Beek, T.F., Kreukels, B.P.C., Cohen-Kettenis, P.T., Steensma, T.D. (2015). Partial gender request and underlying motives of applicants for gender affirming interventions. *Journal of Sexual Medicine*, 12(11), 2201-2205.
- Bouman, W.P., Claes, L., Brewin, N., Crawford, J.R., Millet, N., Fernandez-Aranda, F., & Arcelus, J. (2017). Transgender and anxiety: A comparative study between transgender people and the general population. *International Journal of Transgenderism*, 18(1), 16-26.
- Bouman, W.P., Claes L, Marshall E, Pinner GT, Longworth J, Maddox V, Witcomb G, Jimenez-Murcia S, Fernandez-Aranda F, & Arcelus J. (2016). Socio-demographic variables, Clinical Features and the Role of Pre-assessment Cross-Sex Hormones in older trans people. *Journal of Sexual Medicine*, 13(4), 711-719.
- Bouman, W.P., de Vries, A.L.C, T'Sjoen, G. (2016). Gender Dysphoria and Gender Incongruence: An evolving inter-disciplinary field. *International Review of Psychiatry*, 28(1), 1-4.

- 1
2
3 Bouman, W.P., Suess Schwend, A., Motmans, J., Smiley, A., Safer, J.D., Deutch, M.B.,
4 Adams N.J., & Winter S. (2017). Language and Transgender Health. *International Journal*
5 *of Transgenderism*, 18(1), 1-6.
6
7
8 Bradford, N.J., Nicole Rider, G., Catalpa, J.M., Morrow, Q.J., Berg, D.R., Spencer, K.G.,
9 & McGuire, J.K. (2018). Creating gender: A thematic analysis of genderqueer narratives.
10 *International Journal of Transgenderism*. Advance online publication.
11 doi.org/10.1080/15532739.2018.1474516
12
13
14 Clark, B. A., Veale, J. F., Townsend, M., Frohard-Dourlent, H., & Saewyc, E. (2018). Non-
15 binary youth: Access to gender-affirming primary health care. *International Journal of*
16 *Transgenderism*, 19(2), 158-169.
17
18
19 Claire, C., & Alderson, K. (2013). Living Outside the Gender Binary: A Phenomenological
20 Exploration into the Lived Experience of Female Masculinity. *Canadian Journal of*
21 *Counselling and Psychotherapy*, 47(1), 49–70.
22
23
24 Davis, S. A., & Colton Meier, S. (2014). Effects of testosterone treatment and chest
25 reconstruction surgery on mental health and sexuality in female-to-male transgender
26 people. *International Journal of Sexual Health*, 26(2), 113-128.
27
28
29 de Vries, A. L. C., McGuire, J. K., Steensma, T. D., Eva, C. F., Doreleijers, T. A. H., &
30 Cohen-Kettenis, P. T. (2014). Young adult psychological outcome after puberty
31 suppression and gender reassignment. *Paediatrics*, 134(4), 1–9.
32
33
34 de Vries, A. L. C., Steensma, T. D., Doreleijers, T.A. H., & Cohen-Kettenis, P. T. (2011).
35 Puberty suppression in adolescents with gender identity disorder: A prospective follow-up
36 study. *Journal of Sexual Medicine*, 8(8), 2276–2283.
37
38
39 Dhejne, C., Van Vlerken, R., Heylens, G., & Arcelus, J. (2016). Mental health and gender
40 dysphoria: A review of the literature. *International Review of Psychiatry*, 28(1), 44-57.
41
42 Dhejne, C., Öberg, K., Arver, S., & Landén, M. (2014). An analysis of all applications for sex
43 reassignment surgery in Sweden, 1960–2010: prevalence, incidence, and regrets. *Archives*
44 *of Sexual Behavior*, 43(8), 1535-1545.
45
46
47 Doan, P. L. (2016). To Count or Not to Count: Queering Measurement and the Transgender
48 Community. *WSQ: Women's Studies Quarterly*, 44, 89–110.
49 https://doi.org/10.1353?wsq.2016.0037
50
51
52 Dooley, S., Dawson, M., Zak, L., Ng, C., Effron, L. & Keneally, M. (2015). "Bruce Jenner:
53 'I'm a Woman'". 20/20/ABC News. Retrieved from:
54 <http://abcnews.go.com/Entertainment/bruce-jenner-im-woman/story?id=30570350>
55
56
57
58
59
60

- 1
2
3 Field, A. (2009). *Discovering statistics using IBM SPSS statistics*. Third edition. Thousand
4 Oaks, CA: Sage.
5
6 Government Equalities Office. (2018). National LGBT Survey: Research report. Retrieved
7 from:
8 [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_d](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/722314/GEO-LGBT-Survey-Report.pdf)
9 [ata/file/722314/GEO-LGBT-Survey-Report.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/722314/GEO-LGBT-Survey-Report.pdf)
10
11 Gorin-Lazard, A., Baumstarck, K., Boyer, L., Maquigneau, A., Gebleux, S., Penochet,
12 J.C., ...Bonierbale, M. (2012). Is hormonal therapy associated with better quality of life in
13 transsexuals? A cross sectional study. *Journal of Sexual Medicine*, 9(2), 531-541.
14
15 Gorin-Lazard, A., Baumstarck, K., Boyer, L., Maquigneau, A., Penochet, J. C., Pringuey,
16 D., ... Auquier, P. (2013). Hormonal therapy is associated with better self-esteem, mood,
17 and quality of life in transsexuals. *Journal of Nervous and Mental Disease*, 201(11), 996-
18 1000.
19
20 IBM Corporation (2016). *IBM SPSS Statistics for Windows*. Version 23.0. Armonk, NY: IBM
21 Corporation.
22 Jones, B. A., Bouman, W. P., Haycraft, E., & Arcelus, J. (2018).
23 Development and validation of a new scale to measure outcomes from transgender health
24 services: the Gender Congruence and Life Satisfaction Scale (GCLS). *International*
25 *Journal of Transgenderism*. Advance online publication.
26 doi.org/10.1080/15532739.2018.1453425
27
28 Jones, B.A., Brewin, N., Richards, C., Van Eijk, M., Stephenson-Allen, A. & Arcelus, J.
29 (2017). Investigating the outcome of the initial assessment at a national transgender health
30 service: Time to review the process? *International Journal of Transgenderism*, 18(4), 427-
31 432.
32
33 Jones, B.A., Haycraft, E., Brewin, N., Bouman, W.P., Arcelus, J. (2018). The levels and
34 predictors of physical activity engagement within the treatment seeking transgender
35 population: A matched control study. *Journal of Physical Activity & Health*, 15(2), 99-
36 107.
37
38 Jones, B. A., Haycraft, E., Murjan, S., & Arcelus, J. (2016). Body dissatisfaction and
39 disordered eating in trans people: A systematic review of the literature. *International*
40 *Review of Psychiatry*, 28(1), 81-94.
41
42 Kozee, H. B., Tylka, T. L., & Bauerband, L. A. (2012). Measuring transgender individuals'
43 comfort with gender identity and appearance: Development and Validation of the
44 Transgender Congruence Scale. *Psychology of Women Quarterly*, 36(2), 179–196.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 Kuyper, L. & Wijzen, C. (2014). Gender identities and gender dysphoria in the Netherlands.
4 *Archives of Sexual Behavior*, 43(2), 377–85.
5
- 6 McGuire, J. K., Doty, J. L., Catalpa, J. M., & Ola, C. (2016). Body image in transgender
7 young people: Findings from a qualitative, community based study. *Body Image*, 18, 96-
8 107.
9
- 10 Motmans, J., & Burgwal, A. (2018). Health disparities between binary and non-binary trans
11 respondents: the role of gender identity and socio-economic background characteristics.
12 *International Journal of Transgenderism*. Advance online publication.
13
14 Nicholas, L., (2018). Queer ethics and fostering positive mindsets toward non-binary gender,
15 genderqueer, and gender ambiguity. *International Journal of Transgenderism*. Advance
16 online publication. doi.org/10.1080/15532739.2018.1505576
17
18
19
20
21 Nicole Rider, G.N., Vencill, J.A., Berg, D., Becker-Warner, R., Candelario-Pérez, L., &
22 Spencer, K.G. (2018). The gender affirmative lifespan approach (GALA): A model for
23 competent clinical care with nonbinary clients. *International Journal of Transgenderism*.
24 Advance online publication. doi.org/10.1080/15532739.2018.1485069
25
26
27 Nobili, A., Glazebrook, C. & Arcelus, J. (2018). Quality of life of transgender adults; A
28 systematic review and meta-analysis. *Reviews in Endocrine and Metabolic Disorders*.
29 Advance online publication.
30
31
- 32 Peat, C. M., Peyerl, N. L., Ferraro, F. R., & Butler, M. (2011). Age and body image in
33 Caucasian men. *Psychology of Men & Masculinity*, 12(2), 195-200.
34
- 35 Rankin, S., & Beemyn, G. (2012). Beyond a binary: The lives of gender-nonconforming
36 youth. *About Campus*, 17(4), 2-10.
37
- 38 Richards, C. (2017). In Bouman, W.P. & Arcelus, J. (Eds.). *The Transgender Handbook: A*
39 *Guide for Transgender People, Their Families, and Professionals*. New York, NY: Nova
40 Publishers.
41
42
- 43 Richards, C., & Barker, M-J. (2013). *Sexuality and gender for mental health professionals: A*
44 *practical guide*. London: Sage.
45
- 46 Richards, C., Bouman, W. P., & Barker, M. J. (2018). *Genderqueer and non-binary genders*.
47 London: Palgrave MacMillan..
48
49
- 50 Richards, C., Bouman, W. P., Seal, L., Barker, M. J., Nieder, T.O., & T'Sjoen, G. (2016).
51 Non-binary or genderqueer genders. *International Review of Psychiatry*, 28(1), 95–102.
52
- 53 Rimes, K. A., Goodship, N., Ussher, G., Baker, D., & West, E. (2017). Non-binary and
54 binary transgender youth: Comparison of mental health, self-harm, suicidality, substance
55
56
57
58
59

- 1
2
3 use and victimization experiences. *International Journal of Transgenderism*. Advance
4 online publication. doi: 10.1080/15532739.2017.1370627
5
6 Röder, M., Barkmann, C., Richter-Appelt, H., Schulte-Markwort, M., Ravens-Sieberer, U. &
7
8 Becker, I. (2018) Health-related quality of life in transgender adolescents: Associations
9
10 with body image and emotional and behavioral problems. *International Journal of*
11
12 *Transgenderism*, 19(1), 78-91. doi: 10.1080/15532739.2018.1425649
13
14 Saltzburg, S., & Davis, T. (2010). Co-authoring Genderqueer Youth Identities: Discursive
15
16 tellings and Retellings. *Journal of Ethnic and Cultural Diversity in Social Work*, 19(2),
17
18 87–108.
19
20 Scottish Trans Alliance (2015). Non-binary people's experiences in the UK. Edinburgh:
21
22 Author.
23
24 Thorne, N., Witcomb, G. L., Nieder, T.O., Nixon, E., & Arcelus, J. (2018). A comparison of
25
26 mental health symptomatology and levels of social support in young treatment seeking
27
28 transgender individuals who identify as binary and non-binary. *International Journal of*
29
30 *Transgenderism*. Advance online publication. doi: 10.1080/15532739.2018.1452660
31
32 Tiggemann, M., & McCourt, A. (2013). Body appreciation in adult women: Relationships
33
34 with age and body satisfaction. *Body image*, 10(4), 624-627.
35
36 van Caenegem, E., Wierckx, K., Elaut, E., Buysse, A., Dewaele, A., Van Nieuwerburgh,
37
38 F., De Cuypere, G., & T'Sjoen, G. (2015). Prevalence of gender nonconformity in
39
40 Flanders, Belgium. *Archives of Sexual Behavior*, 44, 1281–1287.
41
42 van de Grift, T. C., Cohen-Kettenis, P. T., Elaut, E., De Cuypere, G., Richter-Appelt, H.,
43
44 Haraldsen, I. R., & Kreukels, B. P. C. (2016). A network analysis of body satisfaction of
45
46 people with gender dysphoria. *Body Image*, 17, 184-190.
47
48 van de Grift, T. C., Elaut, E., Cerwenka, S. C., Cohen-Kettenis, P. T., De Cuypere, G.,
49
50 Richter-Appelt, H., & Kreukels, B. (2017). Effects of medical interventions on gender
51
52 dysphoria and body image: A follow-up study. *Psychosomatic Medicine*, 79(7), 815.
53
54 van de Grift, T.C., Elaut, E., Cerwenka, S.C., Cohen-Kettenis, P.T., & Kreukels,
55
56 B.P.C. (2018). Surgical Satisfaction, Quality of Life, and Their Association After Gender-
57
58 Affirming Surgery: A Follow-up Study. *Journal of Sex & Marital Therapy*, 44(2), 138-
59
60 148. DOI: 10.1080/0092623X.2017.1326190
61
62 Witcomb, G. L., Arcelus, J., & Chen, J. (2013). Can cognitive dissonance methods developed
63
64 in the West for combatting the “thin ideal” help slow the rapidly increasing prevalence of
65
66 eating disorders in non-Western cultures? *Shanghai Archives of Psychiatry*, 25(6), 332–
67
68 340. <http://doi.org/10.3969/j.issn.1002-0829.2013.06.002>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Witcomb, G. L., Bouman, W. P., Brewin, N., Richards, C., Fernandez-Aranda, F., & Arcelus, J. (2015). Body image dissatisfaction and eating-related psychopathology in trans individuals: A matched control study. *European Eating Disorders Review*, 23(4), 287-293.

For Peer Review Only

Table 1. Age and assigned sex for non-binary transgender people, binary transgender people and cisgender people

	Non-binary (n= 97)	Transgender (n=91)	Cisgender (n=338)
Mean (SD) age in years	32.72 (12.17)	35.44 (16.48)	36.32 (12.03)
Sex assigned at birth (n (%))			
Female	66 (67.3)	46 (50.5)	254 (75.1)
Male	31 (31.6)	45 (49.5)	84 (24.9)

Table 2. Mean (SD) and ANCOVA test scores for non-binary and binary transgender people assigned male at birth and cisgender males on measures of body congruence

	Cisgender males (n=84)	Non-binary transgender people (n=31)	Transgender females (n=45)	<i>F</i>
TCS: Appearance congruence	4.46 (.56)	2.19 (.65)	1.90 (.63)	299.41***
HBDS: Body satisfaction	3.58 (.95)	2.65 (1.05)	1.81 (.79)	49.56***
GCLS: Genitalia	4.49 (.32)	3.90 (.64)	2.71 (.88)	120.24***
GCLS: Chest	4.54 (.57)	3.82 (.69)	2.29 (.78)	151.89***
GCLS: Secondary sex characteristics	4.83 (.48)	2.78 (1.01)	1.66 (.62)	348.73***
GCLS: Social gender role recognition	3.94 (1.08)	3.10 (.82)	2.77 (.78)	22.06***

GCLS (Gender Congruence and Life Satisfaction Scale); HBDS (Hamburg Body Drawing Scale); TCS (Transgender Congruence Scale)

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

Table 3. Mean (SD) for non-binary and binary transgender people assigned female at birth and cisgender females on measures of body congruence

	Cisgender females (n=254)	Non-binary transgender people (n=66)	Transgender males (n=46)	F
TCS: Appearance congruence	4.51 (.60)	2.30 (.84)	2.12 (.78)	409.35***
HBDS: Body satisfaction	3.68 (.93)	2.73 (1.31)	2.02 (.93)	58.57***
GCLS: Genitalia	4.54 (.34)	3.80 (.86)	2.52 (.93)	256.83***
GCLS: Chest	4.62 (.47)	2.91 (1.16)	1.54 (.66)	434.39***
GCLS: Secondary sex characteristics	4.78 (.47)	3.53 (1.19)	1.91 (.91)	304.37***
GCLS: Social gender role recognition	4.06 (1.07)	2.62 (.84)	2.63 (.76)	77.57***

GCLS (Gender Congruence and Life Satisfaction Scale); HBDS (Hamburg Body Drawing Scale); TCS (Transgender Congruence Scale)

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$