

Discontinuation of anti-tumour necrosis factor alpha treatment owing to blood test abnormalities, and cost-effectiveness of alternate blood monitoring strategies

Abhishek Abhishek¹, Matthew D. Stevenson,² Georgina Nakafero¹, Matthew J. Grainge,³ Ian Evans,⁴ Oras Alabas,⁴ Tim Card,³ Maarten W. Taal,⁵ Guruprasad P. Aithal,⁶ Christopher P. Fox,⁷ Christian D. Mallen⁸, Danielle A. van der Windt,⁸ Richard D. Riley,⁹ Richard B. Warren^{10,11} and Hywel C. Williams³

¹Academic Rheumatology; ³Lifespan and Population Health; and ⁶Nottingham Digestive Diseases Centre, Translational Medical Sciences, University of Nottingham, Nottingham, UK

²School of Health and Related Research, University of Sheffield, Sheffield, UK

⁴BADBIR, University of Manchester, Manchester, UK

⁵Centre for Kidney Research and Innovation, Translational Medical Sciences, University of Nottingham, Derby, UK

⁷Centre for Cancer Studies, Translational Medical Sciences, School of Medicine, University of Nottingham, Derby, UK

⁸Primary Care Centre Versus Arthritis, Keele University, Keele, UK

⁹Institute of Applied Health Research, College of Medical and Dental Sciences, University of Birmingham, Birmingham, UK

¹⁰Dermatology Centre, Northern Care Alliance NHS Foundation Trust, Manchester, UK

¹¹NIHR Manchester Biomedical Research Centre, Manchester University NHS Foundation Trust, Manchester Academic Health Science Centre, Manchester, UK

Correspondence: Abhishek Abhishek. Email: abhishek.abhishek@nottingham.ac.uk

Linked Article: Paolinell *et al.* *Br J Dermatol* 2024; 190:462–463.

Abstract

Background There is no evidence base to support the use of 6-monthly monitoring blood tests for the early detection of liver, blood and renal toxicity during established anti-tumour necrosis factor alpha (TNF α) treatment.

Objectives To evaluate the incidence and risk factors of anti-TNF α treatment cessation owing to liver, blood and renal side-effects, and to estimate the cost-effectiveness of alternate intervals between monitoring blood tests.

Methods A secondary care-based retrospective cohort study was performed. Data from the British Association of Dermatologists Biologic and Immunomodulators Register (BADBIR) were used. Patients with at least moderate psoriasis prescribed their first anti-TNF α treatment were included. Treatment discontinuation due to a monitoring blood test abnormality was the primary outcome. Patients were followed-up from start of treatment to the outcome of interest, drug discontinuation, death, 31 July 2021 or up to 5 years, whichever came first. The incidence rate (IR) and 95% confidence intervals (CIs) of anti-TNF α discontinuation with monitoring blood test abnormality was calculated. Multivariate Cox regression was used to examine the association between risk factors and outcome. A mathematical model evaluated costs and quality-adjusted life years (QALYs) associated with increasing the length of time between monitoring blood tests during anti-TNF α treatment.

Results The cohort included 8819 participants [3710 (42.1%) female, mean (SD) age 44.76 (13.20) years] that contributed 25 058 person-years (PY) of follow-up and experienced 125 treatment discontinuations owing to a monitoring blood test abnormality at an IR of 5.85 (95% CI 4.91–6.97)/1000 PY. Of these, 64 and 61 discontinuations occurred within the first year and after the first year of treatment start, at IRs of 8.62 (95% CI 6.74–11.01) and 3.44 (95% CI 2.67–4.42)/1000 PY, respectively. Increasing age (in years), diabetes and liver disease were associated with anti-TNF α discontinuation after a monitoring blood test abnormality [adjusted hazard ratios of 1.02 (95% CI 1.01–1.04), 1.68 (95% CI 1.00–2.81) and 2.27 (95% CI 1.26–4.07), respectively]. Assuming a threshold of £20 000 per QALY gained, no monitoring was most cost-effective, but all extended periods were cost-effective vs. 3- or 6-monthly monitoring.

Conclusions Anti-TNF α drugs were uncommonly discontinued owing to abnormal monitoring blood tests after the first year of treatment. Extending the duration between monitoring blood tests was cost-effective. Our results produce evidence for specialist society guidance to reduce patient monitoring burden and healthcare costs.

Accepted: 29 October 2023

© The Author(s) 2023. Published by Oxford University Press on behalf of British Association of Dermatologists. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

What is already known about this topic?

- Hepatic, haematological and renal toxicity due to anti-tumour necrosis factor alpha (TNF α) drugs are uncommon during long-term treatment.
- Despite lack of evidence, all patients established on anti-TNF α treatment in the UK are required to undergo 6-monthly blood tests for the early detection of such toxicity.
- Risk factors for these side-effects are poorly understood and whether it would be cost-effective to reduce the frequency of monitoring blood tests is unknown.

What does this study add?

- It was uncommon for anti-TNF α drugs to be discontinued for hepatic, renal or haematological side-effects; this was even less frequent after 1 year of treatment.
- Age, pre-existing liver disease and diabetes were independent risk factors.
- Assuming a threshold of £20 000/quality-adjusted life years, monitoring 6-monthly, annually and biennially was more cost-effective than monitoring 3-monthly; and monitoring annually and biennially was more cost-effective than monitoring 6-monthly.
- These findings should be used to rationalize monitoring for asymptomatic blood, kidney or liver toxicity during established anti-TNF α treatment.

Anti-tumour necrosis factor alpha (TNF α) drugs are used in the treatment of psoriasis resistant to topical therapies and first-line nonbiologic systemic treatments such as methotrexate.¹ Although effective, they can cause cytopenia and elevated liver enzymes. Concern about these side-effects led the British Association of Dermatology (BAD) to recommend monitoring with full blood count, liver function tests (LFTs), and urea electrolytes and creatinine at least every 6 months during long-term established anti-TNF α treatment.¹ On the contrary, the American Academy of Dermatology does not recommend a fixed monitoring schedule, but leaves the decision on whether to monitor with blood tests – and the frequency of testing – to the patients' dermatologist for all anti-TNF α drugs except for infliximab, for which 6-monthly to annual blood test monitoring is recommended.² This is most likely owing to a higher reported risk of thrombocytopenia, neutropenia and hepatotoxicity with infliximab.^{3,4}

There is no evidence base to support the use of monitoring blood tests for the early detection of toxicity after the first few months of anti-TNF α treatment and the cost-effectiveness of different monitoring strategies has never been evaluated. Previous studies have reported low rates of hepatotoxicity and cytopenia with anti-TNF α drugs in psoriasis, and that the risk of hepatotoxicity was increased in those concomitantly treated with methotrexate.^{5,6} To support evidence-based practice and better use of healthcare resources, we conducted a retrospective cohort study to evaluate the incidence of anti-TNF α discontinuation owing to cytopenia, transaminitis or acute kidney injury; to ascertain the risk factors for this outcome; and to perform a health economic evaluation of alternate intervals between monitoring blood tests.

Materials and methods**Data source**

Data from the BAD Biologic and Immunomodulators Register (BADBIR) were used in this study.⁷ The BAD established

an online pharmacovigilance register (BADBIR) to assess the long-term safety of biologics prescribed for patients with moderate-to-severe psoriasis. Following baseline data acquisition, hospital clinicians recorded changes in therapy, disease activity, changes in treatment (including reasons for stopping treatment) and adverse events. Data on death and malignancy were obtained from linkages with the National Health Service Information Centre system.

Study design

This study protocol was approved by the BADBIR steering committee. It was a retrospective cohort study encompassing the period 1 October 2007 to 31 July 2021.

Study population

Adults (aged ≥ 18 years) with psoriasis with or without arthritis prescribed their first anti-TNF α drugs were eligible. There were no exclusion criteria in terms of comorbidities or concurrent drug prescriptions. In the UK and Republic of Ireland (ROI), anti-TNF α drugs are prescribed from hospital outpatient clinics. The hospital team also organizes monitoring blood tests and acts on any abnormalities.

Follow-up

Patients were followed-up from their first anti-TNF α prescription until the outcome of interest, death, treatment discontinuation, 31 July 2021 or up to 5 years – whichever came first.

Outcome

Anti-TNF α discontinuation owing to a monitoring blood test abnormality was the outcome of interest and was ascertained using the information about reason(s) for stopping treatment specified in BADBIR. Where the reason for stopping treatment was recorded only as an 'adverse event', 'adverse event and inefficacy', 'other' or 'death',

the descriptions of the adverse event in the free text and MedDRA coding were manually reviewed to ascertain outcomes related to side-effects of interest.

Blood test data from the 6-month period before stopping anti-TNF α treatment were used to ascertain whether the treatment was stopped for leukopenia, thrombocytopenia or elevated liver enzymes. The estimated glomerular filtration rate (eGFR) was calculated using the 2021 CKD Epidemiology Collaboration (CKD-EPI) equation, which does not consider race.⁸

Risk factors

Age at registration (years), sex, body mass index (BMI), alcohol intake [not currently drinking alcohol, and self-reporting low (1–14 units weekly), medium (15–21 units weekly) and high alcohol intake (\geq 21 units weekly)], diabetes, high cholesterol and a recorded clinical diagnosis of liver disease before the date of the first anti-TNF α prescription were the risk factors (Table 1). The recorded height and weight at registration were used to calculate BMI. Age, sex, BMI, alcohol intake and diabetes were included as they are associated with drug-induced liver injury.³

Statistical analysis

Mean [standard deviation (SD)] and numbers with percentages were used for descriptive statistics. The incidence rate (IR) and 95% confidence interval (CI) of outcome per 1000 person-years (PY) was calculated for the entire follow-up period, for the first 6 months, for the first 12 months and for the follow-up period after the first 12 months. Multivariable Cox regression was used to examine the association between covariates at baseline and outcome of interest. Missing data for BMI and alcohol were handled by multiple imputation using chained equations. Ten imputations were carried out, and the imputation model included all covariates and the outcome variable. Results from the imputed datasets were combined using Rubin's rule. Data management and analysis were performed in STATA v. 16 (StataCorp, College Station, TX, USA). A *P*-value < 0.05 was considered to be statistically significant.

Health economic methods

Patients were monitored according to each strategy for a period of 5 years. An additional monitoring appointment after the cessation of treatment as a result of an abnormal blood test was assumed. The probability that an abnormal blood test would result in an illness because of the extended monitoring period was estimated by the clinical team members, based on their experience. This involved a remote video-conference followed by asynchronous email discussion. The rates are provided in Table S1 (see [Supporting Information](#)). The clinicians erred toward overestimating the risks rather than underestimating them. The costs and quality-adjusted life year (QALY) losses associated with each condition were estimated following targeted literature reviews (Table S2; see [Supporting Information](#)). A monitoring appointment with blood tests was estimated to cost £24.09; details on this estimation are provided in Appendix S1 (see [Supporting Information](#)). No disutility was assumed for attending a

monitoring appointment. To account for uncertainty in outcome incidence, health economic analyses considered the midpoint and upper and lower 95% CI of the IR from start of treatment, from 1 year after the start of treatment and within the first year of treatment. It was assumed that these risks were distributed exponentially.

For each strategy the costs associated with both monitoring and illness were estimated, as was the loss in QALYs. All values were discounted at a value of 3.5% per annum, as recommended by the National Institute for Health and Care Excellence.⁹ The results were presented in terms of incremental net monetary benefit (iNMB), assuming a cost per QALY threshold of £20 000, compared with monitoring every 3 months. The strategy with the greatest iNMB value was estimated to be the most cost-effective treatment. Sensitivity analysis considered threefold higher risks of illnesses than estimated by the clinicians.

Results

Data for 8819 participants with psoriasis (42.1% female) prescribed their first anti-TNF α agent that contributed 25 058 PY of follow-up were included in this study. Mean (SD) age and BMI were 44.76 (13.20) years and 31.15 (9.44) kg m⁻², respectively. Of the included participants 6744 (76.5%), 1718 (19.5%), 281 (3.2%) and 76 (0.9%) were prescribed adalimumab, etanercept, infliximab and certolizumab or their biosimilars, respectively. Diabetes, high cholesterol, liver disease and renal disease were present in 738 (8.4%), 742 (8.4%), 419 (4.8%) and 131 (1.5%) participants, respectively.

There were 125 anti-TNF α treatment discontinuations for abnormal monitoring blood test results at an IR of 5.85 (95% CI 4.91–6.97)/1000 PY. The IRs of discontinuation of etanercept, infliximab, certolizumab and adalimumab or their biosimilars for abnormal monitoring blood test results were 8.04 (95% CI 5.55–11.65), 18.55 (95% CI 10.53–32.66), 9.79 (95% CI 1.38–69.53) and 4.90 (95% CI 3.96–6.07)/1000 PY, respectively.

Forty-one, 64 and 61 discontinuations owing to abnormal blood test results occurred in the first 6 months, in the first year and after the first year of first prescription during 4111, 7428 and 17 751 PY of follow-up at IRs of 9.97 (95% CI 7.34–13.54), 8.62 (95% CI 6.74–11.01) and 3.44 (95% CI 2.67–4.42)/1000 PY, respectively. The IRs for the discontinuation of etanercept, infliximab, certolizumab and adalimumab or their biosimilars owing to abnormal blood test results after the first year of treatment were 4.63 (95% CI 2.63–8.16), 12.30 (95% CI 5.87–25.81), 23.60 (95% CI 3.32–167.55) and 2.82 (95% CI 2.07–3.83) per 1000 PY.

Blood test results from within the 6 months prior to the stop date were available for 75 patients who stopped treatment. Among them, 51 had an alanine transaminase (ALT) level of > 40 IU L⁻¹ (of these, 29 had an ALT level > 80 IU L⁻¹ and 7 had an ALT level > 200 IU L⁻¹); 5 had a total white blood cell (WBC) count $< 4.0 \times 10^3$ cells mL⁻¹ and none had a WBC count $< 3.0 \times 10^3$ cells mL⁻¹; 8 had a platelet count $< 150 \times 10^3$ cells mL⁻¹ (of these, 4 had a platelet count $< 100 \times 10^3$ cells mL⁻¹ and none had a platelet count $< 50 \times 10^3$ cells mL⁻¹); 2 had an eGFR < 60 mL min⁻¹, with none having an eGFR < 30 mL min⁻¹.

Increasing age, diabetes and liver disease associated with anti-TNF α discontinuation with abnormal blood tests

in the model adjusted for age, sex, BMI, alcohol intake, pre-existing liver disease, diabetes and hypercholesterolaemia [adjusted hazard ratios (HRs) 1.02 (95% CI 1.01–1.04), 1.68 (95% CI 1.00–2.81) and 2.27 (95% CI 1.26–4.07) respectively]. Infliximab was statistically significantly associated with a higher likelihood of stopping treatment owing to abnormal monitoring blood test results than etanercept [unadjusted HR 2.38 (95% CI 1.21–4.67); adjusted HR 2.21 (95% CI 1.12–4.34)]. Patients on adalimumab had a significantly lower risk of stopping treatment owing to abnormal monitoring blood test results compared with etanercept on univariate analysis (unadjusted HR 0.63, 95% CI 0.41–0.97); the data became nonsignificant after adjustment for covariates (adjusted HR 0.66, 95% CI 0.43–1.03).

Health economic results

The iNMB for each strategy is shown in Figure 1. All extended monitoring periods were more cost-effective than 3-monthly monitoring. For all deciles, no monitoring was estimated to be most cost-effective, although annual and biennial monitoring were a considerable improvement on 3-monthly monitoring. Disaggregated results for the base case are shown in Table S3 (see Supporting Information), with combined results in Table S4 (see Supporting Information). Sensitivity analyses using three times the risk of adverse events are

also shown in Figure 1 and showed similar results to the main health economic analysis.

Discussion

In this study of UK and ROI populations we evaluated the incidence and risk factors of clinically significant blood test abnormalities (i.e. ones that required treatment cessation) due to anti-TNF α drugs. We also evaluated the cost-effectiveness of alternate blood test monitoring strategies. Among a cohort of patients with moderate-to-severe psoriasis starting anti-TNF α drugs for the first time, we found that it was uncommon for treatment to be stopped owing to abnormal blood test results, and that this became even less common after the first year of treatment. This is consistent with reports that the median interval between starting anti-TNF α drugs and drug-induced liver injury and neutropenia ranged between 14 to 18 weeks and 12 to 17 weeks, respectively, although late-onset liver enzyme elevations and neutropenia were also reported in these studies.^{10–14} Increasing age and pre-existing liver disease were independent risk factors for stopping anti-TNF α drugs owing to abnormal monitoring blood test results, consistent with previous reports.³ Also consistent with previous reports of a higher risk of drug-induced liver injury with infliximab, we found




Figure 1 Incremental net monetary benefit associated with extended monitoring periods vs. 3-monthly monitoring across different estimated risks (top panel). Sensitivity analyses assuming a triple rate of adverse events (bottom panel). The values on the horizontal axes are the incidence and upper and lower 95% confidence intervals (CIs) of anti-tumour necrosis factor alpha discontinuations owing to abnormal monitoring blood test results during the 5-year follow-up period [5.85 (95% CI 4.91–6.97)/1000 person-years (PY)], during the first year of treatment [8.62 (95% CI 6.74–11.01)/1000 PY] and after the first year of treatment [3.44 (95% CI 2.67–4.42)/1000 PY].

Table 1 Risk factors associated with the discontinuation of anti-tumour necrosis factor alpha (TNF α) drugs owing to abnormal blood test results

	HR (95% CI)	aHR (95% CI)
Age, years	1.03 (1.01–1.04)	1.02 (1.01–1.04)
Sex		
Female	1	1
Male	1.25 (0.88–1.78)	1.31 (0.91–1.89)
BMI (kg m ⁻²) ^a	1.01 (1.00–1.02)	1.01 (0.99–1.02)
Alcohol (units weekly) ^b	1.01 (1.00–1.02)	1.01 (1.00–1.02)
Nondrinker	1	1
≤ 14	0.98 (0.64–1.51)	1.09 (0.71–1.69)
15–21	1.22 (0.67–2.21)	1.39 (0.76–2.56)
≥ 21	0.89 (0.44–1.76)	1.03 (0.50–2.10)
Diabetes ^c		
No	1	1
Yes	2.16 (1.34–3.49)	1.68 (1.00–2.81)
Liver disease ^c		
No	1	1
Yes	2.66 (1.50–4.72)	2.27 (1.26–4.07)
High cholesterol ^c		
No	1	1
Yes	1.01 (0.54–1.87)	0.69 (0.36–1.31)

aHR, adjusted hazard ratio; BMI, body mass index; CI, confidence interval; HR, hazard ratio. ^aBMI was calculated using height in meters/(weight in kg × weight in kg). Data for BMI were missing for 620 participants. ^bData for alcohol intake were missing for 848 participants. ^cComorbidities were defined as present if their year of onset preceded the year of anti-TNF α prescription, else they were defined as absent. No person with kidney disease experienced the outcome of interest.

a significantly higher risk of stopping infliximab treatment associated with abnormal monitoring blood test results.^{11,15}

Our cost-effectiveness analysis showed that monitoring with blood tests every year and every 2 years was more cost-effective than monitoring with blood tests every 3–6 months, and monitoring every 6 months was more cost-effective than monitoring every 3 months, although no monitoring was most cost-effective. This is a novel finding and may be used to reduce the frequency of monitoring blood tests in those established on anti-TNF α treatment (e.g. after the first 6 months). However, such changes in clinical practice should only be implemented once the findings of this study have been considered by the relevant specialist societies.

In this study, among those who stopped treatment as a result of abnormal blood test results, most did so due to abnormal LFT results, and of these the vast majority had mildly elevated liver enzymes. Only a few participants stopped their treatment owing to low WBC and low platelet counts. No participant in this study had either leukopenia or thrombocytopenia that was severe enough to increase the risk of infection and bleeding, respectively. Previous studies have reported occasional patients with severe neutropenia and infection,¹³ although severe thrombocytopenia increasing the risk of bleeding was not reported previously, to the best of our knowledge. Two patients stopped treatment owing to abnormal renal function. This is not unexpected as anti-TNF α drugs are occasionally associated with granulomatous interstitial nephritis.^{16,17}

The strengths of this study included the use of a nationwide cohort of patients with psoriasis from the UK and ROI, increasing the generalizability of the study findings. Furthermore, the reasons for stopping treatment were

systematically documented in BADBIR. Free-text reasons for stopping treatment and MedDRA coding were individually reviewed when there was diagnostic uncertainty. This reduced the chances of missing outcome data. Another strength was the large sample size – to the best of our knowledge this is the largest study of its kind to date. Nevertheless, several limitations need to be considered when interpreting the results. Some patients who stopped treatment owing to inefficacy and mildly abnormal monitoring blood test results were considered to have stopped treatment for abnormal blood test results. This may have increased the event rate and biased the health economic gain from less-frequent monitoring toward null. Lack of information on blood test results is another limitation. Information on disease severity and concurrent medications was not included in the data analysis. Few participants were treated with infliximab and certolizumab, which increased the imprecision around estimates of the incidence of side-effects with these drugs.

In conclusion, this study has shown that anti-TNF α drugs are uncommonly stopped owing to abnormal monitoring blood test results and our health economic analysis showed that it is more cost-effective to undertake monitoring blood tests annually and biennially than the current practice of monitoring every 6 months. The results of this study ought to be considered by guideline writing groups so that the frequency of monitoring blood tests may be reduced.

Funding sources

National Institute for Health and Care Research grant (NIHR130580). BADBIR was supported by the National Institute for Health and Care Research funded Manchester Biomedical Research Centre (NIHR203308).

Conflicts of interest

A.A. reports personal fees from UpToDate (royalty), Springer (royalty), Cadilla Pharmaceuticals (lecture fees), NGM Bio (consulting) and Limbic (consulting) and personal fees from Inflazome (consulting), unrelated to this work. G.P.A. has received consulting fees from Abbott, Albereo, Amryth, AstraZeneca, BenevolentAI, DNDI, GlaxoSmithKline, NuCANA, Pfizer, Roche Diagnostics, Servier Pharmaceuticals and W.L Gore & Associates paid to the University of Nottingham, unrelated to this work. C.P.F. has received consultancy/advisory board fees from AbbVie, GenMab, Incyte, Morphosys, Roche, Takeda, Ono, Kite/Gilead, BMS/Celgene and BTG/Veriton; and departmental research funding from BeiGene unrelated to this work. R.B.W. reports research grants from AbbVie, Almirall, Amgen, Celgene, Janssen, Lilly, LEO, Novartis, Pfizer and UCB, and consulting fees from AbbVie, Almirall, Amgen, Arena, Astellas, Avillion, Biogen, Boehringer Ingelheim, Bristol Myers Squibb, Celgene, DiCE, GSK, Janssen, Lilly, LEO, Novartis, Pfizer, Sanofi, Sun Pharma, UCB and UNION. The other authors declare no conflicts of interest.

Data availability

The data included in this study are available from British Association of Dermatologists Biologic and Immunomodulators Register (BADBIR).

Ethics statement

The British Association of Dermatologists Biologic and Immunomodulators Register (BADBIR) received approval from the North West Research Ethics Committee in March 2007. All participants provided written informed consent in accordance with the Declaration of Helsinki.

Supporting Information

Additional [Supporting Information](#) may be found in the online version of this article at the publisher's website.

References

- Smith CH, Yiu ZZN, Bale T *et al.* British Association of Dermatologists guidelines for biologic therapy for psoriasis 2020: a rapid update. *Br J Dermatol* 2020; **183**:628–37.
- Menter A, Strober BE, Kaplan DH *et al.* Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol* 2019; **80**:1029–72.
- Chalasanani N, Björnsson E. Risk factors for idiosyncratic drug-induced liver injury. *Gastroenterology* 2010; **138**:2246–59.
- Bessissow T, Renard M, Hoffman I *et al.* Review article: non-malignant haematological complications of anti-tumour necrosis factor alpha therapy. *Aliment Pharmacol Ther* 2012; **36**:312–23.
- Hoffmann JHO, Knoop C, Schäkel K *et al.* Long-term safety of combination treatment with methotrexate and tumor necrosis factor (TNF)- α antagonists versus TNF- α antagonists alone in psoriatic patients. *J Dermatol* 2021; **48**:835–43.
- Gottlieb AB, Langley RG, Strober BE *et al.* A randomized, double-blind, placebo-controlled study to evaluate the addition of methotrexate to etanercept in patients with moderate to severe plaque psoriasis. *Br J Dermatol* 2012; **167**:649–57.
- Burden AD, Warren RB, Kleyn CE *et al.* The British Association of Dermatologists' Biologic Interventions Register (BADBIR): design, methodology and objectives. *Br J Dermatol* 2012; **166**:545–54.
- Diao JA, Inker LA, Levey AS *et al.* In search of a better equation – performance and equity in estimates of kidney function. *N Engl J Med* 2021; **384**:396–9.
- National Institute of Health and Care Excellence. NICE health technology evaluations: the manual 2022. Available at: <https://www.nice.org.uk/process/pmg36/resources/nice-health-technology-evaluations-the-manual-pdf-72286779244741> (last accessed 20 November 2023).
- Ghabril M, Bonkovsky HL, Kum C *et al.* Liver injury from tumor necrosis factor- α antagonists: analysis of thirty-four cases. *Clin Gastroenterol Hepatol* 2013; **11**:558–64.
- Björnsson ES, Gunnarsson BI, Gröndal G *et al.* Risk of drug-induced liver injury from tumor necrosis factor antagonists. *Clin Gastroenterol Hepatol* 2015; **13**:602–8.
- Shelton E, Chaudrey K, Sauk J *et al.* New onset idiosyncratic liver enzyme elevations with biological therapy in inflammatory bowel disease. *Aliment Pharmacol Ther* 2015; **41**:972–9.
- Hastings R, Ding T, Butt S *et al.* Neutropenia in patients receiving anti-tumor necrosis factor therapy. *Arthritis Care Res (Hoboken)* 2010; **62**:764–9.
- Rajakulendran S, Gadsby K, Allen D *et al.* Neutropenia while receiving anti-tumour necrosis factor treatment for rheumatoid arthritis. *Ann Rheum Dis* 2006; **65**:1678–9.
- French JB, Bonacini M, Ghabril M *et al.* Hepatotoxicity associated with the use of anti-TNF- α agents. *Drug Saf* 2016; **39**:199–208.
- Sandys V, Moloney B, Lane L *et al.* Granulomatous interstitial nephritis secondary to adalimumab therapy. *Clin Kidney J* 2018; **11**:219–21.
- Plant R, Rafi Ahmed A, McHale T *et al.* A case of adalimumab-induced granulomatous interstitial nephritis. *Cureus* 2021; **13**:e15986.