

Characterising burden of treatment in cystic fibrosis to identify priority areas for clinical trials

Gwyneth Davies^{1**}, Nicola J Rowbotham^{2*}, Sherie Smith^{2*}, Zoe C Elliot³, Katie Gathercole^{4,5}, Oli Rayner⁶, Paul A Leighton⁷, Sophie Herbert^{5,2}, Alistair JA Duff⁸, Suja Chandran⁹, Tracey Daniels¹⁰, Edward F Nash¹¹, Alan R Smyth².

*joint first authors. #author for correspondence (Email: gwyneth.davies@ucl.ac.uk)

Online Supplementary file 2

Table E1. Number of responses to each question (see blank version of survey for questions, in Online Supplementary file 1).

Question	Total no. responses	Applies to Lay (L) / Professional (P)/ All (A)	Question	Total no. responses	Applies to Lay (L) / Professional (P)/ All (A)
1	941	A	25	251	L
2	369	L	26	289	L
3	336	L	27	286	L
4	302	L	28	267	L
5	351	L	29	279	L
6	348	L	30	261	L
7	327	L	31	289	P
8	343	L	32	172	P
9	333	L	33	172	P
10	345	L	34	164	P
11	318	L	35	136	P
12	317	L	36	167	P
13	252	L	37	164	P
14	248	L	38	157	P
15	243	L	39	151	P
16	190	L	40	165	P
17	319	L	41	165	P
18	121	L	42	442	A
19	296	L	43	228	A
20	292	L	44	445	A
21	276*	L	45	442	A
22	279*	L	46	439	A
23	284	L	47	314	A
24	289	L	48	36	A

*Responses to question 21 and 22 were used to generate the Spearman correlation coefficient for number of treatments and time taken for treatments. N=269 respondents answered both questions and therefore were included in this analysis.