

Regularity of $R(X)$ does not pass to finite unions

J. F. Feinstein

Abstract

We show that there are compact plane sets X, Y such that $R(X)$ and $R(Y)$ are regular but $R(X \cup Y)$ is not regular.

1 Introduction

In [11], McKissick gave the first example of a non-trivial, regular uniform algebra. His example was $R(X)$ for a suitable compact plane set X . (We recall the relevant definitions in the next section.) In [5, Corollary 7.7] it was shown that if X and Y are compact plane sets such that $R(X)$ and $R(Y)$ are regular and $X \cap Y$ is countable, then $R(X \cup Y)$ is also regular.

This note addresses the question of whether $R(X \cup Y)$ is regular whenever $R(X)$ and $R(Y)$ are regular, without any additional assumptions on the compact plane sets X and Y . If the answer was positive, the result would clearly also hold for all finite unions. Bearing this in mind, we show that the answer is negative, in Theorem 6 below, by the slightly indirect method of finding **four** compact plane sets X_k , $1 \leq k \leq 4$, such that each $R(X_k)$ is regular, but such that $R\left(\bigcup_{k=1}^4 X_k\right)$ is not regular.

We describe this result by saying that *regularity of $R(X)$ does not pass to finite unions*. Along the way, in Theorem 5, we give a slightly easier example showing that (in an obvious sense) regularity of $R(X)$ does not pass to countable unions.

2 Preliminaries

Throughout this note, by a *compact space* we shall mean a non-empty, compact, Hausdorff topological space; by a *compact plane set* we shall mean a non-empty, compact subset of the complex plane. We shall use the term *clopen* to describe sets which are both open and closed in a given topological space (typically a compact plane set).

Let $a \in \mathbb{C}$ and let $r > 0$. We denote the open disk of radius r and centre a by $D(a, r)$ and the corresponding closed disk by $\bar{D}(a, r)$. We denote the diameter of a non-empty, bounded subset E of \mathbb{C} by $\text{diam}(E)$.

We assume that the reader has some familiarity with uniform algebras. We refer the reader to [1, 9, 12] for further background. For the general theory of commutative Banach algebras, the reader may consult [2, 10].

Let X be a compact space, and let $C(X)$ be the algebra of all continuous complex-valued functions on X . For each function $f \in C(X)$ and each non-empty subset E of X , we denote the uniform norm of the restriction of f to E by $|f|_E$. In particular, we denote by $|\cdot|_X$ the uniform norm on X . When endowed with the norm $|\cdot|_X$, $C(X)$ is a Banach algebra. A *uniform algebra* on X is a closed subalgebra of $C(X)$ that contains the constant functions and separates the points of X . We say that a uniform algebra A on X is *nontrivial* if $A \neq C(X)$, and is *natural* (on X) if X is the character space of A (under the usual identification of points of X with evaluation functionals).

Let A be a natural uniform algebra on X , and let $x \in X$. We denote by J_x the ideal of functions f in A such that x is in the interior of the zero set of f , $f^{-1}(\{0\})$. We denote by M_x the ideal of functions f in A such that $f(x) = 0$.

We say that x is a *point of continuity* (for A) if, for all $y \in X \setminus \{x\}$ we have $J_y \not\subseteq M_x$; we say that x is an *R-point* (for A) if, for all $y \in X \setminus \{x\}$ we have $J_x \not\subseteq M_y$. We say that A is *regular* if, for every closed subset F of X and every $y \in X \setminus F$, there exists $f \in A$ with $f(y) = 1$ and $f(F) \subseteq \{0\}$.

The natural uniform algebra A is regular if and only if every point of X is a point of continuity, and this is also equivalent to the condition that every point of X is an R-point ([6, 7]).

Let X be a compact plane set. By $R(X)$ we denote the set of those functions $f \in C(X)$ which can be uniformly approximated on X by rational functions with no poles on X . It is standard that $R(X)$ is a natural uniform algebra on X .

Let $x \in X$. As we will sometimes need to work with different compact plane sets simultaneously, in this note we will denote the ideals J_x and M_x in $R(X)$ by J_x^X and M_x^X respectively.

We will occasionally use the fact that all idempotents in $C(X)$ are automatically in $R(X)$. This does not require the full force of the Shilov Idempotent Theorem, as for $R(X)$ it follows from Runge's theorem.

Let X be a compact plane set, let Y be a compact subset of X and suppose that $x, y \in Y$ with $x \neq y$. Trivially the restriction $R(X)|_Y$ is contained in $R(Y)$. It follows immediately that if $J_x^X \not\subseteq M_y^X$ then $J_x^Y \not\subseteq M_y^Y$. Thus if x is an R-point for $R(X)$ then x is also an R-point for $R(Y)$. Similarly, if x is a point of continuity for $R(X)$ then x is a point of continuity for $R(Y)$. If $R(X)$ is regular, then so is $R(Y)$.

We now discuss some conditions under which converses to some of these implications hold.

Lemma 1. *Let X be a compact plane set and suppose that E is a non-empty, clopen subset of X . Let $x \in E$. If x is an R-point for $R(E)$ then x is an R-point for $R(X)$. Similarly, if x is a point of continuity for $R(E)$ then x is a point of*

continuity for $R(X)$.

Proof. This is almost trivial in view of the idempotents available in $R(X)$. The only point that may be worth noting is the fact that, for all $y \in E$, each function in J_y^E has an extension in J_y^X . This is because every function in $R(E)$ can be extended to give a function in $R(X)$ which is constantly zero on $X \setminus E$. \square

Let X and Y be compact plane sets, with $Y \subseteq X$, and suppose that X contains no bounded component of $\mathbb{C} \setminus Y$. Then, by Runge's theorem, $R(X)|_Y$ is dense in $R(Y)$. This condition concerning the bounded components of $\mathbb{C} \setminus Y$ is, of course, automatically satisfied if X has empty interior.

Theorem 2. *Let X and Y be compact plane sets, with $Y \subseteq X$ and suppose that $x, y \in Y$ with $x \neq y$. Suppose that X contains no bounded component of $\mathbb{C} \setminus Y$ and that $X \setminus Y$ is the union of a sequence of pairwise disjoint, non-empty relatively clopen subsets Y_n of X whose diameters tend to 0 as $n \rightarrow \infty$. Then Y is a peak set for $R(X)$ and $R(X)|_Y = R(Y)$. Moreover the following implications hold.*

- (i) *If $J_x^Y \not\subseteq M_y^Y$ then $J_x^X \not\subseteq M_y^X$.*
- (ii) *If x is an R -point for $R(Y)$ then x is an R -point for $R(X)$.*
- (iii) *If x is a point of continuity for $R(Y)$ then x is a point of continuity for $R(X)$.*

Proof. In this proof, we work on X when discussing clopen sets or characteristic functions: the characteristic functions of the clopen sets are then precisely the idempotents in $R(X)$.

Note that clopen subsets of X are automatically peak sets (using idempotents in $R(X)$). Clearly Y is an intersection of (a sequence of) clopen subsets of X , so Y is also a peak set for $R(X)$. Thus $R(X)|_Y$ is a uniformly closed subset of $R(Y)$. As X contains no bounded component of $\mathbb{C} \setminus Y$, $R(X)|_Y$ is dense in $R(Y)$, so equality holds.

Using idempotents, we also see that, for each $z \in X \setminus Y$, we have $J_x^X \not\subseteq M_z^X$ and $J_z^X \not\subseteq M_x^X$. Thus (ii) and (iii) follow quickly from (i).

To prove (i), suppose that $J_x^Y \not\subseteq M_y^Y$. Choose $f \in J_x^Y \setminus M_y^Y$. We shall show that f has an extension in J_x^X , from which (i) follows. We know that f has an extension, say F , in $R(X)$. However F might not be in J_x^X , so we modify F as follows.

Choose $r > 0$ such that $f(\bar{D}(x, r) \cap Y) = \{0\}$.

Set

$$S = \{n \in \mathbb{N} : Y_n \cap \bar{D}(x, r) \neq \emptyset\}.$$

If S is finite, then F is already in J_x^X . So we may assume that S is infinite, say $S = \{n_1, n_2, \dots\}$ with $n_1 < n_2 < \dots$. Set $U = \bigcup_{n \in S} Y_n = \bigcup_{k=1}^{\infty} Y_{n_k}$.

We define $\tilde{f} : X \rightarrow \mathbb{C}$ by $\tilde{f} = (1 - \chi_U)F$. We claim that \tilde{f} is the extension of f that we need. Clearly \tilde{f} vanishes on $X \cap D(x, r)$. It remains to show that $\tilde{f} \in R(X)$ (and, in particular, that \tilde{f} is continuous).

For each $m \in \mathbb{N}$, set $U_m = \bigcup_{k=1}^m Y_{n_k}$ and $V_m = \bigcup_{k=m+1}^{\infty} Y_{n_k}$. Then U_m is clopen in X , so $\chi_{U_m} \in R(X)$ and hence $(1 - \chi_{U_m})F \in R(X)$. Since $\text{diam}(Y_{n_k}) \rightarrow 0$ as $k \rightarrow \infty$, we see that the sequence of clopen sets Y_{n_k} can accumulate only at points of $\bar{D}(x, r) \cap Y$, and that $\text{dist}(Y_{n_k}, \bar{D}(x, r) \cap Y) \rightarrow 0$ as $k \rightarrow \infty$. It then follows that $|F|_{V_m} \rightarrow 0$ as $m \rightarrow \infty$, and hence $(1 - \chi_{U_m})F \rightarrow \tilde{f}$ uniformly on X . The result follows. \square

Using Theorem 2 and our earlier observations, we immediately obtain the following corollary concerning the regularity of $R(X)$ for this type of compact plane set X .

Corollary 3. *Let X and Y be compact plane sets, with $Y \subseteq X$. Suppose that X contains no bounded component of $\mathbb{C} \setminus Y$ and that $X \setminus Y$ is the union of a sequence of pairwise disjoint, non-empty relatively clopen subsets Y_n of X whose diameters tend to 0 as $n \rightarrow \infty$. Then $R(X)$ is regular if and only if $R(Y)$ is regular and each $R(Y_n)$ is regular.*

3 Main results

We are almost ready to construct the examples mentioned in the introduction. First we need the following result, which is taken from [3] (see also [4, 8]).

Proposition 4. *There exist a compact plane set X and a non-degenerate closed interval J in \mathbb{R} with $J \subseteq X$ such that $R(X)$ is not regular, but every point of $X \setminus J$ is a point of continuity for $R(X)$.*

Let X, J be as in Proposition 4. Clearly X has empty interior. For every non-empty compact subset K of $X \setminus J$, $R(K)$ is regular, because every point of K is a point of continuity for $R(K)$. We have $R(J) = C(J)$, and so $R(J)$ is regular. However, $R(X)$ is not regular, so not every point of X is a point of continuity for $R(X)$. The points of X which are not points of continuity for $R(X)$ must all be in J . By Lemma 1, J is not clopen in X . In particular, $X \neq J$.

Using this example, we next show that, in a suitable sense, regularity of $R(X)$ does not pass to countable unions.

Theorem 5. *There exist a compact plane set X and a sequence of compact plane sets X_k ($k \in \mathbb{N}$) such that $X = \bigcup_{k \in \mathbb{N}} X_k$ and each $R(X_k)$ is regular, but $R(X)$ is not regular.*

Proof. Let X and J be as in Proposition 4. Scaling X if necessary, we may assume that $\{z \in X : \text{dist}(z, J) \geq 1/2\} \neq \emptyset$. Set $X_1 = J$ and, for $k \geq 2$, set $X_k = \{z \in X : \text{dist}(z, J) \geq 1/k\}$. Then $R(X)$ is not regular, but each $R(X_k)$ is regular, and $X = \bigcup_{k \in \mathbb{N}} X_k$ as required. \square

Thus regularity of $R(X)$ does not pass to countable unions. We finish by showing that regularity of $R(X)$ does not pass to finite unions either.

Theorem 6. *There exist a compact plane set X and four compact plane sets X_k , $k = 1, 2, 3, 4$, such that $X = \bigcup_{k=1}^4 X_k$ and each $R(X_k)$ is regular, but $R(X)$ is not regular.*

Proof. Again, let X and J be as in Proposition 4. Scaling X if necessary, we may assume that $X \subseteq \{z \in \mathbb{C} : \text{dist}(z, J) \leq 1\}$.

We begin by defining subsets K_o and K_e as follows.

Set

$$K_o = J \cup \bigcup_{n \in \mathbb{N}} \{z \in X : 1/(2n+1) \leq \text{dist}(z, J) \leq 1/2n\}$$

and

$$K_e = J \cup \bigcup_{n \in \mathbb{N}} \{z \in X : 1/2n \leq \text{dist}(z, J) \leq 1/(2n-1)\}.$$

Then $X = K_o \cup K_e$, and $R(X)$ is not regular.

It is not clear whether or not $R(K_o)$ or $R(K_e)$ are regular. In the following, we assume that J is clopen in neither K_o nor K_e (otherwise the arguments may be simplified). We now show that each of K_o and K_e is a union of two sets where we can establish regularity using Corollary 3. We may write $K_o = X_1 \cup X_3$ and $K_e = X_2 \cup X_4$, where the sets X_k are defined with the help of vertical bands as follows:

$$X_1 = J \cup \bigcup_{n \in \mathbb{N}} \{z \in X : 1/(2n+1) \leq \text{dist}(z, J) \leq 1/2n \text{ and } \sin(n\text{Re}(z)) \leq 0\};$$

$$X_2 = J \cup \bigcup_{n \in \mathbb{N}} \{z \in X : 1/2n \leq \text{dist}(z, J) \leq 1/(2n-1) \text{ and } \sin(n\text{Re}(z)) \leq 0\};$$

$$X_3 = J \cup \bigcup_{n \in \mathbb{N}} \{z \in X : 1/(2n+1) \leq \text{dist}(z, J) \leq 1/2n \text{ and } \sin(n\text{Re}(z)) \geq 0\};$$

$$X_4 = J \cup \bigcup_{n \in \mathbb{N}} \{z \in X : 1/2n \leq \text{dist}(z, J) \leq 1/(2n-1) \text{ and } \sin(n\text{Re}(z)) \geq 0\}.$$

Then, for $1 \leq k \leq 4$, it is easy to see that X_k is the disjoint union of J with a sequence $(Y_{k,n})_{n=1}^{\infty}$ of pairwise disjoint, non-empty relatively clopen subsets of X_k whose diameters tend to 0. Each of the sets X_k then satisfies the conditions of Corollary 3 with $Y = J$ and $Y_n = Y_{k,n}$ ($n \in \mathbb{N}$). Since $R(J)$ and all of

the $R(Y_{k,n})$ are regular, each $R(X_k)$ is regular by Corollary 3. Certainly $X = \bigcup_{k=1}^4 X_k$, and $R(X)$ is not regular, as required. \square

As noted earlier, it follows that regularity of $R(X)$ does not pass to unions of two sets either. However, we do not know which pair of sets mentioned above demonstrates this. Clearly at least one of the following pairs of sets must provide a suitable example: X_1 and X_3 ; X_2 and X_4 ; K_o and K_e .

We do not know whether or not regularity of $R(X)$ *does* pass to unions of two sets if the sets are assumed to be connected.

We are grateful to the referee for a careful reading and for useful comments and suggestions.

References

- [1] A. Browder, *Introduction to Function Algebras*, Benjamin, New York (1969).
- [2] H. G. Dales, *Banach Algebras and Automatic Continuity*, London Mathematical Society Monographs, New Series, vol. 24, The Clarendon Press Oxford University Press, New York, 2000.
- [3] J. F. Feinstein, *Trivial Jensen measures without regularity*, *Studia Math.*, 148 (2001), 67–74.
- [4] J. F. Feinstein and S. Morley, *Quasianalyticity in certain Banach function algebras*, *Studia Math.*, 238 (2017), 133–153
- [5] J. F. Feinstein, S. Morley and H. Yang, *Abstract Swiss cheese space and classicalisation of Swiss cheeses*, *J. Math. Anal. Appl.*, 438 (2016), 119–141.
- [6] J. F. Feinstein, R. Mortini, *Partial regularity and t -analytic sets for Banach function algebras*, *Math. Z.*, 271 (2012), 139–155.
- [7] J. F. Feinstein and D. W. B. Somerset, *Non-regularity for Banach function algebras*, *Studia Math.*, 141 (2000), 53–68.
- [8] J. F. Feinstein and H. Yang, *Regularity points and Jensen measures for $R(X)$* , *Studia Math.*, 235 (2016), 1–15
- [9] T. W. Gamelin, *Uniform Algebras*, 2nd ed., Chelsea Publishing Company, New York, NY, 1984.
- [10] E. Kaniuth, *A Course in Commutative Banach Algebras*, Graduate Texts in Mathematics 246, Springer, New York, 2009.

- [11] R. McKissick, *A nontrivial normal sup norm algebra*, Bull. Amer. Math. Soc., 69 (1963), 391–395.
- [12] E. L. Stout, *The Theory of Uniform Algebras*, Bogden & Quigley, New York, 1971.