

Short-term integrated rehabilitation for people with newly diagnosed thoracic cancer: a multi-centre randomised controlled feasibility trial

Joanne Bayly joanne.bayly@kcl.ac.uk @jobk2

Lucy Fettes lucy.fettes@kcl.ac.uk

Eleanor Douglas Eleanor.Douglas@nottingham.ac.uk

Maria J Teixeira maria.teixeira@kcl.ac.uk

Nicola Peat Nicola.Peat@gstt.nhs.uk

India Tunnard india.tunnard@kcl.ac.uk

Vishit Patel vishit.patel@kcl.ac.uk

Wei Gao wei.gao@kcl.ac.uk

Andrew Wilcock Andrew.wilcock@nottingham.ac.uk

Irene J Higginson Irene.higginson@kcl.ac.uk

Matthew Maddocks [@MTMaddocks](mailto:matthew.maddocks@kcl.ac.uk)

Abstract

Objectives: Determine the feasibility of recruiting and retaining patients recently diagnosed with thoracic cancer to a trial of short-term integrated rehabilitation; evaluate uptake of theoretically informed components targeting physical function, symptom self-management and participation; estimate sample size requirements for an efficacy trial.

Design: Parallel group randomised controlled feasibility trial

Setting: Three United Kingdom hospitals

Participants: Patients ≤ 8 weeks of thoracic cancer diagnosis, Eastern Cooperative Oncology Group Performance Status 0–3, any cancer stage and treatment plan.

Interventions: Participants randomly allocated (1:1) to short-term integrated rehabilitation and standard care or standard care alone over 30 days.

Main measures: Primary: participant recruitment and retention, targeting $\geq 30\%$ of eligible patients enrolling and $\geq 50\%$ of participants reporting outcomes at 30 days. Secondary: intervention fidelity; missing data and performance of outcome measures for self-efficacy, symptoms, physical activity and health-related quality of life.

Results: Of 159 eligible patients approached, 54 (34%) were recruited. 44 (82%) and 39 (72%) participants reported outcomes at 30 and 60 days. Intervention fidelity was high. Rehabilitation was delivered across 3 (1-3) sessions over 32 (22-45) days (median(range)). Changes in clinical outcomes were modest but most apparent at 60 days for health-related quality of life: Functional Assessment of Cancer Therapy Lung Cancer score median [IQR] change 9.7 (-12.0-16.0) rehabilitation versus 2.3 (-15.0-14.5) standard care.

Conclusion: A trial to examine efficacy of short-term integrated rehabilitation for people newly diagnosed with thoracic cancer is feasible. A sample of 336 participants could detect a meaningful effect on health-related quality of life as the primary outcome.

Keywords: Feasibility Trial, Lung cancer, Mesothelioma, Rehabilitation, Randomised Controlled Trial, Thoracic Cancer

Introduction

Worldwide, more people are diagnosed with lung cancer than any other cancer [1] and the incidence of pleural mesothelioma continues to rise.[2] The prevalence of distressing and disabling symptoms in these thoracic cancers is high.[3, 4] Most people are diagnosed with advanced disease[1, 2] but strive to maintain normality in daily life.[5] Needs based cancer rehabilitation is recommended from diagnosis to reduce the impact of cancer and its treatment on functional well-being and to promote independence.[6] However, many people face difficulties accessing rehabilitation due to poor service provision, lack of clinician recognition of functional need [6] and/or negative views of rehabilitation.[7] Although traditional exercise-based interventions are safe and potentially effective in this population, rates of uptake and completion in supervised programmes are generally low.[8, 9] Symptom management interventions tend to be reactive, targeting patients with well-established symptoms.[10] Individualised pro-active rehabilitation programmes to minimise the impact of thoracic cancer on participation in daily life are lacking. To overcome some of these issues, we have developed a model of short-term integrated rehabilitation drawing on theories of illness,[11, 12] rehabilitation[13] and behaviour change.[14] Rehabilitation processes, components and outcomes were identified via systematic review [15] and focus groups with patients, carers and healthcare professionals.[16] A manual was produced to support delivery of the intervention by a trained rehabilitation practitioner, (e.g. physiotherapist, occupational therapist or dietitian). Short-term integrated rehabilitation aims to support people to self-manage symptoms and immediate functional needs. It also aims to reduce sedentary time to minimise the onset of physical deconditioning. Strategies include maintaining or improving physical activity levels, fitness and participation in daily activities during cancer treatment and beyond.

Before testing the efficacy of short-term integrated rehabilitation in a large-scale trial, we have undertaken this formal feasibility study. We aimed to: (1) determine the feasibility of recruiting and retaining people with thoracic cancer to a trial of short-term integrated

rehabilitation delivered in the period following diagnosis; (2) to evaluate uptake of theoretically informed, individually tailored intervention components and (3) obtain data on the acceptability of selected outcome measures and (4) estimate sample size requirements for an efficacy trial.

Methods

A multi-centre randomised controlled feasibility trial comparing the short-term integrated rehabilitation service plus standard care to standard care took place between February 2018 and April 2019. The protocol was pre-registered (ISRCTN 92666109). Ethical approval was granted by the London South East Research Ethics Committee (Ref 17/LO/1871). Reporting follows the CONSORT guidelines and the Template for Intervention Description and Replication (TIDieR) checklist. No changes were made to the design or methods following trial commencement.

Participants were recruited from thoracic oncology, palliative care and respiratory clinics across three UK hospitals, two in London and one in Nottingham. Eligible participants were adults, within 8 weeks of a clinical or histological diagnosis of primary lung cancer (stage I-IV) or pleural mesothelioma (local or extensive), Eastern Cooperative Oncology Group Performance Status of 0-3 (appendix 1)[17] and the ability to respond to questions in written English (or availability of translators to support this). People already receiving specialist rehabilitation, or who had a co-existing progressive neurological condition (e.g. motor neurone disease), Eastern Cooperative Oncology Group Performance Status 4, not able to complete questionnaires due to cognitive impairment, or with a physician-estimated prognosis of <1 month were excluded. All participants gave written informed consent in accordance with Good Clinical Practice and the Declaration of Helsinki.

Following baseline assessment, conducted in hospital out-patients, in-patients or home settings, participants were randomly allocated (1:1 at the individual level) using an independent web-based randomisation system at the King's Clinical Trials Unit, London, UK.

A minimisation algorithm was used to maintain balance in both the trial groups for recruitment site, performance status (0-1 or 2-3) and disease stage (I-II or III-IV). Following randomisation, the Clinical Trials Unit informed trial staff via secure email. Research nurses, who arranged outcome assessments using self-report mailed questionnaires, were informed of trial entry but not group allocation. Due to the nature of the intervention, it was not possible to maintain allocation blinding for participants and trial physiotherapists (JB, MM, LF, ED). The trial statistician (WG) was also blind to group allocation.

All participants received standard care provided by their hospital for their condition. This included surgical, oncology and supportive and palliative care services. No rehabilitation was routinely provided at two sites. One London site provided multi-professional out-patient based specialist rehabilitation for patients attending oncology follow-up clinics with functional needs identified following screening. We planned to recruit participants who were not scheduled or able to attend these clinics within eight weeks of diagnosis.

Participants allocated to short-term integrated rehabilitation (Figure 1) were offered up to three sessions with a rehabilitation practitioner (physiotherapists) over six weeks (trained by JB). As indicated during model development,[16] a flexible delivery approach was used with appointments lasting up to one hour, in the hospital or home setting, and scheduled according to participant preference. Where possible, first appointments were face to face. Subsequent appointments were face to face or by telephone according to practical considerations and participant preference. The intervention used a psychologically-informed approach[18] to identify immediate functional priorities and concerns, illness understandings and future expectations relating to functional well-being. A intervention manual (available [here](#)) was used to standardise change processes[19] but was unscripted to enable tailoring and adaptation of intervention components to meet the expected variation in function related needs. Intervention components were selected as indicated and included personalised information about the impact of thoracic cancer on function and goal planning for symptom

prevention and/or self-management, graded physical activity, home-based exercise and participation and performance in activities of daily living.

Intervention components were delivered using behaviour change techniques[14] to support receipt and enactment. Information booklets, handheld fans and walking aids were offered and provided to support delivery of intervention components. An individual goal orientated action plan, developed in consultation with each participant and held by them, was reviewed and updated at each session to support enactment. Participants were sign-posted to local hospital, community-based, hospice and charitable services (i.e. local cancer support organisations) as indicated in the manual and where available for on-going support. Those wanting to pursue supervised exercise training to improve fitness were sign-posted to local exercise services. At discharge, a letter summarising the participant's rehabilitation intervention and discharge action plan was sent to the participant and copied to their multi-professional team for follow-up and onward referrals.

Our primary feasibility outcomes were rates of participant enrolment and retention.

Feasibility endpoints, established *a priori*, were defined as $\geq 30\%$ of eligible patients consenting to the trial and $\geq 50\%$ of enrolled participants completing clinical outcome measures at 30 days. Secondary feasibility objectives were number of rehabilitation contacts delivered over 30 days, fidelity of service delivery, including use of intervention components and onwards referral to local rehabilitation services, and missing data on clinical outcome measures at 30 and 60 days. Participant self-reported experiences were assessed by self-reported questionnaire comprising four items on trial processes from a national experience survey[20] and ten items on satisfaction with the intervention from the Functional Assessment of Chronic Illness Therapy Treatment/ Patient Satisfaction (version 4)[21]. Free text comments were also invited.

Clinical outcomes were assessed at baseline, 30 days and 60 days following randomisation via self-reported postal questionnaire. Higher scores are better for all measures except the Integrated Palliative Outcome Scale and the Client Services Receipt Inventory. Symptoms

and concerns were measured by the Integrated Palliative Outcome Scale (17 items, total score 0-68);[22] physical activity level by the Physical Activity Scale for the Elderly (28 items, 0-400);[23] health-related quality of life measured by the Functional Assessment of Cancer Therapy-Lung (total 34 item, 0-136; Functional Assessment of Cancer Therapy-Trial Outcome Index (21 items, 0-84); Functional Assessment of Cancer Therapy-lung cancer subscale 7 items, (0-28);[24] EQ-5D-5L (index values -1 to 1, EQ-VAS 0-100);[25] and confidence measured by the Self-Efficacy Measure for Chronic Disease (6 items, 0-60).[26] The Client Services Receipt Inventory[27] was used to measure resource use relating to hospital and community services, including social care, informal care and equipment provision. No changes were made to outcome assessments during the trial.

As this trial was designed to assess the feasibility of testing the short-term integrated rehabilitation intervention, a formal power calculation around an estimated effect size on clinical outcome was not appropriate. Sample sizes of 24-60 participants have been recommended for feasibility studies.[28, 29] We took a conservative approach and aimed to recruit up to 60 participants which would allow us to estimate feasibility parameters and outcomes with sufficient precision to inform a future sample size calculation.

Statistical analysis

Data were analysed using SPSS (version 24, Chicago, IL, USA) under intention-to-treat principles. Feasibility outcomes and baseline demographics were summarised using descriptive statistics. Data on treatment fidelity in the intervention group were extracted from case report forms using the Behaviour Change Technique Taxonomy (version 1)[30] by a trained member of the research team (JB). Treatment receipt was defined as the participant receiving 1-3 rehabilitation sessions comprising symptom management \pm physical activity or exercise \pm task performance and participation interventions. Reasons for withdrawal or loss-to-follow up from the trial were summarised and classified by Attrition Due to Death (ADD), Attrition Due to Illness (ADI) or Attrition at Random (AaR).[31] The proportions of participants missing each variable were summarised by group at each time point. All serious adverse

events (SAE) and adverse events (AE) relating to the intervention were summarised. This trial did not aim to evaluate effectiveness, therefore clinical outcomes were described by trial group at baseline and both follow-up time points, using standard descriptive statistics, without significance testing.

Results

Figure 2 shows participant flow through the trial. Between February 2018 and January 2019, 374 patients were screened of whom 159 (42%) were eligible and 54 (34%) were enrolled. An overall recruitment rate of 4.5 participants per month was achieved with 1-2 accruals per month at each site. Barriers to additional recruitment included personal absences in the clinical and research teams and changes to site working practices, e.g. availability of lung cancer nurse specialists and allocation of space in oncology clinics. Following randomisation 82% and 72% of participants returned follow-up outcome measures at 30 and 60 days respectively. Attrition was comparable across both groups. In the rehabilitation group, one participant withdrew before receiving the intervention due to rapidly progressing disease, and one was lost to follow-up after receiving one rehabilitation contact. In the standard care group two participants were lost to follow-up, and one withdrew from the trial. Postal questionnaires from four participants who remained in the trial (three rehabilitation group, one standard care group) were not received at 30 days.

Participants had a median (range) age of 67 (44-85) years and 35 (65%) were men. Most were retired or not employed (32/54, 60%) though some were on sick leave (14/54, 27%) or still working (7/54, 14%). Less than one-quarter of participants lived alone. The majority 37/54 (68%) had non-small cell lung cancer (NSCLC), though small cell lung cancer (SCLC), pleural mesothelioma and neuroendocrine tumours were represented. Thirty nine (72%) had stage III-IV disease (27 metastatic), of whom over half were scheduled for treatment with chemotherapy doublet combinations. Group allocation differences in mesothelioma diagnosis and disease stage were noted (Table 1).

Overall engagement with the rehabilitation intervention was high and it was delivered according to protocol (Table 2). Across the 26 participants allocated to the intervention group 69 rehabilitation sessions were delivered with a median of 3 per participant. Twenty-one participants (84%) accessed all three rehabilitation sessions. Two-thirds of rehabilitation sessions were delivered face to face in the home; others occurred in hospital outpatient settings with a single session delivered on a hospital inpatient unit. The median duration of the first session was one hour and follow-up sessions, forty-five minutes. Two participants declined further sessions as they perceived no needs or were too busy with appointments, though both remained in the trial.

Most participants engaged with all core intervention components; physical activity and exercise (24/25, 96%), symptom self-management (20/25, 80%), and task performance/participation (20/25, 80%). Interventions frequently accepted by participants were home-based exercise or physical activity programmes and self-management strategies for fatigue and breathlessness. Few participants reported current problems managing personal activities of daily living, though 20/25 (80%) and 16/25 (64%) were provided with strategies to support work/leisure or domestic activities respectively.

Regarding use of behaviour change techniques,[14] ‘credible source’ and ‘information about health consequences’ were used to support intervention delivery in all participants (supplementary table 1). Behaviour change techniques relating to goal setting, action planning, feedback and self-monitoring, instruction, practice, and ‘verbal persuasion about capability’ were used with over three-quarters of participants. Strategies to reduce negative emotions supported intervention delivery and enactment of rehabilitation plans in 12/25 (48%) participants. Nine interventions delivered (9/26, 36%) included ‘adding objects to the environment’ (handheld fans and assistive devices), though restructuring of the physical and social environment did not take place. Evaluation of participant enactment of rehabilitation plans was clearly documented in most follow up sessions.

No serious adverse events related to the intervention occurred. Two participants died following disease progressions and seven participants were admitted to hospital relating to disease and oncology treatments (disease progression n=3; sepsis, anaemia, pulmonary embolus, blocked stent, all n=1). Six intervention group participants reported transitional worsening of symptoms following exercise or walking activities, two reported delayed onset muscle soreness and one had a non-injurious fall from a bicycle.

Rates of missing data items in returned questionnaires were low (supplementary table 2). Postal questionnaires from four participants who remained in the trial were not received at 30 days. One participant engaged with the intervention, but outcome measures were not received.

The item most frequently missed was Functional Assessment of Cancer Therapy-Lung item 'satisfaction with sex life' (50%) and items in the Client Services Receipt Inventory relating to personal care (45%) and rehabilitation services (physiotherapy, occupational therapy and dietitian, 27-36%). The Physical Activity Scale for the Elderly questionnaire allocates a score of -1 to a weighted domain containing missing items so all obtained questionnaires were scored despite missing items in 6 (14%) and 5 (13%) questionnaires at 30 and 60 days respectively.

Trial participation satisfaction was high in both groups (median 12) with a wider range of scores in the standard care group. Free text comments from fifteen participants described mostly positive experiences. Two intervention and three standard care group participants felt the questionnaires were too long or hard to fill in. Two standard care participants were unhappy with group allocation. Satisfaction with the rehabilitation intervention was also high with a maximum obtainable median score of 27 (range 23-27). Fourteen participants completed free text comments describing the impact of the intervention and the involvement of family members during intervention delivery. Participants valued the practical advice and face to face contact, reporting that the experience had been 'helpful', 'useful,' 'educational' and 'empowering'. Two would have preferred more sessions. Nine commented that they

valued involvement of a family member while three reported feeling able to talk more freely without family member present.

Table 3 displays change in outcomes at 30 and 60 days. Baseline scores (Table 1) were comparable across both groups for all measures except the Physical Activity Scale for the Elderly, which indicated the rehabilitation group were more physically active. Overall, participants were generally sedentary, reported moderate symptom burden (Integrated Palliative Outcome Scale, median [IQR] score 19 [13 - 28]), and reported having confidence to manage their disease (Self-Efficacy Measure for Chronic Disease 6.7 [5.0 - 8.2]). Client Services Receipt Inventory data indicate that about half of all participants were receiving support with personal care.

Changes over time were modest at both time points and a large degree of heterogeneity was evident (Table 3). Change was most apparent for physical activity level and health-related quality of life. Changes in Physical Activity Scale for the Elderly scores were small in both groups at 30 and 60 days, with slightly higher scores in the standard care group at 60 days (median [IQR] 1.6 [-10.7 - 50.6]) and worsening scores for rehabilitation at 60 days (-15.5 [-50.8 - 38.9]). This contrasted with health-related quality of life scores which favoured the rehabilitation group. At 60 days, Functional Assessment of Cancer Therapy-Lung scores improved by a median of 9.7 [-12.0 - 16.0] compared to 2.3 [-15.0 - 14.5] with standard care. Similarly, FACT-Trial Outcome Index rehabilitation group scores improved by 6 [-12 - 15.0] as compared with 4.5 [-13.0 - 9.8] with standard care. Despite comparable reductions in the EQ-5D utility index in both groups, a small worsening in EQ-VAS median score was observed in the standard care group at 30 and 60 days (-2.5 [-20.0 - 5.0] and -6.5 [-13.8 - 8.8]) as compared to no change then a small improvement (0 [0 - 10] and 5.0 [-17.5 - 24.8]) in the rehabilitation group. Symptoms reduced by a small amount at 30 days in both groups though by 60 days, scores had improved by a median [IQR] of -4.5 [-12.5 - 6.0] in the rehabilitation group and -2.1 [-8.0 - 4.0] in the standard care group.

Discussion

This trial has found that it is feasible to recruit and retain people newly diagnosed with thoracic cancer to a randomised controlled trial of short-term integrated rehabilitation versus standard care. Data relating to recruitment, retention and engagement with the rehabilitation intervention suggest the trial was well designed and implemented, and that this model of rehabilitation is acceptable and accessible for patients across a range of clinical and demographic characteristics. The primary feasibility outcomes were achieved: 34% of eligible participants were recruited and 82% were retained on trial for 30 days.

Low baseline levels of physical activity indicate that the sample was representative of the population.[32] Engagement with the rehabilitation intervention and intervention fidelity were high. More than 96% of participants allocated to receive rehabilitation accessed at least one session of rehabilitation and more than 75% accessed each of the three core intervention components. Low levels of missing data in the clinical outcomes suggest that measures were acceptable to participants and have potential to be used in a future trial (see supplementary material). However unclear or missing data in the self-reported Client Services Receipt Inventory domains relating to physiotherapy, occupational therapy, in-patient and out-patient rehabilitation meant it was difficult to quantify and evaluate access to rehabilitation services. Participants rarely completed number of contacts, duration of intervention or provided information to enable assessment of contamination.

The trial did not aim to test effectiveness and inferential statistical tests were not conducted. It is however of interest that changes in median scores at day 30 and day 60 were higher in the rehabilitation group for all measures except EQ-5D-5L index scores and Physical Activity Scale for the Elderly. Modest changes observed in clinical outcome measures were larger in Functional Assessment of Cancer Therapy-Lung and EQ VAS. Changes in Physical Activity Scale for the Elderly scores were less than the minimally important difference for this population.[23] During development work, confidence was identified as a key modifiable

factor regardless of participant disease stage or performance status,[16] however minimal changes were observed in Self-Efficacy Measure for Chronic Disease scores.

This theory-based model of short-term integrated rehabilitation differs from other rehabilitation interventions tested in people with thoracic cancer in key areas. It brings together strategies to optimise physical function, symptom self-management and participation in valued daily activities. Rehabilitation interventions delivered following diagnosis have the potential to minimise the impact of disease and treatment-related symptoms, such as breathlessness and fatigue, on symptom experience, functional independence in activities of daily living and quality of life.[6] The majority of research studies focus on exercise interventions for people on curative treatment pathways.[9] Exercise was prescribed during short-term integrated rehabilitation when participants were willing to engage, but if not, other rehabilitation components, including preventive strategies were used to optimise physical activity levels and to minimise the onset of symptoms developing alongside physical deconditioning. Holistic breathlessness services improve distress relating to refractory breathlessness[10] but have not yet been tested in people mainly limited by exertional breathlessness, as reported by participants in this trial.

The trial has several strengths. Firstly, we were able to recruit and retain participants with a diverse range of demographic characteristics, disease stage and comorbidities on all thoracic cancer treatment pathways. The recruitment and retention rates indicate that the trial design and research personnel were able to address some of the barriers influencing participation in rehabilitation and exercise-based interventions in people with thoracic cancer.[16, 33] Few participants declined for reasons relating to rehabilitation being a burden or unnecessary as found in a large mixed methods study where receptiveness to rehabilitation was low across all participants with mild to severe levels of disability.[7] Most participants in this trial declined as they had too much going on or were not interested, findings congruent with previous studies. Although continuing with normal life following diagnosis is important to patients,[5] it cannot be assumed that they will want to have any

functional concerns met by health providers. Brown et al. found that people with lung cancer may prefer to access functional support from their own family.[34]

The trial has addressed uncertainties surrounding the delivery of rehabilitation interventions at a time when patients are coming to terms emotionally with their new life situation while undergoing a busy and potentially arduous oncology treatment schedule.[35] As prioritised in trial development work,[15, 16] we were able to deliver and test an inclusive, flexible and tailored model of rehabilitation to address the heterogeneity of functional need. The model of short-term integrated rehabilitation echoes effective and valued components of holistic breathlessness services. It was delivered by an expert provider, integrated with busy treatment schedules and focused on participants' immediate functional concerns and priorities.[10]

The comparability of groups at baseline indicates that the randomisation system and minimisation categories worked well. However, all mesothelioma participants were randomised to receive standard care. In view of the changing treatment options and outcomes dependent on histological diagnosis, minimising for diagnosis should be considered in a larger trial. Although the rehabilitation model was designed to be inclusive and accessible for participants with an Eastern Cooperative Oncology Group Performance Status of 0-3, it is disappointing that few disabled participants were recruited with most participants having an Eastern Cooperative Oncology Group Performance Status 0-1. The performance status of those declining to participate was not always available, so it is not known if participants with a performance status of 2-3 were approached and declined or if they were not identified during screening. Recruitment strategies in future trials need to consider how this group may be better represented in the sample, for example, asking people with a performance status of 2-3 to review trial recruitment materials and processes.

The constraints of delivering the intervention in a feasibility trial meant that integration with other oncology services was harder to achieve than was anticipated. To optimise accessibility and to reduce participant burden, we aimed to deliver the intervention alongside

scheduled hospital appointments. This presented challenges as intervention providers were based off site and scheduled hospital appointments were frequently booked or changed at short notice. Attendance at thoracic cancer multi-professional team meetings across three sites was not possible with the available trial personnel. Changes in working practices at two hospital sites meant clinic space was not always available. In addition, some participants had gaps between appointments longer than the 6-week intervention delivery period.

Delivering the intervention in the hospital setting was more achievable when participants were attending for daily radiotherapy or weekly chemotherapy. Some participants treated with surgery lived too far for home visits and had no post-operative appointments within the trial period and were therefore unable to enrol in the trial. Other integration strategies were achieved. These included providing participants with a written rehabilitation plan which they were encouraged to share with other health care professionals involved in their care. A discharge letter summarising the intervention and rehabilitation plan was sent to each participant on discharge and copied to their oncologist and case-notes.

The findings of this feasibility trial have implications for future research and clinical practice. High levels of participant retention, intervention fidelity and rates of data completeness with few non-serious adverse events suggest that the trial processes, intervention and outcome measures were acceptable to participants. Training members of the usual health care team in screening for and delivering the intervention in a full trial should improve capacity to deliver the intervention alongside scheduled appointments and reduce performance bias.

The design of a full trial should consider how to reach participants with Eastern Cooperative Oncology Group Performance Status 2 to 3 and those living further away from the recruiting centres.

Identifying a primary outcome for trials of rehabilitation interventions in this population is challenging, limited in part by the availability of suitable measures.[36] The findings suggest that the more discrete measures (physical activity and confidence) were not relevant to all participants. The inclusive approach to eligibility, heterogeneity of the population, bi-

directionality of functional trajectories, mean that the health-related quality of life measure is more able to capture the impact of the short-term integrated rehabilitation intervention and should be considered as primary outcomes for a future trial. This is in line with the findings of a recent trial of home-based rehabilitation for people with inoperable lung cancer. Edbrooke et al. observed significant improvements in health related quality of life and symptom levels with no changes in the primary outcome, physical function as measured by the 6 minute walking test.[36]

An estimated sample size for a future trial was calculated based on the observed changes in Functional Assessment of Cancer Therapy-Lung. Assuming a difference of 6 points, which is similar to the point estimate of effect we observed, and allowing for 25% attrition, 336 participants would be required to reliably detect this difference (80% power, $p=0.05$) in an efficacy trial.

If shown to be effective, this model of rehabilitation would provide proactive support around the time of diagnosis for people with thoracic cancer. It would provide a route to consecutively screen patients and shift the focus of rehabilitation towards self-management and maintaining rather than regaining function. Further, it can contribute to the integration of rehabilitation into oncology services.

This study has found that it is feasible to deliver and test a new, theory-based model of rehabilitation that differs from exercise interventions typically tested in people newly diagnosed with thoracic cancer. It is an inclusive intervention which addresses the heterogeneity of functional need and barriers to rehabilitation observed in people newly diagnosed with thoracic cancer. We were able to deliver and test the new model in a feasibility trial, and efficacy testing is now warranted.

Clinical Messages

- It is feasible to deliver and test short-term integrated rehabilitation for thoracic cancer in a randomised controlled trial in the period following diagnosis

- A sample size of 336 would be needed to detect a clinically important difference in health-related quality of life in an efficacy trial

Authors Contributions

JB, MM, IJH and AW initiated, designed and obtained funding for the trial; JB had overall responsibility for the conduct of the trial under the supervision of MM, IJH and AW; JB, LF, ED and MJT undertook data collection; JB, MM and WG designed the statistical analysis which was undertaken by JB under the supervision of MM, WG with support from VP. JB wrote the manuscript and all co-authors contributed to critically revising it.

Acknowledgements

The authors would like to thank all the people who participated in the trial and their families. We thank members of the Cicely Saunders Public and Patient Involvement Forum for their involvement during the development and conduct of this trial. In particular, we would like to thank Lesley Smith, for her insightful contributions to the design of the participant information leaflets and for her attendance and input at the research ethics committee meeting. We thank Evelyn Burssens, Cathann Manderson, Sarah Schofield, Param Kaler, Alexander Chan, Pippa Labuc, Faye Dickinson and clinicians at the three sites for their support with screening and recruitment, and Fern Brookes for supporting trial administration.

Funding

JB is funded by a National Institute for Health Research (NIHR) Clinical Doctoral Research Award (ICA-CDRF-2015-01-008) for this research project and NIHR CLAHRC South London. MM is supported by an NIHR Career Development Fellowship (CDF-2017-10-009) and NIHR CLARHC South London. IJH is an NIHR Emeritus Senior Investigator and is supported by NIHR CLARHC South London. This publication presents independent research. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

Declaration of conflicting interests

The author(s) declare that they have no conflicts of interest with respect to the research, authorship and/or publication of this article.

Data

Data will be made available on request.

References

1. McIntyre, A. and A.K. Ganti, *Lung cancer—a global perspective*. Journal of surgical oncology, 2017. **115**(5): p. 550-554.
2. Woolhouse, I., et al., *British Thoracic Society Guideline for the investigation and management of malignant pleural mesothelioma*. Thorax, 2018. **73**(Suppl 1): p. i1-i30.
3. Bayly, J.L. and M. Lloyd-Williams, *Identifying functional impairment and rehabilitation needs in patients newly diagnosed with inoperable lung cancer: a structured literature review*. Supportive Care in Cancer, 2016. **24**(5): p. 2359-2379.
4. Moore, S., L. Darlison, and A.M. Tod, *Living with mesothelioma. A literature review*. European Journal of Cancer Care, 2010. **19**(4): p. 458-68.
5. Salander, P. and S. Lilliehorn, *To carry on as before: A meta-synthesis of qualitative studies in lung cancer*. Lung Cancer, 2016. **99**: p. 88-93.
6. Alfano, C.M. and M. Pergolotti, *Next-Generation Cancer Rehabilitation: A Giant Step Forward for Patient Care*. Rehabil Nurs, 2018. **43**(4): p. 186-194.
7. Cheville, A.L., et al., *How Receptive Are Patients With Late Stage Cancer to Rehabilitation Services and What Are the Sources of Their Resistance?* Archives of Physical Medicine & Rehabilitation, 2017. **98**(2): p. 203-210.
8. Peddle-McIntyre, C.J., et al., *Exercise training for advanced lung cancer*. Cochrane Database of Systematic Reviews, 2019(2).
9. Cavalheri, V., et al., *Exercise training for people following lung resection for non-small cell lung cancer - a Cochrane systematic review*. Cancer Treatment Reviews, 2014. **40**(4): p. 585-94.
10. Brighton, L.J., et al., *Holistic services for people with advanced disease and chronic breathlessness: a systematic review and meta-analysis*. Thorax, 2018: p. thoraxjnl-2018-211589.
11. Leventhal, H., L.A. Phillips, and E. Burns, *The common-sense model of self-regulation (CSM): a dynamic framework for understanding illness self-management*. Journal of behavioral medicine, 2016. **39**(6): p. 935-946.
12. Carel, H., *Phenomenology of Illness*. First ed. 2016, Oxford: Oxford University Press.
13. Wade, D.T., *Describing rehabilitation interventions*. Clinical rehabilitation, 2005. **19**(8): p. 811-818.
14. Michie S, A.L., West R, *The Behaviour Change Wheel A Guide to Designing Interventions*. 2014, Great Britain: Silverback Publishing.
15. Bayly, J., et al., *Changing health behaviour with rehabilitation in thoracic cancer: a systematic review and synthesis*. Psychooncology, 2018.
16. Bayly, J., et al., *Developing an integrated rehabilitation model for thoracic cancer services: views of patients, informal carers and clinicians*. Pilot and Feasibility Studies, 2018. **4**(1): p. 160.
17. Oken, M.M., et al., *Toxicity and response criteria of the Eastern Cooperative Oncology Group*. American journal of clinical oncology, 1982. **5**(6): p. 649-656.
18. Miller, W.R. and S. Rollnick, *Ten things that motivational interviewing is not*. Behavioural and cognitive psychotherapy, 2009. **37**(2): p. 129-140.
19. Hawe, P., A. Shiell, and T. Riley, *Complex interventions: how “out of control” can a randomised controlled trial be?* Bmj, 2004. **328**(7455): p. 1561-1563.

20. Golsorkhi M, S.R., *Report of the Patient Research Experience Survey 2016/2017. National data collection, analysis and recommendations*. 2017, Clinical Research Network Coordinating Centre. National Institute for Health Research: London.
21. Peipert, J.D., et al., *Development and validation of the functional assessment of chronic illness therapy treatment satisfaction (FACIT TS) measures*. Quality of Life Research, 2014. **23**(3): p. 815-824.
22. Murtagh, F.E., et al., *A brief, patient- and proxy-reported outcome measure in advanced illness: Validity, reliability and responsiveness of the Integrated Palliative care Outcome Scale (IPOS)*. Palliative Medicine, 2019. **0**(0): p. 0269216319854264.
23. Granger, C.L., S.M. Parry, and L. Denehy, *The self-reported Physical Activity Scale for the Elderly (PASE) is a valid and clinically applicable measure in lung cancer*. Supportive Care in Cancer, 2015. **23**(11): p. 3211-3218.
24. Cella, D., et al., *What is a clinically meaningful change on the Functional Assessment of Cancer Therapy–Lung (FACT-L) questionnaire?: Results from eastern cooperative oncology group (ECOG) study 5592*. Journal of clinical epidemiology, 2002. **55**(3): p. 285-295.
25. van Hout, B., et al., *Interim scoring for the EQ-5D-5L: mapping the EQ-5D-5L to EQ-5D-3L value sets*. Value in health, 2012. **15**(5): p. 708-715.
26. Ritter, P.L. and K. Lorig, *The English and Spanish self-efficacy to manage chronic disease scale measures were validated using multiple studies*. Journal of clinical epidemiology, 2014. **67**(11): p. 1265-1273.
27. Beecham, J. and M. Knapp, *Costing psychiatric interventions*. Measuring mental health needs, 2001. **2**: p. 200-224.
28. Julious, S.A., *Sample size of 12 per group rule of thumb for a pilot study*. Pharmaceutical Statistics: The Journal of Applied Statistics in the Pharmaceutical Industry, 2005. **4**(4): p. 287-291.
29. Browne, R.H., *ON THE USE OF A PILOT SAMPLE FOR SAMPLE-SIZE DETERMINATION*. Statistics in Medicine, 1995. **14**(17): p. 1933-1940.
30. Michie, S., et al., *The Behavior Change Technique Taxonomy (v1) of 93 Hierarchically Clustered Techniques: Building an International Consensus for the Reporting of Behavior Change Interventions*. Annals of Behavioral Medicine, 2013. **46**(1): p. 81-95.
31. Higginson, I.J., et al., *Evaluating complex interventions in End of Life Care: the MORECare Statement on good practice generated by a synthesis of transparent expert consultations and systematic reviews*. BMC Medicine, 2013. **11**(1): p. 1-11.
32. Granger, C., et al., *Deterioration in physical activity and function differs according to treatment type in non-small cell lung cancer–future directions for physiotherapy management*. Physiotherapy, 2015.
33. Granger, C.L., et al., *Understanding factors influencing physical activity and exercise in lung cancer: a systematic review*. Supportive Care in Cancer, 2017. **25**(3): p. 983-999.
34. Brown, N.M., et al., *Supportive care needs and preferences of lung cancer patients: a semi-structured qualitative interview study*. Supportive Care in Cancer, 2015. **23**(6): p. 1533-9.
35. Ellis, J., et al., *Considerations in developing and delivering a nonpharmacological intervention for symptom management in lung cancer: the views of patients and informal caregivers*. Journal of pain and symptom management, 2012. **44**(6): p. 831-842.

36. Edbrooke, L., et al., *Multidisciplinary home-based rehabilitation in inoperable lung cancer: a randomised controlled trial*. Thorax, 2019: p. thoraxjnl-2018-212996.
37. Excellence, N.I.f.H.a.C. *Improving supportive and palliative care for adults with cancer*. 2004 5th October 2016]; Available from: <https://www.nice.org.uk/guidance/csg4>.




Figure 1 Conceptual Model of Short Term Integrated Rehabilitation for Thoracic Cancer (adapted from Wade 2005[13])




Figure 2 CONSORT flow diagram

*3 participants in the intervention group and one in the standard care group who did not return data at 30 days, did return data at 60 days. ECOG PS = Eastern Cooperative Oncology Group Performance Status.

Table 1 Participant baseline characteristics

Variables		Total N=54	Standard Care N=28	Rehabilitation N=26
Site:	1.	25 (46)	13	12
	2.	18 (33)	9	9
	3.	11 (20)	6	5
Age in years (range)		67 (44-85)	67 (44-85)	66 (46-83)
Gender	Women	19 (35)	8	11
	Men	35 (65)	20	15
Ethnicity	White British	42 (78)	25	17
	Caribbean/Indian/Bangladeshi/	5 (9)	1	4
	African	7 (13)	2	5
	Other white			
Diagnosis	NSCLC (adeno)	24 (44)	10	14
	NSCLC (squamous)	13 (24)	7	6
	SCLC	9 (17)	5	4
	Pleural mesothelioma	5 (9)	5	0
	Neuroendocrine	3 (6)	1	2
Days from diagnosis (range)		44 (2-65)	43 (2-65)	44 (3-60)
Disease by each stage	I	11 (20)	5	6
	II-IIIa	12 (22)	7	5
	IIIb	8 (15)	6	2
	IV	23 (43)	10	13
ECOG PS	0-1	47 (87)	24	23
	2-3	7 (13)	4	3
Metastases	None	27 (50)	16	11
	Local	6 (11)	3	3
	Distant	21 (39)	9	12
Comorbidities	COPD/CVD	18 (34)	7	11
	Diabetes	10 (19)	4	6
	MSK/ Inflammatory arthritis	18 (33)	10	8
Smoking History	Current	5 (9)	2	3
	Ex-smoker	40 (76)	23	17
	Never smoked	8 (15)	2	6
	Missing	1 (2)	1	0
Lives with	Alone	12 (22)	4	8
	Spouse/Partner	31 (57)	18	13
	Other family/ Other	11 (21)	6	5
Employment	Employed	7 (14)	5	2
	Sick leave	14 (27)	4	10
	Retired/not employed	31 (60)	18	13
Self-reported previous physical activity	Sedentary	13 (24)	8	5
	Mild	27 (50)	12	15
	Moderate	12 (22)	6	6
	Vigorous	2 (4)	2	0
Anti-Cancer Treatment (commenced or planned)	Surgery [†]	13 (24)	4	8
	Chemotherapy doublet	22 (41)	15	7
	Radical radiotherapy ± chemotherapy	8(15)	5	3
	Palliative chemoradiotherapy	6 (11)	2	4
	Targeted therapy/immunotherapy [‡]	4 (7)	1	3
	Palliative radiotherapy	1 (2)	0	1
Baseline Outcome Measure Scores	IPOS	19.00 [1.03-28.0]	16.5 [13.0-27.8]	20.0 [12.8-28.3]
	PASE	74.0 [42.9-117.4]	61.8 [30.9-105.7]	97.3 [52.2-170.9]
	FACT-L	93.0 [79.4-104.3]	91.5 [80.6-105.8]	94.5 [76.0-104.0]
	FACT-LCS	17.0 [14.0-23.0]	19.5 [14.0-23.8]	16.5 [13.8-20.8]
	FACT-TOI	52.0 [44.75-63.25]	50.0 [45.3-62.8]	53.5 [39.8-62.8]
	SEMCD	6.7 [5.0-8.2]	6.4 [5.1-8.5]	6.8 [5.0-8.0]
	EQ-5D-5L Index Value	0.76 [0.65-0.84]	0.77 [0.6-0.8]	0.8 [0.65-0.9]
	EQ VAS	67.5 [50.0-80.0]	70.0 [50.0-80.0]	62.5 [48.8-80.0]

Baseline characteristics: values are number (percentage) or median (IQR). NSCLC = non-small cell lung cancer; SCLC= small cell lung cancer; ECOG =Eastern Cooperative Oncology Group; COPD = Chronic Obstructive Pulmonary Disease; CVD =Cardiovascular disease; MSK = musculo-skeletal disease. [†] includes n=2 surgery & chemotherapy [‡] includes n=1 immunotherapy & chemotherapy. IPOS =Integrated Palliative Care Outcome Scale; PASE= Physical Activity Scale for the Elderly; FACT-L/TOI/LCS = Functional Assessment of Cancer Therapy- Lung/trial outcome index/lung cancer subscale; SEMCD = Self-Efficacy Measure for Chronic Disease

Table 2 Intervention Fidelity

Intervention Variable	N (%) unless stated
Rehabilitation providers oncology expertise^a	
Band 7 physiotherapist level 4	2
Band 8 physiotherapist level 3	1
Uptake and mode of rehabilitation	
Days to first session (range, target <14)	7 (5-13)
Days first to last session (range, target <42)	32 (22-45)
Sessions received	
One	25 (96)
Two	23 (88)
Three	21 (84)
Duration of sessions in minutes (range)	
1 st	60 (60-80)
2 nd	48 (35-61)
3 rd	45 (33-60)
Face to face	55 (80)
Home	37 (67)
Hospital outpatient clinic	17 (31)
In-patients	1 (2)
Telephone	14 (20)
Intervention procedures	
Symptom management strategies	20 (80)
Physical activity / fitness strategies (including exercise)	24 (96)
Task performance / participation strategies	20 (80)
Equipment provision	
Handheld fan	7(28)
Walking aids	2(8)
Perching stool	1(4)
Rehabilitation plan	
Agreed and document provided to participant	25 (100)
Shared with oncologist / multi-disciplinary team	24 (96)
Signposting	
Cancer centre rehabilitation	13 (52)
Community based exercise [‡]	5 (20)
Community based rehabilitation	3 (12)
Hospice rehabilitation	3 (12)
Palliative care team	1 (4)
Behaviour Change Techniques^b	
Goals and planning	24 (96)
Feedback and monitoring	23 (92)
Social support	17 (68)
Shaping knowledge	24 (96)
Natural consequences	25 (100)
Comparison of behaviours	23 (92)
Associations	11 (44)
Repetition and substitution	20 (80)
Comparison of outcomes	25 (100)
Regulation	12 (48)
Antecedents	9 (36)
Identity/Self Belief	21 (84)

^aBased on 'Improving supportive and palliative care for adults with cancer. The manual'. NICE 2004.[37] Pg.144-145.

[‡]includes yoga, Pilates, local cancer support organisations and walking groups.

^bcategorised and ordered according to the Behaviour Change Techniques Taxonomy version 1 (Michie 2014)

Table 3 Participant Self-Reported Outcome Data at 30 and 60 days: median (IQR)

Group	Standard Care				Rehabilitation			
	30 days N=28	Change from baseline	60 days N=28	Change from baseline	30 days N=26	Change from baseline	60 days N=26	Change from baseline
Outcome*								
IPOS total	21.0 [12-25]	-2.1 [-5.0-1.0]	18.0 [6.8 -25.0]	-2.1 [-8.0-4.0]	19 [13.5-26.5]	-0.5 [-9.0-5.5]	17.0 [8.8 -23.0]	-4.5 [-12.5-6.0]
PASE total	74.3 [32.6-111.6]	2.0 [-18.5-37.5]	105.7 [62.9-130.8]	1.6 [-10.7-50.6]	75.9 [42.3-153.7]	-4.5[-27.8-32.9]	92.9 [38.9-170.8]	-15.5 [-50.8-38.9]
FACT-L	93.5 [82.8-105.7]	1.3 [-9.7-13.7]	96.0 [78.8-111.9]	2.3[-15.0-14.5]	99.0 [75.3-105.0]	0.0[-8.0-11.0]	97.0 [80.0-103.8]	9.7 [-12.0-16.0]
FACT-LCS	18.0 [16.3-22.3]	0 [-3.0-4.0]	19.0 [14.8-23.8]	1.5 [-5.3-5.0]	19.0 [17.0-23.0]	2.0 [-0.8-5.5]	19.0 [17.0-22.0]	3.0 [-4.0-7.0]
FACT-TOI	50.1 [39.3-57.8]	-1.0 [-10.1-4.5]	53.0 [43.5-66.0]	4.5 [-13.0-9.8]	56.5 [47.0-65.0]	0.0 [-5.0-7.0]	56.2 [46.0-60.0]	6.0 [-12.0-15.0]
SEMCD	5.3 [3.7-8.5]	-0.8 [-1.9-0.7]	6.3 [4.8-8.3]	0.0 [-0.8-0.7]	6.2 [4.8-8.5]	0.3 [-0.8-1.5]	7.0 [5.8-8.3]	0.2 [-1.5-1.5]
EQ-5D-5L Index score	0.65 [0.4-0.8]	-0.09 [-0.23-0.05]	0.67 [0.6-0.9]	-0.04 [-0.18-0.15]	0.7 [0.40-0.9]	-0.03 [-0.24-0.15]	0.7 [0.64-0.8]	-0.02 [-0.28-0.07]
EQ-5D-5L VAS	60.0 [40.0-78.8]	-2.5 [-20.0-5.0]	65.0 [50.0-80.0]	-6.5 [-13.8-8.8]	65.0 [52.5-83.8]	0 [0-10.0]	70.0 [58.8-77.5]	5.0 [-17.5-24.8]
Trial satisfaction	12.0 [9.0-14.0]	n/a	n/a	n/a	12.0 [9.5-14.8]	n/a	n/a	n/a
FACIT-PS-TS	n/a	n/a	n/a	n/a	27 [27-27]	n/a	n/a	n/a
Overall satisfaction					4 [3-4]			

*Higher is better for all measures except IPOS and CSRI where lower scores are better. Median [25th -75th percentile].

IPOS =Integrated Palliative Care Outcome Scale; PASE= Physical Activity Scale for the Elderly; FACT-L/TOI/LCS = Functional Assessment of Cancer Therapy- Lung/trial outcome index/lung cancer subscale; SEMCD = Self-Efficacy Measure for Chronic Disease; FACIT-PS-TS = Functional Assessment of chronic illness therapy treatment/ patient satisfaction. †n=1 participant with 30 day measures lost in mail, completed trial satisfaction at 60 days

Appendix 1

Grade	Eastern Cooperative Oncology Group Performance Status
0	Fully active, able to carry on all pre-disease performance without restriction
1	Restricted in physically strenuous activity but ambulatory and able to carry out work of a light or sedentary nature, e.g. light housework, office work
2	Ambulatory and capable of self-care but unable to carry out work activities; up and about for more than 50% of waking hours
3	Capable of limited self-care, confined to bed or chair for more than 50% of waking hours
4	Completely disabled, cannot carry on any self-care, totally confined to bed or chair

Supplementary Table 1 Details of rehabilitation strategies and behaviour change techniques

Rehabilitation Strategies		N (%)
Symptom management strategies:	Breathlessness	15 (60)
	Fatigue	17 (68)
	Pain	8 (32)
	Nutrition/appetite	8 (32)
	Sleep	7 (28)
	Leaflet	11 (44)
Physical activity/fitness strategies:	Physical activity strategies	24 (96)
	Exercise prescription	22 (88)
	Leaflet	20 (80)
Task performance/participation strategies:	Personal ADL ^a	3 (12)
	Domestic ADL	16 (64)
	Instrumental ADL	20 (80)
Behaviour Change Techniques (BCTs) ^b		
BCT Group	BCT label	
Goals and planning	Goal setting behaviour	23 (92)
	Problem solving	19 (76)
	Goal-setting outcome	20 (80)
	Action planning	24 (96)
	Review behaviour	22 (88)
	Review outcome goals	17 (68)
Feedback and monitoring	Feedback on behaviour	23 (92)
	Self-monitoring of behaviour	21 (84)
	Self-monitoring of outcomes of behaviour	9 (36)
Social support	Unspecified	3 (12)
	Practical	14 (56)
	Emotional	10 (40)
Shaping knowledge	Instruction on how to perform behaviour	24 (96)
	Information about antecedents of behaviour	5 (20)
	Behavioural experiments	1 (4)
Natural consequences	Information about health consequences	25 (100)
	Monitoring of emotional consequences	7 (28)
	Information about emotional consequences	7 (28)
Comparison of behaviour	Demonstration of behaviour	23 (92)
Associations	Prompts and cues	11 (44)
Repetition and substitution	Behavioural practice	21 (84)
	Habits formation	13 (52)
	Graded tasks	15 (60)
Comparison of outcomes	Credible Source	25 (100)
	Comparative imagining of future outcomes	4 (16)
Regulation	Reduce negative emotions	12 (48)
Antecedents	Restructuring the physical environment	0 (0)
	Restructuring the social environment	0 (0)
	Adding objects to the environment	9 (36)
Identity/Self Belief	Framing/reframing	6 (24)
	Verbal persuasion about capability	21 (84)
	Focus on past success	2 (8)
	Self-talk	5 (20)

^a ADL = activities of daily living, ^b classified according to the Behaviour Change Techniques Taxonomy version 1 (Michie 2014)

Supplementary Table 2. Missing data at follow up

Variables	Values*	Total obtained (N)	Missing of obtained N (%)	Total obtained (N)	Missing of obtained N (%)
Patient Reported Measures	Values*	30 days		60 days	
iPOS (17 items) Lower scores = fewer symptoms	0-68	44	1 (2)	39	3 (8)
PASE (12 Items) Higher scores = more active	0-400	44	0	39	0
FACT L (34 items) Higher scores = higher QoL	0-136	44	1 (2)	39	0
FACT TOI (21 items) Higher score = higher QoL	0-84	44	1 (2)	39	0
SEMCD (6 items) Higher score = greater confidence	0-10	44	0	39	1 (3)
EQ-5D-5L (Index values) Lower score = higher QoL	0-1	44	0	39	2 (4)
EQ VAS Higher score = higher QoL	0-100	44	0	39	2 (4)
CSRI ^a - Hospital based care	Yes/no; number/information	44	6 (14)	39	7 (18)
CSRI- Community based care	Yes/no; number/duration/information	44	6(14)	39	4 (10)
CSRI-Investigations	Yes/no; number/duration/information	44	10 (23)	39	6 (15)
CSRI- Personal Care	Yes/no; number/duration/information	44	20 (45)	39	10 (26)
CSRI-Equipment	Yes/no; number/duration/information	44	3 (7)	39	2 (5)
CSRI- Medication	Yes/no; number/duration/information	44	3 (7)	39	5 (13)
Trial Satisfaction (4 items)	0-4	44	1 (6)	n/a	n/a
Treatment satisfaction (10 items)	9 items 0-3 1 item 0-4	15/25 expected	1 (6)	n/a	n/a

IPOS =Integrated Palliative Care Outcome Scale; PASE= Physical Activity Scale for the Elderly; FACT-L/TOI/LCS = Functional Assessment of Cancer Therapy- Lung/trial outcome index/lung cancer subscale; SEMCD = Self-Efficacy Measure for Chronic Disease; FACIT-PS-TS = Functional Assessment of chronic illness therapy treatment/ patient satisfaction. CSRI = Client Services Receipt Inventory. ^aDomain data is recorded missing if contact and frequency items within the domain were missing