

PAPER • OPEN ACCESS

Kerr metric from two commuting complex structures

To cite this article: Kirill Krasnov and Adam Shaw 2025 Class. Quantum Grav. 42 065013

View the article online for updates and enhancements.

You may also like

- <u>Heat capacities and thermodynamic</u> geometry in deformed Jackiw–Teitelboim
- gravity H Babaei-Aghbolagh, Mirmani Mirjalali, Davood Mahdavian Yekta et al.
- Spurious solar-wind effects on acceleration noise in LISA Pathfinder Arnold Tianyi Yang, Indie Desiderio-Sloane and Grant David Meadors
- Quantum curved tetrahedron, quantum group intertwiner space, and coherent states
 Chen-Hung Hsiao and Qiaoyin Pan

Class. Quantum Grav. 42 (2025) 065013 (31pp)

https://doi.org/10.1088/1361-6382/adb531

Kerr metric from two commuting complex structures

Kirill Krasnov* o and Adam Shaw

School of Mathematical Sciences, University of Nottingham, Nottingham NG7 2RD, United Kingdom

E-mail: kirill.krasnov@nottingham.ac.uk and ppyas15@exmail.nottingham.ac.uk

Received 9 August 2024; revised 3 January 2025 Accepted for publication 12 February 2025 Published 3 March 2025



Abstract

The main aim of this paper is to simplify and popularise the construction from the 2013 paper by Apostolov, Calderbank and Gauduchon, which (among other things) derives the Plebański–Demiański family of solutions of GR using ideas of complex geometry. The starting point of this construction is the observation that the Euclidean versions of these metrics should have two different commuting complex structures, as well as two commuting Killing vector fields. After some linear algebra, this leads to an ansatz for the metrics, which is half-way to their complete determination. Kerr metric is a special 2-parameter subfamily in this class, which makes these considerations directly relevant to Kerr as well. This results in a derivation of the Kerr metric that is self-contained and elementary, in the sense of being mostly an exercise in linear algebra.

Keywords: complex geometry, Kerr metric, Plebanski formalism

1. Introduction

Kerr metric [Ker63] is a 2-parameter stationary axisymmetric solution to the vacuum Einstein equations, describing a rotating black hole. Its original derivation was by brute force, assuming algebraic speciality and using the null tetrad formalism¹. Standard textbooks on General Relativity just state the metric and proceed to study its properties. Even a check that the Kerr metric satisfies Einstein equations is a rather non-trivial exercise, even thought doable (by hand) with the right formalism, see below. Books that do derive the Kerr metric are rare and few, and the derivation that can be found in [Cha98] makes it clear why. The derivation is

Original Content from this work may be used under the terms of the Creative Commons Attribution 4.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

^{*} Author to whom any correspondence should be addressed.

¹ See e.g. [AN16] for a description of the very interesting story behind the Kerr's original derivation.

just too complicated. One can give an 'elementary' derivation of the Kerr metric using the the Newman–Janis shift [NJ65], see the recent review [AN16] for this point of view, but this derivation has a flavour of a magic trick. Another 'elementary' derivation is based on the Kerr–Schild form of this metric, see e.g. chapter 8 of [DU18], or similar in spirit [NP12], but there remains some mystery as to why this method should be expected to work. A good further discussion on the issues surrounding the question of simplifying the derivation of the Kerr metric can be found in [BV22].

The Kerr family of solutions is a member of a more general Plebański-Demiański (PD) family [PD76]. All metrics of this family can be analytically continued to produce Euclidean signature metrics in 4D. Being an analytic continuation of a Lorentzian algebraically special metric of type D, the resulting Euclidean metrics are type D with respect to both halves of their Weyl curvature. A beautiful result of [Der83] states that a 4D Ricci-flat metric that has the property that one of its Weyl curvature halves is of type D is conformal to a Kahler metric². In particular, there is an integrable almost complex structure. This theorem can be applied with respect to both halves of the Weyl curvature, and tells us that the analytic continuation of any Lorentzian metric that solves the vacuum Einstein equations and is type D is a Riemannian 4D metric that has two distinct integrable almost complex structures. These are of two different orientations, and complex structures in 4D that are of different orientations commute. This means that the Euclidean Kerr metric is expected to have two different commuting complex structures. By assumption, it also has two commuting Killing vector fields. This is a very rigid geometric setup, and it turns out that to a very large extent the Ricci-flat Riemannian metrics of this type can be described completely. This is the viewpoint taken by the work [ACG13] that re-derived the PD family of metrics as examples of 'ambi-toric' 4D geometries, i.e. metrics that are toric with respect to two different complex structures. We remind the reader that a symplectic toric manifold of dimension 2n is a symplectic manifold endowed with a hamiltonian action of the torus $(S^1)^n$.

The main aim of the present paper is to popularise the results of [ACG13] to the gravitational physics community, and in particular re-derive the (Euclidean) Kerr metric in a rather elementary fashion. The starting point of the analysis is the (motivated above) assumption of two different commuting complex structures, together with two commuting Killing vector fields. One can then introduce a coordinate system adapted to this rigid geometric setup, as well as the corresponding set of one-forms. This leads to an ansatz for the metric. This ansatz is half-way to the derivation of the Kerr metric, and the metric results by substituting the ansatz into the vacuum Einstein equations, in more or less elementary fashion. We believe that the derivation we present is sufficiently simple that it could be included into a GR textbook. Our aim is also to streamline some of the arguments in [ACG13]. This paper is heavy in algebraic geometry, and also its aims are very different from the aim of simplifying the derivation of Kerr. Our aim here is to simplify the arguments in [ACG13] as much as possible and derive everything using as elementary manipulations as possible.

Our other aim is to put to use the Plebanski formalism [Ple77] for 4D GR. In particular, we will see that Derdziński's theorem (quoted above) is an immediate consequence of the Plebanski formalism. This makes this formalism well-adapted to the problem at hand. So, this article can be read as a rephrasal of the derivation in [ACG13] using the Plebanski formalism.

Finally, our main motivation for undertaking this study was to search for a formalism that would be best adapted to the problem of studying perturbations (in particular gravitational

² The theorem in [Der83] is more general, and is to be reviewed in the main text.

perturbations) of the Kerr metric. We believe the geometry we describe here will prove to be useful for this purpose. More remarks on this are contained in the conclusion section.

This article is organised as follows. We begin, in section 2, with a review of works leading to [ACG13]. In section 3 we describe the Plebański formalism. In particular, here we obtain a characterisation of Killing vector fields that is best adapted to this formalism. The material in this section is new. In section 4 we provide a new, Plebanski formalism based proof of the result by Derdziński that type D metrics are conformal to Kähler. We also prove another result by Derdziński, namely that type D metrics possess a Killing vector field. We review the Kerr solution in section 5, and show by an explicit computation how the two commuting complex structures arise. This section motivates our geometric assumptions about the two Killing vector fields that we need to proceed with the derivation of the metric. In section 6 we, more or less, follow [ACG13] and derive the PD family of metrics. The key point of this derivation is that there exists a natural frame consisting of vector fields that all commute with each other. The dual 1-forms are then all closed. This leads to a natural set of geometric coordinates. In terms of these coordinates, the two complex structures can be characterised explicitly, in terms of just two arbitrary functions, each of one variable. This leads to a remarkably simple ansatz for the metric. To impose Einstein equations, we use the chiral Plebanski formalism. After some analysis, Einstein equations are seen to reduce all unknowns to two quartic polynomials, each of one variable.

2. Literature review

Since the discovery of the Kerr metric [Ker63] in 1963 much effort has been put towards understanding the geometry and properties that it carries. The Kerr metric is a 2 parameter family of type D solutions to Einstein's equations. In 1969 Kinnersley parameterised all Lorentzian type D solutions [Kin69]. This was not given in a compact form and many different metrics were presented. Later in 1976 Plebański and Demiański found a single expression for the most general black hole solution [PD76, GP05]. The PD family of metrics is a 7 parameter family of solutions to the Einstein-Maxwell equations. Removing from the PD metric the electric and magnetic charge parameters, as well as the cosmological constant, we are left with a 4 parameter family of solutions. All type D spacetimes have Euclidean analogues, obtained by taking an analytic continuation. An analytic continuation of the 4-parameter PD subfamily gives a double sided type D Euclidean Ricci-flat metric, which is also asymptotically locally Euclidean (ALE). More recently, Chen and Teo generalised the Ricci-flat PD metric to an asymptotically locally flat (ALF) metric [CT15], which becomes ALE in the appropriate limit. However, not all of the Chen-Teo metrics have Lorentzian counterparts. Recently, Biguard and Gauduchon showed [BG21] that the ALF, Ricci-flat, Hermitian, toric metrics fall into one of the 4 types: Kerr, Chen-Teo, Taub-NUT and Taub-bolt.

One of the first realisations of the complex properties of type D spacetimes was through the Newman-Janis shift [NJ65]. The Newman-Janis shift links the Kerr and Schwarzschild solutions of Einstein's field equations through a complex coordinate transformation. It was noticed by Flaherty [Fla74, Fla76] that this shift was related to the existence of an integrable almost complex structure that all type D spacetimes are equipped with. Type D spacetimes were thus known to be locally Hermitian. Later, Derdziński proved [Der83], in Euclidean signature, that all one-sided type D metrics (with an extra condition implied by Einstein's equations) are conformal to Kähler. Soon after Derdziński's result, Przanowski and Baka, using this local Hermiticity, reduced the vanishing of the Ricci tensor for a type D metric to a second order PDE for a single function [PB84]. However, the PDE was written using the complex coordinates

making solutions difficult to find. In 1991 LeBrun derived a related result, with the starting point being a simple form of a Kähler metric with a U(1) symmetry [LeB91]. The metric is parameterised by 2 functions u = u(x, y, z), w = w(x, y, z) and is given by

$$g = e^{u}w\left(dx^{2} + dy^{2}\right) + wdz^{2} + w^{-1}\alpha^{2}, \quad \alpha = dt + \theta, \quad \alpha = J(wdz)$$
 (1)

where $\theta \in \Lambda^1(\mathbb{R}^3)$ and J is the complex structure. The Killing vector is ∂_t , which generates the U(1) symmetry. Vanishing scalar curvature for LeBrun's ansatz becomes the so-called $SU(\infty)$ Toda-Lattice equation

$$u_{xx} + u_{yy} + (e^u)_{zz} = 0. (2)$$

The metric is Ricci flat when $w = cu_z$ for some constant c. This ansatz covers a large class of Kähler metrics in dimension 4, however, the Toda equation is difficult to solve in general. More recently, Tod [Tod20] analysed the case of one-sided type D metrics that have an additional commuting Killing vector field. A more recent related work is [Tod24]. In this case the $SU(\infty)$ Toda equation linearises, as shown by Ward in a different context [War90]. The solutions can then be obtained from axisymmetric solutions of the flat three-dimensional Laplacian. However, even after the linearisation, deriving a general solution from equation (2) remains difficult. Apostolov, Calderbank and Gauduchon took a different approach in [ACG03, ACG13], building upon the fact that many interesting type D metrics are type D with respect to both halves of the Weyl curvature. This is also the approach we follow in the present work.

3. Chiral formalism

In this section, we will briefly describe a well known and very useful projection of the 4 dimensional Einstein action into one of its chiral halves. This leads to what can be called a chiral formalism for 4D GR. For an introduction to chiral and related formulations of General Relativity see [Kra09, Kra20]. We will focus on the Euclidean signature metrics, although the formalism generalises to Lorentzian and split signatures.

3.1. Plebanski action and field equations

The Hodge star operator on 2-form is an endomorphism, $\star:\Lambda^2\to\Lambda^2$, which squares to the identity $\star^2=\mathbb{I}$. The Hodge star splits the space of 2-forms in two eigenspaces $\Lambda^2=\Lambda^+\oplus\Lambda^-$, with eigenvalues ± 1 respectively. We call them self-dual (SD) and anti-SD 2-forms (ASD) for +1 and -1 eigenspaces respectively. Each eigenspace has dimension 3 (as dim $\Lambda^2=6$), meaning we can prescribe bases Σ^i and $\bar{\Sigma}^i$ for Λ^+ and Λ^- . The index i=1,2,3 is an $\mathfrak{so}(3)=\mathfrak{su}(2)$ index, as the Hodge star split is related to the Lie algebra decomposition $\mathfrak{so}(4)=\mathfrak{su}(2)\oplus\mathfrak{su}(2)$. The triple of 2-forms Σ^i is called an SU(2) structure, as the choice of such a triple reduces the structure group $\mathrm{GL}(4,\mathbb{R})$ to the SU(2) preserving each of the 2-forms.

The Riemann curvature can be viewed as another endomorphism acting on the space of 2-forms. It can then be decomposed with respect to the decomposition of the space of 2-forms. As is well-known, the Riemann tensor takes the following form

$$R_{\mu\nu\rho\sigma} = \begin{pmatrix} W^+ + R & \mathcal{R} \\ \mathcal{R} & W^- + R \end{pmatrix} \tag{3}$$

where W^{\pm} are the SD/ASD halves of the Weyl curvature tensor, \mathcal{R} is the traceless part of the Ricci tensor and R is the remaining trace of Ricci. The first row of equation (3) is the SD

projection of the Riemann curvature with respect to one of the pairs of its indices. It is now easy to see that the Einstein condition is equivalent to requiring the ASD part of the first row in equation (3) to be zero. This is useful because one only needs half the of curvature components to enforce the Einstein condition. The action that realises this idea and has Einstein metrics at its critical points is the Plebański action [Ple77]

$$S = \int \Sigma^{i} \wedge F^{i} - \frac{1}{2} \left(\Psi^{ij} + \frac{\Lambda}{3} \delta^{ij} \right) \Sigma^{i} \wedge \Sigma^{j}. \tag{4}$$

Here $F^i=dA^i+\frac{1}{2}\epsilon^{ijk}A^j\wedge A^k$ is the curvature of the $\mathfrak{su}(2)$ connection $A^i\in\Lambda^1\otimes\mathfrak{su}(2)$. The Lagrange multiplier Ψ^{ij} is required to be symmetric and traceless, Λ is the cosmological constant. Varying with respect to Ψ^{ij},A^i and Σ^i one obtains the following Euler–Lagrange equations

$$\Sigma^{i} \wedge \Sigma^{j} \sim \delta^{ij}$$

$$d^{A}\Sigma^{i} = d\Sigma^{i} + \epsilon^{ijk}A^{j} \wedge \Sigma^{k} = 0$$

$$F^{i} = \left(\Psi^{ij} + \frac{\Lambda}{3}\delta^{ij}\right)\Sigma^{j}.$$
(5)

The first of the above equations implies that the 2-forms describe only the 10 components of the metric plus the 3 components defining an SO(3) frame. It is useful to know that given a tetrad basis $e^I = e^0$, e^i one can construct the SD and ASD 2-forms like so

$$\Sigma^{i} = e^{0} \wedge e^{i} - \frac{1}{2} \epsilon^{ijk} e^{j} \wedge e^{k} \tag{6}$$

$$\bar{\Sigma}^i = e^0 \wedge e^i + \frac{1}{2} \epsilon^{ijk} e^j \wedge e^k. \tag{7}$$

The components of the 2-forms satisfy the quaternion algebra

$$\Sigma^{i} = \frac{1}{2} \Sigma^{i}_{\mu\nu} dx^{\mu} \wedge dx^{\nu}, \quad \Sigma^{i}_{\mu}{}^{\alpha} \Sigma^{j}_{\alpha}{}^{\nu} = -\delta^{ij} \delta^{\nu}_{\mu} + \epsilon^{ijk} \Sigma^{k}_{\mu}{}^{\nu}. \tag{8}$$

The second equation in (5) implies that the 1-form A^i corresponds to the SD part of the Levi–Civita connection for the metric defined by the Σ^i 's, see [Kra09, Kra20], and also a more recent [BK24a].

To analyse the final equation we look at the general decomposition of the SD part of the curvature

$$F^{i} = \left(\Psi^{ij} + \frac{R}{3}\delta^{ij}\right)\Sigma^{j} + R^{ij}\bar{\Sigma}^{j}.$$
 (9)

The last line in equation (5) then imposes that $R = \Lambda$ and $R^{ij} = 0$, which are precisely (vacuum) Einstein's equations. This discussion shows that the critical points of the Plebanski action are SU(2) structures that are Einstein, in the sense that the metric defined by the triple of 2-forms according to the formula

$$g(u,v)\nu_g = \frac{1}{6}\epsilon^{ijk}\iota_u \Sigma^i \wedge \iota_v \Sigma^j \wedge \Sigma^k$$
(10)

is Einstein. Here ν_g is the volume form for the metric g, and both sides of the relation are top forms.

3.2. SU(2) structures and Killing vectors

Killing vectors fields are infinitesimal symmetries of the metric $\mathcal{L}_X g_{\mu\nu} = 0$, where \mathcal{L} is the Lie derivative. For our purposes below we need to characterise Killing vectors from the point of view of SU(2) structures. Given that the metric (10) defined by a triple of 2-forms Σ^i is invariant with respect to the SO(3) action on Σ^i , it is suggestive that in order for a vector field X to be infinitesimal isometry at the level of Σ^i , the Lie derivative of Σ^i should be an SO(3) transformation

$$\mathcal{L}_{X}\Sigma^{i} = \epsilon^{ijk}\tilde{\theta}^{j}\Sigma^{k}.\tag{11}$$

We will call a vector field X satisfying this property a Σ -Killing vector field. Here $\tilde{\theta}^i$ is an arbitrary 'internal' vector.

The following lemma describes an equivalent way of stating the Σ -Killing condition

Lemma 1. Σ -Killing vectors $X \in TM$ can be equivalently described as those satisfying

$$d^A \iota_X \Sigma^i = \epsilon^{ijk} \theta^j \Sigma^k, \tag{12}$$

where d^A is the exterior covariant derivative with respect to the Σ -compatible $d^A\Sigma^i=0$ connection A^i .

Proof. To see this we use the Cartan's magic formula

$$\mathcal{L}_{X}\Sigma^{i} = \iota_{X}d\Sigma^{i} + d\iota_{X}\Sigma^{i}. \tag{13}$$

We now require this to be a gauge transformation

$$\mathcal{L}_{X}\Sigma^{i} = \epsilon^{ijk}\tilde{\theta}^{j}\Sigma^{k}. \tag{14}$$

Using $d^A \Sigma^i = 0$ and equation (13) we find

$$\mathcal{L}_{X}\Sigma^{i} = d^{A}\iota_{X}\Sigma^{i} + \epsilon^{ijk}(\iota_{X}A)^{j}\Sigma^{k} = \epsilon^{ijk}\tilde{\theta}^{j}\Sigma^{k} \quad \Rightarrow \quad d^{A}\iota_{X}\Sigma^{i} = \epsilon^{ijk}\theta^{j}\Sigma^{k}$$
 (15)

where $\theta^i = \tilde{\theta}^i - \iota_X A^i$ is the shifted SO(3) gauge transformation parameter.

Next we now link Σ -Killing vector fields with the usual Killing vector fields.

Proposition 1. A Σ -Killing vector field is a Killing vector field for the metric (10) defined by the triple of 2-forms Σ^i .

Before we prove this statement, we need to remind the reader some facts about the decomposition of $\mathfrak{su}(2)$ -valued 2-forms. This construction is from [BK24a], and we will only summarise the main points. Let us denote $E = \mathfrak{su}(2)$, and consider the space of E-valued 2-forms. We introduce an endomorphism acting on this space

$$J_2: \Lambda^2 \otimes E \to \Lambda^2 \otimes E, \quad J_2(B)^i_{\mu\nu} = \epsilon^{ijk} \Sigma^{j\ \alpha}_{[\mu} B^k_{|\alpha|\nu]}, \quad B^i_{\mu\nu} \in \Lambda^2 \otimes E.$$
 (16)

The [BK24a] shows that this operator satisfies

$$J_2(J_2-2)(J_2-1)(J_2+1)=0, (17)$$

and thus has eigenvalues 0, -1, 1, 2. The space $\Lambda^2 \otimes E$ decomposes into the eigenspaces of this operator

$$\Lambda^2 \otimes E = (\Lambda^2 \otimes E)_1 \oplus (\Lambda^2 \otimes E)_3 \oplus (\Lambda^2 \otimes E)_5 \oplus (\Lambda^2 \otimes E)_9. \tag{18}$$

The subscript in $(\Lambda^2 \otimes E)_k$ indicates the dimension of the corresponding space. The spaces $(\Lambda^2 \otimes E)_{1,3,5,9}$ are the eigenspaces of J_2 of eigenvalue 2,1,-1,0 respectively, see section 4.4 of [BK24a] for more details. It can also be shown that $(\Lambda^2 \otimes E)_9 = \Lambda^- \otimes E$. Here Λ^- is the space of anti-self dual 2-forms, with $\Lambda^2 = \Lambda^+ \oplus \Lambda^-$. We also have the following explicit projections to each of the eigenspaces

$$B_{\alpha\beta}^{(i} \Sigma^{j)\alpha\beta} \in S^{2}(E),$$

$$\epsilon^{ijk} B_{\alpha\beta}^{j} \Sigma^{k\alpha\beta} \in E,$$

$$B_{(\mu|\alpha|}^{i} \Sigma^{i\alpha}{}_{\nu)} \in S^{2}(T^{*}M).$$
(19)

The described maps project away some of the irreducible components, while keeping the others. Thus, the kernel of the first map is $(\Lambda^2 \otimes E)_{3+9}$, and so this is a projection onto the $(\Lambda^2 \otimes E)_{5+1}$ component. The second map is a projection onto the $(\Lambda^2 \otimes E)_3$ component. The last is a projection onto the $(\Lambda^2 \otimes E)_{1+9}$ component. The $(\Lambda^2 \otimes E)_1$ component, which contains multiples of Σ^i , is contained in both the first and last projection as the trace part.

Proof. We now prove the proposition. To this end, we consider the *E*-valued 2-form $d^A \iota_X \Sigma^i$, for an arbitrary vector field *X*, and compute its irreducible parts. We have

$$\left(d^{A}\iota_{X}\Sigma^{i}\right)_{\mu\nu} = \partial_{\left[\mu\right.}\left(X^{\rho}\Sigma^{i}_{\left.\left|\rho\right|\nu\right.}\right) + \epsilon^{ijk}A^{j}_{\left.\left[\mu\right.}X^{\rho}\Sigma^{i}_{\left.\left|\rho\right|\nu\right.}\right]}.$$
(20)

We can replace the partial derivative in the first term by the covariant derivative with respect to the metric, because it is applied to a 1-form, and the result is $\mu\nu$ anti-symmetrised. We thus have

$$\left(d^{A}\iota_{X}\Sigma^{i}\right)_{\mu\nu} = \nabla_{\left[\mu\right.}\left(X^{\rho}\Sigma^{i}_{\left.\left|\rho\right|\nu\right.}\right) + \epsilon^{ijk}A^{j}_{\left.\left[\mu\right.}X^{\rho}\Sigma^{i}_{\left.\left|\rho\right|\nu\right.}\right]. \tag{21}$$

We now use the fact that the connection 1-forms A^i_{μ} are the 'intrinsic torsion' components of the SU(2) structure, i.e. we have the following equation

$$\nabla_{\mu} \Sigma_{\rho\nu}^{i} + \epsilon^{ijk} A_{\mu}^{j} \Sigma_{\rho\nu}^{k} = 0. \tag{22}$$

This fact is proven in [BK24a], by considering the *E*-valued 2-forms $X^{\mu}\nabla_{\mu}\Sigma^{i}_{\rho\sigma}$, and showing that only the $(\Lambda^{2}\otimes E)_{3}$ projection is non-vanishing. This is then parametrised by $X^{\mu}A^{i}_{\mu}$. Contracting (22) with X^{ρ} and $\mu\nu$ anti-symmetrising we see that we can rewrite (21) as

$$\left(d^{A}\iota_{X}\Sigma^{i}\right)_{\mu\nu} = \Sigma^{i}_{[\mu}{}^{\rho}\nabla_{\nu]}X_{\rho}.\tag{23}$$

We can now use the algebra equation (8) of Σ 's to compute the projections of $d^A \iota_X \Sigma^i$ onto the irreducible components. We have

$$(\Lambda^2 \otimes E)_1: \quad \Sigma^{i\mu\nu} (d^A \iota_X \Sigma^i)_{\mu\nu} = 3 \nabla_\mu X^\mu$$

$$(\Lambda^{2} \otimes E)_{3} : \quad \epsilon^{ijk} \Sigma^{j\mu\nu} \left(d^{A} \iota_{X} \Sigma^{k} \right)_{\mu\nu} = -2 \Sigma^{i\mu\nu} \nabla_{[\mu} X_{\nu]}$$

$$(\Lambda^{2} \otimes E)_{5} : \quad \Sigma^{\langle i \mid \mu\nu} \left(d^{A} \iota_{X} \Sigma^{[j\rangle} \right)_{\mu\nu} = 0$$

$$(\Lambda^{2} \otimes E)_{9} : \quad \Sigma^{i}_{\langle \mu \mid} {}^{\alpha} \left(d^{A} \iota_{X} \Sigma^{i} \right)_{\alpha \mid \nu\rangle} = -3 \nabla_{\langle \mu} X_{\nu\rangle}. \tag{24}$$

Here $\langle \mu \nu \rangle$ denotes the symmetric traceless part. We now assume that the Σ -Killing equation is satisfied and so $d^A \iota_X \Sigma^i = \epsilon^{ijk} \theta^j \Sigma^k$. The right-hand side here has the only non-vanishing projection

$$\left(\Lambda^2 \otimes E\right)_3: \quad \epsilon^{ijk} \Sigma^{j\mu\nu} \left(\epsilon^{klm} \theta^l \Sigma^m\right)_{\mu\nu} = 8\theta^i. \tag{25}$$

It is then clear that the Σ -Killing equation is equivalent to

$$\nabla_{(\mu} X_{\nu)} = 0, \quad \theta^i = -\frac{1}{4} \Sigma^{i\mu\nu} \nabla_{[\mu} X_{\nu]}. \tag{26}$$

The first is the usual Killing equation, proving the proposition. The last relation gives θ^i in terms of the vector field X. In words, θ^i is the vector giving the SD projection of dX^{\sharp} , where X^{\sharp} is the one-form corresponding to the vector field X.

4. One-sided type D metrics are conformal to Kähler

The aim of this section is to present an alternative (and new) proof of the theorem by Derdziński [Der83]. We use the chiral formalism, which makes the statement almost manifest. We also prove another statement from [Der83], namely that all one-sided type D Einstein manifolds have a Killing vector.

4.1. Derdziński theorem

We start by using the chiral formalism to prove the following theorem

Theorem 1. Let (M, g) be a smooth 4-manifold that is one-sided type D, i.e. with one of the two halves of the Weyl curvature, say SD, having two coinciding eigenvalues. Assume in addition that the SD part of the Weyl curvature is divergence-free $\nabla \cdot W^+ = 0$. Then g is conformal to a Kähler metric.

This was first proved by Derdziński in [Der83]. Note that there is no assumption here that the metric is Einstein, one only assumes that the SD Weyl tensor is divergence-free. This is true for Einstein manifolds (as we will also see later), but is a weaker condition. The proof we present here is an elementary consequence of the chiral Plebanski formalism. In fact, one can motivate the usefulness of the chiral formalism by the fact that this, reasonably non-trivial statement of Riemannian geometry, becomes apparent in this formalism.

We start by translating the devergence-free condition as a condition on objects in the Plebanski formalism. As is clear from equation (3), the SD part of the Weyl curvature is directly related to the matrix Ψ^{ij} , we have $W^+_{\mu\nu\rho\sigma} = \Psi^{ij} \Sigma^i_{\mu\nu} \Sigma^j_{\rho\sigma}$. The condition that the SD Weyl curvature is divergence-free translates to

$$0 = \left(\nabla \cdot W^{+}\right)_{\nu\rho\sigma} = \nabla^{\mu}W^{+}_{\mu\nu\rho\sigma} = \nabla^{A\mu}\left(\Psi^{ij}\Sigma^{i}_{\mu\nu}\Sigma^{j}_{\rho\sigma}\right) = \nabla^{A\mu}\Psi^{ij}\Sigma^{i}_{\mu\nu}\Sigma^{j}_{\rho\sigma}, \quad (27)$$

where ∇^A_μ is the total gauge covariant derivative $\nabla^A_\mu \theta^i_\nu = \nabla_\mu \theta^i_\nu + \epsilon^{ijk} A^j_\mu \theta^k_\nu$ and A^i is the SD connection defined by $d^A \Sigma^i = 0$. As we have already remarked in (22), by construction, this covariant derivative has the property $\nabla^A_\mu \Sigma^i_{\rho\sigma} = 0$, which justifies the last equality.

The object on the right-hand side of (27) is SD with respect to the index pair $\rho\sigma$. It can therefore be converted to an *E*-valued one-form

$$\left(\nabla \cdot W^{+}\right)_{\nu \rho \sigma} \Sigma^{i \rho \sigma} = 4 \nabla^{A \mu} \left(\Psi^{i j} \Sigma^{j}_{\mu \nu}\right) = 4 \nabla^{A \mu} \left(\Psi^{i j}\right) \Sigma^{j}_{\mu \nu}. \tag{28}$$

This can be rewritten in form notations. Considering the 3-form $(\nabla^A \Psi^{ij}) \Sigma^j$, and taking the Hodge star we obtain a multiple of (28). Thus, we see that the following two statements are equivalent

$$(\nabla \cdot W^{+}) = 0 \iff d^{A}(\Psi^{ij}\Sigma^{j}) = 0. \tag{29}$$

We now have the following simple corollary of the Plebanski formalism

Proposition 2. Every Einstein 4-Manifold has divergence free SD Weyl curvature, $\nabla \cdot W^+ = 0$.

Proof of proposition 2. The proof follows from the Bianchi identity $d^A F^i = 0$. In Plebanski formalism, the Einstein equation takes the following form

$$F^{i} = \left(\Psi^{ij} + \frac{\Lambda}{3}\delta^{ij}\right)\Sigma^{j}.\tag{30}$$

Applying the exterior covariant derivative on the left, and using the Bianchi identity, we have

$$0 = d^A F^i = d^A \left(\Psi^{ij} \Sigma^j + \frac{\Lambda}{3} \Sigma^i \right) = d^A \left(\Psi^{ij} \Sigma^j \right) = 0$$
 (31)

where we have used $d^A \Sigma^i = 0$ and that Λ is constant. Using equation (29) we see that Einstein 4-manifolds have divergence free SD Weyl curvature.

Proof of theorem 1. Let us denote the eigenvalues of Ψ^{ij} as α, β, γ . We can assume, without loss of generality that $\alpha \geqslant \beta \geqslant \gamma$. The traceless condition on Ψ^{ij} imposes $\alpha + \beta + \gamma = 0$. This means that $\alpha > 0$ and $\gamma < 0$. There are then two possibilities for two eigenvalues to coincide. One is when the middle eigenvalue coincides with the negative one $\beta = \gamma$. The other is when the two positive eigenvalues coincide. Thus, the matrix Ψ^{ij} is of the two possible forms

$$\Psi^{ij} = \begin{pmatrix} 2\alpha & 0 & 0 \\ 0 & -\alpha & 0 \\ 0 & 0 & -\alpha \end{pmatrix}, \qquad \Psi^{ij} = \begin{pmatrix} \alpha & 0 & 0 \\ 0 & \alpha & 0 \\ 0 & 0 & -2\alpha \end{pmatrix} \tag{32}$$

with α being a positive real function. The two cases differ by the sign of $\det(\Psi)$. See section 2 of [Biq23] for a related discussion. We consider the first case in details. The second case is treated analogously, by observing that it can be mapped to the first by allowing α to be negative and relabelling the basis vectors. In the proof below, we never need to assume that $\alpha>0$, and so the proof covers both of the cases.

We conclude that for every one-sided type D Riemannian metric, one can always pick a basis Σ^i in the space of SD 2-forms, so that Ψ^{ij} is diagonal and of the form of the first matrix in equation (32). With this form of Ψ^{ij} the equation (29) become

$$d(2\alpha\Sigma^{1}) + A^{2}(-\alpha\Sigma^{3}) - A^{3}(-\alpha\Sigma^{2}) = 0$$

$$d(-\alpha\Sigma^{2}) + A^{3}(2\alpha\Sigma^{1}) - A^{1}(-\alpha\Sigma^{3}) = 0$$

$$d(-\alpha\Sigma^{3}) + A^{1}(-\alpha\Sigma^{2}) - A^{2}(2\alpha\Sigma^{1}) = 0.$$
(33)

Using also $d^A \Sigma^i = 0$, these equations can be rewritten as follows

$$2d\alpha \Sigma^1 + 3\alpha d\Sigma^1 = 0 \tag{34}$$

$$d\alpha \Sigma^2 + 3\alpha \left(d\Sigma^2 - A^1 \Sigma^3\right) = 0, \quad d\alpha \Sigma^3 + 3\alpha \left(d\Sigma^3 + A^1 \Sigma^2\right) = 0. \tag{35}$$

The first equation in the above can be written as

$$d\omega = 0, \quad \omega := \alpha^{\frac{2}{3}} \Sigma^{1}. \tag{36}$$

We now introduce a scaled complex linear combinations of the remaining 2-forms

$$\Omega := \alpha^{\frac{2}{3}} \left(\Sigma^2 + i \Sigma^3 \right) \quad \left(\text{resp.} \quad \bar{\Omega} := \alpha^{\frac{2}{3}} \left(\Sigma^2 - i \Sigma^3 \right) \right), \tag{37}$$

where the scaling factor in front is the same as for ω . The last two equations in equation (33) can be rewritten as

$$d\alpha \left(\Sigma^2 + i\Sigma^3\right) + 3\alpha d\left(\Sigma^2 + i\Sigma^3\right) + 3i\alpha A^1\left(\Sigma^2 + i\Sigma^3\right) = 0. \tag{38}$$

For later purposes, we note that using $d^A \Sigma^i = 0$ we can also rewrite the last two equations as

$$d\alpha \Sigma^2 - 3\alpha A^3 \Sigma^1 = 0, \qquad d\alpha \Sigma^3 + 3\alpha A^2 \Sigma^1 = 0, \tag{39}$$

and thus their complex linear combination gives

$$d\alpha \left(\Sigma^2 + i\Sigma^3\right) + 3i\alpha \left(A^2 + iA^3\right)\Sigma^1 = 0. \tag{40}$$

This will be useful later.

We now use (38) to get an equation for Ω

$$d\Omega + \left(iA^{1} - \frac{d\alpha}{3\alpha}\right)\Omega = 0. \tag{41}$$

We will rewrite this equation in a more suggestive way below. Finally, the algebraic equations $\Sigma^i \Sigma^j \sim \delta^{ij}$ imply the following equations

$$\omega\Omega = 0, \quad \Omega\Omega = 0, \quad \omega^2 = \frac{1}{2}\Omega\bar{\Omega}.$$
 (42)

We can now understand where the conformal to Kähler comes from. The second equation in (42) implies that the complex 2-form Ω is decomposable. Its two 1-form factors can be

declared to be the (1,0) forms of an almost complex structure, which defines this almost complex structure. The equation in (41) can then be rewritten in a better way. It is clear that it states

$$d\Omega + \left(iA^{1} - \frac{1}{3\alpha} d\alpha \Big|_{(0,1)} \right) \Omega = 0, \tag{43}$$

because only the projection of $d\alpha$ to the space $\Lambda^{0,1}$ survives the wedge product with Ω . We can then add another term that is valued in $\Lambda^{1,0}$ and rewrite this equation as

$$d\Omega + ia\Omega = 0, \qquad a = A^{1} + \frac{1}{3i\alpha} \left(d\alpha \Big|_{(1,0)} - d\alpha \Big|_{(0,1)} \right). \tag{44}$$

The connection 1-form a is now real-valued. We note that it can also be written as

$$a_{\mu} = A_{\mu}^{1} + \omega_{\mu}^{\nu} \partial_{\nu} \ln \left(\alpha^{\frac{1}{3}} \right). \tag{45}$$

The equation (44) implies that this complex structure is integrable. Indeed, it implies that $d\Omega$ has only a $\Lambda^{2,1}$ part, which is equivalent to integrability. The first of the equations in (42) implies that ω is in $\Lambda^{1,1}$ with respect to this complex structure. It can thus be declared to be the Kähler form. The last equation in (42) is the correct normalisation condition linking the volume form as defined by the Kähler form ω and the volume form defined by Ω . The equation (36) states that the Kähler form is closed. Altogether, we deduce that the metric defined by ω , Ω is Kähler, with Ricci form $\rho = da$. It is also clear that the original metric, defined by the Σ^{i} 's, is conformal to this Kähler metric

$$g_K = \alpha^{\frac{2}{3}} g_{\Sigma}. \tag{46}$$

This proves theorem 1.

4.2. Killing Vectors in Einstein one-sided type D spaces

Another statement proven in [Der83] is about the existence of Killing vectors in every Einstein manifold satisfying the assumptions of theorem 1.

Proposition 3. Let (M, Σ^i) be an Einstein manifold with type D SD Weyl tensor. Let Σ^i be a basis for the triple of SD 2-forms chosen so that the SD part of the Weyl curvature is diagonal, and Σ^I is in the direction of the special (non-repeated) eigenvalue. Then $X^\mu = \Sigma^{1\mu\nu} \nabla_\nu \alpha^{-\frac{1}{3}}$ is a Killing vector field. Here α with $\alpha^2 = \frac{1}{6} \operatorname{Tr} \Psi^2$ is the repeated eigenvalue of the SD Weyl tensor

Proof. Inserting *X* into Σ^1 we find

$$(\iota_{X}\Sigma^{1})_{\mu} = X^{\nu}\Sigma^{1}_{\nu\mu} = \Sigma^{1\nu\rho}\nabla_{\rho}\alpha^{-\frac{1}{3}}\Sigma^{1}_{\nu\mu} = \nabla_{\mu}\alpha^{-\frac{1}{3}} \quad \Rightarrow \quad \iota_{X}\Sigma^{1} = d\alpha^{-\frac{1}{3}} \quad (47)$$

where we have used the quaternion algebra of the 2-forms. The other two 1-forms are computed as

$$\left(\iota_{X}\Sigma^{2}\right)_{\mu} = \Sigma^{1\sigma\rho}\nabla_{\rho}\alpha^{-\frac{1}{3}}\Sigma_{\sigma\mu}^{2} = \Sigma_{\mu}^{1\sigma}\Sigma_{\sigma}^{2\rho}\nabla_{\rho}\alpha^{-\frac{1}{3}},\tag{48}$$

where we have used the algebra of Σ 's to anti-commute $\Sigma^{1,2}$. We similarly have

$$\left(\iota_{X}\Sigma^{3}\right)_{\mu} = \Sigma_{\mu}^{1\sigma}\Sigma_{\sigma}^{3\rho}\nabla_{\rho}\alpha^{-\frac{1}{3}}.\tag{49}$$

Introducing $\Sigma^+ = \Sigma^2 + i\Sigma^3$ we have

$$\left(\iota_{X}\Sigma^{+}\right)_{\mu} = \Sigma_{\mu}^{1\,\sigma}\Sigma_{\sigma}^{+\,\rho}\nabla_{\rho}\alpha^{-\frac{1}{3}}.\tag{50}$$

We now use (40), which we rewrite as

$$d\alpha^{-1/3}\Sigma^{+} = i\alpha^{-1/3}A^{+}\Sigma^{1},\tag{51}$$

where we introduced $A^+ = A^2 + iA^3$. Taking the Hodge dual we get

$$\Sigma_{\mu}^{+\rho} \nabla_{\rho} \alpha^{-1/3} = i \alpha^{-1/3} \Sigma_{\mu}^{1\rho} A_{\rho}^{+}. \tag{52}$$

Using this in (50) we get

$$(\iota_X \Sigma^+)_{\mu} = i\alpha^{-1/3} \Sigma_{\mu}^{1 \sigma} \Sigma_{\sigma}^{1 \rho} A_{\rho}^+ = -i\alpha^{-1/3} A_{\mu}^+.$$
 (53)

We now summarise the above results

$$\iota_{X}\Sigma^{1} = d\alpha^{-1/3}, \qquad \iota_{X}\Sigma^{+} = -i\alpha^{-1/3}A^{+}, \qquad \qquad \iota_{X}\Sigma^{-} = i\alpha^{-1/3}A^{-}.$$
 (54)

Having computed the components of $\iota_X \Sigma^i$, we can compute its exterior covariant derivative and use the characterisation of the Σ -Killing vectors to show that X is a Killing vector field. It will be convenient to rewrite the formulas for the exterior covariant derivative of $\iota_X \Sigma^i$ in terms of the introduced objects Σ^{\pm} , A^{\pm} . We have

$$d_{A}\iota_{X}\Sigma^{1} = d\iota_{X}\Sigma^{1} + \frac{1}{2i} \left(A^{-}\iota_{X}\Sigma^{+} - A^{+}\iota_{X}\Sigma^{-} \right),$$

$$d_{A}\iota_{X}\Sigma^{+} = d\iota_{X}\Sigma^{+} - iA^{+}\iota_{X}\Sigma^{1} + iA^{1}\iota_{X}\Sigma^{+}.$$
 (55)

Substituting (54) into the first line we see it vanishes $d_A \iota_X \Sigma^1 = 0$. For the first term this is manifest from $d^2 = 0$. For the second term there is a cancellation of the two A^+A^- terms. For the second line we have

$$d_{A}\iota_{X}\Sigma^{+} = -\mathrm{i}d\left(\alpha^{-1/3}A^{+}\right) - \mathrm{i}A^{+}d\alpha^{-1/3} + A^{1}\alpha^{-1/3}A^{+}$$
$$= -\mathrm{i}\alpha^{-1/3}\left(dA^{+} + \mathrm{i}A^{1}A^{+}\right). \tag{56}$$

What arises here is the curvature

$$F^{+} = F^{2} + iF^{3} = dA^{+} + iA^{1}A^{+}.$$
 (57)

Using the Einstein equations in the form (30) we have

$$F^{+} = \left(-\alpha + \frac{\Lambda}{3}\right) \Sigma^{+}. \tag{58}$$

This means that all in all

$$d_A \iota_X \Sigma^1 = 0, \qquad d_A \iota_X \Sigma^+ = -i\alpha^{-1/3} \left(-\alpha + \frac{\Lambda}{3} \right) \Sigma^+.$$
 (59)

On the other hand, the Σ -Killing condition with the vector $\theta^i = (\theta, 0, 0)$ becomes

$$d_A \iota_X \Sigma^1 = 0, \qquad d_A \iota_X \Sigma^+ = i\theta \Sigma^+.$$
 (60)

We see that this is indeed satisfied, with

$$\theta = \alpha^{-1/3} \left(\alpha - \frac{\Lambda}{3} \right). \tag{61}$$

This finishes the proof.

5. Two complex structures of Euclidean Kerr

The aim of this section is to exhibit the structures described in the previous section for the example of the Euclidean Kerr metric. This section serves only to motivate the considerations that follow.

5.1. Metric and chiral objects

The Euclidean Kerr metric is given by

$$g = \left(1 - \frac{2Mr}{\rho^2}\right)dt^2 + \frac{\rho^2}{\Delta}dr^2 + \rho^2d\theta^2 + \left(r^2 - a^2 - \frac{2Mra^2}{\rho^2}\sin(\theta)^2\right)\sin(\theta)^2d\phi^2 + \frac{4Mra\sin(\theta)^2}{\rho^2}dtd\phi, \tag{62}$$

where

$$\rho^2 = r^2 - a^2 \cos(\theta)^2$$
, $\Delta = r^2 - 2Mr - a^2$. (63)

This is obtained from the usual Lorentzian Kerr metric in Boyer–Lindquist coordinates through a transformation $t \to it$, $a \to ia$. A useful tetrad basis is given by

$$e^{0} = \frac{\sqrt{\Delta}}{\rho} \left(dt - a \sin(\theta)^{2} d\phi \right), \quad e^{1} = \frac{\rho}{\sqrt{\Delta}} dr, \quad e^{2} = \rho d\theta,$$

$$e^{3} = \frac{\sin(\theta)}{\rho} \left(a dt + \left(r^{2} - a^{2} \right) d\phi \right). \tag{64}$$

5.2. SD objects

The SD 2-forms are readily calculated from the equation (6) using the tetrad basis. We obtain

$$\Sigma^{1} = \left(dt - a\sin(\theta)^{2}d\phi\right) \wedge dr - \sin(\theta)d\theta \wedge \left(adt + \left(r^{2} - a^{2}\right)d\phi\right)$$
 (65)

$$\Sigma^{2} = \sqrt{\Delta} \left(dt - a \sin(\theta)^{2} d\phi \right) \wedge d\theta - \frac{\sin(\theta)}{\sqrt{\Delta}} \left(a dt + \left(r^{2} - a^{2} \right) d\phi \right) \wedge dr \qquad (66)$$

$$\Sigma^{3} = \sqrt{\Delta}\sin(\theta)\,dt \wedge d\phi - \frac{\rho^{2}}{\sqrt{\Delta}}dr \wedge d\theta. \tag{67}$$

The SD connections obtained by solving $d^A \Sigma^i = d\Sigma^i + \epsilon^{ijk} A^j \wedge \Sigma^k = 0$ are the following

$$A^{1} = \frac{M}{z_{\perp}^{2}} \left(dt - a\sin\left(\theta\right)^{2} d\phi \right) + \frac{r\cos\left(\theta\right) - a}{z_{\perp}} d\phi \tag{68}$$

$$A^{2} = -\frac{\sqrt{\Delta}\sin\left(\theta\right)}{z_{+}}d\phi\tag{69}$$

$$A^{3} = -\frac{a\sin(\theta)}{\sqrt{\Delta}z_{+}}dr + \frac{\sqrt{\Delta}}{z_{+}}d\theta \tag{70}$$

where $z_+ = r - a\cos(\theta)$. The SD curvatures can be computed (by hand!) and are given by

$$F^{1} = \frac{2M}{z_{+}^{3}} \Sigma^{1}, \quad F^{2} = -\frac{M}{z_{+}^{3}} \Sigma^{2}, \quad F^{3} = -\frac{M}{z_{+}^{3}} \Sigma^{3}.$$
 (71)

This verifies that Kerr is Ricci-flat, and also gives the components of the SD Weyl tensor $F^i=\Psi^{ij}\Sigma^j$

$$\Psi_{+}^{ij} = \begin{pmatrix} 2\alpha_{+} & 0 & 0\\ 0 & -\alpha_{+} & 0\\ 0 & 0 & -\alpha_{+} \end{pmatrix}, \quad \alpha_{+} = \frac{M}{z_{+}^{3}}.$$
 (72)

The matrix Ψ^{ij} of the SD Weyl is tracefree as expected. Note that we have just used Plebanski formalism to verify that the Kerr metric is Ricci flat. This computation is doable by hand! This once again demonstrates the usefulness and power of the Plebanski formalism.

5.3. ASD objects

The ASD 2-forms can also be calculated using equation (7) to find

$$\bar{\Sigma}^{1} = \left(dt - a\sin(\theta)^{2}d\phi\right) \wedge dr + \sin(\theta)d\theta \wedge \left(adt + \left(r^{2} - a^{2}\right)d\phi\right)$$
 (73)

$$\bar{\Sigma}^{2} = \sqrt{\Delta} \left(dt - a \sin(\theta)^{2} d\phi \right) \wedge d\theta + \frac{\sin(\theta)}{\sqrt{\Delta}} \left(adt + \left(r^{2} - a^{2} \right) d\phi \right) \wedge dr \tag{74}$$

$$\bar{\Sigma}^3 = \sqrt{\Delta}\sin(\theta)\,dt \wedge d\phi + \frac{\rho^2}{\sqrt{\Delta}}dr \wedge d\theta. \tag{75}$$

The corresponding ASD connections are

$$\bar{A}^{1} = -\frac{M}{z_{-}^{2}} \left(dt - a\sin\left(\theta\right)^{2} d\phi \right) + \frac{r\cos\left(\theta\right) + a}{z_{-}} d\phi \tag{76}$$

$$\bar{A}^2 = -\frac{\sqrt{\Delta}\sin\left(\theta\right)}{z_-}d\phi\tag{77}$$

$$\bar{A}^{3} = \frac{a\sin(\theta)}{\sqrt{\Delta}z_{-}}dr + \frac{\sqrt{\Delta}}{z_{-}}d\theta. \tag{78}$$

Here $z_{-} = r + a\cos(\theta)$. The ASD part of the curvature is

$$\bar{F}^1 = -\frac{2M}{z_-^3}\bar{\Sigma}^1, \quad \bar{F}^2 = \frac{M}{z_-^3}\bar{\Sigma}^2, \quad \bar{F}^3 = \frac{M}{z_-^3}\bar{\Sigma}^3.$$
 (79)

Clearly the ASD Weyl tensor is

$$\Psi_{-}^{ij} = \begin{pmatrix} 2\alpha_{-} & 0 & 0\\ 0 & -\alpha_{-} & 0\\ 0 & 0 & -\alpha_{-} \end{pmatrix}, \quad \alpha_{-} = -\frac{M}{z_{-}^{3}}.$$
 (80)

Note that the ASD and SD objects are related by the transformation $t \to -t$, $a \to -a$, which is the Euclidean analogue of the complex conjugation in the Lorentzian signature.

5.4. Conformal to Kähler

Looking at both halves of the Weyl curvature we can see that Euclidean Kerr is of type $D \otimes D$. Then by Derdziński's theorem we know Euclidean Kerr is conformal to two different Kähler metrics. The two conformal factors are

$$\lambda_{\pm} = \frac{1}{z_{+}} \simeq \alpha_{\pm}^{\frac{1}{3}}.\tag{81}$$

Conformally transforming the 2-forms Σ^1 and $\bar{\Sigma}^1$ we find a pair of closed 2-forms

$$\omega_{+} = \frac{1}{z_{+}^{2}} \Sigma^{1} = -dt \wedge d\left(\frac{1}{r - a\cos(\theta)}\right) + d\phi \wedge d\left(\frac{a - r\cos(\theta)}{r - a\cos(\theta)}\right) \tag{82}$$

$$\omega_{-} = \frac{1}{z_{-}^{2}} \bar{\Sigma}^{1} = dt \wedge d\left(\frac{1}{r + a\cos(\theta)}\right) - d\phi \wedge d\left(\frac{a + r\cos(\theta)}{r + a\cos(\theta)}\right). \tag{83}$$

The 2-forms ω_{\pm} are the Kähler forms of the two Kähler metrics.

5.5. Complex Structures

We can extract the complex structures from the metric and the corresponding Kähler forms. It is useful to introduce a basis for TM dual to the tetrad basis in equation (64), these are given by

$$e_{0} = \frac{r^{2} - a^{2}}{\rho \sqrt{\Delta}} \partial_{t} - \frac{a}{\rho \sqrt{\Delta}} \partial_{\phi}, \quad e_{1} = \frac{\sqrt{\Delta}}{\rho} \partial_{r}, \quad e_{2} = \frac{1}{\rho} \partial_{\theta},$$

$$e_{3} = \frac{1}{\rho} \left(a \sin(\theta) \partial_{t} + \frac{1}{\sin(\theta)} \partial_{\phi} \right). \tag{84}$$

These are dual to the tetrads in the sense that $e_I(e^J) = e_I^\mu e_\mu^J = \delta_I^J$. The complex structures can be calculated at the level of the Einstein metric as they are conformally invariant. Hence, we look for operators J_\pm that satisfy $g(\cdot, J_\pm \cdot) = \lambda_\pm^{-2} \omega_\pm$. Written in terms of tetrads and their duals this is simply

$$J_{\pm} = e^1 \otimes e_0 - e^0 \otimes e_1 \mp e^3 \otimes e_2 \pm e^2 \otimes e_3. \tag{85}$$

It is not difficult to see that $J_{\pm}^2 = -\mathbb{I}$. Substituting the expression in Boyer–Lindquist coordinates we find

$$J_{\pm} = \frac{1}{\Delta} dr \otimes \left(\left(r^2 - a^2 \right) \partial_t - a \partial_\phi \right) - \frac{\Delta}{\rho^2} \left(dt - a \sin \left(\theta \right)^2 d\phi \right) \otimes \partial_r \tag{86}$$

$$\mp \frac{\sin(\theta)}{\rho^2} \left(a dt + \left(r^2 - a^2 \right) d\phi \right) \otimes \partial_{\theta} \pm d\theta \otimes \left(a \sin(\theta) \partial_t + \frac{1}{\sin(\theta)} \partial_{\phi} \right). \tag{87}$$

We have explicitly checked that these complex structures are integrable (by computing and checking that their Nijenhuis tensor vanishes) and that they commute.

5.6. Killing vectors

The Euclidean Kerr metric is conformal to two different Kähler metrics with opposite orientations. The general statements of the previous section imply that there should be two Killing vector fields, one coming from each of the two orientations. These are given by

$$X_{+}^{\mu} = J_{+\nu}^{\mu} \left(d\lambda_{+}^{-1} \right)^{\nu}. \tag{88}$$

Plugging in the conformal factors, equation (81), we calculate the intermediate objects

$$d\lambda_{+}^{-1} = dr \pm a\cos(\theta) d\theta. \tag{89}$$

The vector fields dual to the 1-forms are given by

$$\left(d\lambda_{\pm}^{-1}\right)^{\sharp} = \frac{\Delta}{\rho^2} \partial_r \pm \frac{a \sin\left(\theta\right)}{\rho^2} \partial_{\theta}.\tag{90}$$

We get the Killing vectors by applying the complex structures

$$X_{\pm} = J_{\pm} \left(d\lambda_{+}^{-1} \right)^{\sharp} = -\partial_{t}. \tag{91}$$

Thus, we find that the two Killing vectors coming from the two orientations coincide in the case of Kerr. Thus, the Euclidean Kerr is covered by the case (iii) of proposition 11 from [ACG13]. We note that the property that these two Killing vector fields coincide is related to the fact that the Kerr Laplace operator is separable. Indeed, this property holds only when one of the parameters of the PD solution is zero $\epsilon = 0$, see below for the definition of ϵ . This is also when the Laplace operator is separable. For more on these issues see e.g. [GM15].

Even though we see that for Kerr metric there is just one Killing vector field coming from the general construction of proposition 3, there is still another Killing vector field ∂_{ϕ} . The two Killing vector fields $\xi_1 = \partial_t, \xi_2 = \partial_{\phi}$ commute, and they also span an isotropic subspace $\omega_{\pm}(\xi_1, \xi_2) = 0$ with respect to either of the Kähler forms. It is also obvious that each of these two Killing vector fields is J_{\pm} -holomorphic, in the sense that $\mathcal{L}_{\xi_1,2}J_{\pm}=0$. This is manifest in the case of the Kerr metric, because all the geometric objects constructed for it are t,ϕ independent. We will take these properties of ξ_1,ξ_2 as the starting point of the derivation of the Kerr metric in the next section.

For completeness, let us also compute the result of the action of the complex structures J_{\pm} on $\xi_{1,2}$. We have

$$J_{\pm}(\partial_t) = \frac{\Delta}{\rho^2} \partial_r \pm \frac{a \sin(\theta)}{\rho^2} \partial_{\theta}$$
 (92)

$$J_{\pm}(\partial_{\phi}) = -\frac{a\sin(\theta)^{2} \Delta}{\rho^{2}} \partial_{r} \pm \frac{\left(r^{2} - a^{2}\right)\sin(\theta)}{\rho^{2}} \partial_{\theta}.$$
 (93)

It is clear that $\xi_1, \xi_2, J(\xi_1), J(\xi_2)$ span all of TM, for each of the two complex structures. It can also be checked that all four of these vector fields are mutually commuting, for either J. The resulting frame will play an important role in the derivation of the next section.

6. Toric double-sided type D metrics

We now proceed with a derivation of the Euclidean Kerr metric. In fact, our discussion will be more general, and include the more general PD family of metrics. The PD spacetimes [PD76, GP05] are Lorentzian algebraically special (type D) spacetimes that generalise many important GR solutions. In general the PD spacetimes are solutions to the Einstein–Maxwell equations, however, here we will only consider the subclass of metrics that satisfy the vacuum Einstein equations. Our aim is to provide a 'simple' derivation of this class of metrics by building on the conformal to Kähler ideas described in the previous section. Trying to setup Einstein's equation and solve them by brute force is doable but is far from the best strategy, see [Cha98, BV22, NP12] for an example of this. Instead, we follow the approach of [ACG03, ACG13] that is based on the availability of two different commuting complex structures in these spaces.

6.1. Properties of PD spacetimes

We begin by detailing properties of the spacetimes we aim to derive. We are interested in Lorentzian signature spacetimes that are algebraically special (type D), although we will solve for their Euclidean analytic continuations. For example, to obtain a Euclidean metric from the Kerr metric one sends $t \to it$ and $a \to ia$, where t and a are the time coordinate and the angular momentum parameter respectively. The SD and ASD Weyl tensor in Lorentzian spacetimes are complex conjugates of each other, i.e. $W^+ = (W^-)^*$. This is a consequence of the fact that SD 2-forms are complex-valued objects and the ASD 2-forms are complex conjugates of their SD counterparts. Unlike the Euclidean signature case, there are no one sided type D metrics in Lorentzian signature. Thus, analytically continuing a type D Lorentzian metric to Euclidean signature we get a Riemannian metric that is type D with respect to both W^\pm .

We are looking for Einstein metrics, for which we know both W^{\pm} are divergence free. We can then resort to Derdziński's theorem 1 which implies that the metric is conformal to two Kähler metrics, in general not coinciding. We denote these two Kähler metrics by g_{\pm} , so that $g = \lambda_{\pm}^2 g_{\pm}$ where g is the Einstein metric of interest. Being conformal to two Kähler metrics implies the existence of two integrable almost complex structures J_{\pm} . Coming from the SD, ASD sectors respectively, these two complex structures have different orientations, and are not coinciding. Moreover, in 4D, two complex structures of different orientations commute. Having two commuting complex structures is an extremely strong property.

By the discussion in the previous section, one-sided type D Einstein metrics are equipped with a Killing vector, see proposition 3. In our situation of double-sided type D metrics we have two Killing vector fields of this type. However, they may coincide, which, as we have seen in the previous section, is indeed the situation in Kerr. However, the metric we want to reproduce has two commuting Killing vector fields, one related to its stationary property, the other related to it being axisymmetric. So, we will assume that there are two commuting Killing vectors ξ_1, ξ_2 . In the Euclidean setting both of these Killing vectors have compact orbits. This means that we have the two-dimensional torus \mathbb{T}^2 acting on our space by isometries. Such spaces are called toric. Moreover, in our setup with two different Kähler metrics g_{\pm} , we have the action of \mathbb{T}^2 on both g_{\pm} . This is the reason why these spaces are called ambitoric in [ACG13].

The final condition we impose is that the Killing vectors ξ_1, ξ_2 are J-holomoprhic with respect to both J_{\pm} , that is $\mathcal{L}_{\xi_1,2}J_{\pm}=0$, and that $\omega_{\pm}(\xi_1,\xi_2)=0$. We have seen that this is true in the case of Kerr, and so this is a well geometrically-motivated assumption. Having motivated the geometry of the problem, we are ready to convert the assumptions into a concrete metric ansatz.

6.2. A frame of commuting vector fields

Let us consider just one of the two complex structures for the moment. The first statement that we would like to prove is that the collection of vector fields $\xi_1, \xi_2, J(\xi_1), J(\xi_2)$ spans TM and mutually commutes. The first half of the statement follows from the fact that ξ_1, ξ_2 span an isotropic subspace $\omega(\xi_1, \xi_2) = 0$. This means that $J(\xi_1), J(\xi_2)$ is the complementary subspace.

To prove commutativity we need to use both J-holomorphicity, as well as integrability of J. Let us start by giving an equivalent way of stating J-holomorphicity. This is a standard statement in complex geometry.

Lemma 2. A vector field X is J-holomorphic $\mathcal{L}_X J = 0$ if and only if $[X, JY] = J[X, Y], \forall Y \in TM$.

Proof. We have

$$\mathcal{L}_{X}JY = (\mathcal{L}_{X}J)(Y) + J(\mathcal{L}_{X}Y) \qquad \Rightarrow \qquad (\mathcal{L}_{X}J)(Y) = [X,JY] - J[X,Y]. \quad (94)$$

This shows that $(\mathcal{L}_X J) = 0$ if and only if [X, JY] - J[X, Y] = 0 for any Y.

We can now use this to show that most of the vector fields in $\xi_1, \xi_2, J(\xi_1), J(\xi_2)$ commute.

Lemma 3. Let $[\xi_1, \xi_2] = 0$ and $\xi_{1,2}$ be *J*-holomorphic. Then

$$[\xi_1, J(\xi_1)] = 0, \quad [\xi_1, J(\xi_2)] = 0 \quad [\xi_2, J(\xi_1)] = 0 \quad [\xi_2, J(\xi_2)] = 0.$$
 (95)

Proof. This is obvious from the *J*-holomorphicity of $\xi_{1,2}$ and the previous lemma.

To prove that $J(\xi_1)$, $J(\xi_2)$ commute, we need to show that the integrability of J implies that they are also J-holomorphic.

Lemma 4. If J is integrable, and X is a J-holomorphic vector field, then J(X) is also J-holomorphic.

Proof. *J* is integrable if and only if the Nijenhuis tensor

$$N_{I}(X,Y) = [JX,JY] - J[X,JY] - J[JX,Y] - [X,Y]$$
(96)

vanishes. Let us assume that X here is J-holomoprhic. Then the second and the fourth terms annihilate each other, and the vanishing of N_J implies

$$[JX, JY] = J[JX, Y], \tag{97}$$

which is precisely the statement that JX is J-holomorphic.

Now, with both $J(\xi_1), J(\xi_2)$ being J-holomorphic, it is clear that $[J(\xi_1), J(\xi_2)] = J^2[\xi_1, \xi_2] = 0$. We have established that $\xi_1, \xi_2, J(\xi_1), J(\xi_2)$ are mutually commuting.

We end this subsection with another simple lemma.

Lemma 5. Let $\xi_{1,2,3,4}$ span TM and all commute. Let $\theta_{1,2,3,4}$ be the dual 1-forms. Then all these 1-forms are closed.

Proof. This follows from the standard formula

$$d\theta\left(X,Y\right) = X(\theta\left(Y\right)) - Y(\theta\left(X\right)) - \theta\left(\left[X,Y\right]\right), \quad \theta \in \Lambda^{1}, \quad X,Y \in TM. \tag{98}$$

Indeed, evaluating $d\theta$ for any of the 1-forms, on any pair of the vector fields $\xi_{1,2,3,4}$, we see that all the terms on the right-hand side are zero.

6.3. Construction of the frame

This subsection is central. Here we use the availability of the two commuting complex structures to find a convenient parametrisation of the 1-forms dual to $\xi_1, \xi_2, J(\xi_1), J(\xi_2)$. After such a parametrisation is obtained, one can characterise the two commuting complex structures very explicitly. Here we follow [ACG13] rather closely, and even keep some of the notations of these authors. However, unlike this reference, we try to be as elementary as possible, which means without using any algebraic geometry.

First, we note that we can always choose the frame dual to $\xi_1, \xi_2, J(\xi_1), J(\xi_2)$ to be of the form $\theta_1, \theta_2, J(\theta_1), J(\theta_2)$. Since the original vector fields commute, these 1-forms are all closed. This means that there are local coordinates such that

$$\theta_1 = d\tau, \qquad \theta_2 = d\varphi. \tag{99}$$

The condition that the other two 1-forms are closed becomes

$$dJd\tau = 0, \qquad dJd\varphi = 0. \tag{100}$$

We now come to the main statement of this subsection.

Proposition 4. There exists a choice of coordinates x, y, and of two functions F = F(x), G = G(y), such that the two commuting complex structures J_{\pm} are realised as follows

$$J_{\pm}d\varphi = \frac{1}{F}dx \pm \frac{1}{G}dy,$$

$$J_{\pm}d\tau = \frac{x}{F}dx \pm \frac{y}{G}dy,$$

$$J_{\pm}dx = -\frac{F}{y-x}(d\tau - yd\varphi),$$

$$J_{\pm}dy = \pm \frac{G}{y-x}(d\tau - xd\varphi).$$
(101)

Proof. Since $dJ_{\pm}d\tau = 0, dJ_{\pm}d\varphi = 0$, there exist coordinates ξ, η such that

$$d\xi = \frac{1}{2}(J_+ + J_-)d\varphi, \quad d\eta = \frac{1}{2}(J_+ - J_-)d\varphi.$$

$$J_{\pm}d\varphi = d\xi \pm d\eta. \tag{102}$$

The two coordinates introduced (ξ, η) form a good coordinate system together with τ, φ because both complex structures map $(d\tau, d\varphi)$ to a complementary subspace, which we now parametrised by $(d\xi, d\eta)$.

The action of the complex structures on $(d\xi, d\eta)$ gives a linear combination of $(d\tau, d\varphi)$, and can be parametrised as follows.

$$J_{\pm}d\xi = \alpha_{\pm}d\tau + \beta_{\pm}d\varphi, \quad J_{\pm}d\eta = \chi_{\pm}d\tau + \delta_{\pm}d\varphi. \tag{103}$$

Here $\alpha_{\pm}, \beta_{\pm}, \chi_{\pm}, \delta_{\pm}$ are eight at this stage arbitrary functions. However, with $\partial_{\tau}, \partial_{\varphi}$ being Killing vectors, we can assume that these functions depend on ξ, η only. Next we impose that $J_{+}^{2} = -1$ on $d\varphi$

$$J_{\pm}^{2}d\varphi = J_{\pm}d\xi \pm J_{\pm}d\eta = (\alpha_{\pm} \pm \chi_{\pm}) d\tau + (\beta_{\pm} \pm \delta_{\pm}) d\varphi = -d\varphi$$

$$\Rightarrow \quad \alpha_{\pm} \pm \chi_{\pm} = 0, \quad \beta_{\pm} \pm \delta_{\pm} = -1. \tag{104}$$

Next we enforce that J_+ and J_- commute. We have

$$J_{-}J_{+}d\varphi = J_{-}d\xi + J_{-}d\eta = (\alpha_{-} + \chi_{-})d\tau + (\beta_{-} + \delta_{-})d\varphi,$$

$$J_{+}J_{-}d\varphi = J_{+}d\xi - J_{+}d\eta = (\alpha_{+} - \chi_{+})d\tau + (\beta_{+} - \delta_{+})d\varphi.$$
 (105)

The commutativity $[J_+, J_-] = 0$, together with the conditions in equation (104), implies that the action of J_{\pm} on $d\xi, d\eta$ can be parametrised by only two functions

$$J_{\pm}d\xi = -\chi d\tau - (1+\delta) d\varphi, \qquad J_{\pm}d\eta = \pm (\chi d\tau + \delta d\varphi). \tag{106}$$

Lastly, we look at the action on $d\tau$, in general it will have the form

$$J_{\pm}d\tau = a_{\pm}d\xi + b_{\pm}d\eta$$

with 4 arbitrary functions a_{\pm}, b_{\pm} . Again we check

$$J_{\pm}^{2} d\tau = a_{\pm} J_{\pm} d\xi + b_{\pm} J_{\pm} d\eta = (-a_{\pm} \pm b_{\pm}) \chi d\tau + (-a_{\pm} (1 + \delta) \pm b_{\pm} \delta) d\varphi = -d\tau$$

$$\Rightarrow a_{\pm} = -\frac{\delta}{\chi}, \quad b_{\pm} = \mp \frac{1 + \delta}{\chi}. \tag{107}$$

Hence the action on $d\tau$ is

$$J_{\pm}d\tau = -\frac{\delta}{\chi}d\xi \mp \frac{1+\delta}{\chi}d\eta. \tag{108}$$

We have already imposed the closure $dJ_{\pm}d\varphi=0$. It remains to impose $dJ_{\pm}d\tau=0$. We get

$$0 = d(J_{\pm}d\tau) = \left[\left(\frac{\delta}{\chi} \right)_{\eta} \mp \left(\frac{1+\delta}{\chi} \right)_{\xi} \right] d\xi \wedge d\eta$$

$$\Rightarrow \left(\frac{\delta}{\chi} \right)_{\eta} = 0, \quad \left(\frac{1+\delta}{\chi} \right)_{\xi} = 0. \tag{109}$$

These equations are easily solved by introducing two functions $a=a(\xi), b=b(\eta)$ such that $\delta=-\frac{a}{a-b}, \chi=\frac{1}{a-b}$. Summarising the results so far we have

$$J_{\pm}d\varphi = d\xi \pm d\eta$$

$$J_{\pm}d\tau = ad\xi \pm bd\eta$$

$$J_{\pm}d\xi = \frac{1}{a-b}(-d\tau + bd\varphi)$$

$$J_{\pm}d\eta = \pm \frac{1}{a-b}(d\tau - ad\varphi).$$
(110)

From here we make a coordinate transformation $x = a(\xi), y = b(\eta)$. Writing $dx = a'd\xi \equiv Fd\xi, dy = b'd\eta = Gd\eta$, we rewrite the action of the complex structures as

$$J_{\pm}d\varphi = \frac{dx}{F} \pm \frac{dy}{G}$$

$$J_{\pm}d\tau = \frac{xdx}{F} \pm \frac{ydy}{G}$$

$$J_{\pm}dx = -\frac{F}{x-y} (d\tau - yd\varphi)$$

$$J_{\pm}dy = \pm \frac{G}{x-y} (d\tau - xd\varphi),$$
(111)

where F = F(x), G = G(y).

6.4. Product structure

We now consider the operator $j:=-J_+J_-$. This squares to the identity $j^2=\mathbb{I}$. It is also clear that, because both J_\pm are orthogonal (i.e. metric-compatible) complex structures, the operator j is also metric-compatible. Such an operator $j:TM\to TM: j^2=\mathbb{I}$ is known as an orthogonal product structure. The reason for this terminology is that it decomposes the (tangent and) cotangent space into the eigenspaces of eigenvalue ± 1 . It is easy to see that the subspaces of opposite eigenvalue are metric orthogonal. Indeed, let $\eta_1,\eta_2\in\Lambda^1$ be arbitrary eigenforms of j belonging to different eigenspaces

$$j\eta_{+} = +\eta_{+}, \quad j\eta_{-} = -\eta_{-}.$$
 (112)

We then have

$$(\eta_{-}, \eta_{+}) = (j\eta_{-}, j\eta_{+}) = -(\eta_{-}, \eta_{+}) = 0.$$
(113)

Here the bracket denotes the metric pairing.

Now, using one of the complex structures, say J_+ , we can introduce two more 1-forms

$$J_{+}\eta_{-}, J_{+}\eta_{+} \in \Lambda^{1}, \quad s.t. \quad j(J_{+}\eta_{-}) = -J_{+}\eta_{-}, \quad j(J_{+}\eta_{+}) = +J_{+}\eta_{+}.$$
 (114)

It is easy to see that $(J_+\eta_-,\eta_-)=0, (J_+\eta_+,\eta_+)=0$ and so the basis of 1-forms

$$\eta_{+}, J_{+}\eta_{+}, \eta_{-}, J_{+}\eta_{-}$$
 (115)

is metric-orthogonal.

6.5. Metric ansatz

Using the complex structures defined in equation (101), it is now an easy exercise to check that

$$j(dx) = dx, j(dy) = -dy. (116)$$

This means that the 1-forms

$$dx, dy, (d\tau - yd\varphi), (d\tau - xd\varphi), \tag{117}$$

where the last two 1-forms are obtained by applying e.g. J_+ to dx, dy, form a metric orthogonal basis. This means that the metric we are considering is of the form

$$g = A(d\tau - yd\varphi)^{2} + Bdx^{2} + Cdy^{2} + D(d\tau - xd\varphi)^{2}.$$
 (118)

where A, B, C, D are functions of x, y only.

However, metrics of the form (118) are not always compatible with the 2 complex structures. To require this we impose that $g(\cdot,J_{\pm}\cdot)\in\Lambda^2$. Computing the tensor $g(\cdot,J_{\pm}\cdot)$ we find

$$g(\cdot, J_{\pm}\cdot) = \frac{A(x-y)}{F} dx \otimes (dt - yd\varphi) - BF \frac{(dt - yd\varphi)}{x-y} \otimes dx$$
$$\mp \frac{D(x-y)}{G} dy \otimes (dt - xd\varphi) \pm CG \frac{(dt - xd\varphi)}{x-y} \otimes dy. \tag{119}$$

Imposing that this is a 2-form results in the following restrictions to the components of g,

$$\frac{A(x-y)}{F} = \frac{BF}{x-y}, \quad \frac{D(x-y)}{G} = \frac{CG}{x-y}.$$
 (120)

Which are most conveniently solved by introducing two new functions U = U(x, y) and V = V(x, y) so that the solutions become

$$A = \frac{FU}{(x-y)^2}, \quad B = \frac{U}{F}, \quad C = \frac{V}{G}, \quad D = \frac{GV}{(x-y)^2}.$$
 (121)

Thus, finally, the metric and the two compatible 2-forms ($\Sigma_{\pm}^1 = g(\cdot, J_{\pm}\cdot)$) are given by

$$g = FU\left(\frac{d\tau - yd\varphi}{x - y}\right)^2 + \frac{U}{F}dx^2 + \frac{V}{G}dy^2 + GV\left(\frac{d\tau - xd\varphi}{x - y}\right)^2$$
(122)

$$\Sigma_{\pm}^{1} = \frac{U}{x - v} dx \wedge (dt - yd\varphi) \mp \frac{V}{x - v} dy \wedge (d\tau - xd\varphi). \tag{123}$$

It should be appreciated how close this metric ansatz is to the Euclidean Kerr metric. We emphasise that the only information that went into the construction of this ansatz is that there are two commuting Killing vector fields, as well as two commuting complex structures. No field equations have yet been imposed.

6.6. Conformal to two Kähler metrics

We now want to impose the requirement that g is conformal to two different Kähler metrics. That is, we demand that Σ^1_{\pm} are conformal to closed 2-forms

$$d\left(\lambda_{+}^{2}\Sigma_{+}^{1}\right) = 0. \tag{124}$$

with $\lambda_{\pm} = \lambda_{\pm}(x, y)$ being the conformal factors for each Kähler metric. The Lee forms of Σ_{\pm}^1 are 1-forms, θ_{\pm} , defined by

$$d\Sigma_{\pm}^{1} = \theta_{\pm} \wedge \Sigma_{\pm}^{1}. \tag{125}$$

When Σ^1_\pm are conformal to closed 2-forms the Lee forms themselves are closed. In terms of the conformal factors we find that $\theta_\pm = -d\ln(\lambda_\pm^2)$, and hence closed. Using the 2-forms in equation (123) and solving for θ_\pm we find them to be

$$\theta_{\pm} = \left(\frac{V_x}{V} \pm \frac{U}{V(x-y)}\right) dx + \left(\frac{U_y}{U} \mp \frac{V}{U(x-y)}\right) dy, \tag{126}$$

where we write $V_x = \partial_x V$ and similarly for y. We want to impose the condition that both of these are closed 1-forms. It is more efficient to take linear combinations. Firstly, we demand

$$d(\theta_{+} + \theta_{-}) = 0 \quad \Rightarrow \quad \ln(U)_{xy} - \ln(V)_{xy} = 0$$
 (127)

which can be solved by reparameterising by new functions A = A(x), B = B(y) and H = H(x,y), such that

$$U = \frac{x - y}{H^2 A}, \quad V = \frac{x - y}{H^2 R}.$$
 (128)

The introduction of x - y here is to help simplify later formulae. The other linear combination is

$$d(\theta_{+} - \theta_{-}) = 0 \implies -(B^{2})_{y}(x - y) - 2B^{2} = (A^{2})_{x}(x - y) - 2A^{2}.$$
 (129)

Taking derivatives twice with respect to x we get $(A^2)_{xxx} = 0$. Similarly taking the derivative with respect to y twice we get $(B^2)_{yyy} = 0$. This means that these equations can be solved by A^2 and B^2 being quadratic polynomials in their respective variables, with arbitrary coefficients. Further, substituting the quadratic ansatz for A^2 , B^2 back into equation (129) we find that their coefficients coincide, that is

$$A^{2} = R(x), \quad B^{2} = R(y), \quad R(z) = r_{0} + r_{1}z + r_{2}z^{2}.$$
 (130)

We thus find that, given the two functions F(x), G(y) and the 3 constants r_0 , r_1 , r_2 defining A(x), B(y), the metric

$$g = \frac{1}{H^2} \left[\frac{F}{A(x-y)} (d\tau - yd\varphi)^2 + \frac{x-y}{FA} dx^2 + \frac{x-y}{BG} dy^2 + \frac{G}{B(x-y)} (d\tau - xd\varphi)^2 \right]$$
(131)

is an ambitoric metric that is conformal to two Kähler metrics with opposite orientations. This ansatz for the metric is half-way to determining the Kerr metric.

6.7. Solving for the conformal factor

We are interested in a metric that is Einstein, so we now impose Einstein's equations. The easiest way to do this is using the chiral formalism explained in section 3. The derivation we present here is different from that in [ACG13], and, we hope, more straightforward. From equation (131) we choose a convenient basis for the frame

$$e^{0} = \frac{1}{H} \sqrt{\frac{F}{A(x-y)}} \left(d\tau - y d\varphi \right), e^{1} = \frac{1}{H} \sqrt{\frac{x-y}{FA}} dx, e^{2} = \frac{1}{H} \sqrt{\frac{x-y}{BG}} dy,$$

$$e^{3} = \frac{1}{H} \sqrt{\frac{G}{B(x-y)}} \left(d\tau - x d\varphi \right).$$
(132)

Using equation (6) we can build the SD 2-forms

$$\Sigma^{1} = \frac{1}{H^{2}} \left(\frac{1}{A} \left(d\tau - y d\varphi \right) \wedge dx - \frac{1}{B} dy \wedge \left(d\tau - x d\varphi \right) \right)$$
(133)

$$\Sigma^{2} = \frac{1}{H^{2}} \left(\sqrt{\frac{F}{ABG}} \left(d\tau - y d\varphi \right) \wedge dy - \sqrt{\frac{G}{ABF}} \left(d\tau - x d\varphi \right) \wedge dx \right)$$
(134)

$$\Sigma^{3} = \frac{1}{H^{2}} \left(\sqrt{\frac{FH}{AB}} d\varphi \wedge d\tau - \frac{x - y}{\sqrt{ABF}} dx \wedge dy \right). \tag{135}$$

The SD connection, obtained by solving $d^A \Sigma^i = d\Sigma^i + \epsilon^{ijk} A^j \wedge \Sigma^k = 0$, in this basis becomes

$$A^{1} = \left(\frac{BF}{A} - \frac{F^{2}}{AH^{2}} \left(\frac{(x-y)AH^{2}}{F}\right)_{x}\right) \frac{d\tau - yd\varphi}{2(x-y)^{2}} - \left(\frac{AG}{B} + \frac{G^{2}}{BH^{2}} \left(\frac{(x-y)BH^{2}}{G}\right)_{y}\right) \frac{d\tau - xd\varphi}{2(x-y)^{2}}$$

$$A^{2} = \sqrt{\frac{FG}{AB}} \left[\frac{AH_{x} - BH_{y}}{H(x-y)} d\tau + \frac{H(A-B) + 2ByH_{y} - 2AxH_{x}}{H(x-y)} d\varphi\right]$$

$$A^{3} = \frac{1}{\sqrt{AB}} \left[\sqrt{\frac{F}{G}} \frac{H(B-A) + 2B(x-y)H_{y}}{2H(x-y)} dx + \sqrt{\frac{G}{F}} \frac{H(A-B) - 2A(x-y)H_{x}}{2H(x-y)} dy\right]. \tag{136}$$

The curvatures are given by rather long expressions. This is the only place where we needed to resort to algebraic manipulation software. We state the curvature 2-forms by decomposing them into their SD and anti-self dual components. We write

$$F^{i} = M^{ij}\Sigma^{j} + R^{ij}\bar{\Sigma}^{j}. \tag{137}$$

The nonzero components of M^{ij} are

$$M^{11} = \frac{AH^2}{24} \left(\frac{F \log(A)_x}{x - y} \right)_x - \frac{AH^3}{12(x - y)} \left(F \left(\frac{1}{H} \right)_x \right)_x - \frac{AH^2}{24} \left(\frac{F}{x - y} \right)_{xx} - \frac{AFH^2}{48(x - y)^3}$$

$$+ \frac{BH^2}{24} \left(\frac{G \log(B)_x}{x - y} \right)_y - \frac{BH^3}{12(x - y)} \left(G \left(\frac{1}{H} \right)_y \right)_y - \frac{BH^2}{24} \left(\frac{G}{x - y} \right)_{yy} - \frac{BGH^2}{48(x - y)^3}$$

$$+ \frac{ABH^2}{12} \left[\left(\frac{G}{B(x - y)^2} \right)_y - \left(\frac{F}{A(x - y)^2} \right)_x \right] - \frac{H^2}{16(x - y)^3} \left(\frac{A^3G + B^3F}{AB} \right) \qquad (138)$$

$$M^{22} = M^{33} = \frac{ABH^2}{24} \left(\frac{F}{A(x - y)^2} \right)_x - \frac{ABH^2}{24} \left(\frac{G}{B(x - y)^2} \right)_y + \frac{H^2}{48(x - y)^3} \left(\frac{A^3G + B^3F}{AB} \right)$$

$$- \frac{AH^3}{12(x - y)} \left(F \left(\frac{1}{H} \right)_x \right)_x - \frac{AFH^2}{48(x - y)^3} - \frac{AH^2}{24} \left(\frac{F}{(x - y)^2} \right)_x$$

$$- \frac{BH^3}{12(x - y)} \left(G \left(\frac{1}{H} \right)_y \right)_y - \frac{BGH^2}{48(x - y)^3} + \frac{BH^2}{24} \left(\frac{G}{(x - y)^2} \right)_y \qquad (139)$$

$$M^{12} = M^{21} = -\sqrt{\frac{FG}{AB}} \frac{H^2}{24(x - y)^3} \left(\frac{x - y}{2} \left(A^2 \right)_x - A^2 + \frac{x - y}{2} \left(B^2 \right)_y + B^2 \right). \qquad (140)$$

We remind that one of the Einstein equations states that the trace of the matrix M^{ij} coincides with the cosmological constant $M^{11} + M^{22} + M^{33} = \Lambda$. The components of R^{ij} are

$$R^{11} = \frac{AFH}{12} \left(\frac{H}{x-y} \right)_{xx} - \frac{AH^2}{24} \left(\frac{F}{x-y} \right)_{xx} + \frac{AHF_xH_x}{12(x-y)} + \frac{AH^2}{24} \left(\frac{F\log(A)_x}{x-y} \right)_x - \frac{7AFH^2}{48(x-y)^3}$$

$$- \frac{BGH}{12} \left(\frac{H}{x-y} \right)_{yy} - \frac{BH^2}{24} \left(\frac{G}{x-y} \right)_{yy} + \frac{BHG_yH_y}{12(x-y)} + \frac{BH^2}{24} \left(\frac{G\log(B)_y}{x-y} \right)_y - \frac{7BGH^2}{48(x-y)^3}$$

$$+ \frac{H^2}{16(x-y)^3} \left(\frac{B^3F - A^3G}{AB} \right)$$

$$R^{22} = -\frac{FH(AH_x)_x}{24(x-y)} + \frac{FH^2A_x}{24(x-y)^2} - \frac{AFH^2}{48(x-y)^3} + \frac{GH(BH_y)_y}{24(x-y)} + \frac{GH^2B_y}{24(x-y)^2} - \frac{BGH^2}{48(x-y)^3}$$

$$+ \frac{H^2}{48(x-y)^3} \left(\frac{B^3F - A^3G}{AB} \right)$$

$$R^{33} = -\frac{FH(AH_x)_x}{24(x-y)} + \frac{FH^2A_x}{24(x-y)^2} - \frac{AFH^2}{48(x-y)^3} - \frac{GH(BH_y)_y}{24(x-y)} - \frac{GH^2B_y}{24(x-y)} + \frac{BGH^2}{48(x-y)^3}$$

$$- \frac{H^2}{48(x-y)^3} \left(\frac{B^3F - A^3G}{AB} \right)$$

$$R^{12} = \sqrt{\frac{FG}{AB}} \frac{H}{12(x-y)^2} \left[2AB(2(x-y)H_{xy} + H_x - H_y) + \left(2B^2H_y - \frac{H}{2} \left(B^2 \right)_y \right)$$

$$-2A^2H_x + \frac{H}{2} \left(A^2 \right)_x \right).$$

$$(143)$$

$$R^{21} = \sqrt{\frac{FG}{AB}} \frac{H}{12(x-y)^2} \left[2AB(2(x-y)H_{xy} + H_x - H_y) - \left(2B^2H_y - \frac{H}{2} \left(B^2 \right)_y \right)$$

$$-2A^2H_x + \frac{H}{2} \left(A^2 \right)_x \right).$$

We find it useful to first analyse the equations imposed by $R^{12} = 0$, $R^{21} = 0$. Taking the sum and difference of these equations we get

$$2(x - y)H_{xy} + H_x - H_y = 0 ag{146}$$

$$2A^{2}H_{x} - \frac{H}{2}(A^{2})_{x} - 2B^{2}H_{y} + \frac{H}{2}(B^{2})_{y} = 0.$$
 (147)

To solve the first equation we introduce a coordinate substitution t, r = (x + y, x - y) such that is becomes

$$H_{tt} = r \left(\frac{1}{r} H_r\right)_r. \tag{148}$$

Looking for separable solutions of the form H = T(t) + R(r) we obtain the following class of solutions

$$H(t,r) = c_0 r^2 \ln(r) + c_3 r^2 + c_0 t^2 + c_1 t + c_2.$$
(149)

Another class of solutions is obtained by taking H = H(z) and $z = \sqrt{t^2 - r^2}$ the equation reduces to

$$H_{zz} = 0 ag{150}$$

which has a solution of the form

$$H(t,r) = c_4 \sqrt{t^2 - r^2} + \text{const.}$$
 (151)

As the original PDE is linear in H, we can add these solutions. We note that other separable solutions exist in terms of Bessel functions, but these will not be relevant. Substituting the solution into equation (147) we can restrict the allowed values of the constants c_i and r_i , see equation (130) for the definition of the latter. In doing so we find 3 different solutions for H(x, y) and R(z)

1)
$$H(x,y) = 1 + \varepsilon \sqrt{xy}$$
, $R = z$
2) $H(x,y) = \sqrt{xy}$, $R = z^2 + 2\varepsilon z$
3) $H(x,y) = 1 + \varepsilon(x+y)$, $R = 1$ (152)

where we have renamed some constants for later convenience. In the main text we will focus on case 1) as this gives the family that contains the PD metrics. The other cases are described in [ACG13] and briefly in appendix. In case 1), we have that the solutions for H, H, H are

$$H = 1 + \varepsilon \sqrt{xy}, \quad A(x) = \sqrt{x}, \quad B(y) = \sqrt{y}.$$
 (153)

We now perform another change of variables to remove the square roots. We take $x = r^2$, $y = q^2$, taking the positive branch of the square root. The metric in these variables becomes

$$g = \frac{1}{(1 + \varepsilon rq)^2} \left(\frac{C}{r^2 - q^2} \left(d\tau - q^2 d\varphi \right)^2 + \frac{r^2 - q^2}{C} dr^2 + \frac{r^2 - q^2}{D} dq^2 + \frac{D}{r^2 - q^2} \left(d\tau - r^2 d\varphi \right)^2 \right)$$
(154)

where we have introduced C(r) = F/r, D(q) = G/q.

6.8. PD family of solutions

Having chosen a solution for the conformal factor function H^2 , we now calculate the Ricci scalar condition $M^{11} + M^{22} + M^{33} = \Lambda$. The components of the matrix M^{ij} for the metric equation (154) are as follows

$$M^{11} = \frac{1}{6} \frac{(1 + \varepsilon rq)^{2} (r + q)^{3}}{q^{2} - r^{2}} \left[\left(\frac{C}{(r + q)^{3}} \right)_{rr} + \left(\frac{D}{(r + q)^{3}} \right)_{qq} \right] + \frac{(1 + \varepsilon rq)^{5}}{12 (q^{2} - r^{2})} \left[\left(\frac{C}{(1 + \varepsilon rq)^{3}} \right)_{rr} + \left(\frac{D}{(1 + \varepsilon rq)^{3}} \right)_{qq} \right],$$

$$M^{22} = M^{33} = -\frac{1}{12} \frac{(1 + \varepsilon rq)^{2} (r + q)^{3}}{q^{2} - r^{2}} \left[\left(\frac{C}{(r + q)^{3}} \right)_{rr} + \left(\frac{D}{(r + q)^{3}} \right)_{qq} \right] + \frac{(1 + \varepsilon rq)^{5}}{12 (q^{2} - r^{2})} \left[\left(\frac{C}{(1 + \varepsilon rq)^{3}} \right)_{rr} + \left(\frac{D}{(1 + \varepsilon rq)^{3}} \right)_{qq} \right].$$

$$(156)$$

Taking the trace results in the following equation

$$12\varepsilon^{2}\left(q^{2}C+r^{2}D\right)-6\varepsilon\left(1+\varepsilon rq\right)\left(qC_{r}+rD_{q}\right)+\left(1+\varepsilon rq\right)^{2}\left(C_{rr}+D_{qq}\right)=4\left(q^{2}-r^{2}\right)\Lambda. \tag{157}$$

By taking derivatives of both sides 3 times with respect to r and q gives

$$C_{rrrr} = 0, \quad D_{aaaa} = 0. \tag{158}$$

This shows that C and D are now 4th order polynomials of their respective variables. The only remaining nonzero component of the traceless Ricci tensor is then

$$R^{11} = (r^2 - q^2) (1 + \varepsilon rq) D_{qq} + (4q + 6\varepsilon rq^2 - 2\varepsilon r^3) D_q - 4(1 + 3\varepsilon rq) D - (r^2 - q^2) (1 + \varepsilon rq) C_{rr} + (4r + 6\varepsilon r^2 q - 2\varepsilon q^3) C_r - 4(1 + 3\varepsilon rq) C.$$
 (159)

This allows to further restrict C,D. We substitute a generic quartic polynomial ansatz for C,D into $M^{11} + M^{22} + M^{33} = \Lambda$ and $R^{11} = 0$. This gives the following solution

$$C = -b + 2mr + r^2 + 2n\varepsilon r^3 - (\varepsilon^2 b + \Lambda/3) r^4$$
(160)

$$D = b + 2nq - q^2 + 2m\varepsilon q^3 + (\varepsilon^2 b + \Lambda/3)q^4. \tag{161}$$

We have use the freedom of rescaling $r, q \to \lambda r, \lambda q$ to set the coefficients in front of $r^2, -q^2$ to unity.

We have thus derived the Euclidean version of the PD metrics with their 5 parameters, $b, m, n, \varepsilon, \Lambda$ [PD76, GP05]. This is the same result that was obtained in [ACG13], however the approach here was, we believe, more elementary, without any need for algebraic geometry considerations.

To check that the metric we obtained is indeed one sided type $D \otimes D$ we can calculate both parts of the Weyl curvature. We get

$$\Psi = \begin{pmatrix} 2\alpha & 0 & 0 \\ 0 & -\alpha & 0 \\ 0 & 0 & -\alpha \end{pmatrix}, \quad \alpha = \frac{n-m}{6} \left(\frac{1+\varepsilon rq}{r+q} \right)^{3},$$

$$\bar{\Psi} = \begin{pmatrix} 2\bar{\alpha} & 0 & 0 \\ 0 & -\bar{\alpha} & 0 \\ 0 & 0 & -\bar{\alpha} \end{pmatrix}, \quad \bar{\alpha} = \frac{n+m}{6} \left(\frac{1+\varepsilon rq}{r-q} \right)^{3}.$$
(162)

Clearly both halves of the Weyl tensor are of type D. When $m = \pm n$ one of the sides of the Weyl curvature vanishes, in this case the spacetimes are called SD PD [NH15].

6.9. Kerr metric

To obtain the Kerr metric from the more general PD family of metrics we set $\Lambda=0$. We want the metric to be asymptotically flat, requires that $g_{rr}\to 1$ as $r\to\infty$. For this to be the case the polynomial C(r) must be at most quadratic, which necessitates $\varepsilon=0$. Now both C,D are simple quadratic functions

$$C(r) = r^2 - 2Mr - a^2$$
, $D(q) = a^2 - q^2$, (163)

where we renamed the constant terms suggestively, and also set the NUT charge (vanishing in Kerr) n=0. Now $C(r)=\Delta$ of the Kerr metric, and setting $q=a\cos\theta$ makes $D=a^2\sin^2\theta$. These choices can be motivated using the requirement that the metric is an appropriate analytic continuation of a Lorentzian signature metric, and that the Lorentzian metric is axi-symmetric, but we will not attempt this here. The main claim we are making is that the Euclidean Kerr metric is a member of a large family of metrics whose determination reduces to an exercise in linear algebra, determining the metric ansatz, following by a straightforward computation of curvatures using the Plebanski formalism.

With these choices the metric equation (154) becomes

$$g = \frac{\Delta}{\rho^2} \left(d\tau - a^2 \cos^2 \theta d\varphi \right)^2 + \frac{\rho^2}{\Delta} dr^2 + \rho^2 d\theta^2 + \frac{\sin^2 \theta}{\rho^2} \left(a d\tau - a r^2 d\varphi \right)^2. \tag{164}$$

Now further changing the coordinates

$$d\varphi = -\frac{1}{a}d\phi, \qquad d\tau = dt - ad\phi \tag{165}$$

gives the usual form of the Kerr metric in Boyer–Lindquist coordinates, with the frame given by equation (64). The $\Lambda \neq 0$ Kerr metric is another member of the PD family, but we have not attempted to exhibit it explicitly.

7. Conclusion

We have presented an alternative derivation of the (Euclidean) Kerr metric, one that proceeds by deriving the more general PD family. The derivation we described is 'elementary' in the sense that the most non-trivial step of the construction, which is establishing the ansatz (123), proceeds by a calculation in linear algebra. Thus, one gets rather far without imposing any differential equations whatsoever. Some differential equations are imposed at the next step, which requires that the metric is conformal to two different Kähler metrics. This leads to the metric ansatz (131). The rest of the analysis consists in imposing Einstein equations. This is seen to reduce the arbitrariness to two quartic polynomials, each of one variable. One can then recognise the Euclidean Kerr metric in the obtained family of solutions without difficulty. It corresponds to the case of quadratic polynomials.

Our main reason to be interested in the geometry we described is the fact that the (Euclidean) Kerr metric exhibits the beautiful and rich geometry if viewed as a complex 4D manifold. This is of course also one of the motivations of [ACG13]. It can also be seen from the derivation we presented that the Plebanski chiral formalism is best-suited for viewing 4D Euclidean geometry via the prism of complex geometry. Most of what we described is not new for mathematicians, apart from our new characterisation of Killing vector fields in section 3.2 and the new proof of Derdziński theorem in section 4. Some of the reasoning in section 6, in particular in the part where we impose Einstein equations, is also different from [ACG13]. But our main motivation in describing these results is our hope that the adopted here point of view of the Plebanski formalism will make these beautiful ideas more digestible for the gravitational physicists.

Our final remark is that it is very likely that the described here rich ambi-Kähler and ambitoric geometry of the Kerr metric is useful for the problem of gravitational perturbations of the Kerr BH. It is well-known that the problem of scalar, vector and tensor perturbations in the Kerr background is separable, see e.g. [Cha98]. This is usually achieved by considering the Teukolsky equations. On the other hand, it is also well-known that the separability of the Kerr metric wave equation is due to the existence of a non-trivial symmetric rank 2 Killing tensor.

This Killing tensor can be constructed as the square of the anti-symmetric rank 2 Killing-Yano tensor. All this is intimately related to the construction described in this paper, as is explained in [GM15]. Indeed, in this reference the authors show that the existence of a Killing 2-form in 4D is directly related to the metric being ambi-Kähler. All in all, the described here geometry is known to be directly related to the separability of the Kerr wave equation. It thus appears likely that there exists a formalism for gravitational perturbation theory in Kerr that is based on the Plebanski formalism, and the geometry described in this paper. This could be distinct from that given by the Teukolsky equation, and potentially more useful. The basic point is that the Teukolsky equation reduces the problem of describing gravitational perturbations of Kerr to a single scalar. This scalar is an appropriate potential for the metric. It is possible that the geometry we described can lead to a different choice of a potential for the metric. In particular, the pure connection formalism for 4D GR, see [Kra11] may be useful here, in that a potential for the connection rather than the metric may give a more powerful description. We hope to return to such questions in a future publication.

Data availability statement

No new data were created or analysed in this study.

Appendix. Alternative cases solution

Here we briefly explore the Einstein metrics that are solutions to the other cases in equation (152).

A.1. Case 2

In case 2 we can make a slight change of coordinates $r = x + \varepsilon, q = y + \varepsilon$ such that

$$H = \sqrt{(r-\varepsilon)(q-\varepsilon)}, \quad A = \sqrt{r^2 - \varepsilon^2}, \quad B = \sqrt{q^2 - \varepsilon^2}.$$
 (166)

We redefine the functions in the metric C = FA, D = GB such that the metric becomes

$$g = \frac{1}{(r-\varepsilon)(q-\varepsilon)} \left(C \frac{\left(d\tau - (q-\varepsilon)d\phi\right)^2}{\left(r^2 - \varepsilon^2\right)(r-q)} + \frac{r-q}{C} dr^2 + \frac{r-q}{D} dq^2 + D \frac{\left(d\tau - (r-\varepsilon)d\phi\right)^2}{(r-q)(q^2 - \varepsilon^2)} \right). \tag{167}$$

The functions C,D are then quartic polynomials, in which case the Einstein's equations are satisfied when

$$C(r) = -\varepsilon \left(4\Lambda + m\varepsilon^2 + n\varepsilon^3\right) + m\varepsilon^2 r + \frac{2n\varepsilon^3 + m\varepsilon^2 + 4\Lambda}{\varepsilon} r^2 - mr^3 - nr^4$$
(168)

$$D(q) = -C(q) \tag{169}$$

where $m, n, \Lambda, \varepsilon$ are 4 parameters. This gives another class of ambitoric Einstein metrics.

A.2. Case 3

For case 3 we do not need a change of coordinates and can remain in x, y. Similarly to the main text we require the trace of the Ricci tensor to be equal to the cosmological constant,

this becomes

$$(1 + \varepsilon(x + y))^{2} (F_{xx} + G_{yy}) - 6\varepsilon (1 + \varepsilon(x + y)) (F_{x} + G_{y}) + 12\varepsilon^{2} (F + G) = 24(y - x) \Lambda$$
(170)

taking derivatives w.r.t x 3 times and alternatively with y 3 times we obtain the equations

$$F_{xxxx} = 0, \quad G_{yyyy} = 0 \tag{171}$$

which have solutions such that F,G are quartic polynomials in their respective variables. Substituting this back into equation (170) and solving for the constants reveals that

$$F(x) = b + mx + \gamma x^{2} + \left(2\gamma\varepsilon - 2m\varepsilon^{2} - 4\Lambda\right)x^{3} + \frac{\varepsilon\left(2\gamma\varepsilon - 2m\varepsilon^{2} - 4\Lambda\right)}{2}x^{4}$$
 (172)

$$G(y) = -F(y). (173)$$

where $b, m, \gamma, \varepsilon, \Lambda$ are arbitrary constants. This solution for F, G automatically satisfies $R_{\mu\nu} = \frac{\Lambda}{4} g_{\mu\nu}$. The Einstein metric is of the form

$$g = \frac{1}{(1 + \varepsilon(x + y))^2} \left[\frac{F}{x - y} (d\tau - yd\phi)^2 + \frac{x - y}{F} dx^2 + \frac{x - y}{G} dy^2 + \frac{G}{x - y} (d\tau - xd\phi)^2 \right].$$
(174)

ORCID ID

Kirill Krasnov (b) https://orcid.org/0000-0003-2800-3767

References

- [ACG03] Apostolov V, Calderbank D M J and Gauduchon P 2003 The geometry of weakly self-dual Kähler surfaces Compos. Math. 135 279–322
- [ACG13] Apostolov V, Calderbank D M J and Gauduchon P 2016 Ambitoric geometry I: Einstein metrics and extremal ambi-Kähler structures J. Reine Angew. Math. 721 109–47
 - [AN16] Newman E T and Adamo T 2014 The Kerr-Newman metric: a review Scholarpedia 9 31791
- [BG21] Biquard O and Gauduchon P 2023 On toric Hermitian ALF gravitational instantons Commun. Math. Phys. 399 389–422
- [Biq23] Biquard O, Gauduchon P and LeBrun C 2024 Gravitational instantons, Weyl curvature and conformally Kähler geometry Int. Math. Res. Not. 2024 13295–311
- [BK24a] Bhoja N and Krasnov K 2024 SU(2) structures in four dimensions and Plebanski formalism for GR (arXiv:2405.15408 [math.DG])
- [BL67] Boyer R H and Lindquist R W 1967 Maximal analytic extension of the Kerr metric J. Math. Phys. 8 265–81
- [BV22] Baines J and Visser M 2022 Physically motivated ansatz for the Kerr spacetime Class. Ouantum Grav. 39 235004
- [Cha98] Chandrasekhar S 1998 The Mathematical Theory of Black Holes (Oxford University Press)
- [CT15] Chen Y and Teo E 2015 A five-parameter class of solutions to the vacuum Einstein equations Phys. Rev. D 91 124005
- [Der83] Derdzinski A 1983 Self-dual Kähler manifolds and Einstein manifolds of dimension four Compos. Math. 49 405–33
- [DU18] Deruelle N and Uzan J P 2018 Relativity in Modern Physics (Oxford University Press)
- [Fla74] Flaherty E J 1974 An integrable structure for type D spacetimes *Phys. Lett.* A 46 391–2

- [Fla76] Flaherty E J 1976 Hermitian and Kählerian Geometry in Relativity (Springer)
- [GM15] Gauduchon P and Moroianu A 2017 Killing 2-forms in dimension 4 Special Metrics and Group Actions in Geometry (Springer INdAM Series vol 23) ed S Chiossi, A Fino, E Musso, F Podestà and L Vezzoni (Springer) pp 161–205
- [GP05] Griffiths J B and Podolsky J 2006 A new look at the Plebanski-Demianski family of solutions Int. J. Mod. Phys. D15 335–70
- [Ker63] Kerr R P 1963 Gravitational field of a spinning mass as an example of algebraically special metrics Phys. Rev. Lett. 11 237–8
- [Kin69] Kinnersley W 1969 Type D vacuum metrics J. Math. Phys. 10 1195–203
- [Kra09] Krasnov K 2011 Plebanski formulation of general relativity: a practical introduction Gen. Relativ. Gravit. 43 1–15
- [Kra11] Krasnov K 2011 Pure connection action principle for general relativity Phys. Rev. Lett. 106 251103
- [Kra20] Krasnov K 2020 Formulations of General Relativity, Gravity, Spinors and Differential Forms (Cambridge University Press)
- [LeB91] LeBrun C 1991 Explicit self-dual metrics on CP2 # ... # CP2 J. Differ. Geom. 34 223-53
- [NH15] Nozawa M and Houri T 2016 Killing-Yano tensor and supersymmetry of the self-dual Plebanski-Demianski solution Class. Quantum Grav. 33 125008
- [NJ65] Newman E T and Janis A I 1965 Note on the Kerr spinning-particle metric *J. Math. Phys.* **6** 915–7
- [NP12] Nikolic B D and Pantic M R 2012 A possible intuitive derivation of the Kerr metric in orthogonal form based on ellipsoidal metric ansatz (arXiv:1210.5922 [gr-qc])
- [PB84] Przanowski M and Baka B 1984 One-sided type-D gravitational instantons *Gen. Relativ. Gravit.* **16** 797–803
- [PD76] Plebanski J F and Demianski M 1976 Rotating, charged and uniformly accelerating mass in general relativity Ann. Phys., NY 98 98–127
- [Ple77] Plebanski J F 1977 On the separation of Einsteinian substructures J. Math. Phys. 18 2511–20
- [Tod20] Tod P 2020 One-sided type-D Ricci-flat metrics (arXiv:2003.03234 [gr-qc])
- [Tod24] Tod P 2024 Rod structures and patching matrices: a review (arXiv:2411.02096 [math.DG])
- [War90] Ward R S 1990 Einstein-Weyl spaces and SU(∞) Toda fields *Class. Quantum Grav.* 7 L95–L98