

# Atypical presentation of COVID-19 in a frail older person

Hui Sian Tay<sup>1</sup>, Rowan H Harwood<sup>1,2</sup>

<sup>1</sup> Nottingham University Hospitals NHS Trust

<sup>2</sup> School of Health Sciences, University of Nottingham

Correspondence to:

Professor Rowan Harwood  
School of Health Sciences, University of Nottingham,  
Queens Medical Centre,  
Derby Road,  
Nottingham  
NG7 2UJ  
Telephone number +44(0) 115 823 0873  
Mail to: Rowan.Harwood@nottingham.ac.uk

## Abstract:

Common symptoms of pandemic Coronavirus disease (COVID-19) include fever and cough. We describe a 94 year old man with well-controlled schizoaffective disorder, who presented with non-specific and atypical symptoms: delirium, low-grade pyrexia and abdominal pain. He was given antibiotics for infection of unknown source, subsequently refined to treatment for community-acquired pneumonia. Despite active treatment, he deteriorated with oxygen desaturation and tachypnoea. Repeat chest X-ray showed widespread opacification. Post-mortem throat swab identified COVID-19 infection. He was treated in three wards over five days with no infection control precautions. This has implications for the screening, assessment and isolation of frail older people to COVID-specific clinical facilities, and highlights the potential for spread among healthcare professionals and other patients.

## Introduction:

The World Health Organisation characterised Coronavirus disease (COVID-19) as pandemic in March 2020 [1]. In published series 44% of patients had fever on admission and 89% developed it subsequently. Cough was present in 68%, fatigue and sputum production in about 25% [2]. Cardiac complications such as myocardial infarction or heart failure are common. Mortality rates are particularly high in older people [3]. We describe a case highlighting the need to be alert for non-specific or atypical presentations which may delayed testing, diagnosis and isolation.

## Case report:

A 94 year old man with schizoaffective disorder was admitted to hospital in the United Kingdom in March 2020, early in the pandemic. He was found confused by his landlady. His home was untidy and cold. Emergency department assessment concluded: 'not coping at home, possible delirium or mental relapse'.

Further history was given by his daughter by telephone. He had ischemic heart disease, his schizoaffective disorder was well-controlled on valproate semisodium, and he was normally cognitively intact. He was moderately frail (Clinical Frailty Scale 6), and a former smoker. He lived alone and walked with a Zimmer frame. His daughter visited four times a week, the last time three days previously, as she had been unwell with pneumonia.

On arrival, his temperature was 37.4°Celsius, respiratory rate 20/minute, oxygen saturation 98% on air, blood pressure 128/70mmHg and heart rate 96/minute. He was drowsy, disorientated, trying to undress himself and resisting care. He had wheeze and crepitations on chest examination, and abdominal pain and tenderness. Blood test results included white cell count 14 (lymphocytes 0.97, neutrophils 6.3)  $\times 10^9/\text{L}$ , C-reactive protein 258mg/L, urea 15 mmol/L, creatinine 101 micromol/L (baseline 83), lactate 3.4 mmol/L. Blood culture grew *Staphylococcus epidermidis* of doubtful significance. ECG showed sinus rhythm with left bundle branch block. Chest X-ray (CXR) showed no consolidation. Urine culture was negative.

The initial diagnosis was delirium due to infection of unknown source and acute kidney injury. He was treated with intravenous fluids and piperacillin/tazobactam. CT abdomen with contrast showed bibasal lung consolidation. There was hypo-enhancement of the cardiac apex consistent with myocardial ischemia. Echocardiogram showed a dilated left ventricle with severely impaired systolic function. Troponin was raised at 475ng/L (normal 0-59) suggesting type 2 myocardial infarction. A Do Not Resuscitate order was agreed.

Antibiotics were changed to co-amoxiclav and clarithromycin following the CT scan. Three days following admission, he remained unwell: temperature 37.5°, respiratory rate 30/minute, oxygen saturation 91% on 2 litres/minute inspired oxygen. Repeat CXR showed right upper and lower zones airspace opacification in keeping with infection (Figure 1). Oxygen saturation deteriorated to 87% on 15 litres/minute of oxygen; arterial blood gas showed  $\text{pO}_2$  7.8kPa,  $\text{pCO}_2$  5.3kPa. He died 5 days following admission. A nasopharyngeal swab performed post-mortem was positive for SARS-COV-2-RNA.

Seven nursing staff and one doctor subsequently developed symptoms. The patient in the adjacent bed developed confirmed COVID-19.


## Discussion:

81 This man living with frailty presented with non-specific features, which do not appear  
82 in the symptoms listed in published reports [2,3] or guidance [4] on COVID-19. He  
83 developed lower respiratory signs and was treated for bacterial pneumonia. His  
84 temperature was always below 37.8°C, the diagnostic criterion for COVID-19. He  
85 was managed in three different clinical areas in open six-bedded bays without  
86 infection control precautions. Clinicians must be aware of the possibility of COVID-19  
87 presenting non-specifically, including with delirium, particularly with signs suggesting  
88 infection, so cases are not missed when they fall outside current diagnostic and  
89 management guidelines. This has implications for the diagnosis and isolation of  
90 COVID-19 in frail older people, with potential for spread among healthcare  
91 professionals and other patients if the diagnosis is overlooked.

#### 92 93 94 **References:**

- 95 1. World Health Organisation. WHO Director-General's opening remarks at the  
96 media briefing on COVID-19 - 11 March 2020  
97 [https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-](https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020)  
98 [the-media-briefing-on-covid-19---11-march-2020](https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020) Accessed 19 March 2020.
- 99 2. Guan W, Ni Z, Hu Y, et al. Clinical characteristics of coronavirus disease 2019 in  
100 China. NEJM, 2020. doi: 10.1056/NEJMoa2002032
- 101 3. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult  
102 inpatients with COVID-19 in Wuhan, China: a retrospective study. Lancet, 2020. doi:  
103 1016/S0140-6736(20)30566-3.  
104
- 105 4. World Health Organisation. Report of the WHO-China Joint Mission  
106 on Coronavirus Disease 2019 (COVID-19). [https://www.who.int/docs/default-](https://www.who.int/docs/default-source/coronaviruse/who-china-joint-mission-on-covid-19-final-report.pdf)  
107 [source/coronaviruse/who-china-joint-mission-on-covid-19-final-report.pdf](https://www.who.int/docs/default-source/coronaviruse/who-china-joint-mission-on-covid-19-final-report.pdf)

108 Figure1: CXR showing right upper and lower zones airspace opacification and small  
109 bilateral effusion.



110  
111  
112  
113  
114  
115  
116  
117