

Metadata of the chapter that will be visualized online

Chapter Title	The Indian Sundarbans: Biogeochemical Dynamics and Anthropogenic Impacts	
Copyright Year	2021	
Copyright Holder	The Author(s), under exclusive license to Springer Nature Switzerland AG	
Corresponding Author	Family Name	Henderson
	Particle	
	Given Name	Andrew C. G.
	Suffix	
	Division	
	Organization/University	School of Geography, Politics & Sociology, Newcastle University
	Address	Newcastle upon Tyne, UK
	Email	andrew.henderson@ncl.ac.uk
Author	Family Name	Das
	Particle	
	Given Name	Sourav
	Suffix	
	Division	
	Organization/University	School of Oceanographic Studies, Jadavpur University
	Address	Kolkata, India
Author	Family Name	Ghosh
	Particle	
	Given Name	Tuhin
	Suffix	
	Division	
	Organization/University	School of Oceanographic Studies, Jadavpur University
	Address	Kolkata, India
Author	Family Name	Panizzo
	Particle	
	Given Name	Virginia N.
	Suffix	

	Division	
	Organization/University	School of Geography, University of Nottingham
	Address	Nottingham, UK
Author	Family Name	Moorhouse
	Particle	
	Given Name	Heather L.
	Suffix	
	Division	
	Organization/University	Lancaster Environment Centre, Lancaster University
	Address	Lancaster, UK
Author	Family Name	Roberts
	Particle	
	Given Name	Lucy R.
	Suffix	
	Division	
	Organization/University	School of Geography, University of Nottingham
	Address	Nottingham, UK
Author	Family Name	Walton
	Particle	
	Given Name	Richard E.
	Suffix	
	Division	
	Organization/University	School of Geography, Politics & Sociology, Newcastle University
	Address	Newcastle upon Tyne, UK
Author	Family Name	Zheng
	Particle	
	Given Name	Ying
	Suffix	
	Division	
	Organization/University	School of Geographical & Earth Sciences, University of Glasgow
	Address	Glasgow, UK

Author	Family Name	Bass
	Particle	
	Given Name	Adrian M.
	Suffix	
	Division	
	Organization/University	School of Geographical & Earth Sciences, University of Glasgow
	Address	Glasgow, UK
Author	Family Name	McGowan
	Particle	
	Given Name	Suzanne
	Suffix	
	Division	
	Organization/University	School of Geography, University of Nottingham
	Address	Nottingham, UK
Abstract	<p>The Sundarbans region is one of the richest ecosystems in the world and is located on one of the world's largest deltas – the Ganges–Brahmaputra–Meghna system. The Indian Sundarbans have exceptional biodiversity, including rare and globally threatened species, and is made up of a mangrove forest ecosystem with an interconnected network of rivers. The hydrology of the Sundarbans underpin ecosystem health and the potential impact of humans on the region, as the tidal cycle changes water salinity diurnally and freshwater supply changes seasonally with the monsoon. The Indian Sundarbans face multiple pressures with both a reduction in freshwater supply and rising relative sea-level, leading to increased salinization of the mangrove forest. Human-driven alteration of the Sundarbans river catchments is reducing sediment flow, and when coupled with land-use change, is leading to subsidence, deforestation, nutrient enrichment, and heavy metal pollutants impacting the health of the ecosystem. All of these impacts have important ramifications for carbon fluxes that could exacerbate climate change and ecosystem health. In this chapter, we present an overview of our current understanding of biogeochemical dynamics and anthropogenic impacts on the Indian Sundarbans, with a particular focus on water quality, aquatic ecology, and carbon dynamics.</p>	
Keywords (separated by “ - ”)	Indian Sundarbans - Ganges–Brahmaputra–Meghna delta - water quality, biogeochemistry - carbon - pollutants - ecology - sediments	

Chapter 15

The Indian Sundarbans: Biogeochemical Dynamics and Anthropogenic Impacts

Andrew C. G. Henderson, Sourav Das, Tuhin Ghosh, Virginia N. Panizzo,
Heather L. Moorhouse, Lucy R. Roberts, Richard E. Walton, Ying Zheng,
Adrian M. Bass, and Suzanne McGowan

Abstract The Sundarbans region is one of the richest ecosystems in the world and is located on one of the world's largest deltas – the Ganges–Brahmaputra–Meghna system. The Indian Sundarbans have exceptional biodiversity, including rare and globally threatened species, and is made up of a mangrove forest ecosystem with an interconnected network of rivers. The hydrology of the Sundarbans underpin ecosystem health and the potential impact of humans on the region, as the tidal cycle changes water salinity diurnally and freshwater supply changes seasonally with the monsoon. The Indian Sundarbans face multiple pressures with both a reduction in freshwater supply and rising relative sea-level, leading to increased salinization of the mangrove forest. Human-driven alteration of the Sundarbans river catchments is reducing sediment flow, and when coupled with land-use change, is leading to subsidence, deforestation, nutrient enrichment, and heavy metal pollutants impacting the health of the ecosystem. All of these impacts have important ramifications for carbon fluxes that could exacerbate climate change and ecosystem health. In this chapter, we present an overview of our current understanding of biogeochemical dynamics and anthropogenic impacts on the Indian Sundarbans, with a particular focus on water quality, aquatic ecology, and carbon dynamics.

A. C. G. Henderson (✉) · R. E. Walton

School of Geography, Politics & Sociology, Newcastle University, Newcastle upon Tyne, UK
e-mail: andrew.henderson@ncl.ac.uk

S. Das · T. Ghosh

School of Oceanographic Studies, Jadavpur University, Kolkata, India

V. N. Panizzo · L. R. Roberts · S. McGowan

School of Geography, University of Nottingham, Nottingham, UK

H. L. Moorhouse

Lancaster Environment Centre, Lancaster University, Lancaster, UK

Y. Zheng · A. M. Bass

School of Geographical & Earth Sciences, University of Glasgow, Glasgow, UK

© The Author(s), under exclusive license to Springer Nature Switzerland
AG 2021

S. Das, T. Ghosh (eds.), *Estuarine Biogeochemical Dynamics of the East Coast of India*, https://doi.org/10.1007/978-3-030-68980-3_15

Keywords Indian Sundarbans · Ganges–Brahmaputra–Meghna delta · water quality, biogeochemistry · carbon · pollutants · ecology · sediments

[AU1]

15.1 Introduction

The Sundarbans region is one of the richest ecosystems in the world and is located on one of the world's largest deltas – the Ganges–Brahmaputra–Meghna (GBM) system. The Sundarban is located in the estuarine phases of the Rivers Ganga, Brahmaputra, and Meghna between 21°32'N and 21°40'N and 88°05'E and 89°E, spanning regions in both India and Bangladesh (Spalding et al. 1997) and contains arguably the world's largest remaining area of mangroves (an area of ~2529 km², Bhattacharyya 2015). The Indian Sundarbans have exceptional biodiversity, including rare and globally threatened species, for example, the northern river terrapin (*Batagur baska*, Lesson 1831), the Irrawaddy dolphin (*Orcaella brevirostris*, Owen in Gray 1866), the Ganges River dolphin (*Platanista gangetica*, Lebeck 1801), the brown-winged kingfisher (*Pelargopsis amauroptera* Pearson 1841), and the Royal Bengal tiger (*Panthera tigris*, Linnaeus 1785) – the only mangrove tiger on Earth (RAMSAR 2019). The mangrove ecosystem, which makes up the Indian Sundarbans, is an interconnected network of rivers, creeks, rivulets, and semidiurnal tides. The lower delta is dominated by a network of tributary rivers, creeks, and channels, with direct marine influence on the most seaward part of the Indian Sundarbans (Fig. 15.1). As a result, there are a range of hydrological influences (including both freshwater and coastal water) on the mangrove system, and when coupled with its topographic heterogeneity it results in a rich biodiversity (Gopal and Chauhan 2006). This has led to the Sundarbans mangrove forest being designated a World Heritage Site by International Union for Conservation of Nature (IUCN) in 1987; a Biosphere Reserve by United Nations Educational, Scientific and Cultural Organization (UNESCO) in 1989; and a wetland of international importance by RAMSAR in 2019.

[AU2]

Despite its international designation, the Indian Sundarbans face multiple pressures. As the freshwater discharge originating from the Himalayan uplands has decreased in recent decades (Raha et al. 2012), this has led to increased salinization of soil and groundwater within the Sundarbans, leading to the degradation of mangrove ecosystem health (Chowdhury et al. 2019). In addition, anthropogenic activities continue to alter hydrology and sediment flow, while land-use change is leading to deforestation, nutrient enrichment, and heavy metal pollutants causing many mangrove species to become threatened or extinct (Gopal and Chauhan 2006; Sodhi et al. 1987) and triggering an overall degraded ecosystem. This, in turn, has important ramifications for carbon fluxes in the Indian Sundarbans that could further exacerbate climate change and ecosystem health. The following sections aim to explore these different pressures and the impacts they are having on the current and future state of this vital ecosystem.



this figure will be printed in b/w

Fig. 15.1 A Sentinel-2 satellite natural color image taken in March 2018 of the Sundarbans region West Bengal, India, generated through the Sentinel Hub. The main rivers that influence the biogeochemistry and anthropogenic impact of the Sundarbans are labelled and major cities and towns are labelled. Inset map shows the location of the Sundarbans within in India

15.2 Hydrological Regime and Sediment Flow

64

The Indian Sundarbans landscape has evolved from the subduction of the Asian plate under the Burma plate to neotectonic tilting creating a hydrological gradient leading to river discharge from the highlands (Morgan and McIntire 1959). As a result, there are seven major estuarine rivers flowing through the Indian Sundarbans – the Hooghly, the Muriganga, the Saptamukhi, the Thakuran, the Matla, the Gosaba, and the Harinbhanga (also known as Ichamati and Raimangal) (Fig. 15.1). The combination of freshwater and tidal flow shape the deposition and erosion of sediments across the Sundarbans region, creating the dynamic nature of this deltaic environment. The climate of the Sundarbans is subhumid and characterized by hot summers and mild winters (Fig 15.2a). The mean monthly temperature varies between 30 °C to 40 °C in the summer (June to September) and 15 °C to 20 °C in winter (October to March). Precipitation from the annual monsoon during June to September is the major freshwater source to the Indian Sundarbans as it represents 80% of all annual rainfall (1750–1800 mm per annum) for the region. As a result, changes in freshwater inputs from monsoon rains, baseline river discharge during the rest of the year and tidal hydrology strongly influence the Sundarban region.

65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80




Fig. 15.2 (a) Summary of mean monthly temperature and precipitation data from Kolkata, West Bengal, from 1982 to 2012. Data from climate-data.org and is based on an interpolated model of weather station data; (b) Mean monthly discharge of the Bhagirathi and Hooghly River systems, West Bengal. (Data from Rudra (2014) and is derived from a rainfall-runoff model)

15.2.1 Hydrological Regime

15.2.1.1 Freshwater Hydrology

Despite the Indian Sundarbans being part of the GBM system (Chatterjee et al. 2013), year-round continuous flow is limited to a few river channels. The Hooghly River discharges the most freshwater into the Indian Sundarbans and is the western

most branch of the River Ganges reaching the Bay of Bengal (Fig. 15.1) (Rudra 2018). The Raimangal River at the eastern edge of the Indian Sundarbans also brings additional freshwater as a tributary channel of the Ichhamati River, and in turn this influences the discharge of the Gosaba, Harinbhanga, and Jhila Rivers (Chatterjee et al. 2013; Sarkar et al. 2013). While the monsoon seasons create variation throughout the year (Fig 15.2b), the Hooghly River has a more consistent input of freshwater than the Raimangal River (Chatterjee et al. 2013; Ghosh et al. 2013) due to the construction of the Farakka Barrage that diverts 7% of the annual flow of the Ganges to provide a regulated stream of freshwater throughout the dry season to support the operation of the Port of Kolkata (Ghosh et al. 2013).

15.2.1.2 Tidal Influence on Hydrology

The Sundarbans are macrotidal (range: 1.8 to 5.2 m between neap and spring high tides) and it experiences a semidiurnal tide cycle (Gole and Vaidyaraman 1967; Rogers and Goodbred 2014; Sinha et al. 1996). Despite the large volumes of freshwater from the Hooghly and Raimangal Rivers, rising tides still influence the upstream hydrology of the Sundarbans, with tides regularly travelling up to 120 km from the mouth of the Hooghly River during the pre-monsoon season (Gole and Vaidyaraman 1967). In the post-monsoon, the tide can travel 250 km up the Hooghly (Sinha et al. 1996) with the tidal limit at Kalna, West Bengal, during the monsoon (Chatterjee et al. 2013).

As the tides bring saline water with them, they impact both anthropogenic access to freshwater and affect ecological functioning. The incursion of saline waters by flood tides is also controlled by the season in which it happens. For example, the extent of saline waters during the monsoon is low, as the increased freshwater delivered by seasonal rains acts as a barrier to flood tide penetration, with the upper limit typically as far as Nayachar Island in the upper mouth of the Hooghly River (Chatterjee et al. 2013; Ghosh et al. 2013; Sharma et al. 2018). Another effect is the stratification of freshwaters over the saline/brackish waters in the river during the monsoon season (Sadhuram et al. 2005; Chatterjee et al. 2013). In the non-monsoon seasons there is a significant rise in salinity levels within the Hooghly River with 30 ppt (parts per thousand) observed near Diamond Harbour and saline waters reach as far north as Kolkata (Gole and Vaidyaraman 1967), although during ebb tides the limit of saline water moves back down to the mouth of the estuary near Sagar Island (Sinha et al. 1996).

15.2.2 Sediment Flow

All river channels flow into the Indian Sundarbans, including freshwater rivers and tidal inflows, carrying sediments that affect the whole mangrove ecosystem. Sediments carried by the freshwater Hooghly River consist predominantly of sand

and silt (Somayajulu et al. 2002; Massolo et al. 2012) and less than 10% of the sediment consists of clay particles. These sediments are predominantly derived from rain-driven terrestrial erosion to the Ganges (Somayajulu et al. 2002; Rudra 2018), and because of the high discharge of the Ganges and Hooghly Rivers these sediments do not experience much water-column weathering before they reach the Sundarbans and the Bay of Bengal (Somayajulu et al. 2002; Flood et al. 2016). While the Hooghly carries a large volume of sediment, there is notable seasonal variation in sediment loads because of monsoon-driven changes in freshwater discharge (Gole and Vaidyaraman 1967).

Each year flood tides deposit ~12 cm of fresh sediment in to the Indian Sundarbans (Rudra 2018) and tides carry sediments that are more fine-grained than those transported by freshwater rivers (Allison et al. 2003; Flood et al. 2016, 2018). Sediments transported and deposited by flood tides in the Indian Sundarbans also originate from the mouth of the Ganges–Brahmaputra–Meghna River system approximately 275 km to the east (Flood et al. 2016, 2018; Rudra 2018), where >1 billion tons of sediment are discharged each year (Somayajulu et al. 2002; Rogers and Goodbred 2014; Rudra 2018). These sediments are carried by coastal currents westward along the coastline through suspension (Rogers and Goodbred 2014; Flood et al. 2018), where they undergo weathering and degradation in the water column, resulting in fine-grained sediments being transported in suspension by flood tides in to the Sundarbans (Flood et al. 2016, 2018). Sediments are deposited and retained because of lateral accretion along mangrove tree roots (Manna et al. 2012; Flood et al. 2018) and tidal creeks (Rudra 2018).

Resuspension of sediments occurs as a result of bioturbation in intertidal mudflats (Rogers and Goodbred 2014), dredging, winds, and tides. These resuspended sediments are redistributed or carried from the Sundarbans through flooding and wave action. Approximately 430 km² of the Indian Sundarbans were eroded between 1917–2016, which is offset by 220 km² of sediment accumulation over the same period (Rudra 2018). The dynamics of rivers, tides, and sediment movement means these are key processes that drive Sundarbans water quality, ecology, and overall ecosystem health (Gole and Vaidyaraman 1967; Sinha et al. 1996; Rogers and Goodbred 2014).

15.3 Ecology and Water Quality

The Indian Sundarbans are home to a number of endemic enigmatic and globally vulnerable species. By looking at the biology of these fragile Sundarbans ecosystems and the interface with hydrology and biogeochemistry we can document and understand the threats to the Sundarbans wetland ecosystem and its iconic inhabitants.

15.3.1 *Aquatic Ecology* 162

15.3.1.1 **Primary Producers** 163

Aquatic primary production in the Sundarbans is a function of nutrient loading and light penetration, with the latter often constrained by river turbidity (Chaudhuri et al. 2012). Large river and estuarine channels are dominated by the *Bacillariophyceae* algal group – biosiliceous diatoms, followed by *Pyrrophyceae* – dinoflagellates, and *Chlorophyceae* – chlorophytes (Biswas et al. 2010; Manna et al. 2010; De et al. 2011). There are still large gaps in our knowledge about the role of these primary producers in mangrove ecosystems, especially diatoms (Samanta and Bhadury 2018). However, the biovolume of primary produces is highest in the post-monsoon winter months supporting colonies of long-chain diatoms, whereas there are low biovolumes during the monsoon season because of increased total suspended solids (TSS) (derived from rain-driven catchment erosion), reducing light penetration and photosynthesis (Chaudhuri et al. 2012; Bhattacharjee et al. 2013). Prior to the monsoon season the diatom assemblage is dominated by saline-tolerant species (Manna et al. 2010) and this may become a feature of upstream diatom communities as saline intrusion into the delta region becomes more widespread.

15.3.1.2 **Macroinvertebrates** 179

The main consumers of primary producers are the zooplankton, who play an integral role in the transfer of organic matter between trophic levels and export organic carbon to sediments (Bhattacharya et al. 2015a). As macroinvertebrate species occupy distinct trophic levels they respond rapidly to environmental change and are relatively quick and easy to identify, making them effective water-quality indicators (Gannon and Stemberger 1978). Copepods are small cosmopolitan crustaceans, which dominate zooplankton in tidal river systems in the Indian Sundarbans (Bir et al. 2015). Whereas in tidal flats, polychaetes and mollusks are important macro-zoobenthic groups, whose spatiotemporal distribution is driven by salinity, the nature of the substrate (e.g., mudflats exhibit greater diversity than sandflats), and anthropogenic activity (Khan 2003; Roy and Nandi 2012).

In the Sundarbans, compositional changes in zooplankton communities are primarily driven by the quantity and quality of primary producer prey, as well as salinity and water transparency, which can vary seasonally and interannually (Bhattacharya et al. 2015a). Much like primary producers, zooplankton biomass is highest during the post-monsoon season when water currents, salinity, and temperature are at their lowest (Bir et al. 2015). However, extreme climate events, such as cyclone “Aila” in 2009, result in increased suspended particulates and nutrients, reductions in transparency, and primary photosynthesis. As a result there is a decrease in zooplankton diversity, biomass, and abundance (Bhattacharya et al. 2014a). If extreme events across the region worsen, this could modify phytoplankton–

zooplankton interactions and threaten the viability of both open-water and aquaculture fisheries, whose stock require good quantity and quality of these prey organisms. Indeed, continued saltwater intrusion may reduce macrozoobenthic diversity due to reductions in decomposition rate of photosynthetic organic matter following higher sediment salinities, which may modify macrozoobenthic feeding behaviors and consequently impact the higher organisms which they support, for example, wading birds (Bandopadhyay and Burman 2006; Roy and Nandi 2012).

15.3.1.3 Microbial Biodiversity

Mangrove environments are hotspots of microbial diversity because of the complexity of habitats they provide and the fluxes in salinity, nutrients, labile organic compounds, and water levels across daily to seasonal timescales (Chakraborty et al. 2015). Seasonal variations in freshwater flow are an important determinant of community diversity, specifically of the bacterioplankton, where diversity is found to be greater in monsoon seasons compared to post-monsoon (Ghosh and Bhadury 2018). These microbes play a profound role in biogeochemical cycling from metabolizing the considerable allochthonous organic matter inputs of mangrove vegetation (Chakraborty et al. 2015), and therefore the sustenance, productivity, and recovery of this ecosystem (Ghosh et al. 2010; Roy et al. 2002; Santos et al. 2011). While there remains a significant gap in the knowledge of microbial diversity and abundance in the Sundarbans (Ghosh et al. 2010), modifications in microbial abundance, diversity, and community composition have been identified (Ghosh and Bhadury 2018). For example, industrial and boating activity has increased polyaromatic hydrocarbons (PAHs), heavy metal, and nutrient pollution detected by bacterial strains with heavy metal resistance and those involved in hydrocarbon degradation processes (Chakraborty et al. 2015). Eutrophication of these waters has meant bacterial productivity exhibits an exponential relationship to temperature as they are no longer nutrient-limited (Manna et al. 2010, 2012).

15.3.2 Water Quality

15.3.2.1 Nutrients

One of the key factors determining the biodiversity of the Indian Sundarbans is water and the role it plays in transporting nutrients and pollutants in the mangrove ecosystem (Sarkar et al. 2004). In general, phosphorus (P) availability is low in tropical regions where soils have been weathered for millions of years (Yang et al. 2013). Nitrogen (N) can be generated and removed from ecosystems by microbes and so mangroves are important sites for N (and C) cycling with mangrove plants being significant stores of N (Kamruzzaman et al. 2019; Purvaja et al. 2008). In coastal zones, P and N availability changes along the freshwater–marine transition,

because sediments retain less P in marine environments, releasing P to the waters (Blomqvist et al. 2004). Primary production in freshwaters tends to be limited by P, whereas marine waters are generally N-limited and P-replete. Therefore, the tidal cycle in the Sundarbans is a key influence on nutrient distribution in estuaries, and the nutrient status of waters change seasonally to become P-limited after the monsoon when the influence of freshwaters increases, and N-limited during the monsoon and pre-monsoon periods (Chaudhuri et al. 2012). The main source of nutrients are from either freshwater runoff, for example, dissolved silica, nitrate, and phosphate, and/or from intertidal flats, for example, ammonium, nitrate/nitrite, and phosphate (Singh et al. 2016). During low tides, there is an increase in freshwater input into the northern Bay of Bengal, which dilutes nutrient concentration across the continental shelf and the mangrove ecosystem and vice versa during high tides, and these tidal dynamics play a crucial role in regulating short-term variability in nutrient concentrations (Das et al. 2015, 2017). Atmospheric deposition of P also constitutes a major source in the Sundarbans mangroves, comprising >50% of the annual P inputs (Ray et al. 2018a, b). P is hypothesized to be transported from arid regions of western India by pre-monsoonal northwesterly (and westerly) winds (Ray et al. 2018a, b). This seasonal P transport seems likely to either drive or exacerbate the observed seasonal differences in estuaries, but thus far there has been little research into the interplay of monsoonal rainfall, river discharge, and the consequences of desertification in arid regions on nutrient cycling of the Sundarbans.

In addition to natural variability in nutrients, anthropogenic inputs of nutrient-rich effluent have led to the eutrophication of smaller rivers, tidal creeks, and ponds in the Sundarbans, exacerbated by generally reduced flushing rates. However, such phenomena are being more commonly documented within the main estuarine channels such as the Hooghly River where anthropogenic influences has increased at a faster rate (Manna et al. 2010; De et al. 2011). Eutrophication has led to algal blooms, which reduce light penetration for benthic photosynthesis and deplete oxygen for higher trophic species (due to bloom respiration) (Biswas et al. 2014). In addition, harmful algal blooms (HABs) from toxin-producing cyanobacteria (CyanoHABs) such as *Microcystis* species and dinoflagellates have been recorded in Sundarbans aquatic habitats (Manna et al. 2010; Sen et al. 2015). Cyano HABs outcompete other algal groups due to their ability to regulate buoyancy, adaptation to low light, and higher temperatures, and are often able to fix N from the atmosphere (important in systems that are N-limited relative to P typical in these wetlands) (Paerl and Tucker 1995; Walsby and Schanz 2002; Islam et al. 2004; Paerl and Huisman 2008).

15.3.2.2 Heavy Metals

The primary source of heavy metal contamination in coastal areas of West Bengal is the major rivers that run through the Sundarbans (Mitra 1998) and even though these metals can occur naturally in the Sundarbans biogeochemical cycle (Garrett 2000), they predominantly come from industrial and domestic effluents, storm

water runoff, dust, and boating activities. The mineralogy and grain size of sediments of the GBM river system has the potential to trap contaminants with silt and clays, predominantly carrying metal contamination from upstream, with the textural composition and organic matter of sediments critical to the sorption of transition metals (Kumar and Ramanathan 2015; Roy et al. 2018). Consequently, river sediments have become a sink of bioavailable heavy metals, with flooding and dredging leading to the resuspension of sediments, releasing their heavy metal load into the water column. Furthermore, salinity influences the partitioning, physiochemical form, and therefore bioavailability of these metals (Mitra 1998).

The Hooghly River catchment encompasses rural, agricultural, urban, and industrial land uses, including the megacity of Kolkata (population ~ 15 million) before draining into the Bay of Bengal. The metal concentrations of the riverine suspended particulate matter (SPM) ranges 7.9–29 µg/g (mean: 19 ± 5.5 µg/g) for Co, 17–70 µg/g (mean: 49 ± 14 µg/g) for Ni, and 12–55 µg/g (mean: 36 ± 12 µg/g) for Cu, which is higher than the average concentrations for global rivers (Samanta and Dalai 2018). The dissolved concentrations of metals in the Hooghly River estuary range 0.8–24 nM/L (mean: 6.2 ± 6 nM/L) for Co, 3.5–172 nM/L (mean: 50 ± 42 nM/L) for Ni, and 8–178 nM/L (mean: 60 ± 37 nM/L) for Cu. Annually, these contribute up to 1.8% Co, 2.4% Ni, and up to 1.2% Cu of the global riverine metal fluxes (Sumanta and Dalai 2018). The heavy metal concentrations of the Hooghly display seasonal variability with the maximum pollution load pre-monsoon and minimum load during the monsoon (Roy et al. 2018). High concentrations pre-monsoon have been attributed to high temperatures and increased evaporation rates of surface water (Bhattacharya et al. 2014a; Ghosh and Choudhury 1989; Mitra 1998).

Mixing of riverine and marine waters also contributes to changes in the speciation of metals, as well as the resuspension of sediments. Mukherjee et al. (2009) argue physiochemical changes limit the enrichment of heavy metals in river sediments and the high concentrations seen in the Hooghly compared to other regional rivers is because of a large sediment contribution from a bigger catchment area. The elevated concentrations in the Hooghly River are an important mechanism for elevating the amount of dissolved Ba in the river estuary via desorption with mixing of waters (Samanta and Dalai 2018). Similarly, Hg concentrations are positively correlated with pH ($r = 0.58$ – 0.68 , $p < 0.01$) and salinity ($r = 0.52$ – 0.79 , $p < 0.01$) (Bhattacharya et al. 2014b), and some metal concentrations in the waters of the middle and lower Hooghly estuary are significantly higher than other global estuaries in dissolved Ni and Cu (Samanta and Dalai 2018). However, upstream anthropogenic activities are still important in contributing widescale pollution across the Sundarbans.

Anthropogenic pollution within the Sundarbans itself has led to elevated levels of Cd, Cu, Zn, As, Ni, Pb, and Hg, which can cause impacts on biology (Sarkar et al. 2004; Chatterjee et al. 2007, 2009; Chowdhury et al. 2017; Mitra and Ghosh 2014). The source of these contaminants come from a mixture of industrial effluents, boat anti-fouling paint, sewage, fertilizers, and storm water drainage (Chowdhury and Maiti 2016; Mitra and Ghosh 2014; Chatterjee et al. 2007; Mitra et al. 2009; Kumar and Ramanathan 2015). Sediments within the Sundarbans have higher levels of

contamination compared with sediments in the Hooghly estuary because of lower tidal energy and finer-grained sediments (Banerjee et al. 2012). Hooghly River inputs of Cu and Zn are a critical source of heavy metal pollution to the Sundarbans (Chakrabarti et al. 1993; Bhattacharya et al. 2015). Moreover, the metal concentration of fine-grained sediment in the Indian Sundarbans is higher than those in the Bangladesh Sundarbans (Kumar and Ramanathan 2015) (Table 15.1). The industrialization of the upper catchment in India compared to Bangladesh has been suggested as the primary reason for this difference (Rahman et al. 2011).

15.3.2.3 Bioaccumulation and Health

Heavy metal pollution of the Sundarbans has important implications for the health of the ecosystem, aquatic organisms, and the local communities (Bhattacharya et al. 2015). River water in the region is largely unpotable due to the dissolved concentrations of Mn, Pb, and Ni (Bhattacharya et al. 2015). River water is also not suitable for irrigation due to the high concentration of Mn (Bhattacharya et al. 2015) and the large-scale metal pollution in riverine water and sediments is a serious concern as fish, prawns, and crabs have been reported to contain significant level of toxic metals (Dutta et al. 2017a; Mitra et al. 2012). Bioaccumulation of metals in these organisms occurs through the food chain until top level predators accumulate ions at a level that can develop neuronal, abdominal, and cardiovascular diseases. Table 15.2 shows the increase in metal accumulation between water, sediment, and macro benthos. At low concentrations, effects such as diarrhea, vomiting, and skin irritation are common. However, at high concentrations and continued exposure, there are serious health considerations with the International Agency for Research on Cancer (IARC) classifying Cd as a human carcinogen, Pb as possible human carcinogen, and Cr to be the cause of a rare sino-nasal cancer (Dayan and Paine 2001; Järup 2003).

Table 15.1 Comparison of heavy metal concentrations (Fe, Mn, Cu, Zn) across the Indian and Bangladesh Sundarbans

Heavy metal	Indian Sundarban	Bangladesh Sundarban
Fe ($\mu\text{g g}^{-1}$)	38,760–52,829	29,081–45,025
Mn ($\mu\text{g g}^{-1}$)	424–770	342–792
Cu ($\mu\text{g g}^{-1}$)	36–82	12–45
Zn ($\mu\text{g g}^{-1}$)	55–83	29–75

Data from Kumar and Ramanathan (2015)

Table 15.2 Heavy metal concentrations in water, sediments, and macrobenthos from the Sundarbans. Concentrations in the macrobenthos exceed toxic levels

Heavy metal	Water	Sediment	Macrobenthos
Cd ($\mu\text{g g}^{-1}$)	0.04–0.10	6.25–7.38	14.63
Zn ($\mu\text{g g}^{-1}$)	0.01–9.66	24.91–62.0	268.91
Pb ($\mu\text{g g}^{-1}$)	0.03–0.16	33.7–50.33	174.84
Fe ($\mu\text{g g}^{-1}$)	14.3–170.0		
Cr ($\mu\text{g g}^{-1}$)		46.8–78.50	18.76
Cu ($\mu\text{g g}^{-1}$)		20.38–42.01	90.02

Data from Rahman et al. (2009)

15.4 Carbon Biogeochemistry

The Sundarbans contain nearly 3% of the total area of the world’s mangrove ecosystems and have been an important region for understanding carbon cycle dynamics in estuarine delta ecosystems over the past 20 years. In particular, the biogeochemical cycling of different carbon species including dissolved organic carbon (DOC), particulate organic carbon (POC), dissolved inorganic carbon (DIC), and dissolved greenhouse gases (CO_2 ; CH_4) in different environments including estuarine water (e.g., Biswas et al. 2004; Dutta et al. 2019a, b), sediment (e.g. Dutta et al. 2013; Dutta et al. 2017b), mangrove soil and forests (e.g. Rahman et al. 2015; Chanda et al. 2016; Das et al. 2016).

15.4.1 Carbon Fluxes in the Sundarbans

Mangrove estuaries have been recognized as important organic C sources for the ocean and atmosphere (Rosentreter et al. 2018; Ray et al. 2015), with an estimated flux of 55 Tg C yr^{-1} (Sippo et al. 2017) derived from plant litter, phytoplankton, and microphytobenthos (Ray et al. 2015). However, the Sundarbans have a conspicuous lack of data related to its carbon budget. In particular, measurements of POC and DOC have only been taken in the last few years (Ray et al. 2018b; Dutta et al. 2019a, b). Ray et al. (2018b) provide the first baseline data of C export (DOC, POC, and DIC) from the Sundarbans mangroves into the Bay of Bengal, which accounts for $3.03 \text{ Tg C yr}^{-1}$, $0.58 \text{ Tg C yr}^{-1}$, and $3.69 \text{ Tg C yr}^{-1}$, respectively. DIC is the major form of C exported in the Sundarbans region, contributing to >50% of the fluvial C budget (Fig. 15.3), with DIC concentration (DIC) varying between 1.92 to 2.19 mM during a 24-hour period (Dutta et al. 2019a). However, compared to the Hooghly estuary, the major river draining into the Bay of Bengal, the percent contribution and flux of DIC from the Sundarbans is much smaller and it has a greater amount of organic-C flux (as DOC and POC). DOC concentration (DOC) was monitored in different seasons, with similar values observed during the pre- and post-monsoon (pre-monsoon: $294.3 \pm 34 \text{ }\mu\text{M}$; post-monsoon: $262.5 \pm 48.2 \text{ }\mu\text{M}$) (Ray

et al. 2018b), 235 ± 49 (Dutta et al. 2019b). POC is much smaller than DOC, ranging from 28.0 ± 8.6 μM during pre-monsoon to 45.4 ± 7.5 μM post-monsoon (Ray et al. 2018b). When more locations were monitored, a higher post-monsoon POC of 173 ± 111 μM was observed by Dutta et al. (2019b), reflecting the spatial variability of C flux within the Sundarbans region. DIC removal in the Sundarbans is facilitated by phytoplankton uptake, CO_2 outgassing and export to the adjacent continental shelf (Ray et al. 2018b), although significant uncertainty remains.

Much of the work on C biogeochemistry in the Sundarbans has focused on the CO_2 flux from river surface waters (Mukhopadhyay et al. 2002a, b; Biswas et al. 2004; Akhand et al. 2016; Vinh et al. 2019) (Table 15.3). While the mangrove forest is an autotrophic ecosystem and acts as a net C sink (Rodda et al. 2016), more temporal and spatial C flux data is needed to understand its potential to be a large C store. This is important as mangroves can export C to adjacent water bodies, increasing the fraction of CO_2 in water, which can control water-to-air emissions (Akhand et al. 2012) (Table 15.3). A study of the outer part of the Sundarbans found this area to be a CO_2 sink at a rate of $16 \times 10^6 \text{ kg C year}^{-1}$ (Mukhopadhyay et al. 2000), while other studies suggest mangrove estuaries are a net CO_2 source at a rate of $13.8 \text{ kg C ha}^{-1} \text{ year}^{-1}$ (Biswas et al. 2004), or it varies between a net source and sink through the seasons as influenced by the monsoon (Mukhopadhyay et al. 2002a, b) (Table 15.3).

The significance of CH_4 production and export from the Sundarbans has been recently documented (e.g., Mukhopadhyay et al. 2002a, b; Jha et al. 2014; Dutta et al. 2017b). Its importance to the Sundarbans biogeochemical cycle lies in the nature of intertidal mangrove sediments, which are generally anoxic and rich in organic carbon, and therefore creates favorable environments for methanogenesis (Dutta et al. 2017b). While in the riverine and standing waters, the production of CH_4 is linked to the stratification of the water column and anoxic bottom waters




Fig. 15.3 Percentage contribution of the different carbon fractions – dissolved organic carbon (DOC), particulate organic carbon (POC), and dissolved organic carbon (DOC) – (a) riverine C export from the Hooghly River into the Bay of Bengal during the monsoon season with maximum discharge; (b) mangrove-derived C export into the Bay of Bengal. (Date from Ray et al. 2018a, b)

Table 15.3 A summary of CO₂ and CH₄ fluxes and concentration estimates from the Sundarban ecosystem

Place	CO ₂ flux (mmol m ⁻² d ⁻¹)	[CH ₄] (nM)	CH ₄ flux (umol m ⁻² d ⁻¹)	Reference
<i>Sundarbans</i>	-3.65 × 10 ⁹			a.
<i>Hooghly estuary</i>	-2.78–84.4			a.
<i>Sundarbans</i>	0.315			b.
<i>Hooghly-Matla</i>	-0.337			c.
<i>Sundarbans (inner)</i>	0.675			d.
<i>Sundarbans (middle)</i>	0.536			
<i>Sundarbans (outer)</i>	-0.759			
<i>Hooghly estuary</i>	88.8			e.
<i>Matla estuary</i>	6.3			
<i>Saptamukhi estuary (surface water)</i>		69.9		f.
<i>Saptamukhi estuary (sub-surface water)</i>		56.1		
<i>Sundarbans (Muriganga, Saptamukhi, Thakuran)</i>		11.0– 129.0	1.97–134.6	g.
<i>Hooghly estuary</i>		10.3– 59.3	0.88–148.6	

Data sources: a. Mukhopadhyay et al. 2000, 2002b. Biswas et al. 2004; c. Akhand et al. 2012; d. Akhand et al. 2013; e. Akhand et al. 2016; f. Dutta et al. 2017a, b; g. Biswas et al. 2007

(Koné et al. 2010; Borges and Abril 2011). As a result, dissolved CH₄ concentrations ([CH₄]) are 11.0–129.0 nM throughout the year (Biswas et al. 2007) (Table 15.3), with a distinct increase in CH₄ in the post-monsoon period and maximum recorded values in December across all Sundarbans sites. Higher mean concentration for CH₄ are found in surface waters (69.9 nM) compared to their subsurface (56.1 nM) (Dutta et al. 2017b) (Table 15.3).

In mangroves and wetlands, sedimentary-derived CH₄ can escape to the adjacent water and/or atmosphere via plant vascular system-mediated transport, ebullition, and molecular diffusion (Chanton and Dacey 1991), among which ebullition is the dominant pathway (Maher et al. 2019) and is rarely accounted for in the water–air CH₄ budget (Jeffrey et al. 2019). While the [CH₄] in water columns can be partly oxidized to CO₂ via physical and biochemical processes (Hanson and Hanson 1996), this will be limited in well-mixed water bodies, allowing for CH₄ to be emitted (Abril et al. 2007). The CH₄ emission rate from surface waters was between 1.97 and 134.6 μmol m⁻² d⁻¹ in three distributaries in the Sundarbans with clear seasonal variation – minimum during the monsoon and maximum in the post-monsoon (Biswas et al. 2007) (Table 15.3).

15.4.2 Temporal and Spatial Variations of C Flux

422

The biogeochemical processes in the Sundarbans can be significantly different in the monsoon seasons compared to the dry periods of the year. For example, water in the Matla River was found to be marginally oversaturated in CO₂ throughout the year, but transitioned to a CO₂ sink during the post-monsoon season (Akhand et al. 2016). The difference results from the high discharge during the monsoon seasons creating a well-mixed water column, meaning that CO₂ diffusion was limited and there was little organic-rich sediment deposition. Furthermore, the concentrations and fluxes of different forms of C in the Sundarbans are often compared to the Hooghly River estuary, the main artery to the Sundarbans mangroves meaning you are comparing freshwater with coastal saline/brackish waters, which provides a different C dynamic. For DOC, POC, and DIC there is no distinct or consistent spatial pattern between three Sundarbans estuaries (Dutta et al. 2019b), although DIC and DOC were both lower on average than the Hooghly River. Akhand et al. (2013) shows water in the inner and middle Sundarbans regions are oversaturated in CO₂, but undersaturated in the outer region during the summer. As a result, the inner and middle Sundarbans act as a CO₂ source (29.7 and 23.6 mg CO₂ m⁻² day⁻¹, respectively) while the outer Sundarbans is a net sink (−33.4 mg CO₂ m⁻² day⁻¹). This change of carbon sink and source results from higher nutrient availability and chlorophyll *a* concentrations, reflecting primary productivity in the outer mangrove system. Variations in the fluxes of CO₂ also demonstrate the heterotrophic nature of the inner mangrove ecosystems at the land–ocean interface, and C-sink character of the outer mangrove on continental shelves (Chen and Borges 2009).

15.4.3 Source of C in the Indian Sundarbans

445

Very few studies have explored the source of different C species in the Sundarbans water, but a modelling study of the Hooghly–Matla river system by Ray et al. (2015) shows plant litter production and the breakdown of detritus from adjacent Sundarbans mangrove forests are a major source of dissolved inorganic N and C to river waters, and potentially C exports to the continental shelf. In addition, phytoplankton is a leading source of C near Sagar Island, this is not the case for the Saptamukhi estuary in the Sundarbans, where POC is mainly sourced from riverine suspended sediments and soils, but less from marine plankton, as indicated by their C/N ratios (Dutta et al. 2019a). Higher carbon isotope values in POC ($\delta^{13}\text{C}_{\text{POC}}$) in estuarine waters compared to mangroves indicate the modification of POC. Ray et al. (2018b) also demonstrate mangrove forests (including plant litter, eroded soil) are the major source of C exported from Sundarbans to the Bay of Bengal, compared to upstream C-inputs and marine phytoplankton. In addition, the negative relationship between [DIC] and its carbon isotope value ($\delta^{13}\text{C}_{\text{DIC}}$) during low tide, highlights respiration

of marine plankton-derived organic carbon may be an important source of DIC rather than exchange of C-rich porewaters derived from terrestrial sources.

15.4.4 Influence of Salinity and Tide on C

In general, CO₂ flux decreases with increasing salinity toward the open sea (Akhand et al. 2012). In the Matla River, the highest fraction of aqueous CO₂ (fCO₂) coincides with the lowest neap tide, overriding CO₂ uptake by photosynthesis. The hydrological change during the ebb and low tide leads to the mixing of sediment porewater and groundwater with brackish/saline estuary waters. The subsequent biogeochemical interaction that leads to increasing fCO₂ and the extent of CO₂ efflux highlights the role of salinity in C-dynamics over the Sundarbans (Akhand et al. 2016). The importance of tidal stage in controlling dissolved greenhouse gas efflux from water is also demonstrated by Padhy et al. (2020), who show the concentrations of dissolved CH₄ and CO₂ are higher in stagnant water during low tide compared to high tide water. This implies the effect of stagnation and lower salinity and therefore less SO₄²⁻ availability, which increases CO₂ emissions. Apart from high pCO₂ during low tide, Dutta et al. (2019b) also suggest there is a strong influence from estuarine mixing on DIC and δ¹³C_{DIC} during the low tide, both of which correlate with salinity. This can be explained by the impacts of this biogeochemistry on denitrification, sulfate reduction, and aerobic organic matter mineralization to DIC, along with possible organic contributions from porewater.

15.5 Conclusions

This overview of biogeochemical dynamics and anthropogenic impacts on the Indian Sundarbans highlights the importance of the hydrological regime in driving variability in ecosystem health. Diurnal and seasonal changes in salinity, which are driven by the tides and monsoon-driven freshwater availability, influence biological responses, biogeochemical cycling, and carbon dynamics. Also, high concentrations of heavy metals mean they are bioavailable within the major rivers running into the Sundarbans, but there is little evidence of the short- and long-term implications of this pollution for mangrove health, aquatic organisms, and local communities. Overall, there remains a paucity of research into water-quality impacts on aquatic ecology, including nutrient enrichment and heavy metal pollution, carbon cycling through the mangrove system, and how climate change has and will continue to affect the Indian Sundarbans.

Acknowledgments The writing of this chapter would not have been possible without the funding received from the UK Research and Innovation – Global Challenges Research Fund (UKRI GCRF) Living Deltas Hub (NE/S008926/1) www.livingdeltas.org.

References

496

- Abril G, Commarieu M-V, Guérin F (2007) Enhanced methane oxidation in an estuarine turbidity maximum. *Limnol Oceanogr* 52:470–475. <https://doi.org/10.4319/lo.2007.52.1.0470>
- Akhand A, Chanda A, Dutta S, Hazra S (2012) Air – water carbon dioxide exchange dynamics along the outer estuarine transition zone of Sundarban, northern Bay of Bengal, India. *Indian J Geo-Mar Sci* 41:111–116
- Akhand A, Chanda A, Dutta S, Manna S, Sanyal P, Hazra S, Rao KH, Dadhwal VK (2013) Dual character of Sundarban estuary as a source and sink of CO₂ during summer: an investigation of spatial dynamics. *Environ Monit Assess* 185:6505–6515. <https://doi.org/10.1007/s10661-012-3042-x>
- Akhand A, Chanda A, Manna S, Das S, Hazra S, Roy R, Choudhury SB, Rao KH, Dadhwal VK, Chakraborty K, Mostofa KMG, Tokoro T, Kuwae T, Wanninkhof R (2016) A comparison of CO₂ dynamics and air-water fluxes in a river-dominated estuary and a mangrove-dominated marine estuary. *Geophys Res Lett* 43(11):726–11,735. <https://doi.org/10.1002/2016GL070716>
- Allison MA, Khan SR, Goodbred SL, Kuehl SA (2003) Stratigraphic evolution of the late Holocene Ganges–Brahmaputra lower delta plain. *Sediment Geol* 155:317–342. [https://doi.org/10.1016/S0037-0738\(02\)00185-9](https://doi.org/10.1016/S0037-0738(02)00185-9)
- Bandopadhyay B, Burman D (2006) Characterization of soil and water of brackish water fisheries of coastal region of Sundarbans, West Bengal. *J Indian Soc Coast Agric Res* 24(1):78–82
- Banerjee K, Senthilkumar B, Purvaja R, Ramesh R (2012) Sedimentation and trace metal distribution in selected locations of Sundarbans mangroves and Hooghly estuary, Northeast coast of India. *Environ Geochem Health* 34:27–42. <https://doi.org/10.1007/s10653-011-9388-0>
- Bhattacharjee D, Samanta B, Danda A, Bhadury P (2013) Temporal succession of phytoplankton assemblages in a tidal creek system of the Sundarbans mangroves: an integrated approach. *Int J Biodivers*
- Bhattacharya BD, Hwang JS, Tseng LC, Sarkar SK, Rakshit D, Mitra S (2014a) Bioaccumulation of trace elements in dominant mesozooplankton group inhabiting in the coastal regions of Indian Sundarban mangrove wetland. *Mar Pollut Bull* 87:345–351. <https://doi.org/10.1016/j.marpolbul.2014.07.050>
- Bhattacharya S, Dubey SK, Dash JR, Patra PH, Das AK, Mandal TK, Bandyopadhyay SK (2014b) Assemblages of Total mercury in the tropical Macrotidal Bidyadhari estuarine stretches of Indian Sundarban mangrove eco-region. *J Environ Anal Toxicol* 4(6):1
- Bhattacharya BD, Nayak DC, Sarkar SK, Biswas SN, Rakshit D, Ahmed MK (2015) Distribution of dissolved trace metals in coastal regions of Indian Sundarban mangrove wetland: a multivariate approach. *J Clean Prod* 96:233–243. <https://doi.org/10.1016/j.jclepro.2014.04.030>
- Bhattacharya BD, Hwang JS, Sarkar SK, Rakshit D, Murugan K, Tseng LC (2015a) Community structure of mesozooplankton in coastal waters of Sundarban mangrove wetland, India: a multivariate approach. *J Mar Syst* 141:112–121
- Bhattacharyya S (2015) Study of changing facades of Sundarbans – A Remote Sensing & GIS based management approach, PhD Thesis in Department of Geological Sciences, Jadavpur University
- Bir J, Sumon MS, Rahaman S (2015) The effects of different water quality parameters on zooplankton distribution in major river systems of Sundarbans Mangrove. *IOSR J Environ Sci Toxicol Food Technol* 11:56–63
- Biswas H, Mukhopadhyay SK, De TK, Sen S, Jana TK (2004) Biogenic controls on the air-water carbon dioxide exchange in the Sundarban mangrove environment, northeast coast of Bay of Bengal, India. *Limnol Oceanogr* 49:95–101. <https://doi.org/10.4319/lo.2004.49.1.0095>
- Biswas H, Mukhopadhyay SK, Sen S, Jana TK (2007) Spatial and temporal patterns of methane dynamics in the tropical mangrove dominated estuary, NE coast of Bay of Bengal, India. *J Mar Syst* 68:55–64. <https://doi.org/10.1016/j.jmarsys.2006.11.001>

- Biswas H, Dey M, Ganguly D, De TK, Ghosh S, Jana TK (2010) Comparative analysis of phytoplankton composition and abundance over a two-decade period at the land–ocean boundary of a tropical mangrove ecosystem. *Estuar Coasts* 33(2):384–394
- Biswas SN, Rakshit D, Sarkar SK, Sarangi RK, Satpathy KK (2014) Impact of multispecies diatom bloom on plankton community structure in Sundarban mangrove wetland, India. *Mar Pollut Bull* 85(1):306–311
- Blomqvist S, Anneli G, Ragnar E (2004) *Limnol Oceanogr* 49(6):2236–2241 [AU7]
- Borges AV, Abril G (2011) 5.04 – carbon dioxide and methane dynamics in estuaries. In: Wolanski E, McLusky D (eds) *Treatise on estuarine and coastal science*. Academic Press, Waltham, pp 119–161. <https://doi.org/10.1016/B978-0-12-374711-2.00504-0>
- Chakrabarti C, Kundu S, Ghosh P, Choudhury A (1993) A preliminary study on certain trace metals in some plant and animal organisms from mangroves of Sundarbans, India. *Mahasagar* 26:17–20
- Chakraborty A, Bera A, Mukherjee A, Basak P, Khan I, Mondal A, Roy A, Bhattacharyya A, SenGupta S, Roy D, Nag S (2015) Changing bacterial profile of Sundarbans, the world heritage mangrove: impact of anthropogenic interventions. *World J Microbiol Biotechnol* 31(4):593–610
- Chanda A, Mukhopadhyay A, Ghosh T, Akhand A, Mondal P, Ghosh S, Mukherjee S, Wolf J, Lazar AN, Rahman MM, Salehin M, Chowdhury SM, Hazra S (2016) Blue carbon stock of the Bangladesh Sundarban Mangroves: what could be the scenario after a century? *Wetlands* 36:1033–1045. <https://doi.org/10.1007/s13157-016-0819-7>
- Chanton JP, Dacey JWH (1991) Effects of vegetation on methane flux, reservoirs, and carbon isotopic composition. In: Sharkey T, Holland E, Mooney H (eds) *Trace gas emissions from plants*. Academic Press, San Diego, pp 65–92
- Chatterjee M, Silva Filho EV, Sarkar SK, Sella SM, Bhattacharya A, Satpathy KK, Prasad MVR, Chakraborty S, Bhattacharya BD (2007) Distribution and possible source of trace elements in the sediment cores of a tropical macrotidal estuary and their ecotoxicological significance. *Environ Int* 33:346–356. <https://doi.org/10.1016/j.envint.2006.11.013>
- Chatterjee M, Shankar D, Sen GK et al (2013) Tidal variations in the Sundarbans estuarine system, India. *J Earth Syst Sci* 122:899–933. <https://doi.org/10.1007/s12040-013-0314-y>
- Chaudhuri K, Manna S, Sarma KS, Naskar P, Bhattacharyya S, Bhattacharyya M (2012) Physicochemical and biological factors controlling water column metabolism in Sundarbans estuary, India. *Aquat Biosyst* 8(1):26
- Chen CTA, Borges AV (2009) Reconciling opposing views on carbon cycling in the coastal ocean: continental shelves as sinks and near-shore ecosystems as sources of atmospheric CO₂. *Deep Sea Res Part II Top Stud Oceanogr*, Surface Ocean CO₂ Variability and Vulnerabilities 56:578–590. <https://doi.org/10.1016/j.dsr2.2009.01.001>
- Chowdhury A, Maiti SK (2016) Assessing the ecological health risk in a conserved mangrove ecosystem due to heavy metal pollution: a case study from Sundarbans Biosphere Reserve, India. *Hum Ecol Risk Assess Int J* 22:1519–1541. <https://doi.org/10.1080/10807039.2016.1190636>
- Chowdhury R, Favas PJC, Jonathan MP, Venkatachalam P, Raja P, Sarkar SK (2017) Bioremoval of trace metals from rhizosediment by mangrove plants in Indian Sundarban Wetland. *Mar Pollut Bull* 124:1078–1088. <https://doi.org/10.1016/j.marpolbul.2017.01.047>
- Chowdhury R, Sutradhar T, Begam MM, Mukherjee C, Chatterjee K, Basak SK, Ray K (2019) Effects of nutrient limitation, salinity increase, and associated stressors on mangrove forest cover, structure, and zonation across Indian Sundarbans. *Hydrobiologia* 842:191–217
- Das S, Chanda A, Giri S, Akhand A, Hazra S (2015) Characterizing the influence of tide on the physico-chemical parameters and nutrient variability in the coastal surface water of the northern Bay of Bengal during the winter season. *Acta Oceanol Sin* 34(12):102–111
- Das S, Mukherjee A, De TK, De M, Jana TK (2016) Influence of microbial composition on the carbon storage capacity of the mangrove soil at the land-ocean boundary in the Sundarban Mangrove Ecosystem, India. *Geomicrobiol J* 33:743–750. <https://doi.org/10.1080/01490451.2015.1093567>

- Das S, Giri S, Das I, Chanda A, Ghosh A, Mukhopadhyay A, Akhand A, Choudhury SB, Dadhwal VK, Maity S, Kumar TS (2017) Nutrient dynamics of northern Bay of Bengal (nBoB)—emphasizing the role of tides. *Reg Stud Mar Sci* 10:116–134
- Dayan AD, Paine AJ (2001) Mechanisms of chromium toxicity, carcinogenicity and allergenicity: review of the literature from 1985 to 2000. *Hum Exp Toxicol* 20:439–451. <https://doi.org/10.1191/096032701682693062>
- De TK, De M, Das S, Chowdhury C, Ray R, Jana TK (2011) Phytoplankton abundance in relation to cultural eutrophication at the land-ocean boundary of Sunderbans, NE Coast of Bay of Bengal, India. *J Environ Stud Sci* 1(3):169
- Dutta MK, Chowdhury C, Jana TK, Mukhopadhyay SK (2013) Dynamics and exchange fluxes of methane in the estuarine mangrove environment of the Sundarbans, NE coast of India. *Atmos Environ* 77:631–639. <https://doi.org/10.1016/j.atmosenv.2013.05.050>
- Dutta S, Chakraborty K, Hazra S (2017a) Ecosystem structure and trophic dynamics of an exploited ecosystem of Bay of Bengal, Sundarban Estuary, India. *Fish Sci* 83:145–159. <https://doi.org/10.1007/s12562-016-1060-2>
- Dutta MK, Bianchi TS, Mukhopadhyay SK (2017b) Mangrove methane biogeochemistry in the Indian Sundarbans: a proposed budget. *Front Mar Sci* 4. <https://doi.org/10.3389/fmars.2017.00187>
- Dutta MK, Kumar S, Mukherjee R, Sanyal P, Mukhopadhyay SK (2019a) The post-monsoon carbon biogeochemistry of the Hooghly–Sundarbans estuarine system under different levels of anthropogenic impacts. *Biogeosciences* 16:289–307
- Dutta MK, Kumar S, Mukherjee R, Sharma N, Acharya A, Sanyal P, Bhusan R, Mukhopadhyay SK (2019b) Diurnal carbon dynamics in a mangrove-dominated tropical estuary (Sundarbans, India). *Estuar Coast Shelf Sci* 229:106426. <https://doi.org/10.1016/j.ecss.2019.106426>
- Flood RP, Bloemsma MR, Weltje GJ et al (2016) Compositional data analysis of Holocene sediments from the West Bengal Sundarbans, India: geochemical proxies for grain-size variability in a delta environment. *Appl Geochem* 75:222–235. <https://doi.org/10.1016/j.apgeochem.2016.06.006>
- Flood RP, Barr ID, Weltje GJ et al (2018) Provenance and depositional variability of the Thin Mud Facies in the lower Ganges-Brahmaputra delta, West Bengal Sundarbans, India. *Mar Geol* 395:198–218. <https://doi.org/10.1016/j.margeo.2017.09.001>
- Gannon JE, Stemberger RS (1978) Zooplankton (especially crustaceans and rotifers) as indicators of water quality. *Trans Am Microsc Soc*:16–35
- Garrett RG (2000) Natural sources of metals to the environment. *Hum Ecol Risk Assess Int J* 6:945–963. <https://doi.org/10.1080/10807030091124383>
- Ghosh A, Bhadury P (2018) Investigating monsoon and post-monsoon variabilities of bacterioplankton communities in a mangrove ecosystem. *Environ Sci Pollut Res* 25(6):5722–5739
- Ghosh P, Choudhury A (1989) Copper, zinc and lead in the sediment of Hooghly estuary. *Environ Ecol* 7:427–430
- Ghosh A, Dey N, Bera A, Tiwari A, Sathyaniranjan K, Chakrabarti K, Chattopadhyay D (2010) Culture independent molecular analysis of bacterial communities in the mangrove sediment of Sundarban, India. *Saline Syst* 6(1):1
- Ghosh P, Chakrabarti R, Bhattacharya SK (2013) Short-and long-term temporal variations in salinity and the oxygen, carbon and hydrogen isotopic compositions of the Hooghly Estuary water, India. *Chem Geol* 335:118–127. <https://doi.org/10.1016/j.chemgeo.2012.10.051>
- Gole CV, Vaidyaraman PP (1967) Salinity distribution and effect of fresh water flows in the Hooghly River. *Coast Eng* 1966:1412–1434
- Gopal B, Chauhan M (2006) Biodiversity and its conservation in the Sundarban mangrove ecosystem. *Aquat Sci* 68:338–354
- Hanson RS, Hanson TE (1996) Methanotrophic bacteria. *Microbiol Mol Biol Rev* 60:439–471
- Islam MS, Sarkar MJ, Yamamoto T, Wahab MA, Tanaka M (2004) Water and sediment quality, partial mass budget and effluent N loading in coastal brackish water shrimp farms in Bangladesh. *Mar Pollut Bull* 48(5–6):471–485

- Järup L (2003) Hazards of heavy metal contamination. *Br Med Bull* 68:167–182. <https://doi.org/10.1093/bmb/ldg032>
- Jeffrey LC, Maher DT, Johnston SG, Kelaher BP, Steven A, Tait DR (2019) Wetland methane emissions dominated by plant-mediated fluxes: contrasting emissions pathways and seasons within a shallow freshwater subtropical wetland. *Limnol Oceanogr* 64:1895–1912. <https://doi.org/10.1002/lno.11158>
- Jha CS, Rodda SR, Thumaty KC, Raha AK, Dadhwal VK (2014) Eddy covariance based methane flux in Sundarbans mangroves, India. *J Earth Syst Sci* 123:1089–1096. <https://doi.org/10.1007/s12040-014-0451-y>
- Kamruzzaman M, Basak K, Paul SK, Ahmed S, Osawa A (2019) Litterfall production, decomposition and nutrient accumulation in Sundarbans mangrove forests, Bangladesh. *For Sci Technol* 15(1):24–32
- Khan RA (2003) Biodiversity of macrobenthos on the intertidal flats of Sunderban estuarine region, India. *Rec Zool Surv India* 101(3–4):181–205
- Koné YJM, Abril G, Delille B, Borges AV (2010) Seasonal variability of methane in the rivers and lagoons of Ivory Coast (West Africa). *Biogeochemistry* 100:21–37
- Kumar A, Ramanathan A (2015) Speciation of selected trace metals (Fe, Mn, Cu and Zn) with depth in the sediments of Sundarban mangroves: India and Bangladesh. *J Soils Sediments* 15:2476–2486. <https://doi.org/10.1007/s11368-015-1257-5>
- Maher DT, Drexler M, Tait DR, Johnston SG, Jeffrey LC (2019) iAMES: an inexpensive, automated methane ebullition sensor. *Environ Sci Technol* 53:6420–6426. <https://doi.org/10.1021/acs.est.9b01881>
- Manna S, Chaudhuri K, Bhattacharyya S, Bhattacharyya M (2010) Dynamics of Sundarban estuarine ecosystem: eutrophication induced threat to mangroves. *Saline Syst* 6(1):8
- Manna S, Chaudhuri K, Sarma KS, Naskar P, Bhattacharyya S, Bhattacharyya M (2012) Interplay of physical, chemical and biological components in estuarine ecosystem with special reference to Sundarbans, India. In: Voudouris K, Voutsas D (eds) *Ecological water quality—water treatment and reuse*. InTech, Rijeka, Croatia, pp 206–238
- Massolo S, Bignasca A, Sarkar SK, Chatterjee M, Bhattacharya BD, Alam A (2012) Geochemical fractionation of trace elements in sediments of Hooghly River (Ganges) and Sundarban wetland (West Bengal, India). *Environ Monit Assess* 184:7561–7577. <https://doi.org/10.1007/s10661-012-2519-y>
- Mitra A (1998) Status of coastal pollution in West Bengal with special reference to heavy metals. *J Indian Ocean Stud* 5:135–138
- Mitra A, Ghosh R (2014) Bioaccumulation pattern of heavy metals in commercially important fishes on and around Indian Sundarbans 12
- Mitra A, Banerjee K, Sengupta K, Gangopadhyay A (2009) Pulse of climate change in Indian Sundarbans: a myth or reality. *Natl Acad Sci Lett* 32:1–7
- Mitra A, Barua P, Zaman S, Banerjee K (2012) Analysis of trace metals in commercially important crustaceans Collected from UNESCO protected world heritage site of Indian Sundarbans. *Turk J Fish Aquat Sci* 12. https://doi.org/10.4194/1303-2712-v12_1_07
- Morgan JP, McIntire WG (1959) Quaternary geology of the Bengal Basin, East Pakistan and India. *Geol Soc Am Bull* 70:319–342
- Mukherjee D, Mukherjee A, Kumar B (2009) Chemical fractionation of metals in freshly deposited marine estuarine sediments of Sundarban ecosystem, India. *Environ Geol* 58:1757–1767. <https://doi.org/10.1007/s00254-008-1675-4>
- Mukhopadhyay SK, Jana TK, De TK, Sen S (2000) Measurement of exchange of CO₂ in mangrove forest of Sundarbans using micrometeorological method. *Trop Ecol* 41:57–60
- Mukhopadhyay SK, Biswas H, De TK, Sen BK, Sen S, Jana TK (2002a) Impact of Sundarban mangrove biosphere on the carbon dioxide and methane mixing ratios at the NE Coast of Bay of Bengal, India. *Atmos Environ* 36:629–638. [https://doi.org/10.1016/S1352-2310\(01\)00521-0](https://doi.org/10.1016/S1352-2310(01)00521-0)

- Mukhopadhyay SK, Biswas H, De TK, Sen S, Jana TK (2002b) Seasonal effects on the air-water carbon dioxide exchange in the Hooghly estuary, NE coast of Bay of Bengal, India. *J Environ Monit* 4:549–552. <https://doi.org/10.1039/b201614a>
- Padhy SR, Bhattacharyya P, Dash PK, Reddy CS, Chakraborty A, Pathak H (2020) Seasonal fluctuation in three mode of greenhouse gases emission in relation to soil labile carbon pools in degraded mangrove, Sundarban, India. *Sci Total Environ* 705:135909. <https://doi.org/10.1016/j.scitotenv.2019.135909>
- Paerl HW, Huisman J (2008) Blooms like it hot. *Science* 320(5872):57–58
- Paerl HW, Tucker CS (1995) Ecology of blue-green algae in aquaculture ponds. *J World Aquacult Soc* 26(2)
- Purvaja R, Ramesh R, Ray AK, Rixen T (2008) Nitrogen cycling: a review of the processes, transformations and fluxes in coastal ecosystems. *Curr Sci*:1419–1438
- Raha A, Das S, Banerjee K, Mitra A (2012) Climate change impacts on Indian Sunderbans: a time series analysis (1924–2008). *Biodivers Conserv* 21:1289–1307
- Rahman MT, Rahman S, Quraishi SB, Ahmad JU, Choudhury TR, Mottaleb MA (2011) Distribution of heavy metals in water and sediments in Passur River. *Sundarban Mangrove Forest, Bangladesh* 6:11
- Rahman MM, Khan MNI, Hoque AKF, Ahmed I (2015) Carbon stock in the Sundarbans mangrove forest: spatial variations in vegetation types and salinity zones. *Wetl Ecol Manag* 23:269–283. <https://doi.org/10.1007/s11273-014-9379-x>
- Ray R, Rixen T, Baum A, Malik A, Gleixner G, Jana TK (2015) Distribution, sources and biogeochemistry of organic matter in a mangrove dominated estuarine system (Indian Sundarbans) during the pre-monsoon. *Estuar Coast Shelf Sci* 167:404–413
- Ray R, Majumder N, Chowdhury C, Das S, Jana TK (2018a) Phosphorus budget of the Sundarban mangrove ecosystem: box model approach. *Estuar Coasts* 41(4):1036–1049
- Ray R, Baum A, Rixen T, Gleixner G, Jana TK (2018b) Exportation of dissolved inorganic and organic and particulate carbon from mangroves and its implication to the carbon budget in the Indian Sundarbans. *Sci Total Environ* 621:535–547. <https://doi.org/10.1016/j.scitotenv.2017.11.225>
- Rodda SR, Thumaty KC, Jha CS, Dadhwal VK (2016) Seasonal variations of carbon dioxide, water vapor and energy fluxes in tropical Indian mangroves. *Forests* 7:35. <https://doi.org/10.3390/f7020035>
- Rogers KG, Goodbred SL (2014) The Sundarbans and Bengal Delta: the world's largest tidal mangrove and delta system. In: Kale VS (ed) *Landscapes and landforms of India*. Springer Netherlands, Dordrecht, pp 181–187
- Rosentreter JA, Maher DT, Erler DV, Murray R, Eyre BD (2018) Seasonal and temporal CO₂ dynamics in three tropical mangrove creeks—a revision of global mangrove CO₂ emissions. *Geochim Cosmochim Acta* 222:729–745
- Roy M, Nandi NC (2012) Distribution pattern of macrozoobenthos in relation to salinity of Hooghly-Matla estuaries in India. *Wetlands* 32(6):1001–1009
- Roy S, Hens D, Biswas D, Biswas D, Kumar R (2002) Survey of petroleum-degrading bacteria in coastal waters of Sunderban Biosphere Reserve. *World J Microbiol Biotechnol* 18(6):575–581
- Roy D, Pramanik A, Banerjee S, Ghosh A, Chattopadhyay D, Bhattacharyya M (2018) Spatio-temporal variability and source identification for metal contamination in the river sediment of Indian Sundarbans, a world heritage site. *Environ Sci Pollut Res* 25:31326–31345
- Rudra K (2014) Changing river courses in the western part of the Ganga-Brahmaputra delta. *Geomorphology* 227:87–100. <https://doi.org/10.1016/j.geomorph.2014.05.013>
- Rudra K (2018) *Rivers of the Ganga – Brahmaputra – Meghna Delta: a fluvial account of Bengal*. Springer International, pp 1–190
- Sadhuram Y, Sarma VV, Murthy TR, Rao BP (2005) Seasonal variability of physico-chemical characteristics of the Haldia channel of Hooghly estuary, India. *J Earth Syst Sci* 114:37–49. <https://doi.org/10.1007/BF02702007>

- Samanta B, Bhadury P (2018) Study of diatom assemblages in Sundarbans mangrove water based on light microscopy and rbcL gene sequencing. *Heliyon* 4(6):e00663
- Samanta S, Dalai TK (2018) Massive production of heavy metals in the Ganga (Hooghly) river estuary, India: global importance of solute-particle interaction and enhanced metal fluxes to the oceans. *Geochim Cosmochim Acta* 228:243–258
- Santos HF, Carmo FL, Paes JE, Rosado AS, Peixoto RS (2011) Bioremediation of mangroves impacted by petroleum. *Water Air Soil Pollut* 216(1–4):329–350
- Sarkar SK, Frančišković-Bilinski S, Bhattacharya A, Saha M, Bilinski H (2004) Levels of elements in the surficial estuarine sediments of the Hooghly River, Northeast India and their environmental implications. *Environ Int* 30(8):1089–1098
- Sarkar S, Ghosh PB, Das TM et al (2013) Environmental assessment in terms of salinity distribution in the Tropical Mangrove forest of Sundarban, North East Coast of Bay of Bengal, India. *Arch Appl Sci Res* 5:109–118
- Sen A, Saha S, Maita SK, Saha T (2015) A study of planktonic communities along the Bidyadhari River stretch of the Raimangal Estuary of West Bengal, India. *Proceedings of the National Conference on Biodiversity – Issues, Concern & Future Strategies*, pp 49–53
- Sharma M, Chakraborty A, Kuttippurath J, Yadav AK (2018) Potential power production from salinity gradient at the Hooghly Estuary System. *Innov Ener Res* 7:1–7. <https://doi.org/10.4172/2576-1463.1000210>
- Singh G, Ramanathan AL, Santra SC, Ranjan RK (2016) Tidal control on the nutrient variability in Sundarban mangrove ecosystem. *J Appl Geochem* 18(4):495–503
- Sinha PC, Rao YR, Dubey SK et al (1996) Modeling of circulation and salinity in Hooghly estuary. *Mar Geod* 19:197–213. <https://doi.org/10.1080/01490419609388079>
- Sippo JZ, Maher DT, Tait DR, Ruiz-Halpern S, Sanders CJ, Santos IR (2017) Mangrove outwelling is a significant source of oceanic exchangeable organic carbon. *Limnol Oceanogr Lett* 2:1–8. <https://doi.org/10.1002/lo2.10031>
- Somayajulu BLK, Rengarajan R, Jani RA (2002) Geochemical cycling in the Hooghly estuary, India. *Mar Chem* 79:171–183. [https://doi.org/10.1016/S0304-4203\(02\)00062-2](https://doi.org/10.1016/S0304-4203(02)00062-2)
- Spalding M, Blasco F, Field C (1997) World mangrove atlas. International Society for Mangrove Ecosystems, Okinawa
- Vinh TV, Allenbach M, Joanne A, Marchand C (2019) Seasonal variability of CO₂ fluxes at different interfaces and vertical CO₂ concentration profiles within a *Rhizophora* mangrove forest (Can Gio, Viet Nam). *Atmos Environ* 201:301–309. <https://doi.org/10.1016/j.atmosenv.2018.12.049>
- Walsby A, Schanz F (2002) Light-dependent growth rate determines changes in the population of *Planktothrix rubescens* over the annual cycle in Lake Zürich, Switzerland. *New Phytol* 154(3):671–687
- Yang X, Post WM, Thornton PE, Jain A (2013) The distribution of soil phosphorus for global biogeochemical modeling. *Biogeosci Discuss (Online)* 9(4)

Author Queries

Chapter No.: 15 501933_1_En_15_Chapter

Queries	Details Required	Author's Response
AU1	Please check the hierarchy of the section headings and confirm if correct.	
AU2	Please provide details for RAMSAR 2019, Sodhi et al. 1987, Sumanta and Dalai 2018, Chatterjee et al.2009, Rahman et al. 2009 in reference list.	
AU3	Please fix “a” or “b” for Ray et al (2018) here and in other occurrences.	
AU4	Please fix “a” or “b” for Dutta et al. (2017).	
AU5	Please fix “a” or “b” for Mukhopadhyay et al. (2002).	
AU6	Please provide volume number and page range for Bhattacharjee et al. (2013).	
AU7	Please provide article title for Blomqvist et al. (2004).	
AU8	Please provide volume number for Gannon & Stemberger (1978), Purvaja et al. (2008).	
AU9	Please provide page range for Paerl & Tucker (1995), Yang et al. (2013).	
AU10	Please provide publisher location for Rudra (2018).	