

# Identification of *Escherichia coli* *ygaQ* and *rpmG* as novel mitomycin C resistance factors implicated in DNA repair

Edward L Bolt<sup>\*</sup>, Tabitha Jenkins<sup>†</sup>, Valeria Moreira Russo<sup>†</sup>, Sharlene Ahmed<sup>†</sup>, James Cavey and Simon D Cass.

School of Life Sciences, The University of Nottingham. UK

<sup>\*</sup>Correspondence to Ed Bolt; [ed.bolt@nottingham.ac.uk](mailto:ed.bolt@nottingham.ac.uk)

<sup>†</sup>These authors were final year undergraduate honours students

Keywords: DNA repair; mitomycin C; homologous recombination; rpmG; ygaQ

## ABSTRACT

Using the ASKA (A Complete Set of *E. coli* K-12 ORF Archive) library for genome-wide screening of *E. coli* proteins we identified that expression of *ygaQ* and *rpmG* promote mitomycin C resistance (MMC<sup>R</sup>). *YgaQ* mediated MMC<sup>R</sup> was independent of homologous recombination involving RecA or RuvABC, but required UvrD. *YgaQ* is an uncharacterized protein homologous to  $\alpha$ -amylases that we identified to have nuclease activity directed to single stranded DNA of 5' flaps. Nuclease activity was inactivated by mutation of two amino acid motifs, which also abolished MMC<sup>R</sup>. *RpmG* is frequently annotated as a bacterial ribosomal protein, although forms an operon with MutM glycosylase and a putative deubiquitinating enzyme, YicR. *RpmG* associated MMC<sup>R</sup> was dependent on MutM. MMC<sup>R</sup> from *RpmG* resembles DNA repair phenotypes reported for 'idiosyncratic ribosomal proteins' in eukaryotes.

## INTRODUCTION

Chemicals causing covalent modifications to DNA are cytotoxic when their products interfere with biological processes including DNA replication and gene transcription. Mitomycin C (MMC) provokes interstrand DNA crosslinks at 5'-GNC-3' or 5'-CG-3' sequences, and mono-adducts at guanine bases [1-3]. It is a natural antimicrobial synthesized by *Streptomyces caespitosus* that is effective as a treatment for human cancers, and there is continuing interest in mechanisms cells use to overcome genotoxic damage associated with MMC and other cross-linkers [4].

Removal and repair of MMC induced DNA damage, involves interplay between nucleotide excision repair, homologous recombination and repair polymerases. Recent reviews detail the multiple factors implicated in repair of DNA cross-links in human cells and in prokaryotes [4, 5]. In bacteria, UvrA, UvrB and UvrC nucleotide excision repair complexes recognize and eliminate DNA-MMC lesions [5]. DNA molecules generated during and after UvrABC processing can be used as substrates for gap repair by DNA polymerase I, and for homologous recombination initiated by RecA or RecFOR and controlled and completed by helicases (RecG, RuvAB, UvrD, RecQ), and resolvases (RuvC, RecU). The exact events post-excision of the lesion probably depend on the context of repair and the type of lesion being removed.

The importance of homologous recombination for repair of MMC cross-links in *E. coli* is illustrated by the high MMC sensitivity of cells lacking the Holliday junction helicase RuvAB or Holliday junction resolvase RuvC ( $\Delta$ *ruvAB*/ $\Delta$ *ruvC*) [6, 7]. RuvAB and RuvC associate into a 'RuvABC resolvosome' that assists in double strand break repair by branch migrating and resolving Holliday junction DNA into nicked DNA duplexes [8-13]. Similar activities of RuvABC at blocked replication forks can promote repair of blocking lesions and restart of replication [7, 14-16].  $\Delta$ *ruvABC* cells can be rescued from MMC sensitivity by expression of alternative Holliday junction nucleases, the archaeal resolvase Hjc [17], or bacteriophage Rusa [18]. Deletion in *E. coli* of base excision repair enzyme MutM (BER) and nucleotide excision repair (NER) enzymes UvrABC also cause acute sensitivity to MMC [5], highlighting how multiple DNA repair roles are needed to overcome genotoxic effects of MMC.

DNA repair has been intensively studied in *E. coli* to identify DNA repair pathways by genetic analysis, followed by detailed understanding of DNA repair enzyme structure and function [8, 19-24]. DNA repair genes may remain to be identified in *E. coli*, considering the unknown roles for about 30% of *E. coli* genes. A recent genetic screen in *E. coli* unearthed and validated roles for uncharacterized genes in promoting resistance to extreme ionizing radiation [22]. Using protein expression from the ASKA genomic library [25] we screened for MMC resistance of  $\Delta$ *ruvABC* cells, identifying four genes with a validated MMC<sup>R</sup> phenotype. Two of these, *ygaQ* and *rpmG*, are reported in more detail here.

## RESULTS

### Identification of *ygaQ* and *rpmG* as mitomycin C resistance factors in *E. coli*

We searched for uncharacterized genes in *E. coli* whose expression overcame growth inviability associated with mitomycin C (MMC) induced DNA damage. The genetic assay we used exploited the extreme MMC sensitivity of an *E. coli*  $\Delta$ *ruvAB* strain (Figure 1A) resulting from it lacking the RuvABC DNA repair complex. This followed a rationale from previous work identifying that the archaeal Holliday junction resolvase Hjc can restore mitomycin C resistance (MMC<sup>R</sup>) to  $\Delta$ *ruvAB* cells [17] (Figure 1B). An ASKA plasmid library [25] was transformed into *E. coli*  $\Delta$ *ruvAB*, followed by viability tests on MMC agar, resulting in twenty-one colonies with apparent MMC<sup>R</sup> compared to surrounding colonies on replica agar plates, summarized in Figure 1C. Four of these clones were verified for MMC<sup>R</sup> in multiple repeats of the same assay, judged by each growing comparably to pHjc on MMC agar (Supplementary Table S3). Two of these clones (pSTE5 and pDO4) had a strong negative fitness effect on cell viability when expressed

in “wild type” (RuvAB<sup>+</sup>) *E. coli*, and were therefore discarded from the remainder of this study. The two other MMC<sup>R</sup> ASKA clone plasmids, pSA2 and pVM6, were investigated further.

DNA sequencing confirmed that pVM6 and pSA2 contained, respectively, *E. coli* genes *ygaQ* and *rpmG*. *E. coli*  $\Delta$ *ruvAB* cells expressing *ygaQ* or *rpmG* (pYgaQ/pRpmG) were 1000-fold more viable than empty plasmid control, and this effect was dependent on IPTG induction of plasmid gene expression (Figure 1D). Western blotting of proteins from the same IPTG induced MMC<sup>R</sup> cultures detected proteins consistent with predicted sizes consistent with YgaQ and RpmG proteins that were absent from cells containing only empty plasmid vector (Figure 1E). YgaQ also showed multiple protein species of lower than expected molecular mass, probably representing isoforms or protein degradation. MMC<sup>R</sup> of pYgaQ or pRpmG colonies remained dependent on IPTG for plasmid gene expression when sub-cultured as fresh overnight growths, confirming that chromosomal suppressors did not account for the observed phenotype. The same cultures spread onto agar containing rifampicin (0-50  $\mu$ g/mL) did not show evidence of a mutator phenotype, which could promote MMC<sup>R</sup> independently of pYgaQ or pRpmG, compared to wild type cells and a  $\Delta$ *mutS* hyper-mutator control. YgaQ and RpmG were therefore further characterized: Aside from their ability to promote MMC<sup>R</sup> in *E. coli*, they have no obvious relationship to one another in genomic context or predicted protein function, as detailed below, and are therefore dealt with separately.

### Mutagenesis of YgaQ abolishes mitomycin C resistance

The *ygaQ* gene of *E. coli* strain W3110, used for construction of the ASKA library, encodes an uncharacterized protein of 110 amino acids with no conserved domains. *ygaQ* is present in *Escherichia* and *Shigella* species, located next to a predicted  $\alpha$ -amylase gene *ygaR*. In some *E. coli* strains (e.g. MG1655) it is predicted that *ygaQ* and *ygaR* are fused as a single open reading frame; more detailed analysis of YgaQ is presented in supplementary results Figures S2 and S3. Alignment of YgaQ homologues identified many conserved amino acids in YgaQ (Figure 2A). We focused on mutagenesis of individual and combined glutamic and aspartic acid amino acid residues within pYgaQ because of their essential catalytic roles in microbial alpha-amylases. Resulting plasmids were tested for their ability to promote MMC<sup>R</sup> in  $\Delta$ *ruvAB* cells compared to wild type pYgaQ. Combining mutations of Asp-25 and Asp-27 with mutation of a Glu-Arg-Lys triplet at the YgaQ C-terminus (a mutant subsequently referred to as YgaQ<sup>DM</sup>) abolished MMC<sup>R</sup>, similarly to the empty plasmid control (Figure 2B). Western blotting confirmed expression YgaQ<sup>DM</sup> like wild type protein (Figure 2C), confirming that protein mis-folding or instability is unlikely to explain MMC sensitivity from expressing this mutant YgaQ. Furthermore YgaQ<sup>DM</sup> was expressed as soluble protein for purification, as described below. We concluded that MMC<sup>R</sup> associated with YgaQ was specific to this protein, and that the mutagenized residues are important for the biological function of YgaQ when promoting MMC<sup>R</sup> in cells lacking DNA repair by RuvABC.

### YgaQ MMC<sup>R</sup> requires UvrD and acts independently of homologous recombination

Elimination of *ygaQ* ( $\Delta$ *ygaQ*) from RuvABC<sup>+</sup> *E. coli* had no effect on cell viability in mitomycin C agar compared to YgaQ<sup>+</sup> cells.  $\Delta$ *ruvAB* cells were very sick as expected, but combining  $\Delta$ *ygaQ* with  $\Delta$ *ruvAB* caused modest, but reproducible, increased sensitivity to mitomycin C compared to  $\Delta$ *ruvAB* alone (Figure 3A). This is consistent with YgaQ acting independently of RuvABC Holliday junction processing, as expected from the original screening analysis. We returned to pYgaQ to explore this further, testing if it restored MMC<sup>R</sup> to *E. coli*  $\Delta$ *ruvC* cells, in which RuvAB is present. The rationale for this test is based on inability of Hjc and RusA resolvases to restore MMC<sup>R</sup> to  $\Delta$ *ruvC* cells, even though they rescue  $\Delta$ *ruvAB* cells: access of Hjc and RusA to Holliday junctions is blocked by RuvAB. However, unlike Hjc and RusA, YgaQ restored MMC<sup>R</sup> to both  $\Delta$ *ruvC* cells, consistent with it not targeting Holliday junctions (Figure 3B).

We tested if pYgaQ restored MMC<sup>R</sup> to  $\Delta$ *ruvAB* cells that contained additional deletions of genes in DNA repair pathways: *recA*, for recombination dependent repair without Holliday junction formation, *dinG* and *umuD* for translesion synthesis, *recG* and *uvrD* for DNA repair linked to

replication stress, and *uvrB* for excision repair. Interestingly, only deletion of *uvrD* (therefore  $\Delta ruvAB \Delta uvrD$ ) caused pYgaQ to be unable to restore MMC<sup>R</sup> (Figure S1). These data indicate that MMC<sup>R</sup> from YgaQ expression is independent of homologous recombination, and that it might participate in UvrD driven DNA repair processes at blocked replication forks [26]. We purified *E. coli* YgaQ proteins to assay for DNA binding and catalytic activities to gain more understanding of involvement in MMC<sup>R</sup>.

### Purified YgaQ protein is a nuclease that targets single stranded DNA

*E. coli* strain W3110 YgaQ and YgaQ<sup>DM</sup> proteins were purified (Figure S4) and assayed *in vitro* for DNA binding and processing of branched DNA substrates that mimic intermediates formed during DNA repair, replication and recombination. EMSAs mixing purified YgaQ with DNA substrates were inconclusive in determining any substrate binding specificity because YgaQ repeatedly formed in-well aggregates rather than binding complexes. However, YgaQ catalytic activity was identified in similar reactions supplemented with 10 mM Mg<sup>2+</sup> and stopped by treating with proteinase K prior to electrophoresis (Figure 3C and D). Native gels showed YgaQ dependent product formation consistent with nuclease activity on partial and flayed duplex substrates containing ssDNA with a 5' end (Figure 3C). Fully base paired DNA substrates, or substrates with ssDNA terminating at a 3'OH, gave very weak or no activity. A preference for YgaQ targeting 5'-ended ssDNA was confirmed using denaturing gels (Figure 3D); nuclease activity was detected on ssDNA with 5'-terminus, but not on the strand with opposite polarity terminating in 3'OH. No activity was detected on the same strand in a fully based paired fork, confirming that YgaQ targets ssDNA. In the same assays YgaQ<sup>DM</sup> showed greatly reduced activity, in agreement with loss of the MMC<sup>R</sup> phenotype in genetic assays.

### MMC<sup>R</sup> associated with RpmG expression required the presence of MutM

RpmG is conserved widely across bacterial species, encoded within an operon of conserved gene order *yicR-rpmB-rpmG-mutM*. In *E. coli* the operon is transcribed into at least three mRNAs, possibly regulated by *creBC* [27, 28]. YicR, formerly called RadC [29], is a putative JAMM-family de-ubiquitinating (DUB) enzyme (reference [30]), and MutM is a DNA glycosylase required for base excision repair [31]. RpmG is a 53 amino acid protein that is frequently annotated in online databases as ribosomal protein L33, although its deletion in *E. coli* had no discernible effect on ribosome function [32]. We investigated if deleting any of *rpmB*, *yicR* and *mutM* affected pRpmG induced MMC<sup>R</sup> in  $\Delta ruvAB$  cells. MMC<sup>R</sup> associated with expression of pRpmG was lost in  $\Delta mutM$  cells (Figure 4B), but deletions of *yicR* or *rpmB* had no effect. Therefore MMC<sup>R</sup> associated with RpmG is functionally dependent on MutM. We were unable to identify any single or combined amino acid substitutions in RpmG that abolished MMC<sup>R</sup> (summarised in supplementary material). This is possibly because RpmG has a non-catalytic role that facilitates MutM activity through physical interaction involving multiple amino acids.

## DISCUSSION

ASKA libraries have been used to identify genetic factors that influence phenotypes in bacteria [37-39]. We used one version of this resource [25] to screen for novel *E. coli* genes involved in DNA repair, indicated by a mitomycin C resistance (MMC<sup>R</sup>) phenotype. MMC is a potent genotoxic agent by forming inter- and intra-strand cross-links in DNA that block replication and transcription and lead to cell death unless the lesion is repaired. One way to repair MMC lesions is through homologous recombination, which in *E. coli* can involve a "resolvasome" complex called RuvABC. The importance of RuvABC in MMC repair is exemplified by the MMC sensitivity of cells lacking RuvABC (e.g.  $\Delta ruvAB$  in Figure 1A). MMC<sup>R</sup> from expression of *ygaQ* was evident in both a  $\Delta ruvAB$  or  $\Delta ruvC$  strain. This is consistent with YgaQ acting aside from Holiday junction processing by RuvABC, but deletion of YgaQ alone had no MMC sensitivity phenotype, suggesting that at least in the growth conditions we used, YgaQ is subservient to

RuvABC. We speculate that actions of YgaQ in DNA repair or coping with genotoxic stress may become apparent only in response to specific stresses. An interesting observation made from combining  $\Delta ruvAB$  with other gene deletions was that the MMC<sup>R</sup> phenotype from YgaQ expression was lost when cells also lacked UvrD. This suggests that in *E. coli* lacking RuvABC and alternative system for dealing with MMC requires either combined actions of UvrD with YgaQ, or that YgaQ can promote recovery assisted by UvrD. UvrD is a facilitator of DNA repair by exposing lesions for further processing, in some cases by generating ssDNA for removal by nucleases [40]. We observed nuclease activity of YgaQ directed to 5'-ended single stranded DNA (ssDNA), which could therefore potentially degrade ssDNA generated from the 3' to 5' translocation polarity of UvrD. Nuclease activity of YgaQ was abolished or much reduced by introducing several amino acid substitutions (generating YgaQ<sup>DM</sup>), but not by individual amino acid substitutions. Database analyses of the W3110 strain 110 amino acid protein gave significant homology to  $\alpha$ -D-glucosidase, which hydrolyze oligosaccharides into their constituent sugars. It may be plausible that in YgaQ the same kind of fold could be utilized for binding to the DNA backbone leading to hydrolysis of phosphodiester bonds requiring DNA binding and active site chemistry of aspartate, glutamate and arginine residues that were mutated in YgaQ<sup>DM</sup>.

Observation of a role for RpmG in repair of DNA lesions in *E. coli* resembles the reported extra-ribosomal functions of "ribosomal" proteins in prokaryotes and eukaryotes, including in DNA repair [33, 34]. One such "idiosyncratic" ribosomal protein physically interacts with a eukaryotic DNA repair enzyme, stimulating its activity [35, 36]. It is possible that RpmG protein may act in a similar way with MutM.

## METHODS

### Processing the ASKA library for plasmid DNA

The ASKA library of plasmid encoded hexa-histidine tagged *E. coli* W3110 proteins was obtained from NBRP-E.coli at NIG (<http://www.shigen.nig.ac.jp/ecoli/strain/top/top.jsp>), described in [25]. The library contains 4364 open reading frames cloned individually into a plasmid vector for IPTG induced protein expression. Agar plates of up to 96 colonies were flooded with 3-5 mL of sterile LB broth and this was used to extract plasmid DNA by standard methods into 100  $\mu$ L of sterile water, thus generating sub-sections of the library, each containing up to 96 different *E. coli* W3110 genes. For this study, eight undergraduate students were each given aliquots of either five or six sub-sections of the library for transformation into *E. coli* MG1655  $\Delta ruvAB$  to begin the screening process, described below and in Figure 1B.

### Strains and plasmids

Details of the *E. coli* strains and plasmids used in this work are given in Supplementary Tables S1 and S2.

### Viability spot tests and P1 transductions

For viability spot tests of *E. coli* growths, LB cultures were grown to optical density (OD) of 0.4 measured at 600 nm, and then serially diluted 10-fold into M9 salts as indicated in the figures. Typically, 15  $\mu$ L of diluted cells was spotted onto appropriate LB agar.

Transductions were used to move around antibiotic resistance gene deletion cassettes using standard methods from P1 lysates, briefly: An overnight culture of the strain to be transduced was inoculated into 8 mL fresh broth and grown to optical density of 0.8. Pelleted cells were resuspended for 10 minutes at ambient temperature in 1 mL buffer MC (100 mM MgSO<sub>4</sub>, 5 mM CaCl<sub>2</sub>), prior to addition of P1 lysate of various titers prepared to contain the desired selectable gene cassette. Incubation, at typically 37°C for 30 minutes, was followed by addition of sodium citrate to 1 mM, followed by suspension of the P1-*E. coli* mixture in warm liquid 0.6% agar broth and plating onto agar containing the appropriate antibiotic selection. Incubation was at 37°C for

up to 48 hours to allow growth of resistant colonies that were then purified by antibiotic selection and verified for correct insertion of the desired gene cassette.

### Screening Aska plasmids for mitomycin C resistance in *E. coli* $\Delta$ *ruvAB* cells

This is summarized in Figure 1B. A group of 8 undergraduate research project students transformed an apramycin resistant  $\Delta$ *ruvAB* *E. coli* strain N6029 (Supplementary Table S1) with ASKA plasmids and plated cells onto chloramphenicol (15  $\mu$ g/mL) LB agar. In total approximately 11,000 of the resultant colonies were master gridded onto LB agar containing chloramphenicol (15  $\mu$ g/mL), and then replica plated onto LB agar containing either chloramphenicol (15  $\mu$ g/mL), or chloramphenicol plus mitomycin C (0.2  $\mu$ g/mL) and IPTG (0.5 mM). A positive control plasmid that gives MMC<sup>R</sup> in  $\Delta$ *ruvAB* *E. coli* by expressing the resolvase Hjc [17] was included in every stage to compare to ASKA clones. Note that using  $\Delta$ *ruvAB* cells for screening MMC<sup>R</sup> from the ASKA library was appropriate because *ruvA* and *ruvB* genes encoding the RuvAB complex (RuvA<sub>4</sub> or <sub>8</sub>-RuvB<sub>12</sub>), were on separate ASKA 96-well plates, removing the potential for false-positive MMC<sup>R</sup> that could arise if from *ruvA* and *ruvB* were encoded on the same plasmid.

### Mutagenesis of *ygaQ* and purification of *E. coli* YgaQ and YgaQ<sup>DM</sup> proteins

The ASKA plasmid containing *ygaQ* was mutagenized using the Q5 Base-Changer strategy from New England Biolabs. Primer sequences can be provided on request. For protein analysis the gene encoding *E. coli* W3110 YgaQ was synthesized using GeneArt (Life technologies), to include restriction sites for sub-cloning and optimization for codon usage. Sub-cloning of *ygaQ* into pET14b facilitated expression of N-terminally hexa-histidine tagged YgaQ. The same GeneArt process was used to synthesize the gene encoding YgaQ<sup>DM</sup>, with appropriate nucleotide substitutions for the following amino acid substitutions: D23G, D25G, E107G, R108S and K109STOP. YgaQ and YgaQ<sup>DM</sup> were over-expressed and purified in the same way: Briefly, strain BL21 AI harboring the desired plasmid was induced with arabinose at 37°C for 4 hours. Cells lysed in buffer (20 mM Tris. HCl pH 8.0, 500 mM NaCl, 10% (v/v) glycerol, 10 mM imidazole) were passed into a 5 mL Hi-Prep nickel chelation column, with YgaQ proteins luting within a gradient of 0-250 mM imidazole. Fractions containing YgaQ were pooled, dialyzed into a new buffer (20 mM Tris pH 8.0, 100 mM NaCl, 1 mM DTT, 10% (v/v) glycerol) and passed into a 5 mL Hi-Trap heparin column, to which YgaQ proteins did not bind but were collected in the flow-through.

### DNA assays

Base sequences of DNA strands used to construct substrates are given in the Supplementary materials. DNA strands were custom synthesized and HPLC purified by Sigma-Aldrich. DNA strands (300 ng) were <sup>32</sup>P labeled at their 5' ends by incubation with T4 polynucleotide kinase (PNK) and  $\gamma$ -<sup>32</sup>P-ATP (1 hour, 37°C) followed by heat inactivation of PNK. Unincorporated ATP was removed from these reactions using Bio-Spin 6 columns (Bio-Rad). Resulting end-labeled DNA was annealed to other unlabeled DNA strands (900 ng) in buffer (150 mM sodium chloride and 15 mM sodium citrate, pH 7.0) by heating to 95°C for two minutes followed by gradual cooling to room temperature. DNA substrates were then purified, to remove un-annealed oligonucleotide or incomplete DNA structures, by electrophoresis through a 10% acrylamide Tris-Borate-EDTA (TBE) gel followed by autoradiography, excision of gel slice and elution by diffusion at 4°C into 250-500  $\mu$ L of 10 mM Tris.HCl, 50 mM sodium chloride pH 7.5. Nuclease assays were in buffer HB (7 mM Tris.HCl pH 8.0, 9% glycerol, 50 mM NaCl, 100  $\mu$ g/mL BSA) supplemented with 10 mM magnesium chloride at 37°C for 10 minutes. Reactions were stopped by addition of 1 mg/mL proteinase K, 2.5 % w/v SDS prior to electrophoresis through 10% TBE-acrylamide gels 1xTBE buffer.

### ACKNOWLEDGEMENTS

We thank former undergraduate students in The School of Life Sciences for their efforts in analyzing ASKA clones that are not reported in this work: Thiemo Leijeser, Natalie Mettam, Scott Ludford, Elliot McGuinn, Stephen Lawlor, David Omojomolo and Will Hindley.

### Figure legends

**Figure 1. (A).** Viability spot test to illustrate mitomycin C sensitivity of the *E. coli*  $\Delta ruvAB$  strain used for screening the ASKA library for mitomycin C resistance ( $MMC^R$ ). **(B).** The screening procedure. Plasmid DNA isolated from combining typically 96 colonies from an individual ASKA library agar plate was transformed into *E. coli*  $\Delta ruvAB$ . Growth of colonies after plating out on LB agar containing MMC was used to assess  $MMC^R$  when compared to that given by plasmid expression of Hjc resolvase as a positive control, as shown in the panel. Further experimental details, including how *ruvAB* induced false positives were avoided, are given in the methods section. **(C).** Example of a  $MMC^R$  clone arising from the ASKA screen. The panels show details of agar plates after gridding individual colonies in the presence or absence of MMC as indicated. **(D).** Analysis of  $MMC^R$  provided by expression of YgaQ or RpmG, dependent on addition of IPTG to growth media. The graph compares viable colony counts from spot tests in triplicate using  $\Delta ruvAB$  cells transformed by either pHjc (a positive control that restores  $MMC^R$  [17]) and its corresponding empty vector (empty 1, pT7-7), or by ASKA plasmids (Table S2) harbouring *rpmG* (SA2) or *ygaQ* (VM6) and its empty plasmid control (empty 2). A photograph of an example viability spot test for these clones is presented in the panels below. **(E).** Western blot of total cell protein extracted from cultures used to make the viability spot tests shown in (D). YgaQ and RpmG proteins were detected using antibody against their hexa-histidine tag.

**Figure 2. (A).** A ClustalW alignment of YgaQ amino acid sequences from *E. coli* W3110 (the strain used to make the ASKA library [25], *E. coli* MG1655, *Shigella boydii*, and *E. coli* DH1. Highlighted in boxes are the two regions of W3110 YgaQ that when mutagenized in combination gave YgaQ<sup>DM</sup> that could not promote  $MMC^R$  and gave nuclease defective YgaQ protein. **(B).** Example of a viability spot test comparing the  $MMC^R$  of expression from ASKA plasmids YgaQ or YgaQ<sup>DM</sup>. **(C).** Western blot confirming that YgaQ<sup>DM</sup> protein is expressed like YgaQ in cells used for the viability spot test in (B).

**Figure 3. (A).** Graph “killing curves” comparing strains  $\Delta ygaQ$ ,  $\Delta ruvAB$  and  $\Delta ruvAB \Delta ygaQ$  for MMC sensitivity in viability spot tests plotted as a function of MMC concentration as indicated. The assays were done in triplicate with bars representing standard error. **(B).** Graph showing survival of ASKA plasmids expressing YgaQ (pYgaQ) or RpmG (pRpmG) compared to the positive control pHjc and corresponding empty ASKA plasmid vector. Assays were done twice and standard error from the mean is given as bars. **(C).** Non-denaturing TBE acrylamide gel for analysis of products from mixing YgaQ with DNA substrates as indicated. YgaQ was used at 0, 2.5, 25 and 250 nM (lanes 1-12) or 0, 25 and 250 nM (lanes 13-21) in reactions containing 0.6 nM of DNA that was <sup>32</sup>P 5'-end labeled as indicated with (\*). **(D).** Urea denaturing TBE acrylamide gels for analysis of products from mixing YgaQ with forked DNA as indicated; in each substrate the strand presented in grey is labeled at its 5' end. YgaQ and YgaQ<sup>DM</sup> mutant proteins were each used at 0, 2.5, 25 or 250 nM in reactions containing 0.6 nM of DNA.

**Figure 4.** Viability spot test of  $MMC^R$  from expressing RpmG (pRpmG) in  $\Delta ruvAB$  cells compared to cells  $\Delta ruvAB \Delta mutM$ , as indicated.

## REFERENCES

1. Tomasz, M., *Mitomycin C: small, fast and deadly (but very selective)*. Chem Biol, 1995. **2**(9): p. 575-9.
2. Ramos, L.A., et al., *The major mitomycin C-DNA monoadduct is cytotoxic but not mutagenic in Escherichia coli*. Chem Res Toxicol, 1998. **11**(1): p. 64-9.
3. Stevens, C.L., et al., *Chemistry and Structure of Mitomycin C*. J Med Chem, 1965. **8**: p. 1-10.
4. Clauson, C., O.D. Scharer, and L. Niedernhofer, *Advances in understanding the complex mechanisms of DNA interstrand cross-link repair*. Cold Spring Harb Perspect Biol, 2013. **5**(10): p. a012732.
5. Kisker, C., J. Kuper, and B. Van Houten, *Prokaryotic nucleotide excision repair*. Cold Spring Harb Perspect Biol, 2013. **5**(3): p. a012591.
6. Shurvinton, C.E. and R.G. Lloyd, *Damage to DNA induces expression of the ruv gene of Escherichia coli*. Mol. Gen. Genet., 1982. **185**: p. 352-355.
7. Le Masson, M., Z. Baharoglu, and B. Michel, *ruvA and ruvB mutants specifically impaired for replication fork reversal*. Mol Microbiol, 2008. **70**(2): p. 537-48.
8. West, S.C., *The processing of recombination intermediates: mechanistic insights from studies of bacterial proteins*. Cell, 1994. **76**: p. 9-15.
9. West, S.C., *The RuvABC proteins and Holliday junction processing in Escherichia coli*. J. Bacteriol., 1996. **178**: p. 1237-1241.
10. West, S.C., *Processing of recombination intermediates by the RuvABC proteins*. Ann. Rev. Genet., 1997. **31**: p. 213-244.
11. Benson, F.E., et al., *Nucleotide sequencing of the ruv region of Escherichia coli K-12 reveals a LexA regulated operon encoding two genes*. Nucl. Acids Res., 1988. **16**: p. 1541-1549.
12. Whitby, M.C., et al., *Interactions between RuvA and RuvC at Holliday junctions: inhibition of junction cleavage and formation of a RuvA-RuvC-DNA complex*. J. Mol. Biol., 1996. **264**: p. 878-890.
13. Connolly, B., et al., *Resolution of Holliday junctions in vitro requires Escherichia coli ruvC gene product*. Proc. Natl. Acad. Sci. USA, 1991. **88**: p. 6063-6067.
14. McGlynn, P. and R.G. Lloyd, *Action of RuvAB at replication fork structures*. J Biol Chem, 2001. **276**(45): p. 41938-44.
15. Flores, M.-J., et al., *Impairment of lagging strand synthesis triggers the formation of a RuvABC substrate at replication forks*. EMBO J., 2001. **20**: p. 619-629.
16. Michel, B., et al., *Recombination proteins and rescue of arrested replication forks*. DNA Repair (Amst), 2007. **6**(7): p. 967-80.
17. Bolt, E.L., R.G. Lloyd, and G.J. Sharples, *Genetic analysis of an archaeal Holliday junction resolvase in Escherichia coli*. J. Mol. Biol., 2001. **310**(3): p. 577-89.
18. Mahdi, A.A., et al., *Holliday junction resolvases encoded by homologous rusA genes in Escherichia coli K-12 and phage 82*. J. Mol. Biol., 1996. **257**: p. 561-573.
19. Kowalczykowski, S.C., et al., *Biochemistry of homologous recombination in Escherichia coli*. Microbiol. Rev., 1994. **58**: p. 401-465.
20. Eggleston, A.K., A.H. Mitchell, and S.C. West, *In vitro reconstitution of the late steps of genetic recombination in E. coli*. Cell, 1997. **89**(4): p. 607-617.
21. Sancar, A. and W.D. Rupp, *A novel repair enzyme: UVRABC excision nuclease*. Cell, 1983. **33**: p. 249-260.




22. Byrne, R.T., et al., *Escherichia coli* genes and pathways involved in surviving extreme exposure to ionizing radiation. *J Bacteriol*, 2014. **196**(20): p. 3534-45.
23. Clark, A.J. and A.D. Margulies, *Isolation and Characterization of Recombination-Deficient Mutants of Escherichia Coli K12*. *Proc Natl Acad Sci U S A*, 1965. **53**: p. 451-9.
24. Cox, M.M., et al., *The importance of repairing stalled replication forks*. *Nature*, 2000. **404**(6773): p. 37-41.
25. Kitagawa, M., et al., *Complete set of ORF clones of Escherichia coli ASKA library (a complete set of E. coli K-12 ORF archive): unique resources for biological research*. *DNA Res*, 2005. **12**(5): p. 291-9.
26. Flores, M.J., N. Sanchez, and B. Michel, *A fork-clearing role for UvrD*. *Mol Microbiol*, 2005. **57**(6): p. 1664-75.
27. Avison, M.B., et al., *Escherichia coli CreBC is a global regulator of gene expression that responds to growth in minimal media*. *J Biol Chem*, 2001. **276**(29): p. 26955-61.
28. Gifford, C.M. and S.S. Wallace, *The genes encoding formamidopyrimidine and MutY DNA glycosylases in Escherichia coli are transcribed as part of complex operons*. *J Bacteriol*, 1999. **181**(14): p. 4223-36.
29. Attaiech, L., et al., *RadC, a misleading name?* *J Bacteriol*, 2008. **190**(16): p. 5729-32.
30. Ambroggio, X.I., D.C. Rees, and R.J. Deshaies, *JAMM: a metalloprotease-like zinc site in the proteasome and signalosome*. *PLoS Biol*, 2004. **2**(1): p. E2.
31. Jiricny, J., *DNA repair: how MutM finds the needle in a haystack*. *Curr Biol*, 2010. **20**(4): p. R145-7.
32. Maguire, B.A. and D.G. Wild, *The roles of proteins L28 and L33 in the assembly and function of Escherichia coli ribosomes in vivo*. *Mol Microbiol*, 1997. **23**(2): p. 237-45.
33. Wood, I., *Extraribosomal functions of ribosomal proteins*. *Trends Biochem Sci*, 1996. **21**: p. 164-165.
34. Bhavsar, R.B., L.N. Makley, and P.A. Tsonis, *The other lives of ribosomal proteins*. *Hum Genomics*, 2010. **4**(5): p. 327-44.
35. Hegde, V., M. Wang, and W.A. Deutsch, *Human ribosomal protein S3 interacts with DNA base excision repair proteins hAPE/Ref-1 and hOGG1*. *Biochemistry*, 2004. **43**(44): p. 14211-7.
36. Hegde, V., et al., *The high binding affinity of human ribosomal protein S3 to 7,8-dihydro-8-oxoguanine is abrogated by a single amino acid change*. *DNA Repair (Amst)*, 2006. **5**(7): p. 810-5.
37. Yogiara, et al., *Escherichia coli ASKA Clone Library Harboring tRNA-Specific Adenosine Deaminase (tadA) Reveals Resistance towards Xanthorrhizol*. *Molecules*, 2015. **20**(9): p. 16290-305.
38. Fang, X., et al., *GIL, a new c-di-GMP-binding protein domain involved in regulation of cellulose synthesis in enterobacteria*. *Mol Microbiol*, 2014. **93**(3): p. 439-52.
39. Wong, J.L., S.L. Vogt, and T.L. Raivio, *Using reporter genes and the Escherichia coli ASKA overexpression library in screens for regulators of the Gram negative envelope stress response*. *Methods Mol Biol*, 2013. **966**: p. 337-57.
40. Kuper, J. and C. Kisker, *DNA Helicases in NER, BER, and MMR*. *Adv Exp Med Biol*, 2013. **767**: p. 203-24.


ACCEPTED MANUSCRIPT

**Figure 1**

**A.**



**B.**



**Figure 1**




Figure 1

E.




Figure 2

A.


D23-D25

```

MFSIKPGPRNLPIDNPTLLSWNIIDGDLNSKLNLTLEYLNCITNIINSCGVYPQGLKDRREI 60
MFSIKPGPRNLPIDNPTLLSWNIIDGDLNSKLNLTLEYLNCITNIINSCGVYPQGLKDRREI 60
MFSIKPGPGNLPIDNPTLLSWNIIDGDLNSKFNLTLEYLNCITNIINACGVYPQDLKDRREI 60
MFSIKPGPRKLPIDNPSLLSWNIIDGDLNSKLNLTLEYLNCVTNIINACGVYPQDLKDRREI 60
*****:*****:*****:*****:*****:*****:*****
*****:*****:*****:*****:*****:*****:*****
OLE ERK 107-109
ISTFHAEKVINDLLKNDYKISLSPDITYRELNKAQRSITAPDRI GERKT 110
ISTFHAEKVINDLLKNDYKISLSPDITYRELNKAQRSITAPDRI GERKT MFSIKPGPRN 120
ISTFHAEKVINDLLKNDYKISLSPDITYRELNKAQRSITAPDRI GERKT 110
ISTFHAEKVINDLLKNDYKISLSPDITYRELNKAQRSITAPDRI GERKT 110
*****:*****:*****:*****:*****:*****:*****
*****:*****:*****:*****:*****:*****:*****

```

B.



C.






Figure 3



**Figure 3**

