

Glutathione reductase *gsr-1* is an essential gene required for *Caenorhabditis elegans* early embryonic development

José Antonio Mora-Lorca^{a,b}, Beatriz Sáenz-Narciso^c, Christopher J. Gaffney^d, Francisco José Naranjo-Galindo^{a,1}, José Rafael Pedrajas^e, David Guerrero-Gómez^a, Agnieszka Dobrzynska^f, Peter Askjaer^f, Nathaniel J. Szewczyk^d, Juan Cabello^c, Antonio Miranda-Vizuete^{a,*}

^aInstituto de Biomedicina de Sevilla, Hospital Universitario Virgen del Rocío/CSIC/Universidad de Sevilla, 41013 Sevilla, Spain

^bDepartamento de Farmacología, Facultad de Farmacia, Universidad de Sevilla, 41012 Sevilla, Spain

^cCenter for Biomedical Research of La Rioja (CIBIR), 26006 Logroño, Spain

^dMRC/ARUK Centre for Musculoskeletal Ageing Research, University of Nottingham and Medical School Royal Derby Hospital, DE22 3DT Derby, United Kingdom

^eGrupo de Bioquímica y Señalización Celular, Departamento de Biología Experimental, Universidad de Jaén, 23071 Jaén, Spain

^fAndalusian Center for Developmental Biology (CABD), CSIC/JA/Universidad Pablo de Olavide, 41013 Seville, Spain

*To whom correspondence should be addressed: Antonio Miranda-Vizuete, Instituto de Biomedicina de Sevilla (IBIS), Hospital Universitario Virgen del Rocío, 41013 Sevilla, Spain. Tel.: +34 955 923061; Fax: +34 955 923101; E-mail: amiranda-ibis@us.es

¹Present Address: Functional Genomics and Proteomics, Department of Biology, KU Leuven, 3000 Leuven, Belgium

Keywords: *Caenorhabditis elegans*, embryonic development, glutathione reductase, mitochondria, redox

28 HIGHLIGHTS

- 29 • *C. elegans gsr-1* gene encodes cytoplasmic and mitochondria isoforms of glutathione reductase.
- 30
- 31 • *gsr-1* is essential for *C. elegans* embryonic development.
- 32
- 33 • The lethality of *gsr-1* mutants is due to a specific requirement of GSR-1 protein in the cytoplasm.
- 34
- 35 • *gsr-1* embryos have a progressive cell division delay and an aberrant distribution of interphasic
- 36 chromatin.
- 37
- 38 • *gsr-1* worms with maternally contributed GSR-1 are able to reach adulthood but display
- 39 mitochondria-associated phenotypes such as increased fragmentation, decreased mitochondrial
- 40 membrane potential and induction of mitochondrial UPR.

43 ABSTRACT

44 Glutathione is the most abundant thiol in the vast majority of organisms and is maintained in its reduced
45 form by the flavoenzyme glutathione reductase. In this work, we describe the genetic and functional
46 analysis of the *Caenorhabditis elegans gsr-1* gene that encodes the only glutathione reductase protein
47 in this model organism. By using green fluorescent protein reporters we demonstrate that *gsr-1*
48 produces two GSR-1 isoforms, one located in the cytoplasm and one in the mitochondria. *gsr-1* loss of
49 function mutants display a fully penetrant embryonic lethal phenotype characterized by a progressive
50 and robust cell division delay accompanied by an aberrant distribution of interphasic chromatin in the
51 periphery of the cell nucleus. Maternally expressed GSR-1 is sufficient to support embryonic
52 development but these animals are short-lived, sensitized to chemical stress and have increased
53 mitochondrial fragmentation and lower mitochondrial DNA content. Furthermore, the embryonic lethality
54 of *gsr-1* worms is prevented by restoring GSR-1 activity in the cytoplasm but not in mitochondria. Given
55 the fact that the thioredoxin redox systems are dispensable in *C. elegans*, our data support a prominent
56 role of the glutathione reductase/glutathione pathway in maintaining redox homeostasis in the
57 nematode.

58 **INTRODUCTION**


59 Maintenance of thiol redox homeostasis is crucial for survival and the thioredoxin and glutaredoxin
60 systems are the two main pathways that control the redox status in virtually all organisms [1]. The
61 thioredoxin system is composed of thioredoxin reductase (TrxR) and thioredoxins (Trx) while the
62 glutaredoxin system comprises glutathione reductase (GR), glutathione (GSH) and glutaredoxins (Grx),
63 where thioredoxins and glutaredoxins operate as terminal oxidoreductases using the reducing power of
64 NADPH [2]. These two systems are mechanistically and structurally very similar, regulating the
65 formation of disulfides within and between proteins, with the main difference being the use of the
66 tripeptide glutathione (L-γ-glutamyl-L-cysteinyl-glycine) as electron donor for glutaredoxins in the
67 glutaredoxin system. In addition, the glutaredoxin system also catalyzes the formation of disulfides
68 between proteins and GSH, namely glutathionylation, which has been shown to be an important
69 posttranslational modification, modulating the activity of many proteins [3].

70

71 **Thioredoxin system**

72


73



74 **Glutaredoxin system**

75

76



77 GSH is found in cyanobacteria, proteobacteria and some gram-positive bacteria as well as most
78 eukaryotes [4]. Inactivating mutations in the gene encoding γ-glutamylcysteine synthetase, the enzyme
79 that catalyzes the first step in GSH synthesis, are lethal in all organisms studied from yeast to
80 mammals, including plants [5-9] highlighting the physiological relevance of GSH in these organisms
81 Moreover, mutations in the gene encoding glutathione synthetase, that catalyzes the second, final step
82 of GSH synthesis, are also lethal in *Arabidopsis thaliana* [10] and *Mus musculus* [11] while they are

83 viable in *Saccharomyces cerevisiae* [12] and *Drosophila melanogaster* [13]. In these latter organisms, it
84 has been proposed that the intermediate γ -glutamyl-cysteine, which accumulates in glutathione
85 synthetase mutants, substitutes for GSH to allow growth. Of note, although the *C. elegans* glutathione
86 synthetase ortholog, *gss-1*, has not been yet characterized, a *gss-1(tm672)* deletion mutant has been
87 reported as lethal/sterile by the NBRP *C. elegans* Gene Knockout Consortium
88 (<http://www.shigen.nig.ac.jp/c.elegans/>).

89 The lethal phenotype of GSH synthesis mutants contrasts with the dispensability, in the vast
90 majority of eukaryotes, of the glutathione reductase gene that encodes the enzyme that recycles
91 reduced GSH from its oxidized form GSSG. This inessentiality of glutathione reductase for normal
92 growth and development is explained by the thioredoxin system being able to reduce GSSG, a trait that
93 is conserved from bacteria to mammals [14] and also by either accumulating GSSG in yeast vacuole
94 [15] or by excreting GSSG in mammalian cells [16]. Exceptions to glutathione reductase dispensability
95 are the fission yeast *Schizosaccharomyces pombe* [17], the *Plasmodium berghei* parasite in its
96 mosquito oocyst stage but not in the blood stage [18] and the *A. thaliana* GR2 gene, encoding a
97 chloroplastic/mitochondrial glutathione reductase whose inactivation causes early embryonic lethality
98 [19]. The lethality of *A. thaliana* GR2 mutants is most likely due to a deficiency in chloroplasts function,
99 as in this organism mitochondrial thioredoxin reductase TRXR2 is able to reduce GSSG in mitochondria
100 (Meyer AJ, personal communication). In addition, *D. melanogaster* (and probably other insects) lacks a
101 *bona fide* glutathione reductase gene and the reduction of GSSG is performed by the thioredoxin
102 system, which is essential in this organism [20].

103 All eukaryotic organisms relying on glutathione metabolism have two distinct pools of glutathione
104 reductase activity located in cytoplasm and mitochondria, respectively (with the exception of
105 photosynthetic organisms that also have glutathione reductase activity in chloroplasts and peroxisomes
106 [21]). In most cases, one single gene encodes both cytoplasmic and mitochondrial glutathione
107 reductase isoforms by the use of alternative translation initiation sites [22-25]. Whereas in *E. coli* the
108 thioredoxin and glutathione systems are functionally redundant in maintaining redox homeostasis [26],
109 important differences on their respective contribution to the redox status of the different subcellular
110 compartments have been identified in eukaryotic organisms, mainly through work in yeast and
111 mammals. Hence, the yeast thioredoxin system controls the thiol redox status in the cytoplasm where
112 the glutathione system acts merely as a backup. In contrast, the glutathione system is the dominant
113 system to maintain thiol redox control in yeast mitochondria [27-29]. In mammals, knock-out mice
114 lacking glutathione reductase are viable [30] while inactivation of the components of the cytoplasmic
115 and mitochondrial thioredoxin systems causes embryonic lethality in mice [31-34], initially suggesting
116 that the mammalian thioredoxin system may have a predominant role in maintaining redox

117 homeostasis. However, studies on mice harboring conditional alleles of cytosolic thioredoxin reductase
118 TrxR1 to bypass embryonic lethality demonstrate that TrxR1-null mice and cells are robustly viable,
119 relying on the glutathione pathway for survival [35, 36]. Indeed, mice lacking both TrxR1 and glutathione
120 reductase in all hepatocytes sustain hepatic redox homeostasis and organismal survival, through *de*
121 *novo* GSH synthesis via the transulfuration pathway using dietary methionine as cysteine precursor
122 [16]. Together, these data in mammals uphold a more prominent role of the glutathione pathway on
123 maintaining redox homeostasis while the thioredoxin system appears to play key functions during
124 embryonic development.

125 In contrast to mammals, the *C. elegans* thioredoxin system is dispensable for embryonic and
126 postembryonic development, as mutants lacking both cytoplasmic and mitochondrial thioredoxin
127 reductases, *txxr-1* and *txxr-2*, are viable and reach adulthood indistinguishably from wild type controls [37,
128 38], supporting the idea that the glutathione system is also the main responsible for maintenance of
129 redox homeostasis in worms. Lüersen *et al.* have shown that the *C. elegans gsr-1* gene encodes a
130 functional glutathione reductase protein [39], which is required for survival to paraquat and juglone
131 treatments. In this work, we demonstrate that the *C. elegans gsr-1* gene encodes both cytoplasmic and
132 mitochondrial glutathione reductase isoforms and that is required for embryonic development. This
133 lethal phenotype of *gsr-1* mutants arises from a specific requirement of the enzyme in the cytoplasm,
134 the subcellular compartment where GSH is synthesized. In contrast, *gsr-1* mutants maternally
135 expressing GSR-1 are able to develop normally but are sensitized to stress, are short-lived and have
136 compromised mitochondria.

137

RESULTS

***C. elegans* GSR-1 is widely expressed and is targeted to both cytoplasm and mitochondria.**

The *gsr-1* locus is organized into 5 exons and 4 introns and expresses two main mRNA variants, *gsr-1a* and *gsr-1b1* (**Figure 1A**), whose conceptual translation results into two different isoforms, GSR-1a and GSR-1b, with the former having an additional 14 amino acid N-terminal extension (**Figure 1B**). Two other minor mRNA variants have been reported, *gsr-1b2* and *gsr-1b3*, which differ only in the sequence of their respective 5'-UTRs but also generate the GSR-1b isoform (<http://www.wormbase.org/>). *C. elegans* GSR-1 is highly homologous through all protein domains to vertebrate glutathione reductases, including the conserved redox active site CVNVGC (**Figure 1B-C**). The GSR-1a N-terminal extension, mostly encoded by the first *gsr-1* exon, displays characteristics of a mitochondrial targeting sequence (MTS) [40], cleavable by the two matrix protease model (**Figure 1D**) [41]. This predicts that GSR-1a is located in the mitochondrial matrix while GSR-1b, lacking this putative MTS and initiated from a downstream in-frame ATG codon located at the beginning of the second exon (**Figure 1A**), is a cytoplasmic protein.

To demonstrate this dual subcellular localization *in vivo* as well as to describe in detail the tissue and cellular expression pattern of both GSR-1 isoforms, we first generated transgenic worms expressing a translational GFP fusion spanning a 2.2 kb fragment of the *gsr-1* promoter plus the complete *gsr-1* genomic ORF (**Figure 2A**). These transgenic animals showed strong fluorescence in pharynx while weaker labeling was found in hypodermis, intestine, vulva muscle cells, spermatheca, uterine cells, coelomocytes, gonad sheath cells, rectal cells and a couple of unidentified neurons (probably PVPL/R) in the tail (**Figure 2B-E**). Long first introns have been shown to be important for gene regulation by binding a set of transcription factors different to those binding the upstream promoter [42]. As *gsr-1* exons 1 and 2 are separated by a 318 bp intron, considerably larger than the rest of *gsr-1* introns (**Figure 1A**), we set to investigate whether the *gsr-1* first intron also has promoter activity. Interestingly, transgenic worms expressing a translational GFP fusion spanning the *gsr-1* ORF from the first intron (**Figure 2F**) show labeling in some neurons in the head, excretory channel, intestine and coelomocytes (**Figure 2G-K**). This result suggests that the *gsr-1* first intron possess intrinsic promoter activity, independent from that of the *gsr-1* upstream promoter.

To prove that *C. elegans gsr-1* gene encodes both mitochondrial and cytoplasmic isoforms *in vivo*, we generated transgenic worms expressing the full *gsr-1* translational GFP fusion in which either ATG codon is mutated (**Figure 2L**). Thus, inactivation of the second ATG codon forces translation from the first ATG codon resulting in a dotted GFP pattern, consistent with mitochondrial localization (best seen in the pharynx) [43] (**Figure 2M**). In turn, when forcing expression from the first ATG codon by inactivation of the second ATG codon, a diffuse GFP labeling indicating a cytoplasmic localization was

obtained (**Figure 2N**). Similar diffuse fluorescence labeling was found when removing *gsr-1* first exon, which encodes most of GSR-1a MTS (**data not shown**). We further demonstrate that the additional 14 amino acids of GSR-1a constitute a functional MTS on its own as, when expressed in worm muscle cells, the fusion protein MTS*gsr-1*::GFP is found in the typical tubular distribution along the cell longitudinal axis, colocalizing with the TOMM-20::mRFP mitochondrial marker [38] (**Figure 2Q-T**). Together, we conclude that the *C. elegans gsr-1* gene encodes both mitochondrial (GSR-1a) and cytoplasmic (GSR-1b) isoforms. In addition, we further refine and expand the *gsr-1* tissue and cellular expression pattern previously reported [37, 39].

***gsr-1* is essential for *C. elegans* embryonic development and *gsr-1(m+,z-)* mutants are sensitized to chemical and developmental stresses and have compromised mitochondria.**

To investigate GSR-1 function, we used a worm strain carrying the *gsr-1(tm3574)* deletion allele, which spans 383 bp and removes part of *gsr-1* third exon, the third intron and part of the fourth exon (**Figure 1A**). Sequencing of *gsr-1(tm3574)* cDNA demonstrates that it encodes an in frame truncated protein (GSR-1 Δ 193-302, abbreviated as Δ GSR-1) lacking most of the NADPH domain plus a minor fraction of the second FAD domain (**Figure 1B**). Using purified recombinant protein expressed in bacteria, we show that wild type GSR-1b is able to reduce GSSG in a dose-dependent manner while Δ GSR-1b was devoid of enzymatic activity (**Figure 3A**), thus confirming *gsr-1(tm3574)* as a loss of function allele most likely by failure to bind NADPH required for redox cycling. GSR-1b did not function as thioredoxin reductase when using either yeast TRX3 or worm TRX-2 as substrates (**data not shown**).

The *gsr-1(tm3574)* allele is reported as lethal/sterile by the NBRP *C. elegans* Gene Knockout Consortium (<http://www.shigen.nig.ac.jp/c.elegans/>). Six times backcrossed *gsr-1(tm3574)* animals retained the lethal phenotype, so the *tm3574* allele was maintained in heterozygosity using GFP or RFP derivatives of the *qC1* balancer [44]. Homozygous *gsr-1* animals segregating from balanced parents (hereafter referred as *gsr-1(m+,z-)*; *m*, maternal and *z*, zygotic), have a normal embryonic and postembryonic development, reaching reproductive stage indistinguishably of wild type controls. Thus, maternally contributed *gsr-1* mRNA and/or GSR-1 protein is enough to allow *gsr-1* mutants to accomplish a normal life cycle. In turn, the *gsr-1(m-,z-)* embryos generated by these animals invariably arrest during embryogenesis.

RNAi downregulation of *gsr-1* expression has been shown to decrease *C. elegans* lifespan, to sensitize worms to the prooxidant juglone and the superoxide anion generator paraquat [39], to impair worm molting in the absence of cytoplasmic thioredoxin reductase *trxr-1* gene [37] and to induce *skn-1* target genes such as *gst-4* or *gcs-1* without an obvious nuclear translocation of a *skn-1* GFP reporter

[45]. To determine whether the maternal contribution of *gsr-1(m+,z-)* worms is functionally equivalent to the decreased *gsr-1* mRNA levels caused by RNAi, we tested *gsr-1(m+,z-)* individuals in all the above mentioned scenarios. Our data indicate that, indeed, this is the case as *gsr-1(m+,z-)* animals are short-lived (**Figure 3B**), are highly sensitive to juglone and paraquat (**Figure 3C**), have a fully penetrant larval arrest phenotype in a *trxr-1(sv47)* mutant background (**Figure 3D**) and induce GST-4 and GCS-1 reporters without significant SKN-1B/C::GFP nuclear translocation (**Figure 3E**). Therefore, maternally provided *gsr-1* mRNA and/or GSR-1 protein is enough to allow *gsr-1(m+,z-)* mutants to reach reproductive stage under non-stressed growth conditions but it is insufficient to afford effective protection against stress or developmental constrains.

The extreme sensitivity of *gsr-1(m+,z-)* worms to chemicals that impair mitochondrial function prompted us to assess mitochondrial status in these animals. We found that *gsr-1(m+,z-)* worms have a clear mitochondrial fragmentation phenotype in muscle cells (**Figure 4A**) while muscle sarcomere structure is preserved (**Figure 4B**). In addition, the mitochondrial membrane potential and total mitochondria content measured by incorporation of the fluorescent dye JC-10 was significantly decreased (**Figure 4C**). Moreover, *gsr-1(m+,z-)* worms display a strong induction of the mitochondrial UPR reporter HSP-6::GFP [46] (**Figure 4D**), overall indicating that decreased *gsr-1* levels cause mitochondrial stress and dysfunction. As a whole, these results validate the use of *gsr-1(m+,z-)* worms as a tool to investigate the role of GSR-1 in postembryonic development.

***gsr-1(m-,z-)* embryos arrest at the pregastrula/gastrula stage displaying progressive cell division delay and aberrant interphasic chromatin distribution.**

Next, we moved to study in detail the *gsr-1(m-,z-)* embryonic arrest phenotype. For this purpose, we video-recorded several *gsr-1(m-,z-)* embryos (n=26) and found that they all arrest at the pregastrula/gastrula stage ranging from 17 to 103 cells (**Figure 5A-C and Movies 1-3**). This arrest interval is roughly coincidental with the time at which *C. elegans* embryos start expressing a *Pgsr-1::gfp* transcriptional construct (**Figure 5D**). Arrested *gsr-1(m-,z-)* embryonic cells appear normal for at least the length of normal embryogenesis with no signs of necrosis or apoptosis, as it has been reported for most embryonic lethal mutants [47] (**Figure 5A-C and Movies 1-3**). However, cell lineage analysis of *gsr-1(m-,z-)* embryos identifies a dramatic delay of cell divisions timing, already detectable at the first embryonic divisions that is progressively enhanced until complete arrest occurs (**Figure 6A and Supplemental Figure 1A-C**).

We next explored the possible causes of the progressive delay of cell division timing of *gsr-1(m-,z-)* embryos. Among other traits, mutations in genes impairing mitochondrial function or genes encoding chromatin regulators have been shown to retard cell divisions in *C. elegans* embryos [48, 49]. Therefore, we set to investigate whether *gsr-1* mutants impact these two pathways given that

mitochondria-associated phenotypes are found in *gsr-1(m+,z-)* animals and because histone 3 glutathionylation destabilizes nucleosome structure and GSH depletion decreases DNA synthesis rate [50].

To evaluate the integrity of the mitochondrial network in *gsr-1(m-,z-)* embryos, we attempted labeling these organelles with rhodamine 6G followed by *in vivo* image analysis [51]. Despite rhodamine 6G efficiently stained embryo mitochondria, no major differences in the dynamics of the mitochondrial network were observed between young wild type and *gsr-1(m-,z-)* embryos (**Movie 4**). Reduced cell size of older embryos precluded analysis of differentiated embryonic cells. Regarding the other possible cause of cell division delay, we examined chromatin dynamics in *gsr-1(m-,z-)* embryos by time-lapse confocal microscopy. During early development, interphasic chromatin occupied the entire nuclear volume whereas mitotic chromosomes condensed in the nuclear interior (**Figure 6B and Movie 5; 1:10**). However, concomitantly with cessation of cell divisions in *gsr-1(m-,z-)* embryos, a progressive aberrant distribution of chromatin at the nuclear periphery was observed (**Figure 6B and Movie 5; 2:20-5:00**). Collectively, these results suggest that *gsr-1* deficiency impacts nuclear chromatin dynamics pointing to a nuclear dysfunction as a possible cause of *gsr-1(m-,z-)* embryonic arrest.

Cytoplasmic, but not mitochondrial, GSR-1 expression restores viability of *gsr-1(m-,z-)* embryos.

The fully penetrant embryonic arrest phenotype of *gsr-1(m-,z-)* embryos is rescued by transgenes expressing wild type GSR-1 (*vzEx105* transgene) as well as GSR-1::GFP fusion protein (*vzEx93* transgene) (**Figure 7A**), demonstrating that the lethality is due to a *gsr-1* deficiency. In contrast, no rescue was obtained when *gsr-1(m+,z-)* worms and their *gsr-1(m-,z-)* embryos were maintained on an effective external source of different GSH donors or precursors either in solid or liquid medium [5 mM GSH, 40 μ M S-linolenoyl-GSH [52]; 83 mg/ml liposomal GSH [53]; 20 μ M glutathione reduced ethyl ester [54]; 10 mM N-acetyl-L-cysteine [55]; 10 mM L-cystine, 10 mM L-methionine [56]) or 5mM DTT (**data not shown**). This lack of rescue indicates that *gsr-1(m+,z-)* worms do not provide these compounds to their oocytes (or they are rapidly exhausted) and also reflects the inability of these chemicals to cross the *gsr-1(m-,z-)* embryo eggshell, which is non-permeable to most solutes [57].

As mentioned above, both wild type GSR-1 and GSR-1::GFP fusion protein rescue the *gsr-1* embryonic lethal phenotype. In *gsr-1; vzEx105 [Pgsr-1::gsr-1]* animals, untagged GSR-1 produced by the *vzEx105* transgene provides enough maternal load to allow the non-transgenic *gsr-1* progeny to reach the reproductive stage (**Figure 7A, dashed bar**), similarly to *gsr-1(m+,z-)* worms segregating from *gsr-1/qC1* balanced parents. In contrast, in *gsr-1; vzEx93 [Pgsr-1::gsr-1::gfp]* animals, the GSR-

1::GFP fusion protein produced by the *vzEx93* transgene fails to provide maternal rescue, so only the transgenic progeny carrying the *vzEx93* transgene are viable, while all the non-transgenic siblings arrest at early embryogenesis (difference between eggs laid and adults, **Figure 7A**). The failure of the GSR-1::GFP fusion protein to provide maternal rescue could be explained by a lower enzymatic efficiency and/or a faster turnover of the GSR-1::GFP homodimers compared to wild type GSR-1 homodimers.

We next asked whether the embryonic lethal phenotype is due to a specific GSR-1 requirement in cytoplasm or mitochondria or, instead, to the simultaneous absence of the protein in both compartments. To address this, we performed isoform-specific rescue experiments in *gsr-1* mutants expressing, respectively, the mitochondrial GSR-1a::GFP or cytoplasmic GSR-1b::GFP fusion proteins (**Figure 2L-N**). As GSR-1::GFP fusions do not provide maternal load, a qualitative determination of the isoform-specific rescue is possible by simply scoring viable *versus* non-viable progeny. Interestingly, while *gsr-1* mutants expressing GSR-1b::GFP exclusively in the cytoplasm (either from the *vzEx145* [*Pgsr-1::gsr-1b(Δ1st exon)::gfp*] or the *vzEx147* [*Pgsr-1::gsr-1b(1st ATG mutated)::gfp*] transgenes) rescued the embryonic lethal phenotype, no viable progeny was obtained from *gsr-1* mutants expressing the mitochondrial specific GSR-1a::GFP isoform (from the *vzEx158* [*Pgsr-1::gsr-1a(2nd ATG mutated)::gfp*] transgene) (**Figure 7A**). These data imply that cytoplasmic GSR-1 is sufficient to restore normal embryonic development and that alternative redox systems such as the cytoplasmic thioredoxin system (redundant to the glutathione system in other organisms) are not able to substitute GSR-1 when absent from the cytoplasm.

The dispensability of mitochondrial GSR-1a, at least under non-stress normal growth conditions, could be explained by the presence of a redundant GSSG reducing system in mitochondria. In *S. cerevisiae* and *A. thaliana*, mitochondrial thioredoxin reductase is able to reduce GSSG in this organelle in the absence of mitochondrial glutathione reductase [58] (Meyer AJ, personal communication). To test if this genetic redundancy is maintained in *C. elegans*, we generated strains expressing only cytoplasmic GSR-1b::GFP (from *vzEx145* and *vzEx147* transgenes) in a *gsr-1(tm3574) trxr-2(tm2047)* double mutant background. As shown in **Figure 7B**, restoring GSR-1 activity exclusively in the cytoplasm of *gsr-1 trxr-2* double mutants also produced viable progeny, arguing against TRXR-2 as a redundant system for mitochondrial GSR-1a in worms. Collectively, these data demonstrate that in *C. elegans*, cytoplasmic GSR-1b is essential for embryonic development and that mitochondrial GSR-1a might be required for a yet to be identified dispensable function in this organelle.

DISCUSSION

Most eukaryotic organisms have one single glutathione reductase gene encoding both cytoplasmic and mitochondrial isoforms by the use of alternative translation initiation sites [22-25]. In

311 this study, we show that this is also the case for *C. elegans* glutathione reductase *gsr-1* gene, where
 312 translation from the first ATG codon results in the mitochondrial GSR-1a isoform, with an N-terminal
 313 MTS mostly encoded by the first exon, while translation from a second in-frame ATG codon located at
 314 the beginning of the second exon generates the cytoplasmic isoform GSR-1b (**Figure 2L-T**).
 315 Furthermore, *C. elegans* is one of the few examples of an eukaryotic organism with an essential
 316 requirement of glutathione reductase for viability. We show here that it is the cytoplasmic isoform of
 317 GSR-1 that is essential for viability (**Figure 7A**). While in many organisms the thioredoxin system
 318 backups GSSG reduction in the absence of glutathione reductase [14] this appears not to be the case
 319 for *C. elegans*, despite the fact that both systems have been shown to cooperate in other processes
 320 such as worm molting [37]. Therefore, the *C. elegans* thioredoxin and glutathione systems share
 321 common functions but also have specific non-overlapping roles in worm physiology.

322 *gsr-1* mutants with maternal load, *gsr-1(m+,z-)*, are viable and reach the reproductive stage
 323 indiscernibly from wild type controls. In turn, *gsr-1(m-,z-)* animals lacking maternal load arrest during
 324 early embryogenesis, displaying a robust progressive cell division delay phenotype (**Figure 6A**)
 325 concomitant with an aberrant distribution of interphasic chromatin (**Figure 6B**). Several possibilities
 326 might account for this lethal phenotype: **i)** as GSH depletion has been shown to inhibit DNA synthesis
 327 and to compromise cell cycle progression in mammalian cells [59, 60], this dependence on GSH supply
 328 may also apply to *C. elegans* embryos. A possible mechanism could involve histone H3
 329 glutathionylation that leads to a relaxed histone octamer in which DNA is less tightly packed around the
 330 nucleosome, thus opening chromatin [50]. Conversely, deglutathionylation of histone H3 results in a
 331 more compact distribution of chromatin ultimately decreasing gene expression and DNA synthesis.
 332 Thus, the absence of a GSH regenerating system, in *gsr-1(m-,z-)* embryos could compromise nuclear
 333 histone H3 glutathionylation, which may explain the aberrant chromatin distribution and cell division
 334 delay; **ii)** related to this, the thioredoxin and glutathione/glutaredoxin systems are the two enzymatic
 335 pathways providing reducing equivalents to the essential enzyme ribonucleotide reductase (RNR),
 336 responsible for the supply of deoxyribonucleotides for DNA synthesis and repair [61]. While the
 337 glutathione system is much more efficient with the *E. coli* RNR enzyme, in mammals the thioredoxin
 338 system is one order of magnitude more efficient than the glutaredoxin system [62]. If *C. elegans* RNR is
 339 more dependent on the glutathione system, as found in bacteria, failure to reduce RNR could be a
 340 cause of the lethality of *gsr-1(m-,z-)* worms; **iii)** mutations in mitochondrial genes have also been shown
 341 to cause embryonic arrest in *C. elegans* [47, 48]. Although we have not been able to detect abnormal
 342 mitochondrial network distribution in *gsr-1(m-,z-)* embryos (**Movie 4**), the fact that maternally rescued
 343 *gsr-1* mutants have lower amount of mitochondria, increased mitochondrial fragmentation and induced
 344 levels of *hsp-6*, a mitochondrial stress reporter (**Figure 4**), it is possible that impaired mitochondrial

function may also account for the *gsr-1(m-,z-)* embryonic arrest phenotype; **iv)** alternatively, it is possible that *gsr-1(m-,z-)* embryos could die by GSSG poisoning. In mammalian cell cultures with compromised GSSG reducing activity, GSSG is excreted into the cell culture medium to avoid the toxicity of its intracellular buildup [16]. If such mechanism also happens in *C. elegans gsr-1(m-,z-)* embryos, excreted GSSG would accumulate within the non-permeable extracellular peri-embryonic space of the embryo [63], where the progressive buildup of GSSG would ultimately cause embryo poisoning. It will be interesting to test this hypothesis by either microinjecting permeable GSH derivatives, GSH precursors or recombinant GSR-1 into the *gsr-1(m-,z-)* embryos [64] or by transgenically expressing GSR-1 in their peri-embryonic space.

As GSH synthesis is restricted to the cytoplasm, the mitochondrial GSH pool is generated by an active import of cytoplasmic GSH via specific carriers. In turn, it is generally accepted that mitochondrial GSSG is not readily exported from this organelle, being therefore reduced by mitochondrial glutathione reductase [65]. *C. elegans gsr-1* mutants with transgenic GSR-1 expression restricted to cytoplasm are viable, implying that mitochondrial glutathione reductase is dispensable in *C. elegans*, at least under normal non-stress conditions and, therefore, that GSSG does not accumulate in mitochondria. As mitochondrial thioredoxin reductase has been shown to substitute for mitochondrial glutathione reductase in yeast and plants [58] and (Meyer AJ, personal communication), we first tested whether this functional redundancy is maintained in *C. elegans*. However, a double mutant *gsr-1(m-,z-) trxr-2* (with transgenic GSR-1 expression in the cytoplasm to allow development) is also viable (**Figure 7B**), suggesting that *trxr-2* is not functionally redundant with *gsr-1* in worms. We next asked whether mitochondrial GSR-1 is dispensable during normal non-stressed conditions but required under mitochondrial stresses to provide mitochondria with enough GSH to counteract the insults. We tested this possibility and demonstrate that animals lacking mitochondrial GSR-1 are as resistant as wild type controls when exposed to toxic doses of paraquat or juglone, thus ruling out that mitochondrial GSR-1 is needed under mitochondrial stress (**Supplemental Figure 2**). In this context, the dispensability of mitochondrial *gsr-1* (and *trxr-2*) in *C. elegans* poses some interesting hypotheses: **i)** mitochondrial GSSG reduction is carried out by a third, yet unknown, mitochondrial reducing system in the absence of *gsr-1* and *trxr-2*; **ii)** GSSG can be exported from mitochondria to be reduced in the cytoplasm. Although studies in yeast show that the mitochondrial matrix and intermembrane space glutathione pools are maintained separately [66], a GSSG export activity has been described for the inner mitochondrial membrane ATP-binding cassette transporter ATM3 (in *A. thaliana*) and Atm1 (in *S. cerevisiae*) [67]. Thus, it is conceivable that GSSG generated in the mitochondrial matrix could be exported to the intermembrane space by these transporters, from where it can reach the cytoplasm via porins [66] to be reduced by cytoplasmic glutathione reductase as there is no GSSG reducing system in the

mitochondrial intermembrane space. The *C. elegans abtm-1* gene is the ortholog of ATM3/Atm1 and its mutation causes embryonic lethality. Instead, worms with downregulated levels of *abtm-1* by RNAi are viable, although have increased oxidative stress and ferric iron accumulation [68]. This lethality of *abtm-1* mutants is presumably originated by a failure to transport Fe-S clusters to cytoplasm rather than a GSSG accumulation in mitochondria, as these animals are wild type for *gsr-1*. We performed *abtm-1* RNAi in *gsr-1(m+,z-)* worms and did not observe any compound phenotype, suggesting that GSSG is not accumulating in their mitochondria, even in the absence of this putative GSSG exporter (**data now shown**); **iii**) alternatively, mitochondrial GSR-1 could be needed for yet unknown additional non-essential roles in this organelle, other than reducing GSSG. An example of a novel additional function of glutathione reductase is the reduction of the redox-regulated mitoNEET protein, an outer mitochondrial membrane protein that has been implicated in energy homeostasis and mitophagy [69, 70].

In summary, we describe here the phenotypes that the absence of glutathione reductase causes in the model organism *C. elegans*. The availability in this organism of cytoplasmic and mitochondrial thioredoxin reductase mutants combined with that of *gsr-1* now characterized offers an ideal scenario where to study the respective contribution of the different compartmentalized redox systems in the context of a multicellular organism. Although our results clearly show that GSR-1 is dispensable in mitochondria, it is yet intriguing why *gsr-1(m+,z-)* mutants display mitochondria-associated phenotypes. It is plausible that under limiting GSR-1 availability as in *gsr-1(m+,z-)* mutants, resources are derived to maintain the essential cytoplasmic function thus provoking mitochondrial susceptibility. Future studies aimed to specifically inactivate *gsr-1* mitochondrial or cytoplasmic isoforms by CRISPR technology will be instrumental to further refine their respective contribution to cell function and redox homeostasis.

400 MATERIALS AND METHODS

402 *C. elegans* strains and culture conditions

403 The standard methods used for culturing and maintenance of *C. elegans* were as previously
404 described [71]. The strains used in this work are described in **Supplemental Table I**. All experiments
405 were performed at 20°C unless otherwise noted. All VZ strains are 6x backcrossed with N2 wild type.

407 *gsr-1* expression constructs, transgenesis and image analysis of transgenic animals.

408 For fluorescent reporter constructs, DNA fragments containing the 2.2 kb upstream promoter
409 region of the *gsr-1* gene (including the first 15 nucleotides of the coding region), the 2.2 kb promoter
410 region plus the complete *gsr-1* genomic ORF and the *gsr-1* genomic ORF except the first exon were
411 amplified from *C. elegans* wild type genomic DNA and cloned into the PstI and Ball sites of the
412 pPD95.81 vector (Fire Lab *C. elegans* Vector Kit 1995 (unpublished)) to generate *gsr-1* transcriptional
413 and translational GFP reporters, respectively. Mutagenesis reactions (1st exon deletion and 1st and 2nd
414 ATG codon substitutions) were performed on the 2.2 kb promoter *gsr-1* translational GFP reporter
415 construct using the QuickChange II XL Site Directed Mutagenesis Kit (Stratagene) and confirmed by
416 sequencing. For the reporter construct targeting GFP to muscle cell mitochondria directed by *gsr-1*
417 MTS, we amplified a 2.4 kb region of the *myo-3* gene promoter with primers that included the *gsr-1* MTS
418 sequence from the pBN42 plasmid (*[Pmyo-3::gfp::3'-UTR_{unc-54}]* [72]) and cloned it into the BamHI site
419 of the pPD95.81 vector. Correct orientation of the insert was confirmed by sequencing. For expression
420 of wild type GSR-1, we amplified a fragment containing the *gsr-1* 2.2 kb promoter region, the genomic
421 ORF and 3'-UTR from *C. elegans* wild type genomic DNA and cloned it into the PstI and SpeI sites of
422 the pSPARK I vector (Canvax Biotech, Córdoba, Spain). The sequences of all primers used for cloning
423 are available upon request. Transgenic worms were generated by DNA microinjection as previously
424 described [73]. The pRF4 [*rol-6(su1006)*] [73] and coel::GFP [*Punc-122::gfp*] [74] plasmids were used
425 as coinjection markers at 50 ng/μl. The *gsr-1* expression constructs described above were injected into
426 N2 wild type animals (either as circular plasmids or linear fragments) at varying concentrations ranging
427 from 10 to 50 ng/μl.

428 For image analysis of fluorescent transgenic worms, animals were mounted in a 5 μl drop of 10
429 mM levamisole on a 3% agarose pad covered with a coverslip. Differential interference contrast (DIC)
430 and fluorescence imaging was performed on a Zeiss AxioImager M2 with ApoTome Unit fluorescence
431 microscope and images were captured with the AxioVision 4.8 Software (Zeiss) (**Figure 2A-K**).

433 **Imaging of mitochondria, sarcomeres and measures of mitochondrial function.**

434 For **Figure 2M-N and 2R-T**, animals were picked into 25 ml M9 buffer and were immobilized only
435 by the pressure from the cover slip. Images in M and N were taken using a Nikon H600L microscope
436 (Nikon Corporation, Tokyo, Japan) and a Nikon Digital Sight DS-Fi1 digital camera with proprietary
437 software. Images R – T were taken on a Zeiss AX10 microscope (Carl Zeiss AG, Oberkochen,
438 Germany) with an Axiocam MRC digital camera and Axiovision LE software. Images R – T were taken
439 using GFP, RFP, and triple-pass filter sets, respectively. Images in **Figure 4A** were taken in n = 60
440 animals and **Figure 4B-C** in n = 20 animals at young adulthood with a fixed exposure of 1 s and the
441 fluorescence was quantified using ImageJ. Corrected total animal fluorescence was calculated using
442 data from integrated density and area, however, mean fluorescence minus background fluorescence
443 produced a similar result. Data were non-parametric and were therefore analysed using a Mann
444 Whitney test and significance was set at P <0.05.

445

446 **RNA extraction and RT-PCR**

447 For total RNA extraction, gravid hermaphrodites were washed off the plates with M9 buffer and
448 dissolved in 5M NaOH bleaching solution. Embryos were collected and washed several times with M9
449 buffer. RNA was extracted from embryos using the NucleoSpin RNA II (Macherey-Nagel) kit following
450 the manufacturer's instructions. Total RNA was DNase-treated using the Amplification Grade Dnase I
451 (Sigma) and 1µg of Dnase-treated RNA was reverse transcribed in a 20 µl reaction mixture. cDNA was
452 generated using the iScript™ cDNA Synthesis Kit (Biorad). 1µg of cDNA was used for *gsr-1* RT-PCR
453 reactions using MBL-Taq DNA Polymerase (Dominion-MBL).

454

455 **Recombinant protein expression, purification and glutathione reductase enzymatic activity**

456 *gsr-1* cDNA from N2 wild type and *gsr-1(tm3574)* mutants was amplified with the forward primer
457 5'-ACTGCATATGTCTGGCGTCAAG-3' and the reverse primer 5'-
458 CATGCTCGAGTTATTATTCCGGCTTCACAC-3' and cloned into the NdeI and XhoI restrictions sites of
459 the pET-15b vector (Novagen) to generate the constructs His-GSR-1b and His-ΔGSR-1b, respectively.
460 These constructs were used to transform the *E. coli* BL21(DE3) strain and recombinant protein
461 expression was induced by adding 0.5 mM IPTG to a 500 ml LB medium bacteria culture of 0.5-0.7 OD
462 supplemented with 0.1 mg/ml ampicillin and further incubating the cells at 25°C and 200 rpm during 5
463 hours. Cells were collected by centrifugation, immediately resuspended in 25 ml Tris-HCl 20 mM pH 8,
464 0.1 M NaCl buffer supplemented with 15 mg de lysozyme and 30 mg de Dnase I. After 5 min incubation

465 at room temperature the preparation was sonicated for 30 min on ice and the cell free extract was
466 obtained by centrifugation at 10000 x g during 30 min at 4°C. Recombinant His-GSR-1b and His-ΔGSR-
467 1b proteins were purified from the cell free extract using a TALON column (Clontech) equilibrated with
468 Tris-HCl 20 mM pH 8, 0.1 M NaCl buffer and eluted with 100 mM imidazol. Finally, the purified proteins
469 were dialyzed against the same buffer and concentrated using Centricon YM-10 filter devices
470 (Millipore). The glutathione reductase enzymatic activity was determined as previously described [75].
471 Briefly, a standard assay mixture (0.5 ml) containing potassium phosphate buffer 0.1 M, pH 7.2, EDTA 1
472 mM, NADPH 200 μM, with different concentrations of His-GSR-1b and His-ΔGSR-1b was prepared and
473 the reaction was initiated by addition of 0.5 mM GSSG. The decrease of absorbance at 340 nm
474 (indicating NADPH consumption) was recorded spectrophotometrically at 25°C.

475

476 **Embryonic cell lineage analysis**

477 For embryonic cell lineage analysis, 4D-microscopy was carried out using standard live-animal
478 mounting techniques on a Leica DM6000 microscope fitted with DIC optics. The use of DIC optics
479 allows cells tracing without using any dye or fluorescent marker that might alter the cell cycle
480 progression. Embryonic cell lineage was determined as described [76]. In summary, gravid
481 hermaphrodites were dissected and 2- to 4-cell stage embryos were mounted on 4% agar pads in
482 water, and sealed with Vaseline. Imaging was performed at 25°C. The multi-focal time-lapse
483 microscopy of the samples was controlled with the open source software Micro-manager ([www.micro-](http://www.micro-manager.org)
484 [manager.org](http://www.micro-manager.org)). Pictures on 30 focal planes (1micron/section) were taken every 30 seconds for 12 h.
485 Embryo lineages were analyzed with the software SimiBiocel (SIMI GmbH, www.simi.com).

486

487 **Brood size determination**

488 L4 larvae were singled onto OP50 seeded plates, allowed to lay eggs at 20°C and transferred to
489 new plates every 12 hours until the animals stopped reproducing. Eggs were counted and then
490 incubated further at 20°C to quantify adult progeny four days later.

491

492 **Longevity assay**

493 Lifespan assays were performed at 25°C as previously described [77] with slight modifications.
494 Tightly synchronized embryos from gravid adult hermaphrodites were allowed to develop through the L4
495 larval stage and then transferred to fresh NMG plates in groups of 20 worms per plate for a total of 80
496 individuals per experiment. The day animals reached the L4 larval stage was used as $t = 0$. Animals

497 were transferred to fresh plates daily until progeny production ceased and after that they were
498 transferred every second to third day but monitored daily for dead animals. Nematodes that did not
499 respond to gentle prodding and displayed no pharyngeal pumping were scored as dead. Animals that
500 crawled off the plate or died due to internal hatching or extruded gonad were censored and incorporated
501 as such into the data set.

502

503 **Paraquat and Juglone acute treatment**

504 Plates containing the appropriate amount of the respective chemical were prepared freshly prior
505 the experiment. To minimize the number of animals crawling off the agar, OP50 bacteria (from seeded
506 plates) was directly spread in the center of the plates using an inoculating loop. L4 larvae were
507 transferred to plates and survival was determined after 16 hours incubation at 20°C. Nematodes that did
508 not respond to gentle prodding and displayed no pharyngeal pumping were scored as dead. Animals
509 that crawled off the plate were censored.

510

511 **Embryonic/larval arrest assays**

512 10 gravid *trxr-1(sv47); gsr-1(tm3574)/qC1::rfp* hermaphrodites were allowed to lay eggs during 3
513 hours at 20°C and subsequently removed. Embryos were counted and then allowed to develop for 4
514 days at 20°C after which non RFP *trxr-1(sv47); gsr-1(m+,z-)* adults, larvae and unhatched embryos
515 were quantified.

516

517 **SKN-1 dependent gene expression quantification and nuclear translocation of SKN-1 fluorescent** 518 **reporter**

519 Worms at their third day of adulthood expressing *dvEx166 [Pgst-4::gfp]* and *svEx741 [Pgcs-1::gfp]*
520 extrachromosomal arrays were mounted in a 5 ml drop of 10 mM levamisole on a 3% agarose pad
521 covered with a coverslip. Fluorescence imaging was performed on a Olympus BX61 fluorescence
522 microscope equipped with a Olympus DP72 camera and images were captured with the
523 CellSensDimension 1.12 Software (Olympus). All micrographs were taken with identical image capture
524 settings and quantification of GFP expression (measured as the fluorescence mean of 25 worms
525 divided by the selected area and normalized by the background adjacent to the selected worm in the
526 same image) was performed using the ImageJ Software (NIH). When needed, equal adjustment of
527 brightness and contrast on control and matched problem images was implemented using Adobe
528 Photoshop 10 Software (Adobe Systems). To visualize SKN-1B/C::GFP nuclear translocation, young

adults were washed from plates, anesthetized with 1 mM levamisole and mounted on 2% agarose pads. Visualization and imaging was performed using an Olympus IX81 automated inverted microscope and Slidebook (version 5.0) software. SKN-1B/C::GFP localization quantification, the percent intestinal SKN-1 nuclear localization was categorically scored as follows: none: no localization, low: posterior or anterior intestinal localization, medium: posterior and anterior intestinal localization, high: localization throughout the entire intestine [78].

Mitochondrial related phenotypes

To determine if *gsr-1* deficiency had any effect on mitochondria related phenotypes, age-synchronized, young adult animals (n = 60) were assessed for mitochondrial networking in *gsr-1(m+,z-); ccls4251 [Pmyo-3::mito::gfp]* versus control *ccls4251 [Pmyo-3::mito::gfp]* animals. There was significantly greater mitochondrial fragmentation in the *gsr-1* animals than controls at 20°C (P <0.001) and 25°C (P <0.05). Determination of sarcomere structure in *gsr-1(m+,z-); jls01[Pmyo-3::myo-3::gfp]* were compared to control *jls01[Pmyo-3::myo-3::gfp]* animals (n = 20 per group) both at 20°C and at 25°C. There was no significant difference between groups (P >0.05). For mitochondrial content determination, we loaded animals with JC-10, a dye that accumulates proportionally to a negative mitochondrial membrane potential as previously described [79]. Wild type control animals had significantly greater fluorescence of the gut mitochondria than did *gsr-1(m+,z-)* animals (P <0.001). Images were taken with a fixed exposure of 1 s on the Nikon H600L microscope and the fluorescence was quantified using ImageJ. Corrected total animal fluorescence was calculated as follows: Integrated density – area of the worm x mean fluorescence of three background readings. Data were analyzed using a Mann-Whitney test as data were non-parametric.

Embryonic chromatin dynamics analysis.

BN323 animals were maintained at 16°C and shifted to 25°C 2 hours prior to microscopy. Heterozygous and homozygous *gsr-1* mutants were dissected and early embryos were mounted together in M9 buffer on 2% agarose pads. Coverslips were placed over the embryos and sealed with VALAP (1:1:1 mixture of Vaseline, lanolin and paraffin). Confocal epifluorescence and DIC images from three focal planes were acquired at 25°C every 10 minutes on a Nikon A1R microscope through a Plan Apo VC 60x/1.4 objective (Nikon, Tokyo, Japan) using a pinhole of 39.7 µm (1.2 Airy Units).

560 **ACKNOWLEDGMENTS**

561

562 Some strains were provided by the CGC, which is funded by NIH Office of Research
563 Infrastructure Programs (P40 OD010440) and by the Japanese National Bioresource Project. We
564 thank Cristina Cecchi, Amir Shapir, Paul Sternberg, Bart Braeckman, Chris Link, Simon Tuck,
565 Keith Blackwell, José López-Barneo and LivOn Labs for strains and chemicals, Katie McCallum
566 and Danielle Garsin for their help with *skn-1* experiments and Elizabeth Veal and Michel Toledano
567 for critical reading of the manuscript. Prof. Rafael Fernández-Chacón is deeply acknowledged for
568 his continuous support. AMV was supported by grants from the Spanish Ministry of Economy and
569 Competitiveness (BFU2015-64408-P) and the Instituto de Salud Carlos III (PI11/00072,
570 cofinanced by the Fondo Social Europeo) and is a member of the GENIE and EU-ROS Cost
571 Action of the European Union. NJS was supported by a grant from the US National Institutes of
572 Health National Institute for Arthritis and Musculoskeletal and Skin Diseases (AR-054342). CG
573 was funded by a Doctoral Training Studentship provided by the University of Nottingham. PA was
574 supported by the Spanish Ministry of Economy and Competitiveness (BFU2013-42709P). JC is a
575 member of the GENIE Cost action and was funded by Rioja Salud Foundation.

576

577

577 FIGURES

578

579 **Figure 1: *gsr-1* mRNA and protein analysis. A)** Schematic representation of the two main *gsr-1*
 580 mRNA variants. Boxes represent exons and lines show spliced introns. White boxes indicate 5'-UTR
 581 and 3'-UTR, respectively, and grey boxes indicate the ORF. The region of the *gsr-1* mRNA missing
 582 in the *gsr-1(tm3574)* deletion allele is underlined. **B)** Protein domain organization of GSR-1a and
 583 GSR-1b isoforms as well as that of the truncated Δ GSR-1 protein resulting from translation of the
 584 *gsr-1(tm3574)* allele. Numbers indicate the amino acid residues flanking the different protein
 585 domains, based on the domain organization reported for the human GSR orthologue [80]. **C)**
 586 Sequence alignment of *C. elegans* GSR-1 with the corresponding human and zebrafish GSR
 587 orthologues. The red rectangle indicates the conserved redox active site CVNVGC. The yellow
 588 boxes highlight the residues involved in GSSG binding [81]. The NADPH and FAD binding sites are
 589 underlined. Identical residues are shown in black. The numbers on the left indicate the respective
 590 amino acid residue. The different domains boundaries are indicated above the sequences and are
 591 inferred from comparison with those reported for human GSR protein [80]. **D)** Sequence of the GSR-
 592 1a mitochondrial targeting sequence (MTS). Amino acid residues matching the two matrix protease
 593 model (in blue. X, any amino acid residue) [41] are highlighted in red. The MTS cleavage site is
 594 indicated by the upside-down arrow. The residues encoded by *gsr-1* first and second exons are
 595 separated by an hyphen.

596

597 **Figure 2: GSR-1 tissue, cellular and subcellular expression analysis. A)** Scheme of the GSR-1
 598 translational GFP fusion construct used to determine expression from *gsr-1* upstream promoter.
 599 Exons are in grey boxes and introns in white boxes. **B-E)** Differential interference contrast and **B'-E')**
 600 fluorescence images of transgenic worms expressing the *Pgsr-1::gsr-1::gfp* construct: ai, anterior
 601 intestine; coe, coelomocytes; gsc, gonad sheet cell; hy, hypodermis; ph, pharynx; rc, rectal cells; sp,
 602 spermatheca; tn, tail neurons; uc, uterine cells; vmc, vulva muscle cells. **F)** Scheme of the GSR-1
 603 translational GFP fusion construct used to determine expression from *gsr-1* first intron. Exons are in
 604 grey boxes and introns in white boxes. **G-K)** Differential interference contrast and **G'-K')**
 605 fluorescence images of transgenic worms expressing the *gsr-1 first intron::gsr-1::gfp* construct: ai,
 606 anterior intestine; coe, coelomocytes; ec, excretory cell; hn, head neurons; int, intestine; pi, posterior
 607 intestine. **L)** Schemes of the GSR-1 translational GFP fusion constructs used to determine
 608 subcellular localization of GSR-1a and GSR-1b isoforms. Red X indicates the mutated ATG codon.
 609 Exons are in grey boxes and introns in white boxes. **M)** Dotted fluorescence labeling in pharynx of
 610 transgenic worms expressing the *Pgsr-1::gsr-1::gfp* construct with the second ATG codon mutated.

N) Diffuse cytoplasmic fluorescence labeling in pharynx of transgenic worms expressing the construct the *Pgsr-1::gsr-1::gfp* construct with the first ATG codon mutated. **Q)** Scheme of the construct used to determine GSR-1a MTS functionality when fused to GFP under the control of the *myo-3* muscle promoter. GSR-1a MTS is shown in a grey box. Fluorescence images of muscle cells of transgenic worms expressing **R)** the *Pmyo-3::MTS_{gsr-1}::gfp* construct, **T)** the *Pmyo-3::tomm-20::mrfp* construct and **S)** merged picture showing colocalization in muscle tubular mitochondria. Scale bar 20 μ m in all images except 100 μ m in B. Note Images B-K and M-T were taken with different microscopes and settings (see Materials and Methods).

Figure 3: Glutathione reductase enzymatic activity and phenotypes of *gsr-1(m+,z-)* mutants.

A) GSSG reductase enzymatic activity of different concentrations of recombinant His-GSR-1b (straight lines) and His- Δ GSR-1b (dashed lines) in the presence of 0.5 mM GSSG and 0.25 mM NADPH. **B)** The longevity was assayed at 20°C in *E. coli* OP50. Kaplan-Meier plot was used to show the fraction of animals that survive over time. Longevity was performed twice, obtaining similar results, and the composite data is shown. The survival rate of *gsr-1(m+,z-)* animals was compared to that of wild type using the log-rank (Mantel-Cox) test and the differences were found significant (** $p < 0.001$). **C)** Sensitivity of *gsr-1(m+,z-)* animals to acute paraquat and juglone exposure. Data are the mean $\pm$ SD of four independent experiments with three biological replicates each ($n \geq 200$; ** $p < 0.001$, by unpaired, two-tail Student *t*-test). **D)** Percentage of developmental stages of single and double mutants combining the *txxr-1(sv47)* and the *gsr-1(m+,z-)* mutation. Data are the mean of three independent experiments (n = total number of individuals scored). **E)** Induction of GST-4 and GCS-1 transcriptional GFP reporters and percentage of nuclear localization of a SKN-1B/C translational GFP reporter in a *gsr-1(m+,z-)* background. For GST-4 and GCS-1 reporters, the data are the mean $\pm$ SD of 25 individuals for each genotype (** $p < 0.001$, * $p < 0.01$ by unpaired, two-tail Student *t*-test). For SKN-1 reporter the data are from two independent experiments (n = total number of individuals scored).

Figure 4: *gsr-1(m+,z-)* mutants display mitochondrial phenotypes. A) Increased mitochondrial fragmentation is observed in *gsr-1(m+,z-)* animals expressing the mitochondrial reporter *ccIs4251 [Pmyo-3::mito::gfp]* in muscle cells. Data are the total $\pm$ SEM of three independent experiments with 20 animals per experiment at 20°C and 25°C (n = 60 total at each temperature) (* $p < 0.05$, ** $p < 0.001$, by a Mann Whitney test). Representative images of each genotype are shown. Scale bar 20 μ m. **B)** Muscle cell integrity is not affected in *gsr-1(m+,z-)* worms as demonstrated by the maintenance of sarcomere myotubular structure visualized using the *jIs01 [Pmyo-3::myo-3::gfp]*

reporter. Data are the total $\pm$ SEM of one experiment with 20 animals ($p > 0.05$ by a Mann Whitney test). Representative images of each genotype are shown. Scale bar 20 μm . **C)** Quantification of the total mitochondrial content measured by JC-10 dye incorporation. Data are the total $\pm$ SEM of 20 individuals per genotype ($*** p < 0.001$ by a Mann Whitney test). Representative images of each genotype are shown. Scale bar 100 μm . **D)** Mitochondrial UPR is induced in *gsr-1(m+,z-)* worms expressing the reporter *zcls13 [Phsp-6::gfp]*. Data are the mean $\pm$ SD of 40 individuals per genotype ($*** p < 0.001$, by unpaired two-tail Student *t*-test). Representative images of each genotype are shown. Scale bar 200 μm .

Figure 5: Identification of the cell-stage time of arrest of *gsr-1(m-,z-)* embryos and determination of GSR-1 expression initiation. A-C) Differential interference contrast still images of developing *wild type* and *gsr-1(m-,z-)* embryos analyzed by 4D-microscopy. Three representative independent experiments are provided, illustrating the cases at which *gsr-1(m-,z-)* embryos arrest at the earliest, the latest and an intermediate cell-stage. **A)** Arrest at 17 cell-stage; **B)** Arrest at 33 cell-stage; **C)** Arrest at 103 cell-stage. *Gsr-1* in the images refers to *gsr-1(m-,z-)* embryos. **D)** Fluorescence and differential interference contrast still images of developing embryos from transgenic animals expressing the transcriptional construct *Pgsr-1::gfp*. Three independent embryos are shown. *Gsr-1* expression begins approximately at 100 min since embryos start developing. In these initial stages, *gsr-1* stronger expression is found in E blastomere descendants (intestine precursors) and MS blastomeres descendants (neurons and mesodermic tissues precursors). Scale bar 10 μm . All analyses were performed at 25°C.

Figure 6: Cell division delay and progressive perinuclear localization of interphasic chromatin in *gsr-1(m-,z-)* embryos. A) Cell lineage timing analysis of the EMS blastomere descendants of the *gsr-1(m-,z-)* embryo depicted in **Figure 5B** compared to that of wild type embryos. For complete cell lineage analysis, see **Supplemental Figure 1B**. **B)** *gsr-1(m+,z+)* and *gsr-1(m-,z-)* embryos expressing mCherry::HIS-58 (magenta) and GFP::TBB-2 were observed by live confocal microscopy. Time is indicated in hours:minutes from beginning of recording (see Movie 5). At 2:20 and more evidently at 5:00 chromosomes are condensed at the nuclear periphery of *gsr-1(m-,z-)* embryos. Scale bar 10 μm .

Figure 7: Rescue of *gsr-1(tm3574)* mutants embryonic lethality. A) The *gsr-1(tm3574)* embryonic lethal phenotype is rescued when GSR-1 activity is restored in the cytoplasm (GSR-1b, transgenes *vzEx145* and *vzEx147*) but not in mitochondria (GSR-1a, transgene *vzEx158*). The

679 dashed bar in *gsr-1; vzEx105 [Pgsr-1::gsr-1]* animals indicates the fraction of viable, maternally
680 rescued, non-transgenic progeny. Data are the mean $\pm$ SD of the progeny from at least 10 animals of
681 each genotype. **B)** Viable progeny from *gsr-1; trxr-2* double mutant worms expressing GSR-1b::GFP
682 in cytoplasm. Data are the mean $\pm$ SD of the progeny from at least 10 animals of each genotype.

683
684

684
685 **MOVIES**

686
687 **Movies 1-3: Movie recordings of *C. elegans* wild type vs *gsr-1(m-,z-)* embryos.** Developmental
688 arrest of the *gsr-1(m-,z-)* embryo occurs at 17-cell stage (movie 1), 33-cell stage (movie 2) and 103-
689 cell stage (movie 3). Records were performed at 25°C over 12h as explained in materials and
690 methods. Movie 1 corresponds to **Figure 5A**, Movie 2 corresponds to **Figure 5B** and Movie 3
691 corresponds to **Figure 5C**

692
693 **Movie 4: Movie recording of *C. elegans* wild type vs *gsr-1(m-,z-)* embryos stained with**
694 **rhodamine 6G.** The day before the recording wild type and *gsr-1(m-,z-)* L4 worms were placed on
695 NGM agar plates supplemented with rhodamine 6G at a final concentration of 2.5 ug/ml and
696 incubated at 20°C. Two hours before the recording plates were transferred to 25°C. Recordings were
697 performed with a NIKON A1R confocal microscope through a Plan Apo VC 60x/1.4 objective. Time
698 is indicated in hours : minutes from beginning of recording.

699
700 **Movie 5: Movie recording of *C. elegans* *gsr-1(m+,z+)* and *gsr-1(m-,z-)* embryos expressing**
701 **mCherry::HIS-58 (magenta) and GFP::TBB-2.** Time is indicated in hours:minutes from beginning of
702 recording. Corresponds to **Figure 6B**.

703
704

REFERENCES

- [1] Hanschmann, E. M.; Godoy, J. R.; Berndt, C.; Hudemann, C.; Lillig, C. H. Thioredoxins, glutaredoxins, and peroxiredoxins--molecular mechanisms and health significance: from cofactors to antioxidants to redox signaling. *Antioxid Redox Signal* **19**:1539-1605; 2013.
- [2] Meyer, Y.; Buchanan, B. B.; Vignols, F.; Reichheld, J. P. Thioredoxins and glutaredoxins: unifying elements in redox biology. *Annu Rev Genet* **43**:335-367; 2009.
- [3] Xiong, Y.; Uys, J. D.; Tew, K. D.; Townsend, D. M. S-glutathionylation: from molecular mechanisms to health outcomes. *Antioxid Redox Signal* **15**:233-270; 2011.
- [4] Fahey, R. C.; Sundquist, A. R. Evolution of glutathione metabolism. *Adv Enzymol Relat Areas Mol Biol* **64**:1-53; 1991.
- [5] Grant, C. M.; MacIver, F. H.; Dawes, I. W. Glutathione is an essential metabolite required for resistance to oxidative stress in the yeast *Saccharomyces cerevisiae*. *Curr Genet* **29**:511-515; 1996.
- [6] Romero-Aristizabal, C.; Marks, D. S.; Fontana, W.; Apfeld, J. Regulated spatial organization and sensitivity of cytosolic protein oxidation in *Caenorhabditis elegans*. *Nat Commun* **5**:5020; 2014.
- [7] Fraser, J. A.; Kansagra, P.; Kotecki, C.; Saunders, R. D.; McLellan, L. I. The modifier subunit of *Drosophila* glutamate-cysteine ligase regulates catalytic activity by covalent and noncovalent interactions and influences glutathione homeostasis in vivo. *J Biol Chem* **278**:46369-46377; 2003.
- [8] Cairns, N. G.; Pasternak, M.; Wachter, A.; Cobbett, C. S.; Meyer, A. J. Maturation of arabidopsis seeds is dependent on glutathione biosynthesis within the embryo. *Plant Physiol* **141**:446-455; 2006.
- [9] Dalton, T. P.; Dieter, M. Z.; Yang, Y.; Shertzer, H. G.; Nebert, D. W. Knockout of the mouse glutamate cysteine ligase catalytic subunit (Gclc) gene: embryonic lethal when homozygous, and proposed model for moderate glutathione deficiency when heterozygous. *Biochem Biophys Res Commun* **279**:324-329; 2000.
- [10] Pasternak, M.; Lim, B.; Wirtz, M.; Hell, R.; Cobbett, C. S.; Meyer, A. J. Restricting glutathione biosynthesis to the cytosol is sufficient for normal plant development. *Plant J* **53**:999-1012; 2008.
- [11] Winkler, A.; Njalsson, R.; Carlsson, K.; Elgadi, A.; Rozell, B.; Abraham, L.; Ercal, N.; Shi, Z. Z.; Lieberman, M. W.; Larsson, A.; Norgren, S. Glutathione is essential for early embryogenesis--analysis of a glutathione synthetase knockout mouse. *Biochem Biophys Res Commun* **412**:121-126; 2011.
- [12] Grant, C. M.; MacIver, F. H.; Dawes, I. W. Glutathione synthetase is dispensable for growth under both normal and oxidative stress conditions in the yeast *Saccharomyces cerevisiae* due to an accumulation of the dipeptide gamma-glutamylcysteine. *Mol Biol Cell* **8**:1699-1707; 1997.
- [13] Logan-Garbisch, T.; Bortolazzo, A.; Luu, P.; Ford, A.; Do, D.; Khodabakhshi, P.; French, R. L. Developmental ethanol exposure leads to dysregulation of lipid metabolism and oxidative stress in *Drosophila*. *G3 (Bethesda)* **5**:49-59; 2014.
- [14] Kanzok, S. M.; Schirmer, R. H.; Turbachova, I.; Iozef, R.; Becker, K. The thioredoxin system of the malaria parasite *Plasmodium falciparum*. Glutathione reduction revisited. *J Biol Chem* **275**:40180-40186; 2000.

749 [15] Morgan, B.; Ezerina, D.; Amoako, T. N.; Riemer, J.; Seedorf, M.; Dick, T. P. Multiple
750 glutathione disulfide removal pathways mediate cytosolic redox homeostasis. *Nat Chem Biol*
751 **9**:119-125; 2013.

752 [16] Eriksson, S.; Prigge, J. R.; Talago, E. A.; Arner, E. S.; Schmidt, E. E. Dietary methionine can
753 sustain cytosolic redox homeostasis in the mouse liver. *Nat Commun* **6**:6479; 2015.

754 [17] Lee, J.; Dawes, I. W.; Roe, J. H. Isolation, expression, and regulation of the pgr1(+) gene
755 encoding glutathione reductase absolutely required for the growth of *Schizosaccharomyces*
756 *pombe*. *J Biol Chem* **272**:23042-23049; 1997.

757 [18] Pastrana-Mena, R.; Dinglasan, R. R.; Franke-Fayard, B.; Vega-Rodriguez, J.; Fuentes-
758 Caraballo, M.; Baerga-Ortiz, A.; Coppens, I.; Jacobs-Lorena, M.; Janse, C. J.; Serrano, A. E.
759 Glutathione reductase-null malaria parasites have normal blood stage growth but arrest
760 during development in the mosquito. *J Biol Chem*; 2010.

761 [19] Tzafrir, I.; Pena-Muralla, R.; Dickerman, A.; Berg, M.; Rogers, R.; Hutchens, S.; Sweeney,
762 T. C.; McElver, J.; Aux, G.; Patton, D.; Meinke, D. Identification of genes required for embryo
763 development in *Arabidopsis*. *Plant Physiol* **135**:1206-1220; 2004.

764 [20] Kanzok, S. M.; Fechner, A.; Bauer, H.; Ulschmid, J. K.; Muller, H. M.; Botella-Munoz, J.;
765 Schnewly, S.; Schirmer, R.; Becker, K. Substitution of the thioredoxin system for glutathione
766 reductase in *Drosophila melanogaster*. *Science* **291**:643-646; 2001.

767 [21] Meyer, Y.; Belin, C.; Delorme-Hinoux, V.; Reichheld, J. P.; Riondet, C. Thioredoxin and
768 Glutaredoxin Systems in Plants: Molecular Mechanisms, Cross Talks and Functional
769 Significance. *Antioxid Redox Signal*; 2012.

770 [22] Kelner, M. J.; Montoya, M. A. Structural organization of the human glutathione reductase
771 gene: determination of correct cDNA sequence and identification of a mitochondrial leader
772 sequence. *Biochem Biophys Res Commun* **269**:366-368; 2000.

773 [23] Tamura, T.; McMicken, H. W.; Smith, C. V.; Hansen, T. N. Gene structure for mouse
774 glutathione reductase, including a putative mitochondrial targeting signal. *Biochem Biophys*
775 *Res Commun* **237**:419-422; 1997.

776 [24] Outten, C. E.; Culotta, V. C. Alternative start sites in the *Saccharomyces cerevisiae* GLR1
777 gene are responsible for mitochondrial and cytosolic isoforms of glutathione reductase. *J Biol*
778 *Chem* **279**:7785-7791; 2004.

779 [25] Song, J. Y.; Cha, J.; Lee, J.; Roe, J. H. Glutathione reductase and a mitochondrial
780 thioredoxin play overlapping roles in maintaining iron-sulfur enzymes in fission yeast.
781 *Eukaryot Cell* **5**:1857-1865; 2006.

782 [26] Prinz, W. A.; Aslund, F.; Holmgren, A.; Beckwith, J. The role of the thioredoxin and
783 glutaredoxin pathways in reducing protein disulfide bonds in the *Escherichia coli* cytoplasm.
784 *J Biol Chem* **272**:15661-15667; 1997.

785 [27] Kumar, C.; Igbaria, A.; D'Autreaux, B.; Planson, A. G.; Junot, C.; Godat, E.; Bachhawat, A.
786 K.; Delaunay-Moisan, A.; Toledano, M. B. Glutathione revisited: a vital function in iron
787 metabolism and ancillary role in thiol-redox control. *EMBO J* **30**:2044-2056; 2011.

788 [28] Toledano, M. B.; Delaunay-Moisan, A.; Outten, C. E.; Igbaria, A. Functions and cellular
789 compartmentation of the thioredoxin and glutathione pathways in yeast. *Antioxid Redox*
790 *Signal* **18**:1699-1711; 2013.

791 [29] Gostimskaya, I.; Grant, C. M. Yeast mitochondrial glutathione is an essential antioxidant
792 with mitochondrial thioredoxin providing a back-up system. *Free Radic Biol Med*; 2016.

793 [30] Rogers, L. K.; Tamura, T.; Rogers, B. J.; Welty, S. E.; Hansen, T. N.; Smith, C. V. Analyses of
794 glutathione reductase hypomorphic mice indicate a genetic knockout. *Toxicol Sci* **82**:367-
795 373; 2004.

796 [31] Matsui, M.; Oshima, M.; Oshima, H.; Takaku, K.; Maruyama, T.; Yodoi, J.; Taketo, M. M.
 797 Early embryonic lethality caused by targeted disruption of the mouse thioredoxin gene. *Dev*
 798 *Biol* **178**:179-185; 1996.
 799 [32] Conrad, M.; Jakupoglu, C.; Moreno, S. G.; Lippl, S.; Banjac, A.; Schneider, M.; Beck, H.;
 800 Hatzopoulos, A. K.; Just, U.; Sinowatz, F.; Schmahl, W.; Chien, K. R.; Wurst, W.; Bornkamm, G.
 801 W.; Brielmeier, M. Essential role for mitochondrial thioredoxin reductase in hematopoiesis,
 802 heart development, and heart function. *Mol Cell Biol* **24**:9414-9423; 2004.
 803 [33] Jakupoglu, C.; Przemeck, G. K.; Schneider, M.; Moreno, S. G.; Mayr, N.; Hatzopoulos, A. K.;
 804 de Angelis, M. H.; Wurst, W.; Bornkamm, G. W.; Brielmeier, M.; Conrad, M. Cytoplasmic
 805 thioredoxin reductase is essential for embryogenesis but dispensable for cardiac
 806 development. *Mol Cell Biol* **25**:1980-1988; 2005.
 807 [34] Nonn, L.; Williams, R. R.; Erickson, R. P.; Powis, G. The absence of mitochondrial
 808 thioredoxin 2 causes massive apoptosis, exencephaly, and early embryonic lethality in
 809 homozygous mice. *Mol Cell Biol* **23**:916-922; 2003.
 810 [35] Mandal, P. K.; Schneider, M.; Kolle, P.; Kuhlencordt, P.; Forster, H.; Beck, H.; Bornkamm,
 811 G. W.; Conrad, M. Loss of thioredoxin reductase 1 renders tumors highly susceptible to
 812 pharmacologic glutathione deprivation. *Cancer Res* **70**:9505-9514; 2010.
 813 [36] Suvorova, E. S.; Lucas, O.; Weisend, C. M.; Rollins, M. F.; Merrill, G. F.; Capecchi, M. R.;
 814 Schmidt, E. E. Cytoprotective Nrf2 pathway is induced in chronically txnrd 1-deficient
 815 hepatocytes. *PLoS One* **4**:e6158; 2009.
 816 [37] Stenvall, J.; Fierro-Gonzalez, J. C.; Swoboda, P.; Saamarthy, K.; Cheng, Q.; Cacho-Valadez,
 817 B.; Arner, E. S.; Persson, O. P.; Miranda-Vizuete, A.; Tuck, S. Selenoprotein TRXR-1 and GSR-1
 818 are essential for removal of old cuticle during molting in *Caenorhabditis elegans*. *Proc Natl*
 819 *Acad Sci U S A* **108**:1064-1069; 2011.
 820 [38] Cacho-Valadez, B.; Munoz-Lobato, F.; Pedrajas, J. R.; Cabello, J.; Fierro-Gonzalez, J. C.;
 821 Navas, P.; Swoboda, P.; Link, C. D.; Miranda-Vizuete, A. The characterization of the
 822 *Caenorhabditis elegans* mitochondrial thioredoxin system uncovers an unexpected
 823 protective role of thioredoxin reductase 2 in beta-amyloid peptide toxicity. *Antioxid Redox*
 824 *Signal* **16**:1384-1400; 2012.
 825 [39] Luersen, K.; Stegehake, D.; Daniel, J.; Drescher, M.; Ajonina, I.; Ajonina, C.; Hertel, P.;
 826 Woltersdorf, C.; Liebau, E. The glutathione reductase GSR-1 determines stress tolerance and
 827 longevity in *Caenorhabditis elegans*. *PLoS One* **8**:e60731; 2013.
 828 [40] Habib, S. J.; Neupert, W.; Rapaport, D. Analysis and prediction of mitochondrial
 829 targeting signals. *Methods Cell Biol* **80**:761-781; 2007.
 830 [41] Hendrick, J. P.; Hodges, P. E.; Rosenberg, L. E. Survey of amino-terminal proteolytic
 831 cleavage sites in mitochondrial precursor proteins: leader peptides cleaved by two matrix
 832 proteases share a three-amino acid motif. *Proc Natl Acad Sci U S A* **86**:4056-4060; 1989.
 833 [42] Fuxman Bass, J. I.; Tamburino, A. M.; Mori, A.; Beittel, N.; Weirauch, M. T.; Reece-Hoyes,
 834 J. S.; Walhout, A. J. Transcription factor binding to *Caenorhabditis elegans* first introns reveals
 835 lack of redundancy with gene promoters. *Nucleic Acids Res* **42**:153-162; 2014.
 836 [43] Petit, E.; Michelet, X.; Rauch, C.; Bertrand-Michel, J.; Terce, F.; Legouis, R.; Morel, F.
 837 Glutathione transferases kappa 1 and kappa 2 localize in peroxisomes and mitochondria,
 838 respectively, and are involved in lipid metabolism and respiration in *Caenorhabditis elegans*.
 839 *FEBS J* **276**:5030-5040; 2009.
 840 [44] Edgley, M. L.; Baillie, D. L.; Riddle, D. L.; Rose, A. M. Genetic balancers. *WormBook*:1-32;
 841 2006.

842 [45] Wang, J.; Robida-Stubbs, S.; Tullet, J. M.; Rual, J. F.; Vidal, M.; Blackwell, T. K. RNAi
843 screening implicates a SKN-1-dependent transcriptional response in stress resistance and
844 longevity deriving from translation inhibition. *PLoS Genet* **6**; 2010.

845 [46] Yoneda, T.; Benedetti, C.; Urano, F.; Clark, S. G.; Harding, H. P.; Ron, D. Compartment-
846 specific perturbation of protein handling activates genes encoding mitochondrial
847 chaperones. *J Cell Sci* **117**:4055-4066; 2004.

848 [47] Ahringer, J. Embryonic tissue differentiation in *Caenorhabditis elegans* requires dif-1, a
849 gene homologous to mitochondrial solute carriers. *EMBO J* **14**:2307-2316; 1995.

850 [48] Asencio, C.; Navas, P.; Cabello, J.; Schnabel, R.; Cypser, J. R.; Johnson, T. E.; Rodriguez-
851 Aguilera, J. C. Coenzyme Q supports distinct developmental processes in *Caenorhabditis*
852 *elegans*. *Mech Ageing Dev* **130**:145-153; 2009.

853 [49] Kruger, A. V.; Jelier, R.; Dzyubachyk, O.; Zimmerman, T.; Meijering, E.; Lehner, B.
854 Comprehensive single cell-resolution analysis of the role of chromatin regulators in early *C.*
855 *elegans* embryogenesis. *Dev Biol* **398**:153-162; 2015.

856 [50] Garcia-Gimenez, J. L.; Olasso, G.; Hake, S. B.; Bonisch, C.; Wiedemann, S. M.; Markovic, J.;
857 Dasi, F.; Gimeno, A.; Perez-Quilis, C.; Palacios, O.; Capdevila, M.; Vina, J.; Pallardo, F. V. Histone
858 h3 glutathionylation in proliferating mammalian cells destabilizes nucleosomal structure.
859 *Antioxid Redox Signal* **19**:1305-1320; 2013.

860 [51] Badrinath, A. S.; White, J. G. Contrasting patterns of mitochondrial redistribution in the
861 early lineages of *Caenorhabditis elegans* and *Acroboloides* sp. PS1146. *Dev Biol* **258**:70-75;
862 2003.

863 [52] Cascella, R.; Evangelisti, E.; Zampagni, M.; Becatti, M.; D'Adamio, G.; Goti, A.; Liguri, G.;
864 Fiorillo, C.; Cecchi, C. S-linolenoyl glutathione intake extends life-span and stress resistance
865 via Sir-2.1 upregulation in *Caenorhabditis elegans*. *Free Radic Biol Med* **73**:127-135; 2014.

866 [53] Shibamura, A.; Ikeda, T.; Nishikawa, Y. A method for oral administration of hydrophilic
867 substances to *Caenorhabditis elegans*: Effects of oral supplementation with antioxidants on
868 the nematode lifespan. *Mech Ageing Dev* **130**:652-655; 2009.

869 [54] Levy, E. J.; Anderson, M. E.; Meister, A. Transport of glutathione diethyl ester into
870 human cells. *Proc Natl Acad Sci U S A* **90**:9171-9175; 1993.

871 [55] Yang, W.; Hekimi, S. A mitochondrial superoxide signal triggers increased longevity in
872 *Caenorhabditis elegans*. *PLoS Biol* **8**:e1000556; 2010.

873 [56] Edwards, C.; Canfield, J.; Copes, N.; Brito, A.; Rehan, M.; Lipps, D.; Brunquell, J.;
874 Westerheide, S. D.; Bradshaw, P. C. Mechanisms of amino acid-mediated lifespan extension in
875 *Caenorhabditis elegans*. *BMC Genet* **16**:8; 2015.

876 [57] Wharton, D. Nematode egg-shells. *Parasitology* **81**:447-463; 1980.

877 [58] Trotter, E. W.; Grant, C. M. Overlapping roles of the cytoplasmic and mitochondrial
878 redox regulatory systems in the yeast *Saccharomyces cerevisiae*. *Eukaryot Cell* **4**:392-400;
879 2005.

880 [59] Dethlefsen, L. A.; Lehman, C. M.; Biaglow, J. E.; Peck, V. M. Toxic effects of acute
881 glutathione depletion by buthionine sulfoximine and dimethylfumarate on murine mammary
882 carcinoma cells. *Radiat Res* **114**:215-224; 1988.

883 [60] Markovic, J.; Mora, N. J.; Broseta, A. M.; Gimeno, A.; de-la-Concepcion, N.; Vina, J.;
884 Pallardo, F. V. The depletion of nuclear glutathione impairs cell proliferation in 3t3
885 fibroblasts. *PLoS One* **4**:e6413; 2009.

886 [61] Sengupta, R.; Holmgren, A. Thioredoxin and glutaredoxin-mediated redox regulation of
887 ribonucleotide reductase. *World J Biol Chem* **5**:68-74; 2014.

888 [62] Zahedi Avval, F.; Holmgren, A. Molecular mechanisms of thioredoxin and glutaredoxin
889 as hydrogen donors for Mammalian S phase ribonucleotide reductase. *J Biol Chem* **284**:8233-
890 8240; 2009.

891 [63] Olson, S. K.; Greenan, G.; Desai, A.; Muller-Reichert, T.; Oegema, K. Hierarchical assembly
892 of the eggshell and permeability barrier in *C. elegans*. *J Cell Biol* **198**:731-748; 2012.

893 [64] Brennan, L. D.; Roland, T.; Morton, D. G.; Fellman, S. M.; Chung, S.; Soltani, M.; Kevek, J.
894 W.; McEuen, P. M.; Kempfues, K. J.; Wang, M. D. Small molecule injection into single-cell *C.*
895 *elegans* embryos via carbon-reinforced nanopipettes. *PLoS One* **8**:e75712; 2013.

896 [65] Ribas, V.; Garcia-Ruiz, C.; Fernandez-Checa, J. C. Glutathione and mitochondria. *Front*
897 *Pharmacol* **5**:151; 2014.

898 [66] Kojer, K.; Bien, M.; Gangel, H.; Morgan, B.; Dick, T. P.; Riemer, J. Glutathione redox
899 potential in the mitochondrial intermembrane space is linked to the cytosol and impacts the
900 Mia40 redox state. *EMBO J* **31**:3169-3182; 2012.

901 [67] Schaedler, T. A.; Thornton, J. D.; Kruse, I.; Schwarzlander, M.; Meyer, A. J.; van Veen, H.
902 W.; Balk, J. A conserved mitochondrial ATP-binding cassette transporter exports glutathione
903 polysulfide for cytosolic metal cofactor assembly. *J Biol Chem* **289**:23264-23274; 2014.

904 [68] Gonzalez-Cabo, P.; Bolinches-Amoros, A.; Cabello, J.; Ros, S.; Moreno, S.; Baylis, H. A.;
905 Palau, F.; Vazquez-Manrique, R. P. Disruption of the ATP-binding cassette B7 (ABTM-
906 1/ABCB7) induces oxidative stress and premature cell death in *Caenorhabditis elegans*. *J Biol*
907 *Chem* **286**:21304-21314; 2011.

908 [69] Landry, A. P.; Cheng, Z.; Ding, H. Reduction of mitochondrial protein mitoNEET [2Fe-2S]
909 clusters by human glutathione reductase. *Free Radic Biol Med*; 2015.

910 [70] Lazarou, M.; Narendra, D. P.; Jin, S. M.; Tekle, E.; Banerjee, S.; Youle, R. J. PINK1 drives
911 Parkin self-association and HECT-like E3 activity upstream of mitochondrial binding. *J Cell*
912 *Biol* **200**:163-172; 2013.

913 [71] Sulston J, H. J. *Methods. In: The Nematode Caenorhabditis elegans.* . Cold Spring Harbour,
914 New York: Cold Spring harbour Laboratory Press, Cold Spring Harbour; 1998.

915 [72] Morales-Martinez, A.; Dobrzynska, A.; Askjaer, P. Inner nuclear membrane protein LEM-
916 2 is required for correct nuclear separation and morphology in *C. elegans*. *J Cell Sci*
917 **128**:1090-1096; 2015.

918 [73] Mello, C. C.; Kramer, J. M.; Stinchcomb, D.; Ambros, V. Efficient gene transfer in
919 *C.elegans*: extrachromosomal maintenance and integration of transforming sequences. *EMBO*
920 *J* **10**:3959-3970; 1991.

921 [74] Miyabayashi, T.; Palfreyman, M. T.; Sluder, A. E.; Slack, F.; Sengupta, P. Expression and
922 function of members of a divergent nuclear receptor family in *Caenorhabditis elegans*. *Dev*
923 *Biol* **215**:314-331; 1999.

924 [75] Muller, S.; Walter, R. D.; Fairlamb, A. H. Differential susceptibility of filarial and human
925 erythrocyte glutathione reductase to inhibition by the trivalent organic arsenical melarsen
926 oxide. *Mol Biochem Parasitol* **71**:211-219; 1995.

927 [76] Nieto, C.; Almendinger, J.; Gysi, S.; Gomez-Orte, E.; Kaech, A.; Hengartner, M. O.;
928 Schnabel, R.; Moreno, S.; Cabello, J. ccz-1 mediates the digestion of apoptotic corpses in *C.*
929 *elegans*. *J Cell Sci* **123**:2001-2007; 2010.

930 [77] Larsen, P. L.; Albert, P. S.; Riddle, D. L. Genes that regulate both development and
931 longevity in *Caenorhabditis elegans*. *Genetics* **139**:1567-1583; 1995.

932 [78] Inoue, H.; Hisamoto, N.; An, J. H.; Oliveira, R. P.; Nishida, E.; Blackwell, T. K.; Matsumoto,
933 K. The *C. elegans* p38 MAPK pathway regulates nuclear localization of the transcription
934 factor SKN-1 in oxidative stress response. *Genes Dev* **19**:2278-2283; 2005.

935 [79] Etheridge, T.; Rahman, M.; Gaffney, C. J.; Shaw, D.; Shephard, F.; Magudia, J.; Solomon, D.
 936 E.; Milne, T.; Blawzdziejewicz, J.; Constantin-Teodosiu, D.; Greenhaff, P. L.; Vanapalli, S. A.;
 937 Szewczyk, N. J. The integrin-adhesome is required to maintain muscle structure,
 938 mitochondrial ATP production, and movement forces in *Caenorhabditis elegans*. *FASEB J*
 939 **29**:1235-1246; 2015.
 940 [80] Krauth-Siegel, R. L.; Blatterspiel, R.; Saleh, M.; Schiltz, E.; Schirmer, R. H.; Untucht-Grau,
 941 R. Glutathione reductase from human erythrocytes. The sequences of the NADPH domain and
 942 of the interface domain. *Eur J Biochem* **121**:259-267; 1982.
 943 [81] Urig, S.; Lieske, J.; Fritz-Wolf, K.; Irmeler, A.; Becker, K. Truncated mutants of human
 944 thioredoxin reductase 1 do not exhibit glutathione reductase activity. *FEBS Lett* **580**:3595-
 945 3600; 2006.
 946
 947







FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5





FIGURE 6





FIGURE 7

Supplemental Table 1: *C. elegans* strains used in this work.

Strain name	Genotype	Reference/Source
N2	Wild type, DR subclone of CB original (Tc1 pattern I)	CGC ^a
FX3574	<i>gsr-1(tm3574)</i> III	NBRP ^b
JK2533	<i>qC1 [dpy-19(e1259) glp-1(q339) qIs26 [Plag-2::gfp; rol-6(su1006)]] III/eT1 (III;V)</i>	CGC
MT20108	<i>dpy-17(e164) unc-32(e189)/qC1 [dpy-19(e1259) glp-1(q339) nls281 [Pmyo-2::rfp]] III</i>	CGC
VZ64	<i>gsr-1(tm3574)/qC1 [dpy-19(e1259) glp-1(q339) qIs26 [Plag-2::gfp; rol-6(su1006)]] III</i>	This study
VZ454	<i>gsr-1(tm3574)/qC1 [dpy-19(e1259) glp-1(q339) nls281 [Pmyo-2::rfp]] III</i>	This study
VZ291	<i>gsr-1(tm3574) III; vzEx105 [Pg_{gsr-1}::gsr-1::3'-UTR_{gsr-1}; Punc-122::gfp]</i>	
VZ292	<i>gsr-1(tm3574) III; vzEx106 [Pg_{gsr-1}::gsr-1::3'-UTR_{gsr-1}; Punc-122::gfp]</i>	This study
VZ232	<i>vzEx76 [Pg_{gsr-1}::gfp; rol-6 (su1006)]</i>	This study
VZ233	<i>vzEx77 [Pg_{gsr-1}::gfp; rol-6 (su1006)]</i>	This study
VZ234	<i>vzEx78 [Pg_{gsr-1}::gfp; rol-6 (su1006)]</i>	This study
VZ257	<i>vzEx93 [Pg_{gsr-1}::gsr-1::gfp; rol-6(su1006)]</i>	This study
VZ258	<i>vzEx94 [Pg_{gsr-1}::gsr-1::gfp; rol-6(su1006)]</i>	This study
VZ341	<i>gsr-1(tm3574) III; vzEx93 [Pg_{gsr-1}::gsr-1::gfp; rol-6(su1006)]</i>	This study
VZ307	<i>vzEx117 [gsr-1 1st intron::gsr-1::gfp]</i>	This study
VZ308	<i>vzEx118 [gsr-1 1st intron::gsr-1::gfp]</i>	This study
VZ309	<i>vzEx119 [gsr-1 1st intron::gsr-1::gfp]</i>	This study
VZ342	<i>gsr-1(tm3574)/qC1 [dpy-19(e1259) glp-1(q339) qIs26 [Plag-2::gfp; rol-6(su1006)]] III; vzEx119 [gsr-1 1st intron::gsr-1::gfp]</i>	This study
VZ375	<i>gsr-1(tm3574) III; vzEx144 [Pg_{gsr-1}::gsr-1(Δ1st exon)::gfp]</i>	This study
VZ378	<i>gsr-1(tm3574) III; vzEx145 [Pg_{gsr-1}::gsr-1(Δ1st exon)::gfp]</i>	This study
VZ393	<i>gsr-1(tm3574) III; vzEx146 [Pg_{gsr-1}::gsr-1(Δ1st exon)::gfp]</i>	This study
VZ395	<i>gsr-1(tm3574) III; vzEx147 [Pg_{gsr-1}::gsr-1(1st ATG->CCT)::gfp]</i>	This study
VZ396	<i>gsr-1(tm3574) III; vzEx148 [Pg_{gsr-1}::gsr-1(1st ATG->CCT)::gfp]</i>	This study
VZ397	<i>gsr-1(tm3574) III; vzEx149 [Pg_{gsr-1}::gsr-1(1st ATG->CCT)::gfp]</i>	This study
VZ445	<i>vzEx158 [Pg_{gsr-1}::gsr-1(2nd ATG->GGT)::gfp]</i>	This study
VZ446	<i>vzEx159 [Pg_{gsr-1}::gsr-1(2nd ATG->GGT)::gfp]</i>	This study
VZ447	<i>vzEx160 [Pg_{gsr-1}::gsr-1(2nd ATG->GGT)::gfp]</i>	This study
VZ610	<i>gsr-1(tm3574)/qC1 [dpy-19(e1259) glp-1(q339) nls281 [Pmyo-2::rfp]] III; vzEx158 [Pg_{gsr-1}::gsr-1(2nd ATG->GGT)::gfp]</i>	This study
PS6187	<i>unc-119 (ed3) III; syEx1155 [Pmyo-3::tomm-20::rfp::3xMyc; unc-119(+)]</i>	[1]
VZ427	<i>vzEx155 [Pmyo-3::MTS_{gsr-1}::gfp]</i>	This study
VZ428	<i>vzEx156 [Pmyo-3::MTS_{gsr-1}::gfp]</i>	This study
VZ429	<i>vzEx157 [Pmyo-3::MTS_{gsr-1}::gfp]</i>	This study
VZ436	<i>vzEx155 [Pmyo-3::MTS_{gsr-1}::gfp]; syEx1155 [Pmyo-3::tomm-20::rfp::3xMyc; unc-119(+)]</i>	This study
VZ12	<i>trxr-2(tm2047)</i>	[1]
VZ594	<i>gsr-1(tm3574) trxr-2(tm2047)/qC1 [dpy-19(e1259) glp-1(q339) nls281 [Pmyo-2::rfp]] III</i>	This study
VZ603	<i>gsr-1(tm3574) trxr-2(tm2047); vzEx147 [Pg_{gsr-1}::gsr-1(1st ATG->CCT)::gfp]</i>	This study
VZ604	<i>gsr-1(tm3574) trxr-2(tm2047); vzEx145 [Pg_{gsr-1}::gsr-1(Δ1st exon)::gfp]</i>	This study

VB1414	<i>trxr-1(sv47) IV</i>	[2]
VZ78	<i>gsr-1(tm3574)/qC1 [dpy-19(e1259) glp-1(q339) qIs26 [Plag-2::gfp; rol-6(su1006)]] III; trxr-1(sv47) IV</i>	This study
VZ492	<i>gsr-1(tm3574)/qC1 [dpy-19(e1259) glp-1(q339) nIs281 [Pmyo-2::rfp]] III; trxr-1(sv47) IV</i>	This study
CL1166	<i>dvEx166 [Pgst-4::gfp::NLS]</i>	[3]
VZ498	<i>gsr-1(tm3574)/qC1 [dpy-19(e1259) glp-1(q339) nIs281 [Pmyo-2::rfp]] III; dvEx166 [Pgst-4::gfp::NLS]</i>	This study
VB2317	<i>svEx741 [Pgcs-1::gfp]</i>	[2]
VZ468	<i>gsr-1(tm3574)/qC1 [dpy-19(e1259) glp-1(q339) nIs281 [Pmyo-2::rfp]] III; svEx741 [Pgcs-1::gfp]</i>	This study
LD001	<i>ldIs007 [Pskn-1::skn-1b/c::gfp; rol-6(su1006)] X</i>	[4]
VZ452	<i>gsr-1(tm3574)/qC1 [dpy-19(e1259) glp-1(q339) nIs281 [Pmyo-2::rfp]] III; ldIs007 [Pskn-1::skn-1b/c::gfp; rol-6(su1006)] X</i>	This study
PD4251	<i>ccls4251 [Pmyo-3::nucgfp::LacZ::NLS; Pmyo-3::mitogfp]; dpy-20(+)] I; dpy-20(e1282) IV</i>	CGC
VZ508	<i>ccls4251 [Pmyo-3::nucgfp::LacZ::NLS; Pmyo-3::mitogfp]; dpy-20(+)] I; gsr-1(tm3574)/qC1 [dpy-19(e1259) glp-1(q339) nIs281 [Pmyo-2::rfp]] III</i>	This study
PJ707	<i>jIs01[Pmyo-3::myo-3::gfp; rol-6(su1006)]</i>	[5]
VZ507	<i>gsr-1(tm3574)/qC1 [dpy-19(e1259) glp-1(q339) nIs281 [Pmyo-2::rfp]] III; jIs01[Pmyo-3::myo-3::gfp; rol-6(su1006)]</i>	This study
SJ4100	<i>zcls13 [Phsp-6::gfp] V</i>	[6]
VZ495	<i>gsr-1(tm3574)/qC1 [dpy-19(e1259) glp-1(q339) nIs281 [Pmyo-2::rfp]] III; zcls13 [Phsp-6::gfp] V</i>	This study
BN245	<i>ltIs37 [Ppie-1::mCherry::his-58; unc-119 (+)] IV; oJIs1 [Ppie-1::gfp::tbb-2; unc-119(+)] V; ltIs24 [Ppie-1::gfp::tba-2; unc-119(+)]</i>	[7]
BN323	<i>gsr-1(tm3574)/qC1 [dpy-19(e1259) glp-1(q339) qIs26 [Plag-2::gfp; rol-6(su1006)]] III; ltIs37 [Ppie-1::mCherry::his-58; unc-119 (+)] IV; oJIs1 [Ppie-1::gfp::tbb-2; unc-119(+)] V</i>	This study

^a Caenorhabditis Genetics Center (<http://www.cbs.umn.edu/CGC/>)

^b National BioResource Project C. elegans (<http://www.shigen.nig.ac.jp/c.elegans/index.jsp>)

SUPPLEMENTAL FIGURES

Supplemental Figure 1: Complete cell lineage analysis of *wild type* and *gsr-1(m-,z-)* embryos.

Comparison of cell lineage and cell divisions timing of *wild type* control embryos (grey lines) with *gsr-1(m-,z-)* embryos (black lines with red dots) arresting at **A)** 17 cell-stage; **B)** 33 cell-stage; **C)** 103 cell-stage (same as depicted in **Figure 5A-C**). Green dots show cell mitosis.


Supplemental Figure 2: Sensitivity of *gsr-1(m-,z-)* animals with restricted expression of GSR-

1b::GFP in cytoplasm to acute paraquat and juglone exposure. *gsr-1(m-,z-)* worms with GSR-1 activity restricted to cytoplasm are as resistant as wild type control to paraquat and juglone treatments. Therefore, mitochondrial glutathione reductase isoform is not required for resistance to these chemicals. Data are the mean $\pm$ SD of three independent experiments with three biological replicates each ($n \pm 200$).


SUPPLEMENTAL REFERENCES

- [1] Cacho-Valadez, B.; Munoz-Lobato, F.; Pedrajas, J. R.; Cabello, J.; Fierro-Gonzalez, J. C.; Navas, P.; Swoboda, P.; Link, C. D.; Miranda-Vizuete, A. The characterization of the *Caenorhabditis elegans* mitochondrial thioredoxin system uncovers an unexpected protective role of thioredoxin reductase 2 in beta-amyloid peptide toxicity. *Antioxid Redox Signal* **16**:1384-1400; 2012.
- [2] Stenvall, J.; Fierro-Gonzalez, J. C.; Swoboda, P.; Saamarthi, K.; Cheng, Q.; Cacho-Valadez, B.; Arner, E. S.; Persson, O. P.; Miranda-Vizuete, A.; Tuck, S. Selenoprotein TRXR-1 and GSR-1 are essential for removal of old cuticle during molting in *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A* **108**:1064-1069; 2011.
- [3] Link, C. D.; Johnson, C. J. Reporter transgenes for study of oxidant stress in *Caenorhabditis elegans*. *Methods Enzymol* **353**:497-505; 2002.
- [4] An, J. H.; Blackwell, T. K. SKN-1 links *C. elegans* mesendodermal specification to a conserved oxidative stress response. *Genes Dev* **17**:1882-1893; 2003.
- [5] Szewczyk, N. J.; Kozak, E.; Conley, C. A. Chemically defined medium and *Caenorhabditis elegans*. *BMC Biotechnol* **3**:19; 2003.
- [6] Yoneda, T.; Benedetti, C.; Urano, F.; Clark, S. G.; Harding, H. P.; Ron, D. Compartment-specific perturbation of protein handling activates genes encoding mitochondrial chaperones. *J Cell Sci* **117**:4055-4066; 2004.
- [7] Morales-Martinez, A.; Dobrzynska, A.; Askjaer, P. Inner nuclear membrane protein LEM-2 is required for correct nuclear separation and morphology in *C. elegans*. *J Cell Sci* **128**:1090-1096; 2015.

A



B



C

