

Quantitative analysis of ruminal bacterial populations involved in lipid metabolism in dairy cows fed different vegetable oils

E. Vargas-Bello-Pérez^{1a}, N. Cancino-Padilla¹ J. Romero² and P. C. Garnsworthy³.

¹*Departamento de Ciencias Animales, Facultad de Agronomía e Ingeniería Forestal, Pontificia Universidad Católica de Chile, Santiago, Chile. Casilla 306. C.P. 6904411.*

²*Laboratorio de Biotecnología, Instituto de Nutrición y Tecnología de los Alimentos (INTA), Universidad de Chile, Santiago, Chile.*

³*The University of Nottingham, Sutton Bonington Campus, Loughborough, LE12 5RD, United Kingdom.*

^a *Corresponding author: evargasb@uc.cl. Departamento de Ciencias Animales. Facultad de Agronomía e Ingeniería Forestal. Pontificia Universidad Católica de Chile. Telephone: +56 (2) 3544239 / +56 (2) 3544142.*

Short title: Bacterial populations involved in lipid metabolism

Abstract

Vegetable oils are used to increase energy density of dairy cow diets, although they can provoke changes in rumen bacteria populations and have repercussion on the biohydrogenation process. The aim of this study was to evaluate the effect of two sources of dietary lipids; soybean oil (SO; an unsaturated source) and hydrogenated palm oil (HPO; a saturated source) on bacterial populations and the fatty acid (FA) profile of ruminal digesta. Three non-lactating Holstein cows fitted with ruminal cannulae were used in a 3x3 Latin square design with 3 periods consisting of 21 d. Dietary treatments consisted of a basal diet (Control; no fat supplement), and the basal diet supplemented with SO (2.7 % of DM) or HPO (2.7 % of DM). Ruminal digesta pH, NH₃-N and VFA were not affected by dietary treatments. Compared with control and HPO, total bacteria measured as copies of

16S rDNA/ml by qPCR was decreased ($P<0.05$) by SO. *Fibrobacter succinogenes*, *Butyrivibrio proteoclasticus*, and *Anaerovibrio lipolytica* loads were not affected by dietary treatments. In contrast, compared with control, load of *Prevotella bryantii* was increased ($P<0.05$) with HPO diet. Compared with control and SO, HPO decreased ($P<0.05$) C18:2 cis n-6 in ruminal digesta. Contents of C15:0 iso, C18:11 trans-11 and C18:2 cis-9, trans-11 were increased ($P<0.05$) in ruminal digesta by SO compared with control and HPO. In conclusion, supplementation of SO or HPO do not affect ruminal fermentation parameters whereas HPO can increase load of ruminal *Prevotella bryantii*. Also, results observed in our targeted bacteria may have depended on the saturation degree of dietary oils.

Keywords: Soybean oil, rumen fermentation, vegetable oil, palm oil

Implications

A better knowledge of the rumen microbiome may help us to understand, and eventually modulate, the effect of nutrition on milk fat production and quality. This work was conducted to evaluate the effect of two sources of dietary lipids; soybean oil (SO; an unsaturated source) and hydrogenated palm oil (HPO; a saturated source) on bacterial populations and the fatty acid profile of ruminal digesta. Contents of C15:0 iso, C18:11 trans-11 and C18:2 cis-9, trans-11 were increased by SO. Supplementation with SO or HPO (2.7 % DM) did not affect ruminal fermentation parameters whereas HPO can increase loads of ruminal *Prevotella bryantii*.

Introduction

Dietary polyunsaturated fatty acids (PUFA) have toxic effects on ruminal microorganisms, therefore, lipid supplementation often leads to changes in ruminal microbial populations and shifts in ruminal fermentation parameters (Zhang *et al.*, 2008). Rumen microbes attempt to detoxify PUFA by biohydrogenation (Maia *et al.*, 2010). Biohydrogenation pathways require an initial hydrolysis of ingested dietary glyceride by microbial lipases/esterases causing the release of FA (Prive *et al.*, 2015) at this stage; *Anaerovibrio lipolytica* is recognized as one of the major species involved in lipid hydrolysis in ruminants (Prive *et al.*, 2013). Wallace *et al.* (2006) proposed that *Butyrivibrio* genus contained the main bacterial species involved in the biohydrogenation process. However, Huws *et al.* (2011) demonstrated that as yet uncultured bacteria belonging to the genera *Prevotella* and *Anaerovoax*, and unclassified *Ruminococcaceae* and *Clostridiales* may play more important roles in ruminal biohydrogenation.

It is known that cellulolytic bacteria (*Fibrobacter succinogenes*, *Ruminococcus flavefaciens*, *Ruminococcus albus*, and *Butyrivibrio fibrisolvens*) are important in the biohydrogenation process of dietary sources of PUFA (Potu *et al.*, 2011). Also, *Butyrivibrio proteoclasticus* has been reported to be the principal rumen bacteria involved in biohydrogenation of C18:1 FA (Boeckaert *et al.*, 2008). On the other hand, *Prevotella bryantii* has been described as a ruminal bacterium that is involved in oligosaccharolytic and xylanolytic activities (Tajima *et al.*, 2001) and also *Prevotella* spp. has been reported as resistant to inhibitory effects of dietary PUFA (Huws *et al.*, 2010).

81

82 Supplementing dairy cow diets with soybean oil (SO) can increase milk bioactive
83 FA such as C18:1 trans-11 (Allred *et al.*, 2006; Vargas-Bello-Perez *et al.*, 2015a).
84 Also, SO has been shown to reduce cellulolytic bacteria, protozoa populations and
85 total concentration of volatile fatty acids (Yang *et al.*, 2009). On the other hand,
86 hydrogenated vegetable oils have been used to increase the energy content of
87 dairy cow diets in housed (Kargar *et al.*, 2012) and pasture systems (Schroeder *et*
88 *al.*, 2002) without effect on milk composition (Vargas-Bello-Perez *et al.*, 2015b).

89

90 To our knowledge, no study on the effect of dietary hydrogenated palm oil on
91 ruminal bacterial populations in dairy cows has been published. Also, animal trials
92 reporting use of oils and their effect on rumen microbiome have less risk of bias
93 compared with *in vitro* studies. Therefore, the aim of this study was to make a
94 quantitative analysis of bacterial populations involved in ruminal biohydrogenation
95 (*Fibrobacter succinogenes*, *Butyrivibrio proteoclasticus* and *Anaerovibrio lipolytica*)
96 and *Prevotella bryantii* (one of the most predominant ruminal bacteria) in dairy
97 cows fed different vegetable oils (soybean oil as an unsaturated source and
98 hydrogenated palm oil as a saturated source). The effect of fat supplements on the
99 FA profile of ruminal digesta was another objective. Our hypothesis was that
100 supplementation with saturated versus unsaturated oils would have different
101 effects on bacterial populations that were or were not involved in biohydrogenation.

102

103 **Materials and methods**

104 *Animals and treatments*

Three non-lactating Holstein cows (684.7 ± 84.7 kg BW) fitted with ruminal cannulae # 3C (Bar Diamond, Inc., Boise, Idaho, USA) were used in a 3×3 Latin square design with 3 periods consisting of 21 d. Cows were fed to satisfy the requirements of a dry cow on the last trimester of gestation consuming 10 kg DM daily (NRC, 2001). Dietary treatments (Table 1) were a basal diet (C) containing 56% forage and 44% concentrate ratio with no fat supplement, and fat-supplemented diets containing soybean oil (SO; 170 g/d/cow = 2.7% DM) and hydrogenated palm oil (HPO; 170 g/d/cow = 2.7% DM). The amounts of oils used were similar to those reported to alter rumen FA in previous studies (Yang et al., 2009, Vargas-Bello-Perez et al., 2015a). The most important FA in dietary oils were: SO contained (g/100g) 25 of C18:1 cis-9 and 51 of C18:2 cis n-6, whereas HPO contained 47 of C16:0 and 43 of C18:0. Oils were administrated separately and mixed manually into the daily TMR for each cow. Animals were housed in individual stalls (2.4 × 6 m) and had free access to fresh water. Animal care and procedures were carried out according to the guidelines of the Animal Care and Use Committee of the Pontificia Universidad Católica de Chile.

Samples

On the last day of each 21-d period, samples of whole ruminal digesta were collected from the anterior, dorsal and mid-ventral regions of the rumen at 0900 h (2 h post feeding) and were squeezed through three layers of cheesecloth. Ten ml of residual ruminal fluid was immediately used to determine pH by using a pH meter (PP-201 GOnDO Electronic, Taipei, Taiwan), 10 ml were kept for NH₃-N analysis (Bal et al., 2000) and another 10 ml were preserved for volatile fatty acid

(VFA) determination by adding 1 ml of 25% metaphosphoric acid. Samples were frozen (-20°C) for later analysis. The VFA measurement were performed by gas chromatograph (GC-2010) equipped with a 30-m wall-coated open tubular-fused silica capillary column (Stabilwax-DA; 30 m × 0.32 mm i.d., 0.25 µm film thickness, Restek, Bellefonte, PA). Oven temperature was programmed for 145°C for 2 min and then increased from 145 to 220°C at 4°C/min. The injector and flame-ionization detector were 250 and 300°C, respectively. Following pH determination, the strained ruminal fluid was centrifuged for 10 min at 3,000 × g at room temperature. The supernatant was discarded and the residue was stored at -20°C until microbiology analysis.

DNA extraction

Samples from each cow and every period were weighed (240 ± 12 µg) and deposited in 1.5 ml Eppendorf tubes. Subsequently, 300 µl of phosphate-buffered saline (PBS) solution were added and mixed to homogenize the sample. DNA was obtained by incubating the sample for 30 min at 37 °C with lysozyme (1 µg/µl) and then for 30 min at 37 °C with proteinase K (0.1 mg / mL). DNA extraction was performed using the Power Soil DNA Isolation Kit (Mo-Bio Laboratories, Inc.), according to manufacturer's recommendations.

qPCR conditions

Primers (forward and reverse) used to target bacterial species of interest are described in Table 2. Primers for *Anaerovibrio lipolytica* (Tajima *et al.*, 2001), *Fibrobacter succinogenes* (Tajima *et al.*, 2001), *Butyrivibrio proteoclasticus* (Huws

et al., 2010), and *Prevotella bryantii* (Tajima et al., 2001) were those reported in previous research. Once obtained, the primers were tested for specificity using the probe match function at the Ribosomal Database Project (RDP; <https://rdp.cme.msu.edu/probematch/search.jsp>) as described Huws et al. (2007). The oligonucleotides from each target bacteria were synthesized for Integrated DNA Technologies (IDT, Coralville, IA). These primers were also analyzed for the requirements necessary for real-time PCR.

Real-time PCR quantification (qPCR) of total ruminal bacteria and bacterial species of interest was performed on a Rotor Gene 6000 (Corbett Life Science, Brisbane, Australia). Quantification of total ruminal bacteria was accomplished by qPCR amplifying the V3-V4 region of the 16S rRNA gene using the conserved bacterial domain-specific primers 341f (5'-CCTACGGGAGGCAGCAG-3') and 788r (5'-GGACTACCAGGGTATCTAA-3'). PCR reactions were carried out in quadruplicate and in 10 µl final volume containing 1 µl of extracted DNA (1: 1000 dilution), 25 pmol/µL of each primer, DNase-free water and 2x LightCycler® 480 DNA Master SYBR Green I (Roche Applied Science). PCR conditions started with an initial denaturation at 95°C for 5 min, followed by 50 cycles of denaturation at 95°C for 10 s, annealing at 60°C for 10 s and extension at 72°C for 15 s. The reaction mixture for quantification of specific bacteria consisted of 1 µl of DNA template, 20 pmol/µL of each specific primers described in Table 2, DNase-free water and 2x LightCycler® 480 DNA Master SYBR Green I (Roche Applied Science). The PCR program was similar to total bacterial quantification, except for annealing temperature. Annealing for *Anaerovibrio lipolytica* and *Butyrivibrio proteoclasticus*

was performed at 62°C, and *Fibrobacter succinogenes* and *Prevotell*
annealing was performed at 60°C. Specificity of qPCR reactions was confirmed by
analyzing the temperature characteristics of melting curves – increase of
temperature from 72 to 95°C, holding 1 s on the first step and 5 s on next steps.

The number of copies of the target bacterial 16S rDNA were determined by the
serially dilution of purified genomic DNA extracted from ruminal samples, with the
objective to construct specific calibration curves, and thus calculate the
concentration of total and target bacteria in samples. The bacterial concentrations
were calculated considering the rRNA operon copy number of each bacterial
genome described in Genbank as follow: *Fibrobacter succinogenes*, 3 copies
(Accession number CP001792.1); *Prevotella bryantii*, 4 copies (Accession number
NZ_AUKF000000000.1); *Butyrivibrio proteoclasticus*, 6 copies (Accession number
NZ_JHWL000000000.1) and *Anaerovibrio lipolytica*, 1 copy (Accession number
NZ_JHYA000000000.1). The qPCR efficiencies for bacterial species of interest were
obtained using standard dilution curves in quadruplicate of *Anaerovibrio lipolytica*,
Fibrobacter succinogenes, *Butyrivibrio proteoclasticus* and *Prevotella bryantii* 16S
rDNA, respectively. The qPCR efficiencies were calculated according to the
equation: $E = [(10^{(1 / \text{slope})}) - 1]$. Standard curves were generated using relative
concentration vs. the threshold cycle (Ct). The qPCR efficiencies (E) were
calculated from the given slopes (M) in a RotorGene 6000 software. Based on the
slopes of the standard curves, the qPCR efficiencies ranged from 80% to 97%. The
transcripts studied showed high linearity: $R^2 > 0.99$.

201 Additionally, to check the expected sizes of each PCR product, the amplicons were
202 visualized by electrophoresis on a 1% (w/v) agarose gel was stained using
203 ethidium bromide and Lambda DNA/ HindIII marker was used to compare the 16S
204 rDNA amplification fragments.

206 *Sequence analysis*

207 To verify the correct amplification in the qPCR assays of specific bacteria, the PCR
208 products were sequenced using the Macrogen USA sequencing service. The 16S
209 rDNA sequences were compared to the available databases using the basic Local
210 Alignment Search Tool (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) and aligned with
211 reference sequences using Sequence Match function at the Ribosomal Database
212 Project (https://rdp.cme.msu.edu/seqmatch/seqmatch_intro.jsp) to determine the
213 approximate phylogenetic affiliations.

215 *Fatty acid analysis*

216 Lipids from oils, diets and ruminal digesta were extracted with chloroform/methanol
217 (2:1, v/v) by the method of Bligh and Dyer (1959) and trans-esterified with sodium
218 methoxide according to the method of Christie (1982) using a methylation reagent
219 (1.75 mL methanol:0.4 mL of 5.4 mol/L sodium methylate) and a termination
220 reagent (1 g oxalic acid/30 mL diethyl ether) according to Chouinard *et al.* (1999).
221 All chemicals and solvents used for this method were of analytical grade. A GC
222 system (GC-2010, Shimadzu Scientific Instruments) equipped with a 100-m
223 column (Rt-2560 column 100 m × 0.32 mm × 0.20 µm column, Restek, Bellefonte,
224 PA) was used. The GC conditions were as follows: the oven temperature was

initially set at 110°C for 4 min after injection, and then increased to 240°C (20°C/min) with equilibration time of 2 min. The inlet and flame-ionization detector temperatures were 260°C, the split ratio was 15:1 and a 2 µl injection volume was used. The hydrogen carrier gas flow to the detector was 40 mL/min, airflow was 400 mL/min, and the flow of nitrogen makeup gas was 25 mL/min. Fatty acid peaks were identified by using a fatty acid methyl ester standard (FAME; Supelco 37 Component FAME mix, Bellefonte, PA, USA).

Statistical analysis

Bacterial qPCR data were log₁₀-transformed to attain normality. Data were analyzed as a 3×3 Latin square design using the GenStat (12th Edition) statistical package (VSN International Ltd, Oxford, UK). Fixed effects were experimental periods and treatments and the random effect was the cow. When significant treatment effects were detected, means were separated using Tukey test. Probability of $P < 0.05$ was used to determine significant differences among means.

Results

Ruminal fermentation parameters and ruminal bacteria quantification

Cows consumed all their individual allocation of TMR (10 kg DM per cow per day) with no feed refusals. Rumen digesta pH, NH₃-N and total VFA were similar for the three dietary treatments and averaged 6.9, 13.3 mg/dL and 57.5 mmol/L, respectively. Molar proportions (mol/100 mol) of individual VFA were comparable across dietary treatments and averaged 63.8 for acetate, 22.7 for propionate, 11.2 for butyrate and 2.4 for valerate (Table 3).

In this study, ruminal bacterial populations involved in lipid metabolism were quantified by qPCR (Table 2). The obtained PCR products were checked by expected size and sequenced. All the PCR products corresponded to the expected size: *Fibrobacter succinogenes* (500 bp), *Butyrivibrio proteoclasticus* (200 bp), *Prevotella bryantii* (550 bp) and *Anaerovibrio lipolytica* (600 bp) (not shown) and the sequences corresponded to the target organism. Compared with control, total bacteria (copies of 16S rDNA / ml) was decreased ($P < 0.05$) by SO and increased ($P < 0.05$) by HPO. The load of target bacteria (bacteria / ml) was similar for all dietary treatments and averaged: 4.52 for *Fibrobacter succinogenes*, 2.92 for *Butyrivibrio proteoclasticus* and 4.19 for *Anaerovibrio lipolytica* (Table 4). However, the load of *Prevotella bryantii* was increased ($P < 0.05$) by HPO but not by SO.

Fatty acid composition of ruminal digesta

Data from the FA composition of ruminal digesta is shown in Table 5. The most abundant FA in ruminal digesta regardless of dietary treatment were (g/100g):

C14:0 (4.4), C15:0 (5.7), C16:0 (36.5), C18:0 (21.7), C18:1 cis-9 (5.4) and C18:3 cis-9, 12, 15 (4.3). Compared with control and SO, HPO decreased ($P < 0.05$) C18:2 cis n-6 (1.28 and 1.64 vs. 0.75 g/100g) and total contents of monounsaturated (15.61 and 17.9 vs. 7.66 g/100g) and unsaturated (22.54 and 25.79 vs. 11.22 g/100g) FA in ruminal digesta. Also, compared with control and HPO, SO increased ($P < 0.05$) contents (g/100g) of C15:0 iso (1.92 and 1.04 vs. 2.76), C18:1 trans-11 (0.96 and 0.23 vs. 1.68) and C18:2 cis-9, trans-11 (1.42 and 0.42 vs. 1.65). Dietary treatments did not affect contents (g/100g) of the following FA: C10:0, C11:0, C12:0, C13:0, C14:0, C14:1, C15:0, C15:1 cis-10, C16:0, C16:0 iso, C16:1 trans-9 + C17:0 iso, C16:1 cis-9, C17:0, C17:1 cis-10, C18:0, C18:1 cis-9 and C18:3 cis-9, 12, 15.

Discussion

In this study, ruminal pH and $\text{NH}_3\text{-N}$ were not affected by dietary treatments, this partly agrees with studies (Yang *et al.*, 2009) who did not report ruminal pH changes when cows were fed soybean oil and linseed oil, but did observe increases in ruminal $\text{NH}_3\text{-N}$ concentration. Benchaar *et al.* (2012) reported no effect on pH, VFA and $\text{NH}_3\text{-N}$ when dairy cows were supplemented with linseed oil at 2, 3 and 4% DM. In the current study, lack of difference in ruminal fermentation parameters may be due to the amount (almost 3% of DM) of oil incorporated into the basal diet. Differences from other studies such as Yang *et al.* (2009), on the effect dietary oils on ruminal fermentation parameters in dairy cows may be explained by the amount of dietary oil and the forage source used, for example, VFA patterns were not affected when cows were supplemented with linseed oil

(3% DM) on a hay-base diet (Ueda et al., 2003) whereas on a corn silage-based diet they were changed (Doreau et al., 2009).

The chemical configuration of dietary lipids is associated with their effects on ruminal microorganisms. For example, PUFA are more toxic for biohydrogenating bacteria (e.g., *Butyrivibrio fibrisolvens*) than monoenoic FA (Lourenco et al., 2010). Consequently, SO which is a rich source of C18:2 cis n-6 is expected to have strong negative effects on ruminal bacterial populations; this agrees in part with the reduction of total bacteria (copies of 16S rDNA / ml) caused by SO treatment observed in this study. During rumen biohydrogenation, C18:2 cis n-6 yields several intermediate compounds until reduction to C18:0 (Castagnino et al., 2015). In the present study, C18:1 trans-11 and C18:2 cis-9, trans-11 (biohydrogenation intermediate isomers) were increased in rumen contents with SO compared to control and HPO. This is important for milk production because those FA can escape from the rumen and be secreted in milk as shown by Bu et al., (2007) who observed increases in the C18:1 trans-11 and C18:2 cis-9, trans-11 concentrations of milk fat when dairy cows were supplemented with vegetable oils and oilseeds rich in C18:2 cis n-6.

Increases of C15:0 iso provoked by SO are particularly interesting, since branched-chain FA have been suggested to reflect rumen function (e. g., ruminal fermentation pattern) and also contribute to the formation of the main odd- and branched-chain FA in milk (Vlaeminck et al., 2006). The odd- (C15:0 and C17:0) and branched (C13:0 iso, C14:0 iso, C15:0 iso, C16:0 iso, C17:0 iso, C18:0 iso,

C13:0 anteiso, C15:0 anteiso, C17:0 anteiso) chain fatty acids (OBCFA) profile of the rumen bacteria appears to be largely determined by the FA synthase activity of the microorganism rather than by the precursor availability (Vlaeminck et al., 2006). Consequently, variation in the OBCFA profile leaving the rumen is expected to mirror changes in the relative abundance of specific bacterial populations in the rumen rather than an altered bacterial FA synthesis. In this study, supplementation with SO may have influenced the FA synthase activity of ruminal microorganisms, specifically from *Prevotella* spp. and *Butyrivibrio fibrisolvens* (Fievez et al., 2012). It has been suggested that higher proportions of iso-fatty acids in solid associated bacteria reflect their enrichment in cellulolytic bacteria (e. g., *Butyrivibrio fibrisolvens*), whereas higher proportions of anteiso-C15:0 in liquid associated bacteria might indicate their enrichment in pectin and sugar fermenting bacteria (e. g., *Prevotella* spp.) (Bessa et al., 2009).

Normally in dairy cow diets, ruminal biohydrogenation of C18:2 cis n-6 varies between 70% and 95%, indicating that with the exception of diets containing marine lipids C18:0 is the major FA escaping from rumen (Shingfield *et al.*, 2013). In the present study this was corroborated by the FA profile of rumen digesta where C16:0 and C18:0 were the most predominant saturated FA (especially in HPO). Also, in the current study, HPO decreased ruminal C18:2 cis n-6, which may be explained by the levels of C18:2 cis n-6 in the HPO diet which was notably lower than control and SO.

It has been recognized that cellulolytic bacteria can be affected by dietary supplementation of lipid with high concentrations of PUFA (Paillard *et al.*, 2007). This is explained by factors such as disruption of microbial cell membranes and cell function caused by PUFA and lipid coating of feed particles (especially fibrous components) and bacteria (Yang *et al.*, 2009). The antimicrobial effect of lipids in the rumen is related to the cytotoxic effects of FA on membrane function of eukaryotic cells (Maia *et al.*, 2010). Long chain unsaturated FA appear to be more toxic to ruminal bacteria since they can attach to lipid bilayers in bacterial membranes (because of their hydrophobic and amphiphilic nature). The longer the chains, and the more double bonds, the easier it is for FA to attach and destroy membranes of bacteria (Zheng *et al.*, 2005).

Although, *Prevotella* spp. has been reported to be resistant to dietary PUFA (Huws *et al.*, 2010), in this study, *Prevotella bryantii* load was increased by HPO (a saturated source), which agrees in part with Choi *et al.* (2013) who reported that C16:0 and C18:0 have less antibacterial effect than PUFA (HPO diet contained 46 g of C16:0 and 36 g of C18:0 per 100g total FA). In concordance with that, it has been reported that consumption of animal fats (mainly saturated FA) has been associated with the presence of *Prevotella* and *Bacteroides* (Tremaroli and Bäckhed, 2012). Another possible explanation for increased *Prevotella* with the HPO diet may be the interaction of a saturated lipid source and a substrate (our basal diet comprised of 56% forage and 44% concentrate ratio). The *Prevotella* spp. are the dominant bacteria in the rumen (Stevenson and Weimer, 2007) and their ruminal populations vary according to different substrates, for example; on a

hay diet, *Prevotella ruminicola* is the predominant whereas on a grain diet *Prevotella bryantii* is the most numerous among these species (Tajima *et al.*, 2001). Our results are similar to those reported by Rico and Harvatine (2013) who fed dairy cows with a control diet composed by 60% forage and 40% concentrate and a low-fiber diet supplemented with 3 g/100g of SO, later, the authors (Rico *et al.*, 2015), studied the ruminal microbiome and found that the abundance of *Prevotella bryantii* was lowered in the control diet.

This study used qPCR to analyze bacterial population's concentrations using their DNA and did not use isolation from pure cultures. Compared to culture dependent studies, our results may be more precise because we avoided 'plate count anomaly' and the use of laborious protocols to isolate the target bacterial populations (Amann *et al.*, 1995). The primers used in this study were previously validated (Huws *et al.*, 2010; Tajima *et al.*, 2001). Furthermore, the primers were checked using the probe match tool in the Ribosomal Database Project (Huws *et al.*, 2007; Cole *et al.*, 2014). One interesting point is that the sum of the selected bacterial populations corresponded to a half of the total bacterial, and this observation was independent of the diet used. Therefore, more studies should be performed to obtain a clear picture of the changes on ruminal bacterial populations; a metagenomic approach could provide a deeper composition of ruminal populations.

Conclusions

In conclusion, supplementation with SO or HPO (2.7 % DM) did not affect ruminal fermentation parameters whereas HPO can increase loads of ruminal *Prevotella bryantii*. Also, results observed in our target bacteria may have depended on the degree of saturation of dietary oils.

Acknowledgments

This study was funded by a research grant from FONDECYT 11121142 and 1140734 (Fondo Nacional de Desarrollo Científico y Tecnológico, Chile). We would like to thank Juanita Clavijo, Jorge Manzor and Pamela Alvarez for technical assistance and Fundación Agro U.C. for the animal facilities and assistance in obtaining research data.

References

Amann RI, Ludwig W, Schleifer KH 1995. Phylogenetic identification and in situ detection of individual microbial cells without cultivation. Microbiological Research 59, 143–169.

Allred SL, Dhiman TR, Brennand CP, Khanal RC, McMahon DJ, Luchini ND 2006. Milk and cheese from cows fed calcium salts of palm and fish oil alone or in combination with soybean products. Journal of Dairy Science 89, 234–248.

Bal MA, Shaver RD, Jirovec AG, Shinnors KJ, Coors JG 2000. Crop processing and chop length of corn silage: Effects on intake, digestion, and milk production by dairy cows. Journal of Dairy Science 83, 1264–1273.

415

416

417 Benchaar C, Romero-Perez GA, Chouinard PY, Hassanat F, Eugene M, Petit HV,
418 Cortes C 2012. Supplementation of increasing amounts of linseed oil to dairy cows
419 fed total mixed rations: Effects on digestion, ruminal fermentation characteristics,
420 protozoal populations, and milk fatty acid composition. *Journal of Dairy Science* 95,
421 4578-4590.

422

423 Bligh EG, Dyer WJ 1959. A rapid method of total lipid extraction and purification.
424 *Canadian Journal of Biochemistry and Physiology* 37, 911-917.

425

426 Boeckaert C, Vlaeminck B, Fievez V, Maignien L, Dijkstra J, Boon N 2008.
427 Accumulation of trans C18:1 fatty acids in the rumen after dietary algal
428 supplementation is associated with changes in the *Butyrivibrio* community. *Applied*
429 *and Environmental Microbiology* 74, 6923–6930.

430

431 Bessa RJB, Maia MRG, Jeronimo E, Belo AT, Cabrita ARJ, Dewhurst RJ, Fonseca
432 AJM 2009. Using microbial fatty acids to improve understanding of the contribution
433 of solid associated bacteria to microbial mass in the rumen. *Animal Feed Science*
434 *and Technology* 150, 197–206.

435

436 Bu DP, Wang JQ, Dhiman TR, Liu SJ 2007. Effectiveness of oils rich in linoleic and
437 linolenic acids to enhance conjugated linoleic acid in milk from dairy cows. *Journal*
438 *of Dairy Science* 90, 998–1007.

439

440 Castagnino PS, Messana JD, Fiorentini G, De Jesus RB, Vito ES, Carvalho IP,
441 Berchielli TT 2015. Glycerol combined with oils did not limit biohydrogenation of
442 unsaturated fatty acid but reduced methane production in vitro. *Animal Feed*
443 *Science and Technology* 201, 14-24.

444

445 Choi, J. S., N. H. Park, S. Y. Hwang, J. H. Sohn, I. Kwak, K. K. Cho, and I. S. Choi.
446 2013. The antibacterial activity of various saturated and unsaturated fatty acids
447 against several oral pathogens. *Journal of Environmental Biology* 34, 673-676.

448

449 Chouinard PY, Corneau L, Saebo A, Bauman DE 1999. Milk yield and composition
450 during abomasal infusion of conjugated linoleic acids in dairy cows. *Journal of*
451 *Dairy Science* 82, 2737-2745.

452

453 Christie WW 1982. A simple procedure for rapid transmethylation of glycerolipids
454 and cholesteryl esters. *The Journal of Lipid Research* 23,1072-1075.

455

456 Cole JR, Wang Q, Fish JA, Chai B, McGarrell DM, Sun Y, Brown CT, Porras-Alfaro
457 A, Kuske CR, Tiedje JM 2014. Ribosomal Database Project: data and tools for high
458 throughput rRNA analysis. *Nucleic Acids Research* 42, D633–642.

459

460 Doreau M, Aurousseau E, Martin C 2009. Effects of linseed lipids fed as rolled
461 seeds, extruded seeds or oil on organic matter and crude protein digestion in cows.
462 *Animal Feed Science and Technology* 150, 187-196.

463

464 Fievez V, Colman E, Castro-Montoya JM, Stefanov I, Vlaeminck B 2012 Milk odd-
465 and branched-chain fatty acids as biomarkers of rumen function-An update. *Animal*
466 *Feed Science and Technology* 172, 51–65.

467

468 Huws SA, Edwards JE, Kim EJ, Scollan ND 2007. Specificity and sensitivity of
469 eubacterial primers utilized for molecular profiling of bacteria within complex
470 microbial ecosystems. *Journal of Microbiological Methods* 70, 565–569.

471

472 Huws S, Lee M, Muetzel S, Scott M, Wallace R, Scollan N 2010. Forage type and
473 fish oil cause shifts in rumen bacterial diversity. *FEMS Microbiology Ecology* 73,
474 396-407.

475

476 Huws SA, Kim EJ, Lee MRF, Scott MB, Tweed JKS, Pinloche E, Wallace RJ,
477 ,Scollan ND 2011. As yet uncultured bacteria phylogenetically classified as
478 *Prevotella*, *Lachnospiraceae* incertae sedis and unclassified *Bacteroidales*,
479 *Clostridiales* and *Ruminococcaceae* may play a predominant role in ruminal
480 biohydrogenation. *Environmental Microbiology* 13, 1500-1512.

481

482 Kargar S, Ghorbani GR, Alikhani M, Khorvash M, Rashidi L, Schingoethe DJ 2012.
483 Lactational performance and milk fatty acid profile of Holstein cows in response to
484 dietary fat supplements and forage: Concentrate ratio. *Livestock Science* 150, 274-
485 283.

486

487 Llorente B, Bravo-Almonacid F, Cvitanich C, Orlowska E, Torres HN, Flawiá MM,
488 Alonso GD 2010. A quantitative real-time PCR method for in planta monitoring of
489 *Phytophthora infestans* growth. *Letters in Applied Microbiology* 51, 603–610.
490

491 Lourenco M, Ramos-Morales E, Wallace R 2010. The role of microbes in rumen
492 lipolysis and biohydrogenation and their manipulation. *Animal* 4,1008-1023.
493

494 Maia M., Chaudhary L, Bestwick C, Richardson A, McKain N, Larson T, Graham I,
495 Wallace R 2010. Toxicity of unsaturated fatty acids to the biohydrogenating ruminal
496 bacterium, *Butyrivibrio fibrisolvens*. *BMC Microbiology* 10, 1-10.
497

498 NRC. Nutrient Requirements of Dairy Cattle. 7th rev. ed. Natl. Acad. Sci.,
499 Washington, DC. 2001.
500

501 Paillard D, McKain N, Chaudhary LC, Walker ND, Pizette F, Koppova I, McEwan
502 NR, Kopecký J, Vercoe PE, Louis P, Wallace RJ 2007. Relation between
503 phylogenetic position, lipid metabolism and butyrate production by different
504 *Butyrivibrio*-like bacteria from the rumen. *Antonie Van Leeuwenhoek* 91,417-422.
505

506 Potu R, AbuGhazaleh A, Hastings D, Jones K, Ibrahim S 2011. The effect of lipid
507 supplements on ruminal bacteria in continuous culture fermenters varies with the
508 fatty acid composition. *Journal of Microbiology* 49, 216-223.
509

Prive F, Kaderbhai NN, Girdwood S, Worgan HJ, Pinloche E, Scollan ND, Huws SA, Newbold CJ 2013. Identification and Characterization of Three Novel Lipases Belonging to Families II and V from *Anaerovibrio lipolyticus* 5ST. Plos One 8, e69076.

Prive F, Newbold CJ, Kaderbhai NN, Girdwood SG, Golyshina OV, Golyshin PN, Scollan ND, Huws SA 2015. Isolation and characterization of novel lipases/esterases from a bovine rumen metagenome. Applied Microbiology and Biotechnology 99, 5475-5485.

Rico DE, Harvatine KJ 2013. Induction of and recovery from milk fat depression occurs progressively in dairy cows switched between diets that differ in fiber and oil concentration. Journal of Dairy Science 96, 6621-6630.

Rico DE, Preston SH, Risser JM, Harvatine KJ 2015. Rapid changes in key ruminal microbial populations during the induction of and recovery from diet-induced milk fat depression in dairy cows. British Journal of Nutrition 114, 358-367.

Schroeder GF, Gagliostro GA, Becu-Villalobos D, Lacau-Mengido I 2002. Supplementation with partially hydrogenated oil in grazing dairy cows in early lactation. Journal of Dairy Science 85, 580-594.

Shingfield KJ, Bonnet M, Scollan ND 2013. Recent developments in altering the fatty acid composition of ruminant-derived foods. Animal 7, 132-162.

534

535 Stevenson DM, Weimer PJ 2007. Dominance of *Prevotella* and low abundance of
536 classical ruminal bacterial species in the bovine rumen revealed by relative
537 quantification real-time PCR. *Applied Microbiology and Biotechnology* 75,165-174.

538

539 Tajima K, Aminov R, Nagamine T, Matsui H, Nakamura M, Benno Y 2001. Diet-
540 dependent shifts in the bacterial population of the rumen revealed with real-time
541 PCR. *Applied and Environmental Microbiology* 67, 2766-2774.

542

543 Tremaroli V, Backhed F 2012. Functional interactions between the gut microbiota
544 and host metabolism. *Nature* 489,242-249.

545

546 Ueda K, Ferlay A, Chabrot J, Loo JJ, Chilliard Y, Doreau M 2003. Effect of linseed
547 oil supplementation on ruminal digestion in dairy cows fed diets with different
548 forage: Concentrate ratios. *Journal of Dairy Science* 86, 3999-4007.

549

550 Vargas-Bello-Perez E, Fehrmann-Cartes K, Iniguez-Gonzalez G, Toro-Mujica P,
551 Garnsworthy PC 2015a. Short communication: Chemical composition, fatty acid
552 composition, and sensory characteristics of Chanco cheese from dairy cows
553 supplemented with soybean and hydrogenated vegetable oils. *Journal of Dairy*
554 *Science* 98, 111-117.

555

556 Vargas-Bello-Perez E, Iniguez-Gonzalez G, Fehrmann-Cartes K, Toro-Mujica P,
557 Garnsworthy PC 2015b. Influence of fish oil alone or in combination with

558 hydrogenated palm oil on sensory characteristics and fatty acid composition of
559 bovine cheese. *Animal Feed Science and Technology* 205, 60-68.

560

561 Vlaeminck B, Fievez V, Cabrita ARJ, Fonseca AJM, Dewhurst R 2006. Factors
562 affecting odd- and branched-chain fatty acids in milk: a review. *Animal Feed*
563 *Science and Technology* 131, 389–417.

564

565 Wallace RJ, Chaudhary LC, McKain N, McEwan NR, Richardson AJ, Vercoe PE,
566 Walker ND, Paillard D 2006. *Clostridium proteoclasticum*: a ruminal bacterium that
567 forms stearic acid from linoleic acid. *FEMS Microbiology Letters* 265,195-201.

568

569 Yang S, Bu D, Wang J, Hu Z, Li D, Wei H, Zhou L, Loo J 2009. Soybean oil and
570 linseed oil supplementation affect profiles of ruminal microorganisms in dairy cows.
571 *Animal* 3, 1562-1569.

572

573 Zhang CM, Guo YQ, Yuan ZP, Wu YM, Wang JK, Liu JX, Zhu WY 2008. Effect of
574 octadeca carbon fatty acids on microbial fermentation, methanogenesis and
575 microbial flora in vitro. *Animal Feed Science and Technology* 146, 259–269.

576

577 Zheng HC, Liu JX, Yao JH, Yuan Q, Ye HW, Ye JA, Wu YM 2005. Effects of
578 dietary sources of vegetable oils on performance of high-yielding lactating cows
579 and conjugated linoleic acids in milk. *Journal of Dairy Science* 88, 2037-2042.

580

581

582

Table 1 *Ingredients and chemical composition of control, soybean oil (SO), and hydrogenated palm oil (HPO) diets*

	Diet ²		
	Control	SO	HPO
	(% of DM)		
Ingredient composition			
Alfalfa hay	17	17	17
Corn silage	18	18	18
High-moisture corn	10	10	10
Soybean hulls	34	34	34
Wheat bran	19	19	19
Vitamin and mineral premix ¹	2	2	2
Soybean oil	0	2.7	0
Hydrogenated palm oil	0	0	2.7
Chemical composition, %			
DM	53.6	53.6	53.6
CP	16.6	16.6	16.6
Ether extract	2.3	5.1	6.3
NDF	39.2	39.2	39.2
ADF	21.0	21.0	21.0
Lignin	3.6	3.6	3.6
Ash	6.0	6.0	6.0
Fatty acid composition, g/100g of FA			
C4:0	0.03	0.09	0.73
C6:0	0.05	0.04	0.01
C8:0	0.03	0.03	0.07
C10:0	1.63	0.15	0.10
C12:0	0.16	0.13	2.08
C14:0	0.26	0.15	1.70
C16:0	15.6	13.7	45.9
C18:0	18.7	18.8	36.3
C18:1 cis-9	0.42	1.78	0.04
C18:2 cis n-6	46.9	49.5	5.03
C18:3 cis-6, 9, 12	0.17	0.10	0.19
C18:3 cis-9, 12, 15	7.44	6.38	0.55

¹Contained per kg: 25, 000 mg of P; 80,000 mg of Ca; 25,000 mg of Mg; 1,612 mg of S; 300,000 IU of vitamin A; 50,000 IU of vitamin D₃ and 1,600 IU of vitamin E.

²Control = basal diet / no fat supplement; SO = basal diet + 170 g/d/ cow of SO; HPO = basal diet + 170 g/d/cow of HPO.

591 **Table 2** PCR primers and template DNA for detection of ruminal bacteria

Target bacterium		Primer ¹	Primer concentration (μM)	Purified template of DNA (ng)	Product size (bp)
<i>Fibrobacter succinogenes</i>	Forward	GGTATGGGATGAGCTTGC	20	30	500
	Reverse	GCCTGCCCCTGAACTATC			
<i>Butyrivibrio proteoclasticus</i>	Forward	TCCGGTGGTATGAGATGGGC	20	30	200
	Reverse	GTCGCTGCATCAGAGTTTCCT			
<i>Prevotella bryantii</i>	Forward	ACTGCAGCGCGAACTGTCAGA	20	26	550
	Reverse	ACCTTACGGTGGCAGTGTCTC			
<i>Anaerovibrio lipolytica</i>	Forward	TGGGTGTTAGAAATGGATTC	20	28	600
	Reverse	CTCTCCTGCACTCAAGAATT			

592 ¹*Fibrobacter succinogenes*, *Prevotella bryantii*, and *Anaerovibrio lipolytica* primers were described by Tajima et al (2001)
593 whereas *Butyrivibrio proteoclasticus* primers were described by Huws et al. (2010).
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608

Table 3 *Ruminal pH, NH₃-N and VFA from cows fed control, soybean oil (SO), and hydrogenated palm oil (HPO)*

	Diet ¹			SED	P-value
	Control	SO	HPO		
pH	6.90	6.88	6.89	0.10	0.78
NH ₃ -N (mg/dL)	13.6	11.9	14.3	1.22	0.32
Total VFA (mmol/L)	50.5	59.6	62.5	4.33	0.32
Molar proportion (mol/100 mol)					
Acetate	63.9	63.9	63.7	0.96	0.97
Propionate	22.5	22.8	22.8	1.30	0.99
Butyrate	11.2	11.2	11.3	0.30	0.97
Valerate	2.4	2.4	2.3	0.20	0.94

¹ Control = basal diet / no fat supplement; SO = basal diet + 170 g/d/ cow of SO; HPO = basal diet + 170 g/d/cow of HPO.

Table 4 Quantification of ruminal bacteria by PCR from cows fed control, soybean oil (SO), and hydrogenated palm oil (HPO)

	Diet ¹			SED	P-value
	Control	SO	HPO		
Total bacteria (copies 16S rDNA / ml) ²	11.84 ^b	11.75 ^c	12.06 ^a	0.08	<0.01
Target bacterium (bacteria/ml) ^{2, 3}					
<i>Fibrobacter succinogenes</i>	4.96	4.25	4.36	0.32	0.26
<i>Butyrivibrio proteoclasticus</i>	3.04	2.74	2.99	0.37	0.73
<i>Prevotella bryantii</i>	3.41 ^b	3.51 ^b	3.90 ^a	0.08	0.04
<i>Anaerovibrio lipolytica</i>	4.10	4.20	4.28	0.23	0.76

¹ Control = basal diet / no fat supplement; SO = basal diet + 170 g/d/ cow of SO; HPO = basal diet + 170 g/d/cow of HPO.

² Log₁₀

³ Based on ribosomal operon copy number

Means in the same row with different superscripts (a, b, c) are different (P<0.05)

Table 5 *Fatty acid composition of ruminal digesta from cows fed control, soybean oil (SO), and hydrogenated palm oil (HPO)*

Fatty acid (g/100g of fatty acid)	Diet ¹			SED	P-value
	Control	SO	HPO		
C10:0	2.35	1.77	1.63	0.90	0.71
C11:0	0.30	0.39	0.29	0.24	0.92
C12:0	0.57	0.63	0.21	0.17	0.23
C13:0	2.54	1.30	2.00	0.84	0.44
C14:0	4.10	5.47	3.69	1.53	0.53
C14:1	2.81	2.43	1.73	1.05	0.61
C15:0	7.10	6.17	3.94	2.21	0.42
C15:1 cis-10	3.41	2.21	1.13	0.79	0.10
C15:0 iso	1.92 ^b	2.76 ^a	1.04 ^c	0.43	0.04
C16:0	35.06	32.84	41.62	3.89	0.17
C16:0 iso	2.40	1.35	0.28	2.23	0.52
C16:1 trans-9 + C17:0 iso	0.93	0.63	0.46	0.26	0.28
C16:1 cis-9	0.66	0.52	0.39	0.10	0.24
C17:0	1.59	1.74	0.79	0.48	0.21
C17:1 cis-10	0.61	0.78	0.69	0.30	0.85
C18:0	17.00	16.42	31.61	6.24	0.11
C18:1 trans-11	0.96 ^b	1.68 ^a	0.23 ^c	0.06	<0.01
C18:1 cis-9	5.65	8.23	2.30	2.88	0.23
C18:2 cis n-6	1.28 ^a	1.64 ^a	0.75 ^b	0.38	<0.01
C18:2 cis-9, trans-11	1.42 ^b	1.65 ^a	0.42 ^c	0.06	0.03
C18:3 cis-9, 12, 15	4.81	5.41	2.53	1.50	0.24
Σ Saturated fatty acids	70.44	66.54	85.64	7.03	0.07
Σ Monounsaturated fatty acids	15.61 ^a	17.90 ^a	7.66 ^b	3.26	0.04
Σ Polyunsaturated fatty acids	6.92	7.88	3.56	1.87	0.16
Σ Unsaturated fatty acids	22.54 ^a	25.79 ^a	11.22 ^b	4.63	0.04

¹ Control = basal diet / no fat supplement; SO = basal diet + 170 g/d/ cow of SO; HPO = basal diet + 170 g/d/cow of HPO.

Means in the same row with different superscripts (a, b, c) are different (P<0.05)