

Introduction

Medications prescribed, dispensed and held in a patient's home in anticipation of need are becoming increasingly used in the UK (Wilcock, 2011). These 'just in case' or 'anticipatory' medications can be prescribed in advance to manage common symptoms at the end of life including pain, nausea and vomiting, anxiety and excessive secretions. They are usually prescribed by a GP, dispensed and then held in the patient's home or nursing home for use if and when needed. Having immediate access to necessary medications in the home is regarded as a key means of improving symptom management and enabling better end of life care in the community (Carney, 2011, Lawton et al., 2012, Faull et al., 2013, National Institute for Health and Care Excellence, 2015).

Anticipatory prescribing is seen as enabling professional staff such as community and district nurses to respond immediately to patients' needs (Carney, 2011 p13). Yet little is known about nurses' roles and experiences in anticipatory prescribing (Wilson and Seymour, 2013, Eisenhauer et al., 2007) and it is cited as a recommended area of research in the new NICE guideline on caring for the dying (National Institute for Health and Care Excellence, 2015). A synthesis of the existing related research suggests that nurses may be challenged by a lack of resources, limited knowledge and experience with the relevant medications and power differentials between themselves and medical practitioners (Wilson and Seymour, 2013). Although local audits have been carried out to assess the implementation of anticipatory prescribing (Carney, 2011, Carney and MacRobbie, 2008, Scott-Aiton, 2009, Amass and Allen, 2005, Lawton et al., 2012), there are no larger scale studies in existence.

Insights from a qualitative study by Faull et al. (2013) suggest that anticipatory prescribing is complex and that health care staff have a range of interpretations of what is meant by prescribing in advance. The authors conclude that anticipatory prescribing is a process rather than a one-off event and highlight the need to build relationships between professionals to ensure good communication between teams and across organisational boundaries, including out-of-hours care.

This paper reports the findings of a survey that elicited the views of 575 community/district, nursing home, and palliative care nurses from two UK regions: the East Midlands and Lancaster/South Cumbria. The aim of the survey was to gain insight into the roles and experiences of a wide range of community nurses in end of life medication decisions. This paper draws on both quantitative and qualitative aspects of the survey to give an overview of the key findings.

Methods

The survey was part of a three phase mixed methods study to examine community nurses' experiences of working with anticipatory prescriptions. Phase one was a literature review of 26 studies from 10 countries (Wilson and Seymour, 2013). Phase two was a qualitative ethnographic study using interviews and observations (Wilson et al., 2015, Wilson et al., 2014). The survey was the third phase of the study, informed by the data gathered in the literature review and ethnography. The survey questionnaire was developed by the research team supported by an advisory panel with expertise in community and end of life care nursing. A face validity exercise was undertaken with a group of district nurses from the East Midlands. The nurses completed the survey and gave feedback in a focus group on the wording, style, order and appropriateness of the questions. In light of their comments, adjustments were made to the questionnaire.

Survey sample

The survey was conducted in two regions: East Midlands and Lancaster/South Cumbria. The Binley's database of NHS staff was used to select 500 district and community nurses at random in each area. In addition all specialist palliative care nurses, Macmillan and Marie Curie nurses¹ and managers and matrons of nursing homes were also selected in the two areas. It was recognised that the database would not be completely up to date and search terms might identify some nurses who were not working in the fields required. In order to minimise the number of questionnaires sent to

¹ Grouped and referred to a 'Palliative care nurses'

inappropriate people, the lists were all examined by hand by the study team. During this process job and institutional titles were inspected to exclude individuals who were not likely to be involved in caring for adults who had died at home or in a nursing home. For example, we excluded all those located in hospitals or with job titles that would not have a primary focus on palliative care for adults.

Data collection and management

From February to May 2013 a postal questionnaire and two reminders were sent to 1739 individuals. 181 mailings were returned as the individual no longer worked at the institution concerned, leaving a total valid sample of 1558. As questionnaires were returned the data were logged in a database managed through the programme Statistical Package for the Social Sciences (SPSS)®.

Data analysis

Survey data were analysed to produce a series of descriptive statistics supported by the SPSS software. Responses were compared across the three categories of nurses (nursing home nurses, community/district nurses and palliative care nurses). The free text sections were analysed using a thematic analysis (Braun and Clarke, 2006) to allocate codes to the comments made in order to identify patterns. These codes were then grouped and distilled to generate themes. Themes were then compared with the quantitative findings and are used in this paper to add depth. As not all respondents answered all questions, total response rates for individual questions are reported and vary throughout the paper.

Ethical approval

We obtained ethical approval from the National Research Ethics Service (11/EM/0213) and governance approval from the NHS Trusts at each research site.

Findings

The overall response rate was 37% (n=575/1558). While nursing home nurses (49.6%; n=231/466) and palliative care nurses (47.4%; n=151/318) responded well, a lower response from

community/district nurses (24.9%; n=193/774) reduced the overall response rate. Comparing the two regions, the East Midlands had a slightly better response rate (39.7%; n=380/957) compared to Lancaster/South Cumbria (32.4%; n=195/601). Respondents were predominantly female (95%; n=540/569), had worked as a nurse for 20 years or more (64%; n=366/572) and were likely to have some kind of additional training in palliative (70%; n=404/575) and/or end of life care (78%; n=440/575). Respondents reported attending a range of courses and training therefore it was not possible to compare the level or quality of the training reported. From the written comments those with additional qualifications appeared to value this training vis à vis the anticipatory prescribing process. Some of those with a prescribing qualification acknowledged that they could take on the prescription writing role in the process:

Community/district nurses know more about end of life drugs and care than GPs.

Consequently the decision to prescribe generally is dictated by the nurse and the GP will take their advice. Hopefully all nurses will hold NISP (V300) [Nurse Independent and Supplementary Prescribers] qualification soon and prescribe for patients. (184: District nurse team leader)

I feel that as a NISP [Nurse Independent and Supplementary Prescribers] I am extremely confident in prescribing for palliative care patients, specifically in end of life care. (1204: Palliative care nurse)

Caring for patients at the end of life

Respondents were asked how many patients they cared for at the end of life in the past year. Overall 83.7% (n=481/575) reported providing care to at least one person who had died in the past year. Of these 481 nurses, 471 (98.1%) reported that an 'an anticipatory prescription for drugs aimed at relieving symptoms and distress' was in place for at least one patient. 98.3% (n=463/471) of respondents reporting this said that, in at least one case, these prescriptions were used. Respondents reporting that in no cases were they used (just 8 respondents) worked in care homes or in 'other' roles.

Characteristics of patients where an anticipatory prescription was used

Having answered some general questions about anticipatory prescriptions, respondents were then asked to recall details of one specific patient for whom an anticipatory prescription had been used and to answer a series of further questions about that particular case. Table 1 gives a breakdown of the characteristics of these individual patient cases. Where patients' age category was reported (n=412), 63.8% (n=263) were said to be aged 70 or over. The majority of the patients for whom the use of an anticipatory prescription was recalled were reported by nurses as dying in their own home or care home (96.4%; n=432/448). A primary cause of death was provided for 434 patient cases and in 79.3% (n=344/434) of these, cancer was reported by nurses as the registered cause of death. Fewer respondents from nursing homes reported that cancer was a cause of death among the patients whom they recalled as receiving anticipatory medications.

Table 1 - Characteristics of patient cases reported by nurses to have received anticipatory prescription medications

Characteristic		Number of patient cases as reported by nurse respondents (%)
Gender	Male	194/442 (43.9%)
	Female	248/442 (56.1%)
Age	Under 50	32/412 (7.7%)
	50-69	117/412 (28.4%)
	70-89	263/412 (63.9%)
Place of death	Hospital	4/448 (0.9%)
	Hospice	12/448 (2.7%)
	Nursing/care home	198/448 (44.2%)
	Own home	234/448 (52.2%)
Cause of death	Cancer	344/434 (79.3%)
	Heart disease	27/434 (6.2%)
	Stroke	12/434 (2.8%)
	Other	51/434 (11.8%)

The writing and dispensing of the medications

In general, nurses reported working well with GPs and perceived that they had good access to the medications needed. Figure 1 shows that 79.2% (n=427/539) of nurses reported that they 'infrequently or never' found doctors reluctant to prescribe anticipatory medication and 11.1% (n=60/539) reported this as 'neither frequent nor infrequent'. However, it was evident there was

some variation. A small proportion, 9.6% (n=52/539), agreed that some doctors were reluctant to provide anticipatory prescriptions (see Figure 1) and this point was reflected in the qualitative comments:

I work with different GP practices. All very individual, some more knowledgeable than others. Some happy to listen to nursing home staff and families and take our opinions on board. I have to get assistance from palliative care nurse at times to help me get GP to prescribe. (1273: Matron in a nursing home)

The difficulty I found is that not all practices in my location are happy to prescribe anticipatory medications as they feel this is a waste on their budget and feel that out of hours doctors are there to be called upon for the prescribing of anticipatory drugs. In the main the reluctance of GPs in prescribing is improving (1636: Manager in a nursing home)

Similarly, a few nurses (8.6%; n=45/525) said that they 'always or frequently' experienced significant difficulties in obtaining the drugs specified in anticipatory prescriptions used in end of life care. For a further 11.2% (n=59/525) this was reported as 'neither frequent nor infrequent'. However the majority of nurses (80.2%; n=421/525) said they encountered these difficulties 'infrequently or never' (see figure 2).




Figure 1 - Prescription writing (539 responses)




Figure 2 – Dispensing (525 responses)

Across the three categories of nurses slightly less than one fifth (18.4%; n=97/527) agreed with the statement 'anticipatory prescriptions are incorrectly written up by doctors'. However, when looking at the categories of respondents this was reported more frequently by palliative care nurses (24.3%; n=28/115) and community/district nurses (25.7%; n=45/175) than by nursing home nurses (9.7% n=19/195). Comments about these issues were also added in the written section of the survey:

I have real problems with GPs prescribing [end of life] drugs, they are usually unable to estimate doses correctly, have no idea how to prescribe when a patient has a Fentanyl patch or on a slow release morphine. ...This weekend on one day 2 prescriptions were written incorrectly. (1019: Palliative care nurse)

Anticipatory medications used

Where respondents reported on the care of one specific patient who had been in receipt of anticipatory medications, Midazolam was the drug most commonly reported to have been used in the last month of life of this specific patient. This applied to 81.9% (n=367/448) of the patients where a use of an anticipatory prescription was recalled by nurses. This was followed by Levomepromazine (32.5%; n=143/440) and then Haloperidol (23.9%; n=104/435). Figure 3 shows the percentage of drugs used by category of respondent; nurses in all three categories reported similar patterns of usage.




Figure 3 - Drugs used by respondent category

Managing the use of anticipatory medications

Having the confidence to make decisions relating to the use of drugs specified in anticipatory prescriptions was considered to be important by 92.0% (n=483/525) of nurses responding to the survey. For example, when reporting the use of Midazolam in specific patient cases, 82.2% (n=287/349) of the responding nurses felt they had the responsibility of the assessing that patients' response to the drug. 34.1% (n=119/349) of nurses responding to this question about a specific patient care felt this responsibility lay with the GP, and 3.2% (n=11/349) with the specialist palliative care doctor (for this question respondents had the opportunity to select more than one option).

Table 2 shows that for these patient cases medications were either 'not increased' or just 'increased gradually' over the last three days of life, with a 'strong increase' on the last day being rare.

Table 2 – Reported changes in dosages for individual patient cases in the last three days of life

	Midazolam	Haloperidol	Levomopromazine
No increase	51.1% (n=165)	54.0% (n=47)	55.2% (n=69)
Gradual increase	42.1% (n=136)	41.4% (n=36)	40.0% (n=50)
Strong increase last day	6.8% (n=22)	4.6% (n=4)	4.8% (n=6)
TOTAL (=100%)*	323	87	125

*Patients may have received more than one of these drugs

Nurses reported that the anticipatory medications successfully controlled those symptoms they were intended to relieve in 89.6% of the patient cases they recalled. In a more general sense, most

respondents (96.0%; n=504/525) agreed that when anticipatory prescriptions were in place they 'always or frequently' enabled them to improve the quality of end of life care they provided:

It helps to provide better quality of care for the patients, especially out of hours. (482: District nurse)

Anticipatory prescribing is essential- whilst medications are not always needed it gives peace of mind to the resident, relatives and staff that, should symptoms develop, medication is at hand to deal with them. Proactive decision making is essential in this area. (1473: Matron in a nursing home)

Discussion

This paper has reported on one aspect of a wider study to explore the nurses' roles in the use of anticipatory prescriptions. A survey was used to elicit views from 575 community/district, nursing home, and palliative care nurses across two regions in England. Initially nurses were asked to report their views about anticipatory prescribing in general; they were then asked to refer to a specific patient whom they recalled had received anticipatory medications and to answer a series of further questions about that patient. The survey shows that nurses report the presence of anticipatory prescriptions as very common indeed. Moreover, when in place these prescriptions were normally used. In line with Carney's (2011) audit of the use of 'just in case' boxes in Scotland, the majority of nurses felt that having anticipatory medications in place enabled them to provide better care and could improve the quality of death.

Our findings indicate that nurses have a number of pivotal and complex roles throughout the 'process' of anticipatory prescribing (Faull et al., 2013). The survey findings support those of the qualitative work undertaken in the earlier part of the study and reported elsewhere (Wilson et al., 2015), and was designed to indicate the prevalence of the issues raised by nurses.

Our findings give some insight into the types of medication used for symptom management and the pattern of their use. Of the 441 specific cases where nurses recalled that a patient had received anticipatory medications, Midazolam was reported as the most commonly used drug, with Levomepromazine and then Haloperidol used less often. These are often considered to be three of the 'core' drugs commonly used in anticipatory prescriptions (Wilcock, 2011). For the individual cases reported it was rare for nurses to recall any sudden increase in dosage of the medications used in the last three days of life; any increases that were recalled were considered by respondents as 'gradual'.

Nurses responding to the survey reported a particular responsibility for monitoring the results of giving the medications, with GPs and other doctors were less often involved in this. This echoed narratives from nurses in the ethnographic element in phase two of the study, recounting that decisions about the use of anticipatory medications were primarily regarded to be the responsibility of the health care team, and often that of the nurse (Wilson et al., 2015).

Although for the most part nurses reported that relationships with doctors were working well, a small proportion of respondents reported difficulties, again echoing findings from the ethnography in phase two of the study (Wilson et al., 2015). Some encountered reluctance in some doctors to provide anticipatory prescriptions. In addition a small number of respondents stated the effectiveness of the process could be challenged by prescriptions that were incorrectly written up by doctors, or by difficulties in obtaining the drugs. These findings resonate with observations by Faull et al. (2013) that there is potential for the process to stall at any stage. Recent guidance from NICE (National Institute for Health and Care Excellence, 2015) suggests that when considering the need for anticipatory medications clinicians should not only weigh up benefits and harms of prescribing in advance, but also recognise the possibility of sudden deterioration and time it might take to obtain

medications. Hence the recommendation is that prescriptions be put in place as early as possible, yet remain under review (National Institute for Health and Care Excellence, 2015).

A small number of qualitative comments by responding nurses indicated that they valued being a nurse prescriber and that this had enhanced their confidence and capacity to provide good care. Webb and Gibson's (2011) audit of nurse independent prescribers supports this, concluding that prescribing by nurses is effective, timely and appropriate, especially during out-of-hours periods.

Limitations

Since 181/1739 (10.4%) of the questionnaires sent to people listed by Binley's were returned as the person had moved on since the list was compiled, it seems likely that the response was somewhat skewed towards nurses with more experience, as this statistic suggests a time lag before Binley's picks up a name and adds it to their database. If true, there would tend to be an under-representation of people recently starting in posts, and this group may be more likely to contain nurses who had not worked in the role before. A high proportion of respondents with twenty years or more experience may also indicate that the questionnaire was passed on to nurses with more experience to complete and/or that more experienced nurses were more likely to deal with patients with anticipatory prescriptions in place.

The survey was sent to nurses at a time when end of life care was under a great deal of scrutiny due to a media furore and the subsequent repeal of the Liverpool Care Pathway (<https://www.gov.uk/government/publications/review-of-liverpool-care-pathway-for-dying-patients>); this may have led to some nurses being reluctant to share their experiences of using anticipatory medications at the end of life, resulting in a lower response rate, particularly from community/district nurses.

It is important for the reader to note that we did not have direct access to patient data. Nurses were asked to report on one specific case with which they had been involved, we therefore have no knowledge of how representative this case may have been of that respondent's caseload, or how accurate was their reporting of medication use.

Conclusion

The survey was able to gain insight into the roles and experiences of a substantial number, and range of nurses working with end of life medications in the community. Survey responses show that anticipatory prescriptions are very common, and once in place are often used. The comments in the survey show that nurses value the provision of anticipatory prescriptions and in general believe them to improve the care they provide for their patients. Although for the most part the relationships with doctors were reported by the nurse respondents as working well, a small proportion recorded difficulties when some doctors were reluctant to provide prescriptions, when prescriptions were incorrectly written up by doctors, or in obtaining drugs. This suggests pharmacists, nurses and GPs need to work together to establish the most appropriate ways of facilitating this process. In qualitative comments several nurses indicated that they valued being a nurse prescriber and that this enhanced their confidence and capacity to provide good care. Increasing the number of Nurse Independent and Supplementary Prescribers (NISP) may help to support the process of anticipatory prescribing.

References

- Amass, C. & Allen, M. 2005. How a 'just in care' approach can improve out-of-hours palliative care. *Pharmaceutical Journal*, 275, 22-23.
- Carney, J. 2011. *Analysis of 'just in care' anticipatory audit forms 1st December 2010-28th February 2011*. Inverness, Palliative Care Network and NHS Highland (report).
- Carney, J. & MacRobbie, A. 2008. *Analysis of 'Just in Case' anticipatory prescribing pilot*. Inverness, Palliative Care Network and NHS Highland (report).
- Eisenhauer, L. A., Hurley, A. C. & Dolan, N. 2007. Nurses' Reported Thinking During Medication Administration. *Journal of Nursing Scholarship*, 39, 82-87.

- Faull, C., Windridge, K., Ockleford, E. & Hudson, M. 2013. Anticipatory prescribing in terminal care at home: what challenges do community health professionals encounter? *BMJ Supportive and Palliative Care*, 3, 91-97.
- Lawton, S., Denholm, M., Macaulay, L., Grant, E. & Davie, A. 2012. Timely symptom management at end of life using 'just in case' boxes. *British Journal of Community Nursing*, 17, 182-183.
- National Institute for Health and Care Excellence 2015. *Caring for the Dying Adult: NICE guideline*. London: NICE.
- Scott-Aiton, A. 2009. Creating a tool to improve anticipatory prescribing in palliative care. *Pharmaceutical Journal*, 283, 283-284.
- Webb, W. A. & Gibson, V. 2011. Evaluating the impact of nurse independent prescribing in a weekend clinical nurse specialist service. *International Journal of Palliative Nursing*, 17, 537-543.
- Wilcock, A. 2011. Pre-emptive prescribing in the community. In: Twycross, R. & Wilcock, A. (eds.) *Palliative Care Formulary (PCF4)*. 4th Edition ed. <http://www.palliativedrugs.com/SamplePDF/PCF4/Chapter17.pdf>: Palliativedrugs.com.
- Wilson, E., Morbey, H., Brown, J., Payne, S., Seale, C. & Seymour, J. 2015. Administering anticipatory medications in end-of-life care: A qualitative study of nursing practice in the community and in nursing homes. *Palliative Medicine*, 29, 60-70.
- Wilson, E. & Seymour, J. 2013. Understanding the role of nurses in the management of symptoms and distress in the last days of life. In: Sterckx, S., Raus, K. & Mortimer, F. (eds.) *Continuous sedation at the end of life*. Cambridge: Cambridge University Press.
- Wilson, E., Seymour, J., Morbey, H., Payne, S., Brown, J. & Seale, C. 2014. Anticipatory prescribing in end-of-life care. *Nursing Times*, 110, 16-17.

Braun, V. & Clarke, V. 2006. Using thematic analysis in psychology. *Qualitative Research in Psychology*, 3, 77-101.