



RESEARCH PAPER

Natural variation in *Arabidopsis thaliana* reveals shoot ionome, biomass, and gene expression changes as biomarkers for zinc deficiency tolerance

Ana Carolina A.L. Campos^{1,2,*}, Willem Kruijer³, Ross Alexander^{1,4}, Robert C. Akkers¹, John Danku^{2,*}, David E. Salt^{2,*} and Mark G.M. Aarts^{1,†}

¹ Laboratory of Genetics, Wageningen University, Droevendaalsesteeg 1, 6708 PB Wageningen, The Netherlands

² Institute of Biological and Environmental Sciences, University of Aberdeen, Cruickshank Building, Aberdeen AB24 3UU, UK

³ Biometris, Wageningen University and Research, PO Box 100, 6700AC Wageningen, The Netherlands

⁴ School of Life Sciences, John Muir Building, Heriot Watt University, Edinburgh EH14 4AS, UK

* Present address: School of Biosciences, University of Nottingham, Sutton Bonington Campus, Loughborough, Leicestershire LE12 5RD, UK

† Correspondence: mark.aarts@wur.nl

Received 16 November 2016; Editorial decision 17 May 2017; Accepted 20 May 2017

Editor: Christine Raines, University of Essex

Abstract

Zinc (Zn) is an essential nutrient for plants, with a crucial role as a cofactor for many enzymes. Approximately one-third of the global arable land area is Zn deficient, leading to reduced crop yield and quality. To improve crop tolerance to Zn deficiency, it is important to understand the mechanisms plants have adopted to tolerate suboptimal Zn supply. In this study, physiological and molecular aspects of traits related to Zn deficiency tolerance were examined in a panel of 19 *Arabidopsis thaliana* accessions. Accessions showed a larger variation for shoot biomass than for Zn concentration, indicating that they have different requirements for their minimal Zn concentration required for growth. Accessions with a higher tolerance to Zn deficiency showed an increased expression of the Zn deficiency-responsive genes *ZIP4* and *IRT3* in comparison with Zn deficiency-sensitive accessions. Changes in the shoot ionome, as a result of the Zn treatment of the plants, were used to build a multinomial logistic regression model able to distinguish plants regarding their Zn nutritional status. This set of biomarkers, reflecting the *A. thaliana* response to Zn deficiency and Zn deficiency tolerance, can be useful for future studies aiming to improve the performance and Zn status of crop plants grown under suboptimal Zn concentrations.

Key words: Biofortification, biomarker, mineral concentration, plant ionome, shoot growth, zinc usage index.

Introduction

Zinc (Zn) is an essential micronutrient required for plant growth and development. Many agricultural soils in the Middle East, India, and parts of Australia, America, and Central Asia are Zn deficient, often due to poor Zn availability caused by high pH in calcareous soils. Zn-deficient soils affect both crop yield and quality, and can also result in

human malnutrition through the intake of food containing low concentrations of Zn and other micronutrients (Cakmak, 2007; Alloway, 2009). The World Health Organization (WHO) and the Food and Agriculture Organization (FAO) of the United Nations estimate that about one-third of the world's population suffers from some form of Zn deficiency

(Allen *et al.*, 2006). Since plants are often the main source of dietary Zn, improving plant Zn concentration and tolerance to Zn deficiency is an important goal in fighting this so-called 'hidden hunger' (www.harvestplus.org).

Plants exposed to Zn deficiency show reduced growth. Severe deficiency results in extensive leaf chlorosis, wilting, stunting, leaf curling, and reduced root elongation, while mild stress results in chlorosis in young leaves and early senescence of older leaves (Marschner, 1995). In *Arabidopsis thaliana*, all of these symptoms, as well as delayed flowering, are observed when plants are grown under Zn deficiency (Talukdar and Aarts, 2007). Zn deficiency also affects the function of enzymes such as copper/zinc superoxide dismutase (Cu/Zn SOD) and carbonic anhydrase (CA), leading to an accumulation of reactive oxygen species (ROS), which cause oxidative damage and a reduction in photosynthesis (Clemens, 2010; Ibarra-Laclette *et al.*, 2013).

The threshold Zn concentration below which plants are considered to be Zn deficient is ~15–20 $\mu\text{g g}^{-1}$ dry biomass. This can vary from species to species and between plants of the same species (Marschner, 1995; White and Broadley, 2011). The ability of a plant to grow and yield under Zn-limiting conditions compared with ideal growth conditions is defined as Zn efficiency. It is based on the difference in relative growth or yield between plants grown under control and Zn-deficient conditions. Another parameter used is the Zn usage index (ZnUI), which quantifies the amount of dry matter produced per mg of Zn in the plant. The ZnUI is useful for the comparison of plant genotypes which do not show significant differences in Zn concentration, but differ in biomass production under Zn deficiency (Siddiqi and Glass, 1981; Cakmak *et al.*, 1998; Good *et al.*, 2004; Genc *et al.*, 2006).

To avoid problems associated with inappropriate Zn supply, plants have developed an efficient homeostasis mechanism. Different genes act in the uptake of Zn from soil, distribution over different organs, tissues, cells, and organelles, and (re) mobilization through the plant, to control Zn homeostasis (Sinclair and Krämer, 2012). While the actual Zn deficiency sensor is not yet known, the Zn deficiency response in *A. thaliana* seems to start with the activation of the transcription factors bZIP19 and bZIP23, the function of which is essential for plants to survive Zn deficiency (van de Mortel *et al.*, 2006; Assunção *et al.*, 2013;).

Zn is among the essential elements which make up the plant ionome (Salt *et al.*, 2008). Previous studies have shown that the plant ionome profile reflects the physiological state of plants under various genetic, developmental, and environmental backgrounds, and can be used as a biomarker for a particular physiological condition (Huang and Salt, 2016). Ionome-based biomarker models have been used to determine differences in the plant nutritional status among large sets of different genotypes and experimental batches (Baxter *et al.*, 2008a). Natural variation for the concentration of elements comprising the plant ionome has been studied in *A. thaliana*, revealing important mineral homeostasis mechanisms in plants (Rus *et al.*, 2006; Loudet *et al.*, 2007; Baxter *et al.*, 2008a, 2010; Kobayashi *et al.*, 2008; Morrissey *et al.*, 2009; Chao *et al.*, 2012; Pineau *et al.*, 2012; Koprivova *et al.*, 2013).

To improve efficiently the performance of crops grown under suboptimal Zn conditions and increase the Zn content in their edible parts, it is of paramount importance to understand the physiological and molecular mechanisms underlying plants' tolerance to Zn deficiency. Aspects of natural variation for Zn deficiency tolerance have been described for several plant species, including *A. thaliana* (Graham *et al.*, 1992; Rengel and Graham, 1996; Cakmak *et al.*, 1998; Hacisalihoglu *et al.*, 2004; Genc *et al.*, 2006; Ghandilyan *et al.*, 2012; Karim *et al.*, 2012). However, to date, a detailed study on natural variation of plants' tolerance to Zn deficiency involving both physiological and molecular mechanisms has not been performed. In this study, we evaluated natural variation among 19 diverse *A. thaliana* accessions to identify physiological and molecular traits involved in the tolerance to Zn deficiency. It shows that high-throughput screening of genetic variation for Zn deficiency tolerance can be simplified by focusing on the combination of changes in the ionome profile, the minimum Zn concentration required for growth, and the expression level of Zn deficiency-responsive genes.

Materials and methods

Plant material and hydroponic growth

A set of 19 *A. thaliana* accessions was selected based on their diverse site of origin (Supplementary Table S1 at *JXB* online). Seeds were surface-sterilized with chlorine vapour and sown in Petri dishes on wet filter paper followed by a 4 d stratification treatment at 4 °C in the dark, to promote uniform germination. Seeds were transplanted to 0.5% (w/v) agar-filled tubes, the bottoms of which were cut off, and placed in a modified half-strength Hoagland nutrient solution for hydroponic growth (Assunção *et al.*, 2003): 3 mM KNO₃, 2 mM Ca(NO₃)₂, 1 mM NH₄H₂PO₄, 0.5 mM MgSO₄, 1 μM KCl, 25 μM H₃BO₃, 2 μM MnSO₄, 0.1 μM CuSO₄, 0.1 μM (NH₄)₆Mo₇O₂₄, and 20 μM Fe(Na)EDTA. The solution pH was set at 5.5 using KOH and buffered with 2 mM MES. Plants were grown hydroponically in two experiments performed separately. In experiment one, referred to as the mild Zn deficiency experiment, we compared plants grown for 41 d under control (2 μM ZnSO₄) and mild Zn deficiency (0.05 μM ZnSO₄). In experiment two, referred to as the severe Zn deficiency experiment, we compared plants grown for 31 d under control (2 μM ZnSO₄) and severe Zn deficiency (no Zn added). Plants were grown in a climate-controlled chamber set at 70% relative humidity, with a 12 h day (120 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$) and 20 °C/15 °C day/night temperatures. The hydroponic system consisted of plastic trays (46 × 31 × 8 cm) holding 9 litres of nutrient solution, covered with a non-translucent 5 mm thick plastic lid with evenly spaced holes in a 7 × 10 format holding the agar-filled tubes with plantlets. The nutrient solution was refreshed once a week. Shoot fresh weight (SFW) was measured in all samples during harvesting. Some samples were immediately frozen in liquid nitrogen and stored at –80 °C for gene expression and element concentration analysis. The shoot dry weight (SDW) of these samples was calculated based on a fresh weight/dry weight correction factor obtained from additional plants which were dried for 72 h at 60 °C. For each trait, the treatment versus control relative values were determined as $\text{Rel_trait} = (\text{trait Zn deficiency} / \text{trait control}) \times 100$. The ZnUI was calculated based on the following formula:

$$\text{ZnUI} = \left(\frac{\text{shoot biomass (mg)}}{\text{shoot Zn concentration (ppm)}} \right)$$

Mineral elemental analysis

For each treatment, the shoot ionome profile was determined for five biological replicates of each *A. thaliana* accession. Samples were first dried for 72 h at 60 °C, transferred to 96-well plates with tubes containing one 5 mm glass bead, and homogenized for 5 min at 30 Hz with a Qiagen 96-well plate mixer mill. A 2–4 mg aliquot of leaf tissue was transferred to Pyrex test tubes (16 × 100 mm) and digested with 0.9 ml of concentrated nitric acid (Baker Instra-Analyzed; Avantor Performance Materials; <http://www.avantormaterials.com>) for 5 h at 115 °C. Samples were diluted to 10 ml with 18.2 MΩcm Milli-Q water. Elemental analyses were performed with inductively coupled plasma mass spectrometry, ICP-MS (Elan DRC II; PerkinElmer, <http://www.perkinelmer.com>) for Li, B, Na, Mg, P, S, K, Ca, Mn, Fe, Co, Ni, Cu, Zn, As, Se, Rb, Sr, Mo, and Cd. A reference, composed of pooled samples of digested leaf material, was prepared and included every ninth sample in all sets of 70 samples to correct for variation between and within ICP-MS analysis runs. Seven samples from each sample set were weighed and used during the iterative weight normalization process to estimate the weight of the remaining 63 samples from the set (Danku *et al.*, 2013). The following elements were not added to the nutrient solution: Li, Co, Ni, As, Se, Rb, Sr, and Cd and, except for Cd, their concentrations are not shown.

Gene expression

Gene expression analysis was performed for eight accessions with different ZnUI values selected from the 19 accessions grown under mild Zn deficiency conditions. Frozen leaf material from plants grown under mild and severe Zn deficiency and their respective control treatments was used, in three biological replicates, each consisting of material from three plants. Total RNA was extracted using the method of Onate-Sanchez and Vicente-Carbajosa (2008). cDNA was synthesized from 1 µg of total RNA using the iScript cDNA synthesis kit from BioRad as per the manufacturer's instructions. Following synthesis, cDNA was diluted 10-fold. Quantitative real-time PCRs (qRT-PCRs) were performed in triplicate with iQ SYBR Green Supermix (BioRad) using an iQ Real Time PCR machine (BioRad). Relative transcript levels of selected genes were determined by qRT-PCR. The expression of *IRT3* (At1g60960), *ZIP3* (At2g32270), *ZIP4* (At1g10970), *bZIP19* (At4g35040), *CSD2* (At2g28190), and *CA2* (At5g14740) was measured. The oligonucleotides used for each gene are shown in Supplementary Table S2. Amplicon lengths were between 80 bp and 120 bp, and all primer combinations had at least 95% efficiency. Reaction volumes were 10 µl (5 µl of SYBR green qPCR mix, 300 nmol of each primer, and 4 µl of cDNA template). Cycling parameters were 4 min at 95 °C, then 40 cycles of 15 s at 95 °C and 30 s at 55 °C. Gene expression values were normalized to the reference genes *PEX4* (At5g25760), *SAND* (At2g28390), and *18S*. Gene expression levels relative to the average of the reference genes for each accession under mild and severe Zn deficiency and their respective control treatments were calculated based on Δ CT values. Gene expression levels of accessions exposed to mild and severe Zn deficiency, relative to their respective control treatment, were calculated based on $\Delta\Delta$ CT values (Livak and Schmittgen, 2001).

Statistical analysis

For all shoot traits and gene expression levels relative to reference gene expression, a two-way ANOVA was performed to test for significant differences between treatments, accessions, and the interaction between treatments and accessions. A one-way ANOVA was performed to test for significant differences between accessions for relative gene expression values, relative change in SDW, Zn concentration, and Zn content. A one-way ANOVA was also performed to test for significant differences in element concentrations between the four treatments used (mild and severe Zn deficiency and their respective controls). Element concentration values were log₁₀-transformed

and a Benjamini–Hochberg correction of the *P*-values was performed. When significant differences were found, a Tukey's HSD post-hoc test with a significance level of 0.05 was performed. Broad-sense heritability (H_2) was calculated as the ratio between estimated genetic variance and total phenotypic variance (Kruijer *et al.*, 2015).

Multivariate analysis and classification

To predict the Zn deficiency nutritional status of accessions based on their ionic profile, various multinomial logistic regression (MLR) models were used, similar to the model described by Baxter *et al.* (2008b). In all cases, 11 elements (B, Mg, P, S, K, Ca, Mn, Fe, Cu, Zn, and Mo) were considered, the concentrations of which were reliably measured. At first, element concentrations were log₁₀-transformed and the transformed element concentration values in the severe or mild Zn-deficient plants were normalized to their respective control treatment by subtracting the means of the control group. Thereafter, plants from the control treatment of the two experiments were considered to have the same 'control' status. Hence, plants can be in either a control, mild, or severe Zn deficiency state. These states have different probabilities, which were modelled as a linear function of the element concentrations. The prediction for the state of a new plant was defined as the state with the highest probability. Finally, the prediction performance of the following MLR models were compared: (i) univariate MLR models, for each element; (ii) a multivariate MLR model, including all elements; and (iii) a multivariate MLR model with all elements except Zn. The multivariate models included a LASSO penalty, which is a multiple of the absolute values of the regression coefficients. The level of penalization was chosen by 10-fold cross-validation. The prediction performance of all models was assessed by drawing 100 times a training set of 199 plants from the total of 398 plants, while the remaining 199 plants were used as a validation set. Each training set was drawn in a stratified manner, respecting the number of plants in the Zn sufficiency (2 × 100), mild (99), and severe Zn deficiency treatment (99) categories. A penalized logistic regression model was fit for each training set using the R package 'glmnet' (Friedman *et al.*, 2010), and used to predict the status of the 199 plants in the validation set. Prediction performance was estimated by averaging the proportion of correctly classified plants over the 100 validation sets.

Results

Natural variation in Zn deficiency response for physiological and morphological traits

Arabidopsis thaliana accessions were grown hydroponically under control conditions (2 µM ZnSO₄) and either mild (0.05 µM ZnSO₄) or severe Zn deficiency (no Zn added). After 31 d of exposure to severe Zn deficiency, plants showed clear deficiency symptoms compared with plants in the control treatment. This was primarily visible as reduced growth, leaf curling, and the presence of chlorotic and necrotic spots (Fig. 1A, B). After 31 d in the mild Zn deficiency treatment, accessions did not show any sign of Zn deficiency, hence they were grown for an additional 10 d. Even then, only a few accessions had visible symptoms of Zn deficiency, mainly slight chlorosis in leaves and reduced growth (Fig. 1C, D), confirming that the treatment was indeed mild.

Accessions showed significant phenotypic variation for most traits analysed which varied according to the trait and Zn treatment (Supplementary Tables S3, S5). Plants in the severe Zn deficiency treatment had shoot Zn concentrations




Fig. 1. Comparison of *A. thaliana* accessions grown under control and severe or mild Zn-deficient conditions. Representative examples of *A. thaliana* accessions grown in hydroponic medium under Zn-sufficient control conditions ($2 \mu\text{M ZnSO}_4$) (A and C) or severe (no Zn added) (B) and mild Zn deficiency ($0.05 \mu\text{M ZnSO}_4$) (D). Plants in (A) and (B) are grown for 31 d; plants in (C) and (D) are grown for 41 d. Accessions from left to right in rows from top to bottom: C24, Per-1, Tsu-0, Mc-0, Hau-0, Mt-0, Shah, Kas-2, Bor-4, Wag-3, Ors-1, Pa-2, Li-5:2, Ge-0, Can-0, Var 2-1, Ler-1, Cvi-0, Bur-0, and Col-0. Scale bars indicate 2 cm.

close to the reported minimum required for growth, which is $\sim 15\text{--}20 \mu\text{g g}^{-1}$ dry biomass (Marschner, 1995). Shoot Zn concentrations under mild Zn deficiency were approximately twice as high than under severe Zn deficiency (Fig. 2). In addition, plants in the mild Zn deficiency experiment had a higher SDW than plants in the severe Zn deficiency experiment, as they were grown for 10 d longer. From all shoot traits, only Zn concentration was significantly correlated between the controls of the two Zn deficiency experiments, indicating that during the additional 10 d of growth between experiments, other factors such as the growth rate of accessions affected their shoot biomass and Zn content in a different manner (Supplementary Table S8). Accession Cvi-0 had to be excluded from further analysis as it had established poorly and too many plants were lost, especially from the mild Zn deficiency experiment.

In both Zn deficiency treatments, most accessions showed reduced SDW relative to their respective control treatments, while few had a higher SDW and apparently were not affected by the reduced Zn supply (Fig. 3A, B). Nearly all accessions had a reduction in shoot Zn concentration of $\sim 80\%$ in both Zn deficiency treatments relative to their respective controls (Fig. 3C, D). Also, accessions with high shoot Zn concentrations were not always among

the ones with a high shoot total Zn content, due to differences in SDW. Tsu-0, Col-0, and Mt-0 were the best performing accessions under mild Zn deficiency in terms of having similar Zn concentrations to the other accessions and higher SDW across the Zn deficiency and control treatments. Thus, these accessions seem to be able to maintain growth under Zn deficiency, albeit with some reduction in shoot Zn concentration. Conversely, Pa-2, C24, and Li-5:2 performed poorly under mild Zn deficiency, with a strong reduction in growth in comparison with the other accessions though with a small reduction in shoot Zn concentrations in both Zn deficiency treatments. These accessions appear to have a poor ability to take up Zn under both control and Zn-deficient conditions, which results in a limited capacity to grow and to maintain cellular Zn levels. Only accession Bor-4 showed an increase in SDW under severe Zn deficiency relative to its control treatment, even though this was not statistically different from most of the other accessions (Fig. 3A; Supplementary Table S8). Bor-4 also showed an increase in SDW under mild Zn deficiency, as did Shah. However, it is important to note that these two accessions were among the ones with the lowest SDW in their respective control treatments, which could explain their lower sensitivity to Zn deficiency.




Fig. 2. Relationships between shoot dry weight (SDW) and Zn concentration of 19 *A. thaliana* accessions grown under Zn deficiency. SDW is expressed in mg and Zn concentration in $\mu\text{g g}^{-1}$ DW. See Supplementary Table S1 for the list of accessions. Data for plants grown under severe Zn deficiency (no Zn added; A) or mild Zn deficiency ($0.05 \mu\text{M ZnSO}_4$; B) are indicated with grey dots, and plants grown under their respective control conditions ($2 \mu\text{M ZnSO}_4$) with black dots. Plants used for (A) grew for 31 d and plants used for (B) grew for 41 d.

Accessions with contrasting tolerance to Zn deficiency show differences in the expression of Zn deficiency-responsive genes

The ZnUI was used to determine the amount of biomass produced per unit of tissue Zn concentration (Fig. 4). In accordance with the results previously shown for SDW and Zn concentration, the accessions Mt-0 and Tsu-0 had the highest ZnUI values for both Zn deficiency treatments, and C24 and Pa-2 had the lowest values. Even though only in the mild Zn deficiency treatment, these accessions had significantly higher or lower ZnUI values when compared with the other accessions (Supplementary Table S6). Eight accessions with different ZnUI values in the mild Zn deficiency treatment were then selected to examine whether natural variation for Zn deficiency tolerance is reflected at the gene expression level. Mild Zn deficiency was favoured over the severe treatment as the variation between accessions for SDW was larger in the mild treatment. In addition, mild Zn deficiencies are more likely to be found in nature. The accessions Tsu-0 and Col-0 had high ZnIU values, accessions Ge-0, Bur-0, and Can-0 were intermediate, and Pa-2, C24, and Per-1 had low ZnUI values. Accessions with higher ZnUI values were considered to be more tolerant to Zn deficiency (Fig. 4).

The expression of six genes involved in the plant Zn deficiency and oxidative stress response was determined in Zn deficiency-tolerant and -sensitive accessions (Fig. 5; Supplementary Fig. S1). *bZIP19* encodes one of the two redundant bZIP transcription factors which control the Zn deficiency response in *A. thaliana*. We also looked at the expression of the *IRT3*, *ZIP4*, and *ZIP3* transcriptional targets genes of bZIP19, all encoding ZIP-like Zn transporter proteins, strongly induced following Zn deficiency (Assunção *et al.*, 2010). The expression of the *CSD2* gene, encoding a Cu/Zn SOD which needs Zn as a structural component to function (Sharma *et al.*, 2004), and the *CA2* gene, encoding a

carbonic anhydrase (CA) requiring Zn as cofactor, was also determined. *CSD2* is needed for detoxification of superoxide radicals, while *CA2* facilitates the diffusion of CO_2 through the liquid phase of the cell to the chloroplast, important for photosynthesis (Randall and Bouma, 1973; Li *et al.*, 2013). Both *CSD2* and *CA2* are expected to decrease in expression under Zn deficiency exposure due to the reduced concentration of Zn in the cells (Ibarra-Laclette *et al.*, 2013).

There was a significant effect of both the mild and severe Zn deficiency treatments on the expression level of most studied genes. The exceptions were *bZIP19* and *CA2* in the severe Zn deficiency treatment (Supplementary Table S3). The Zn deficiency-responsive genes *IRT3*, *ZIP4*, and *ZIP3* were up-regulated in all accessions under both Zn deficiency treatments, confirming that the plants sensed Zn deficiency (Fig. 5; Supplementary Fig. S1). *ZIP4* and *IRT3* in particular were in general more highly expressed in the more Zn deficiency-tolerant accessions than in the more Zn deficiency-sensitive accessions, with Tsu-0 especially showing strong induction of these genes under severe Zn deficiency. The expression of *ZIP3*, which is predominantly expressed in roots (van de Mortel *et al.*, 2006), is the least prominent of the three Zn transporter genes in shoots. The expression levels of *CSD2* and *CA2* were generally low and variable in both Zn deficiency treatments, but these genes are down-regulated especially under mild Zn deficiency. The Zn deficiency-tolerant accessions Ge-0 and Bur-0 had the highest induction of the *CA2* and *CSD2* genes under severe Zn deficiency (Fig. 5). Significant accession by treatment interaction was found for all genes tested, except for *bZIP19*, in at least one of the Zn deficiency experiments (Supplementary Table S3), indicating that gene expression differences between the response of accessions to Zn deficiency are pronounced.

To understand further the relationship between the expression levels of Zn deficiency-responsive genes and




Fig. 3. Relative changes in shoot dry weight and Zn concentration of 19 *A. thaliana* accessions grown under severe (A and C) and mild (B and D) Zn deficiency, compared with their respective control treatments. Relative changes are expressed as percentages of the control (%). One-way ANOVA of these data and pairwise comparisons between accessions are provided in Supplementary Tables S5 and S6. See Supplementary Table S1 for the list of accessions. Plants were grown in hydroponic medium under Zn-sufficient control conditions ($2 \mu\text{M ZnSO}_4$) and severe (no Zn added) or mild Zn deficiency ($0.05 \mu\text{M ZnSO}_4$). Plants in (A) and (C) are grown for 31 d and plants in (B) and (D) are grown for 41 d.

Zn deficiency tolerance traits, a correlation analysis was performed. Under severe Zn deficiency, we found a significant positive correlation between the expression levels of *IRT3* and *CSD2* with ZnUI and of *ZIP4* with SFW (Supplementary Table S9).

Zn deficiency affects the shoot ionomic profile of A. thaliana accessions

The shoot ionome of the 19 *A. thaliana* accessions was then determined. Box plots of the combined results per element showed a substantial variation between treatments for almost all the elements measured (Fig. 6; Supplementary Table S7). Significant differences between treatments were observed for Zn, Mg, Mo, Cu, and Cd concentrations in both the mild and severe Zn deficiency experiments. B, Na, and Ca concentrations were significantly different between treatments only in the mild Zn deficiency experiment, and Mn and Fe concentrations only in the severe Zn deficiency experiment. When

comparing Zn concentrations across the four treatments, there was a significant difference between severe and mild deficiency but not between their respective control treatments.

H^2 values were calculated to estimate the genetic contribution to the observed phenotypic variation (Table 1). H^2 values were generally higher in the mild compared with the severe Zn deficiency experiment and in plants exposed to Zn deficiency in comparison with their control treatments. The heritability for ZnUI was highest in the mild Zn deficiency treatment, suggesting that under those conditions a large part of the observed variation is due to genetic differences between accessions. Fe concentration had the lowest heritability in both control treatments, whereas Mo concentration had the highest heritability across the treatments. Even though the Zn concentrations of plants grown under severe Zn deficiency were very low, there was substantial heritability for both Zn concentration and Zn content, with values of 0.49 and 0.41, respectively, indicating that the minimal Zn concentration/content levels are subject to genetic variation.




Fig. 4. Shoot Zn usage index (ZnUI) of *A. thaliana* accessions grown in severe (A) and mild (C) Zn deficiency and their respective control treatments (B and D). The letters above each bar indicate if the accession was already bolting (B) or flowering (F) when harvested. The ZnUI is defined as shoot biomass (in mg)/shoot Zn concentration (in ppm). Plants were grown in hydroponic medium under Zn-sufficient control conditions ($2 \mu\text{M ZnSO}_4$) and severe (no Zn added) or mild Zn deficiency ($0.05 \mu\text{M ZnSO}_4$). Plants in (A) and (C) are grown for 31 d and plants in (B) and (D) are grown for 41 d. One-way ANOVA of these data and pairwise comparisons between accessions are provided in Supplementary Tables S3 and S4.

Classification of the plant Zn deficiency state using multinomial logistic regression

The univariate model (i.e. with a single element as the only predictor) performed poorly as a predictor of plant nutritional status, for most elements, and often mistakenly identified plants under Zn deficiency as being control (Table 2). As expected, only the Zn concentration was able to separate the three classes very well, with prediction accuracies ranging from 0.92 for the plants under severe Zn deficiency to 0.99 for the control plants. Cu also had a good prediction performance for severe Zn deficiency, while Ca was the only element (apart from Zn) that identified a substantial number of the plants under mild Zn deficiency (Table 2). Mg, Mn, Fe, and Mo performed only marginally well, having some ability to identify plants under severe and mild Zn deficiency. For the other elements (B, P, S, and K), the univariate model performed no better than a naïve classifier that would always predict control conditions.

The penalized multivariate model, fitted on all elements except Zn, performed much better than the univariate model: the predicted accuracy for mild (0.6596) and severe Zn deficiency (0.7750) was far higher than with any element alone (except Zn), and the accuracy for the control treatments (0.8738) was still very good. When this model was fitted on

all elements (including Zn), it performed similarly to the univariate model fitted with Zn alone, the latter having a higher accuracy for the controls and mild Zn deficiency treatments and less for the severe Zn-deficient plants (Table 2).

Discussion

The natural variation in the response of *A. thaliana* to two levels of Zn deficiency was examined, with a focus on physiological and molecular traits. Analysis of genetic variation indicated that: (i) accessions vary for the minimum Zn requirement for growth; (ii) tolerance to Zn deficiency seems to be related to an increased expression of genes encoding Zn transmembrane transporter proteins (*ZIP4* and *IRT3*); and (iii) Zn deficiency results in changes in the plant ionome which can be used as a biomarker to predict the plant's physiological condition.

Natural variation of growth and Zn concentration in response to severe and mild Zn deficiencies

The tested *A. thaliana* accessions showed substantial diversity for all traits studied in both Zn deficiency experiments (Figs 1–3). Extreme accessions were identified for all traits, confirming the existing large natural variation in the *A.*




Fig. 5. Normalized gene expression levels of *bZIP19*, *IRT3*, *ZIP3*, *ZIP4*, *CSD2*, and *CA2* in rosette leaves of eight *A. thaliana* accessions under Zn deficiency (Zn⁻) and control treatments (Zn⁺ control) in the severe (left) and mild Zn deficiency experiments (right). Accessions are ranked from left to right according to decreasing Zn usage index values under mild Zn deficiency (see Fig. 4). Plants were grown in hydroponic medium under Zn-sufficient control conditions (2 μ M ZnSO₄) and severe (no Zn added) or mild Zn deficiency (0.05 μ M ZnSO₄), for 31 d or 41 d, respectively. Error bars represent the SE; one-way ANOVA and pairwise comparisons between accessions are provided in Supplementary Tables S3 and S4.





Fig. 6. Box plots comparing mineral element concentrations in shoots of 19 *A. thaliana* accessions grown under severe and mild Zn deficiency and their respective control treatments. Plants were grown in hydroponic medium under Zn-sufficient control conditions ($2 \mu\text{M ZnSO}_4$) and severe (no Zn added) or mild Zn deficiency ($0.05 \mu\text{M ZnSO}_4$). Plants in the severe Zn deficiency condition were grown for 31 d and plants in the mild Zn deficiency condition were grown for 41 d. For each concentration, the box represents the interquartile range (IQR), the bisecting line represents the median, the whiskers indicate 1.5 times the IQR, and the open circles indicate outlier points. Lower case letters denote statistically different groups when comparing the four treatments using a one-way ANOVA with groupings by Tukey's HSD test with a significance level of $P \leq 0.05$. The results of this ANOVA are shown in Supplementary Table S7.

thaliana response to Zn deficiency conditions and endorsing this panel of representative accessions as a valuable resource to study the plant response and tolerance to Zn deficiency. The response of *A. thaliana* to Zn deficiency has previously been examined in the Ler \times Cvi recombinant inbred line (RIL) population, in which a large variation in SDW and Zn concentration was observed (Ghandilyan *et al.*, 2012).

The mild Zn deficiency treatment is more suitable to reveal genetic variation underlying a plant's response to Zn nutrition, with higher heritability for most traits in comparison with the other treatments (Table 1). The disadvantage of using this mild treatment was that plants were 10 d older than in the severe Zn deficiency treatment, resulting in the initiation of flowering in some accessions. Such a change in development could include remobilization of minerals

Fig. 6. *Continued*

from older to younger organs (e.g. from rosette leaves to developing fruits); however, Waters and Grusak (2008) previously showed that the contribution of remobilization is probably <10% of the seed mineral content, so we considered this not much of a disturbing factor. In addition, this treatment seems better in representing Zn-deficient conditions likely to be encountered by *A. thaliana* in nature, with an average Zn concentration in leaves of 26 ppm in comparison with 18 ppm in the severe Zn deficiency treatment. To support this, Ghandilyan *et al.* (2012) observed leaf average Zn concentration of 40 ppm when using a Zn-deficient and nutrient-poor soil originating from Eskisehir, Central Anatolia in Turkey to grow the *A. thaliana* Ler×Cvi RIL population. Furthermore, the harshness of the severe Zn deficiency treatment seems to be beyond the genetic capacity of most accessions to tolerate, based on the extensive chlorosis displayed by nearly all accessions in this treatment and their very low average leaf Zn concentration, which was within or below the minimum Zn concentration range of 15–20 ppm required for growth as suggested by Marschner (1995).

Heritabilities of most traits were higher in the severe and mild Zn deficiency treatments than in their respective controls, further supporting the observed large genetic variation for all traits in response to the Zn deficiency treatments. Contrary to these observations, Ghandilyan *et al.* (2012) reported lower heritability values for shoot biomass and

most element concentrations in *A. thaliana* plants grown in Zn-deficient soil compared with control conditions. Yet, other studies show that heritabilities for the same trait can change according to the growth conditions (Ghandilyan *et al.*, 2009; Richard *et al.*, 2011; Baxter *et al.*, 2012), hence the importance of taking heritability into account when selecting growth conditions most amenable to detect genetic variation for a specific trait.

The control treatments of the two Zn deficiency experiments were significantly correlated with respect to the Zn concentration, but not for SDW and Zn content (Supplementary Table S8). This is probably due to differences in growth rate between the *A. thaliana* accessions during the additional 10 d of growth in the mild Zn deficiency experiment. Previous studies have shown that growth rate is highly variable among plants, being affected by both internal and external factors such as developmental processes and environmental conditions (El-Lithy *et al.*, 2004; Zhang *et al.*, 2012). Differences in growth rate between accessions in the mild and severe Zn deficiency experiments are probably caused by differences in the initiation of flowering. Most accessions in the control treatment of the mild Zn deficiency experiment were flowering or bolting at the harvesting day; while only three accessions, of the control treatment, were flowering in the severe Zn deficiency experiment at the harvesting day, which was 10 d earlier than in the mild Zn deficiency experiment (Fig. 1A, C).

Table 1. Broad-sense heritability (H^2) values for the traits measured in *A. thaliana* accessions grown under severe and mild Zn deficiency and their respective Zn sufficiency (control) conditions

Traits	Mild		Severe	
	Control	Zn deficiency	Control	Zn deficiency
SFW	0.44	0.62	0.41	0.66
SDW	0.68	0.78	0.40	0.48
ZnUI	0.65	0.81	0.40	0.57
SZnC	0.60	0.62	0.50	0.41
[Zn]	0.63	0.65	0.60	0.49
[Mn]	0.68	0.69	0.60	0.64
[Fe]	0.36	0.53	0.32	0.83
[Cu]	0.50	0.75	0.59	0.38
[Mo]	0.91	0.97	0.86	0.75
[Cd]	0.59	0.73	0.49	0.76
[B]	0.67	0.51	0.63	0.78
[Na]	0.48	0.37	0.55	0.60
[Mg]	0.59	0.71	0.55	0.46
[P]	0.62	0.71	0.44	0.72
[S]	0.45	0.59	0.53	0.58
[K]	0.51	0.65	0.46	0.48
[Ca]	0.72	0.69	0.42	0.52

SFW, shoot fresh weight (g); SDW, shoot dry weight (mg); ZnUI, Zn usage index; SZnC, shoot total Zn content (μg); and [X], mineral element concentrations ($\mu\text{g g}^{-1}$ DW).

Table 2. Estimated prediction performance values for elements used in the logistic regression model to predict plant nutritional Zn status

Univariate models	Control	Zn deficiency		Average
		Severe	Mild	
B	0.914	0.002	0.0837	0.4804
Mg	0.901	0.364	0	0.5442
P	1	0	0	0.5025
S	0.993	0	0.0061	0.5005
K	0.991	0	0.0102	0.5005
Ca	0.949	0	0.2673	0.5427
Mn	0.882	0.412	0	0.5467
Fe	0.977	0.278	0	0.5608
Cu	0.877	0.716	0.0286	0.6276
Zn	0.996	0.92	0.9857	0.9744
Mo	0.911	0.204	0	0.509
<i>Multivariate models</i>				
All elements except Zn	0.8738	0.7750	0.6596	0.7962
All elements	0.9921	0.9332	0.9549	0.9681

Physiological and molecular mechanisms of Zn deficiency tolerance in *A. thaliana*

Arabidopsis thaliana accessions showed a larger variation for relative change in SDW than in Zn concentration under both Zn deficiency treatments (Fig. 3). This indicates the presence of genetic variation for their minimum Zn requirement and for the ability to tolerate low Zn concentrations. This is not

unique for *A. thaliana* though. Also for barley, bread and durum wheat, common bean, and rice, different genotypes are reported to have similar shoot Zn concentrations with different levels of Zn deficiency tolerance (Cakmak *et al.*, 1998; Rengel, 2001; Genc *et al.*, 2002; Hacısalihoglu *et al.*, 2003; Wissuwa *et al.*, 2006; Sadeghzadeh *et al.*, 2009). Further indications that *A. thaliana* accessions vary for the minimum Zn requirement is shown by a few accessions with slightly higher SDW in the Zn-deficient treatment relative to its control.

The ability to enhance the root Zn uptake and the root to shoot Zn transport are among the proposed mechanisms underlying tolerance to Zn deficiency (Rengel, 2001; Broadley *et al.*, 2007), but the Zn deficiency signal may come from shoots. Indeed, accessions considered tolerant to Zn deficiency had a higher expression of the Zn deficiency-responsive genes *ZIP4* and *IRT3* in shoots (Fig. 5). These genes, encoding Zn transmembrane transporters (Grotz *et al.*, 1998), are transcriptionally responsive to Zn deficiency and mainly expressed in roots, but are also expressed in shoot tissue in response to low Zn, suggesting a role in both Zn uptake and distribution (Lin *et al.*, 2009; Jain *et al.*, 2013). Our findings indicate that higher tolerance to Zn deficiency may be the result of an increased, or more efficient, shoot Zn re-allocation capacity, and that natural variation for it may reflect variation in the expression of these and other Zn transport genes in *A. thaliana*.

Previous studies have shown that tolerance to Zn deficiency can also be affected by the plant's capacity to deal with the high levels of ROS produced under low Zn conditions (Rengel, 2001; Sinclair and Krämer, 2012). In this study, a relationship was found between the expression of *CA2* and ZnUI (Supplementary Table S9). Further studies examining the ability of plants to tolerate ROS under Zn deficiency and other mechanisms not included in this study, but thought to contribute to tolerance to Zn stress, will be useful for a more complete understanding of the mechanisms involved in plant tolerance to Zn deficiency (Cakmak *et al.*, 1996; Rengel, 2001; Gao *et al.*, 2005; Genc *et al.*, 2006; Hoffland *et al.*, 2006; Wissuwa *et al.*, 2006; Chen *et al.*, 2009; Impa *et al.*, 2013a, b). This should include examining the ability of plants to increase the bioavailability of Zn^{2+} ions in the soil, to improve the root system architecture to scavenge larger soil volumes, and a more efficient utilization, compartmentalization, and remobilization of Zn.

Model to predict Zn deficiency status based on the concentration of other elements

Exposing *A. thaliana* plants to different levels of Zn deficiency also affects the homeostasis of other elements, which made it possible to develop an MLR model able to predict the Zn deficiency status of a plant based on changes in other elements (Table 2). This approach is analogous to the model used by Baxter *et al.* (2008b) to predict the physiological status of *A. thaliana* plants exposed to Fe or P deficiency. Contrary to the MLR model developed for Zn deficiency, Baxter *et al.* (2008b) found that changes in Fe concentration alone had no power to detect Fe deficiency, and detection was

totally dependent on analysis of other elements. This difference could be caused by the two different Zn deficiency treatments used in this study which incorporated more data points in the model, while only one deficiency treatment was used in the Fe deficiency study (Baxter *et al.*, 2008b), but it could also be because in that study the Fe concentrations in leaves of plants grown under low and normal Fe did not differ, while in our study, the shoot Zn concentrations of plants grown under severe and mild Zn deficiency were significantly different, having extremely low Zn concentrations in comparison with control conditions. The Zn concentration thus appears to be much less tightly controlled in *A. thaliana* than Fe. In that respect, Zn corresponds more with P, for which their model did incorporate P concentration (Baxter *et al.*, 2008b). This analysis provides strong evidence that elements do not behave independently upon Zn deficiency and it shows the power of using a combination of elements as a phenotype of interest to detect a plant's nutritional status. The use of these traits to evaluate the tolerance of crops to Zn deficiency has the potential to simplify and shorten the process of identification of Zn deficiency-tolerant varieties. However, further studies confirming the application of biomarkers comparable with those found for *A. thaliana* in the evaluation of Zn deficiency tolerance in crops will be needed.

Conclusion

This study demonstrates that several physiological and molecular mechanisms underlie differences in Zn deficiency tolerance in *A. thaliana*. These include the minimum Zn concentration required for growth and the ability to take up and translocate Zn by inducing the expression of Zn deficiency-responsive genes. ZnUI, the reduction in SDW, and the expression level of Zn deficiency-responsive genes, such as *ZIP4* and *IRT3*, are useful proxies to evaluate plant tolerance to Zn deficiency in future studies. A mild Zn deficiency condition is more amenable for genetic studies than a severe stress, with higher heritability values for most studied traits and providing a more natural condition, at least for *A. thaliana*. Finally, the shoot ionome profile is a useful predictor of the plant Zn deficiency status. Changes in Zn concentration alone or in combination with other elements have an excellent capacity to detect physiological plant Zn deficiency in the absence of other visible symptoms. While we have shown this now for *A. thaliana*, a model plant species, the application of our findings will be in crops. Although it will be more difficult to establish this, we expect our research to inspire others to test the applicability of the described biomarkers in crops, under experimental and field conditions.

Supplementary data

Supplementary data are available at *JXB* online.

Table S1. Detailed information of the *A. thaliana* accessions used in this study.

Table S2. Oligonucleotide PCR primer sequences.

Table S3. Two-way ANOVA of shoot dry weight, Zn concentration, total Zn content, and gene expression of *A. thaliana* accessions exposed to control and Zn deficiency.

Table S4. Tukey pairwise multiple comparison between accessions for shoot dry weight, shoot Zn concentration, shoot Zn content, and gene expression levels (relative to reference genes) in the severe and mild Zn deficiency experiments.

Table S5. One-way ANOVA of differences between *A. thaliana* accessions exposed to control and Zn deficiency for relative changes in shoot dry weight, Zn concentration, total Zn content, Zn usage index, and gene expression.

Table S6. Tukey pairwise multiple comparison between accessions for relative change in shoot dry weight, shoot Zn concentration, shoot Zn content, and gene expression values in the severe and mild Zn deficiency treatments relative to their respective control treatments.

Table S7. One-way ANOVA of the log₁₀-transformed shoot element concentrations to determine significant differences between treatments.

Table S8. Pearson correlation coefficients for the comparison of shoot traits measured in the 19 *A. thaliana* accessions grown under severe or mild Zn deficiency and their respective control treatments.

Table S9. Pearson correlation coefficients for the comparison of gene expression with shoot traits measured in eight *A. thaliana* accessions grown under severe or mild Zn deficiency and their respective control treatments.

Fig. S1. Relative gene expression in leaves of eight *A. thaliana* accessions grown under severe and mild Zn deficiency, compared with their respective control treatments.

Acknowledgements

We gratefully acknowledge Maarten Koornneef for his critical comments on earlier versions of this manuscript. During the review and publication process of this manuscript Dr John Danku died. He will be greatly missed by his family, friends and colleagues and his untimely death is a significant loss to the field of ionomics. This work was supported by the Centre for BioSystems Genomics and grant 93512008 of the ZonMWZenith program, both initiatives under the auspices of the Netherlands Genomics Initiative, and by the EU-COST Action FA0905.

References

- Allen L, Benoist B, Dary O, Hurrell R. 2006. Guidelines on food fortification with micronutrients. Geneva Switzerland: World Health Organization and Food and Agricultural Organization of the United Nations.
- Alloway BJ. 2009. Soil factors associated with zinc deficiency in crops and humans. *Environmental Geochemistry and Health* **31**, 537–548.
- Assunção AGL, Herrero E, Lin Y-F, *et al.* 2010. *Arabidopsis thaliana* transcription factors bZIP19 and bZIP23 regulate the adaptation to zinc deficiency. *Proceedings of the National Academy of Sciences, USA* **107**, 10296–10301.
- Assunção AGL, Persson DP, Husted S, Schjørring JK, Alexander RD, Aarts MGM. 2013. Model of how plants sense zinc deficiency. *Metallomics* **5**, 1110–1116.
- Assunção AGL, Schat H, Aarts MGM. 2003. *Thlaspi caerulescens*, an attractive model species to study heavy metal hyperaccumulation in plants. *New Phytologist* **159**, 351–360.
- Baxter I, Brazelton JN, Yu D, *et al.* 2010. A coastal cline in sodium accumulation in *Arabidopsis thaliana* is driven by natural variation of the sodium transporter AthKT1;1. *PLoS Genetics* **6**, e1001193.

- Baxter I, Hermans C, Lahner B, Yakubova E, Tikhonova M, Verbruggen N, Chao DY, Salt DE.** 2012. Biodiversity of mineral nutrient and trace element accumulation in *Arabidopsis thaliana*. *PLoS One* **7**, e35121.
- Baxter I, Muthukumar B, Park HC, et al.** 2008a. Variation in molybdenum content across broadly distributed populations of *Arabidopsis thaliana* is controlled by a mitochondrial molybdenum transporter (MOT1). *PLoS Genetics* **4**, e1000004.
- Baxter IR, Vitek O, Lahner B, Muthukumar B, Borghi M, Morrissey J, Guerinot ML, Salt DE.** 2008b. The leaf ionome as a multivariable system to detect a plant's physiological status. *Proceedings of the National Academy of Sciences, USA* **105**, 12081–12086.
- Broadley MR, White PJ, Hammond JP, Zelko I, Lux A.** 2007. Zinc in plants. *New Phytologist* **173**, 677–702.
- Cakmak I.** 2007. Enrichment of cereal grains with zinc: agronomic or genetic biofortification? *Plant and Soil* **302**, 1–17.
- Cakmak I, Sari N, Marschner H, Kalayci M, Yilmaz A, Eker S, Gulut KY.** 1996. Dry matter production and distribution of zinc in bread and durum wheat genotypes differing in zinc efficiency. *Plant and Soil* **180**, 173–181.
- Cakmak I, Torun B, Erenoglu B, Öztürk L, Marschner H, Kalayci M, Ekiz H, Yilmaz A.** 1998. Morphological and physiological differences in the response of cereals to Zn deficiency. *Euphytica* **100**, 349–357.
- Chao DY, Silva A, Baxter I, Huang YS, Nordborg M, Danku J, Lahner B, Yakubova E, Salt DE.** 2012. Genome-wide association studies identify heavy metal ATPase3 as the primary determinant of natural variation in leaf cadmium in *Arabidopsis thaliana*. *PLoS Genetics* **8**, e1002923.
- Chen WR, He ZL, Yang XE, Feng Y.** 2009. Zinc efficiency is correlated with root morphology, ultrastructure, and antioxidative enzymes in rice. *Journal of Plant Nutrition* **32**, 287–305.
- Clemens S.** 2010. Zn—a versatile player in plant cell biology. *Plant Cell Monographs* **17**, 281–298.
- Danku JM, Lahner B, Yakubova E, Salt DE.** 2013. Large-scale plant ionomics. *Methods in Molecular Biology* **953**, 255–276.
- Ei-Lithy ME, Clercx EJ, Ruys GJ, Koornneef M, Vreugdenhil D.** 2004. Quantitative trait locus analysis of growth-related traits in a new *Arabidopsis* recombinant inbred population. *Plant Physiology* **135**, 444–458.
- Friedman J, Hastie T, Tibshirani R.** 2010. Regularization paths for generalized linear models via coordinate descent. *Journal of Statistical Software* **33**, 1–22.
- Gao X, Zou C, Zhang F, van der Zee SATM, Hoffland E.** 2005. Tolerance to zinc deficiency in rice correlates with zinc uptake and translocation. *Plant and Soil* **278**, 253–261.
- Genc Y, McDonald GK, Graham RD.** 2002. Critical deficiency concentration of zinc in barley genotypes differing in zinc efficiency and its relation to growth responses. *Journal of Plant Nutrition* **25**, 545–560.
- Genc Y, McDonald GK, Graham RD.** 2006. Contribution of different mechanisms to zinc efficiency in bread wheat during early vegetative stage. *Plant and Soil* **281**, 353–367.
- Ghandilyan A, Ilk N, Hanhart C, et al.** 2009. A strong effect of growth medium and organ type on the identification of QTLs for phytate and mineral concentrations in three *Arabidopsis thaliana* RIL populations. *Journal of Experimental Botany* **60**, 1409–1425.
- Ghandilyan A, Kutman UB, Kutman BY, Cakmak I, Aarts MGM.** 2012. Genetic analysis of the effect of zinc deficiency on *Arabidopsis* growth and mineral concentrations. *Plant and Soil* **361**, 227–239.
- Good AG, Shrawat AK, Muench DG.** 2004. Can less yield more? Is reducing nutrient input into the environment compatible with maintaining crop production? *Trends in Plant Science* **9**, 597–605.
- Graham RD, Asher JS, Hynes SC.** 1992. Selecting zinc-efficient cereal genotypes for soils of low zinc status. *Plant and Soil* **146**, 241–250.
- Grotz N, Fox T, Connolly E, Park W, Guerinot ML, Eide D.** 1998. Identification of a family of zinc transporter genes from *Arabidopsis* that respond to zinc deficiency. *Proceedings of the National Academy of Sciences, USA* **95**, 7220–7224.
- Hacisalihoglu G, Hart JJ, Wang YH, Cakmak I, Kochian LV.** 2003. Zinc efficiency is correlated with enhanced expression and activity of zinc-requiring enzymes in wheat. *Plant Physiology* **131**, 595–602.
- Hacisalihoglu G, Öztürk L, Cakmak I, Welch RM, Kochian LV.** 2004. Genotypic variation in common bean in response to zinc deficiency in calcareous soil. *Plant and Soil* **259**, 71–83.
- Hoffland E, Wei C, Wissuwa M.** 2006. Organic anion exudation by lowland rice (*Oryza sativa* L.) at zinc and phosphorus deficiency. *Plant and Soil* **283**, 155–162.
- Huang XY, Salt DE.** 2016. Plant ionomics: from elemental profiling to environmental adaptation. *Molecular Plant* **9**, 787–797.
- Ibarra-Laclette E, Lyons E, Hernández-Guzmán G, et al.** 2013. Architecture and evolution of a minute plant genome. *Nature* **498**, 94–98.
- Impa SM, Gramlich A, Tandy S, Schulin R, Frossard E, Johnson-Beebout SE.** 2013a. Internal Zn allocation influences Zn deficiency tolerance and grain Zn loading in rice (*Oryza sativa* L.). *Frontiers in Plant Science* **4**, 534.
- Impa SM, Morete MJ, Ismail AM, Schulin R, Johnson-Beebout SE.** 2013b. Zn uptake, translocation and grain Zn loading in rice (*Oryza sativa* L.) genotypes selected for Zn deficiency tolerance and high grain Zn. *Journal of Experimental Botany* **64**, 2739–2751.
- Jain A, Sinilal B, Dhandapani G, Meagher RB, Sahi SV.** 2013. Effects of deficiency and excess of zinc on morphophysiological traits and spatiotemporal regulation of zinc-responsive genes reveal incidence of cross talk between micro- and macronutrients. *Environmental Science and Technology* **47**, 5327–5335.
- Karim MR, Zhang YQ, Tian D, Chen FJ, Zhang FS, Zou CQ.** 2012. Genotypic differences in zinc efficiency of Chinese maize evaluated in a pot experiment. *Journal of the Science of Food and Agriculture* **92**, 2552–2559.
- Kobayashi Y, Kuroda K, Kimura K, Southron-Francis JL, Furuzawa A, Kimura K, Iuchi S, Kobayashi M, Taylor GJ, Koyama H.** 2008. Amino acid polymorphisms in strictly conserved domains of a P-type ATPase HMA5 are involved in the mechanism of copper tolerance variation in *Arabidopsis*. *Plant Physiology* **148**, 969–980.
- Koprivova A, Giovannetti M, Baraniecka P, Lee BR, Grondin C, Loudet O, Kopriva S.** 2013. Natural variation in the ATPS1 isoform of ATP sulfurylase contributes to the control of sulfate levels in *Arabidopsis*. *Plant Physiology* **163**, 1133–1141.
- Kruijer W, Boer MP, Malosetti M, Flood PJ, Engel B, Kooke R, Keurentjes JJ, van Eeuwijk FA.** 2015. Marker-based estimation of heritability in immortal populations. *Genetics* **199**, 379–398.
- Li Y, Zhang Y, Shi D, et al.** 2013. Spatial-temporal analysis of zinc homeostasis reveals the response mechanisms to acute zinc deficiency in *Sorghum bicolor*. *New Phytologist* **200**, 1102–1115.
- Lin YF, Liang HM, Yang SY, Boch A, Clemens S, Chen CC, Wu JF, Huang JL, Yeh KC.** 2009. *Arabidopsis* IRT3 is a zinc-regulated and plasma membrane localized zinc/iron transporter. *New Phytologist* **182**, 392–404.
- Livak KJ, Schmittgen TD.** 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* **25**, 402–408.
- Loudet O, Saliba-Colombani V, Camilleri C, Calenge F, Gaudon V, Koprivova A, North KA, Kopriva S, Daniel-Vedele F.** 2007. Natural variation for sulfate content in *Arabidopsis thaliana* is highly controlled by APR2. *Nature Genetics* **39**, 896–900.
- Marschner H.** 1995. Mineral nutrition of higher plants. Amsterdam: Academic Press.
- Morrissey J, Baxter IR, Lee J, Li L, Lahner B, Grotz N, Kaplan J, Salt DE, Guerinot ML.** 2009. The ferroportin metal efflux proteins function in iron and cobalt homeostasis in *Arabidopsis*. *The Plant Cell* **21**, 3326–3338.
- Oñate-Sánchez L, Vicente-Carbajosa J.** 2008. DNA-free RNA isolation protocols for *Arabidopsis thaliana*, including seeds and siliques. *BMC Research Notes* **1**, 93.
- Pineau C, Loubet S, Lefoulon C, Chalies C, Fizames C, Lacombe B, Ferrand M, Loudet O, Berthomieu P, Richard O.** 2012. Natural variation at the FRD3 MATE transporter locus reveals cross-talk between Fe homeostasis and Zn tolerance in *Arabidopsis thaliana*. *PLoS Genetics* **8**, e1003120.
- Randall PJ, Bouma D.** 1973. Zinc deficiency, carbonic anhydrase, and photosynthesis in leaves of spinach. *Plant Physiology* **52**, 229–232.
- Rengel Z.** 2001. Genotypic differences in micronutrient use efficiency in crops. *Communications in Soil Science and Plant Analysis* **32**, 1163–1186.
- Rengel Z, Graham RD.** 1996. Uptake of zinc from chelate-buffered nutrient solutions by wheat genotypes differing in zinc efficiency. *Journal of Experimental Botany* **47**, 217–226.

- Richard O, Pineau C, Loubet S, Chalies C, Vile D, Marquès L, Berthomieu P.** 2011. Diversity analysis of the response to Zn within the *Arabidopsis thaliana* species revealed a low contribution of Zn translocation to Zn tolerance and a new role for Zn in lateral root development. *Plant, Cell and Environment* **34**, 1065–1078.
- Rus A, Baxter I, Muthukumar B, Gustin J, Lahner B, Yakubova E, Salt DE.** 2006. Natural variants of AtHKT1 enhance Na⁺ accumulation in two wild populations of *Arabidopsis*. *PLoS Genetics* **2**, e210.
- Sadeghzadeh B, Rengel Z, Li C.** 2009. Differential zinc efficiency of barley genotypes grown in soil and chelator-buffered nutrient solution. *Journal of Plant Nutrition* **32**, 1744–1767.
- Salt DE, Baxter I, Lahner B.** 2008. Ionomics and the study of the plant ionome. *Annual Review of Plant Biology* **59**, 709–733.
- Sharma PN, Kumar P, Tewari RK.** 2004. Early signs of oxidative stress in wheat plants subjected to zinc deficiency. *Journal of Plant Nutrition* **27**, 451–463.
- Siddiqi MY, Glass ADM.** 1981. Utilization index: a modified approach to the estimation and comparison of nutrient utilization efficiency in plants. *Journal of Plant Nutrition* **4**, 289–302.
- Sinclair SA, Krämer U.** 2012. The zinc homeostasis network of land plants. *Biochimica et Biophysica Acta* **1823**, 1553–1567.
- Talukdar S, Aarts MGM.** 2007. *Arabidopsis thaliana* and *Thlaspi caerulescens* respond comparably to low zinc supply. *Plant and Soil* **306**, 85–94.
- van de Mortel JE, Almar Villanueva L, Schat H, Kwekkeboom J, Coughlan S, Moerland PD, Ver Loren van Themaat E, Koornneef M, Aarts MGM.** 2006. Large expression differences in genes for iron and zinc homeostasis, stress response, and lignin biosynthesis distinguish roots of *Arabidopsis thaliana* and the related metal hyperaccumulator *Thlaspi caerulescens*. *Plant Physiology* **142**, 1127–1147.
- Waters BM, Grusak MA.** 2008. Whole-plant mineral partitioning throughout the life cycle in *Arabidopsis thaliana* ecotypes Columbia, Landsberg erecta, Cape Verde Islands, and the mutant line ysl1ysl3. *New Phytologist* **177**, 389–405.
- White PJ, Broadley MR.** 2011. Physiological limits to zinc biofortification of edible crops. *Frontiers in Plant Science* **2**, 80.
- Wissuwa M, Ismail AM, Yanagihara S.** 2006. Effects of zinc deficiency on rice growth and genetic factors contributing to tolerance. *Plant Physiology* **142**, 731–741.
- Zhang X, Hause RJ Jr, Borevitz JO.** 2012. Natural genetic variation for growth and development revealed by high-throughput phenotyping in *Arabidopsis thaliana*. *G3* **2**, 29–34.