

Autonomous Magnetic Labelling of Functional Mesenchymal Stem Cells for Improved Traceability and Spatial Control in Cell Therapy Applications.

Richard Harrison^{1#}, Hareklea Markides^{2#}, Robert. H. Morris³, Paula Richards², Alicia J. El Haj², Virginie Sottile^{1}.*

¹ Wolfson Centre for Stem Cells, Tissue Engineering and Modelling (STEM), School of Medicine, The University of Nottingham

² Institute of Science and Technology in Medicine, Keele University

³ School of Science and Technology, Nottingham Trent University

These authors contributed equally to this study.

* Corresponding author:

Dr. Virginie Sottile

Wolfson Centre for Stem Cells, Tissue Engineering and Modelling (STEM)

School of Medicine, The University of Nottingham

CBS Building, University Park, Nottingham NG7 2RD, UK

Telephone: +44 115 823 1235

E-mail: virginie.sottile@nottingham.ac.uk

Running Title: Magnetic Mesenchymal Stem Cells Labelling for Imaging and Improved Cell Control.

Abstract

Mesenchymal stem cells (MSCs) represent a valuable resource for regenerative medicine treatments for orthopaedic repair and beyond. Following developments in isolation, expansion and differentiation protocols, efforts to promote clinical translation of emerging cellular strategies now seek to improve cell delivery and targeting. This study shows efficient live MSC labelling using silica-coated magnetic particles (MPs), which enable 3D tracking and guidance of stem cells. A procedure developed for the efficient and unassisted particle uptake was shown to support MSC viability and integrity, while surface marker expression and MSC differentiation capability were also maintained. In vitro, MSCs showed a progressive decrease in labelling over increasing culture time, which appeared linked to the dilution effect of cell division rather than particle release, and did not lead to detectable secondary particle uptake. Labelled MSC populations demonstrated magnetic responsiveness in vitro through directed migration in culture and when seeded onto a scaffold, supporting MP-based approaches to cell targeting. The potential of these silica-coated MPs for MRI-cell tracking of MSC populations was validated in 2D and in a cartilage repair model following cell delivery. These results highlight silica-coated magnetic particles as a simple, safe and effective resource to enhance MSC targeting for therapeutic applications and improve patient outcomes.

Keywords: Mesenchymal stem cell, cell labelling, magnetic microparticle, cell targeting, regenerative medicine

1. Introduction

Over the past decades, a range of iron oxide-based magnetic particles (MPs) have been developed for clinical applications in the field of magnetic resonance imaging (MRI) (Gilchrist *et al.*, 1957). Superparamagnetic iron oxide nanoparticles (SPIOs) are a specific class of magnetic particles known for their application as T₂-weighted negative MRI contrast agents designed to overcome the inherent low sensitivity associated with MRI (Pooley 2005, Bulte and Kraitchman 2004). Magnetic particles composed of either a magnetite (Fe₃O₄) or maghemite (γ-Fe₂O₃) core (Berman *et al.*, 2011, Gupta and Gupta 2005) surrounded by a biocompatible polymer such as silica and dextran have been used for the labelling and identification of cell populations (Kunzmann *et al.*, 2011). FDA-approved iron-based particles such as Endorem (also referred to as Feridex) and Resovist have been used as MRI contrast agents in recent years (Berman *et al.*, 2011, Jasmin *et al.*, 2011), however since these products are no longer clinically used there is a need for validated products offering low toxicity, biocompatibility, and chemical stability in physiological conditions (Hofmann-Antenbrink *et al.*, 2010, Mahmoudi *et al.*, 2011).

While the literature on MPs has largely focused on developing particle design, synthesis and characterization (McBride *et al.*, 2013), recent studies have also investigated MPs for cell-based applications beyond MRI imaging, as their applied magnetic fields have been used to develop new approaches to enhance transfection (Pickard *et al.*, 2011), induce hyperthermia (Kobayashi 2011), force in vitro aggregation (Fayol *et al.*, 2013), enable regenerative therapies (El Haj *et al.*, 2012) and activate cell receptor signalling on the cell membrane (Henstock *et al.*, 2014). Their small size and magnetic properties, coupled with versatile surface coatings (Gupta and Gupta 2005) open a range of new approaches which could see MPs

enhance existing and future regenerative cell therapies. Such cell-based approaches require the targeted delivery of functional populations such as mesenchymal stem cells (MSCs), which have become a resource of prime importance for their skeletal regeneration ability (Caplan 2007, Quarto *et al.*, 2001), but also for their properties of immune modulation (Le Blanc *et al.*, 2003), anti-inflammation (Uccelli 2008) or trophic secretion (Caplan and Dennis 2006). MSC-based therapies for tissue repair require auxiliary approaches which enable *in vivo* tracking, delivery and targeting, in order to monitor and improve the retention of functional cells at the intervention site (Wimpenny *et al.*, 2012).

In this study, the suitability of MPs presenting a silica surface with negatively charged silanol groups was investigated for use in human mesenchymal stem cells as a labelling, imaging and manipulation agent. The labelling dynamics and cellular response were analyzed with a particular emphasis on markers of cell health, identity and functional potential of the target population, as well as their suitability for cell tracking purposes in an articular model. Observations presented here can help refine novel applications of MP labelling and evaluate the resulting health considerations of future MP-assisted stem cell therapies.

2. Materials and Methods

All reagents were purchased from Life Technologies unless otherwise stated.

2.1. Human Mesenchymal Stem Cell Cultures: A human bone marrow derived mesenchymal stem cell line (hMSCs) (Okamoto *et al.*, 2002, France *et al.*, 2014) was cultured and expanded under standard cell culturing conditions (37.5°C, 5% CO₂) in standard culture medium consisting of Dulbecco's Modified Eagle's Medium

(DMEM) supplemented with 10% (v/v) Fetal Bovine Serum (FBS), 1% (v/v) non-essential amino acids, 1 mM L-Glutamine, 1mM Pyruvate and 1% Penicillin/streptomycin. Cells were passaged using Trypsin/EDTA. For some experiments, hMSCs stably transfected to constitutively express GFP (gMSCs) following an established protocol (Peister *et al.*, 2004) were used under standard cell culturing conditions in standard culture medium to enable fluorescence microscopy.

Primary human mesenchymal stem cells (pMSCs) were isolated from human bone marrow aspirate (Lonza, UK). In brief, the bone marrow aspirate was seeded in fibronectin coated flasks at a mononuclear cell density of 1.5×10^3 cell/cm² and cultured for one week (37°C, 5% CO₂) in pMSC isolation medium containing low glucose DMEM (Lonza Biowhittaker, UK) supplemented with 10 % FBS (Lonza Biowhittaker), 1 % L-Glutamine (Sigma-Aldrich, UK) and 1 % Penicillin/Streptomycin (Sigma-Aldrich). A 50% medium change with fresh pMSC isolation medium was performed after one week, followed by a switch one week later to hMSC proliferation media (high glucose DMEM supplemented with 10 % FBS, 1 % L-Glutamine and 1 % Penicillin/Streptomycin). pMSCs were identified as those which had adhered to the tissue culture vessel after 14 days in culture.

2.2. Cell Labelling With Magnetic Particles (MPs): hMSCs and pMSCs were labelled with 1000 nm particles composed of a maghemite core with a solid unmodified silica surface as previously described (Markides *et al.*, 2013), using standard (SiMAG) or fluorescently tagged (ScreenMAG-Silanol) particles as specified (Chemicell, Germany). In brief, adherent cell populations were incubated with MPs (1-10 µg/mL) in medium for 24 h, using serum-containing or serum-free medium (MRI

experiments) as specified (for cell labelling experiments, standard medium containing 10% FBS was used unless otherwise stated). The next day, cells were thoroughly washed with phosphate buffered saline (PBS) in order to remove excess particles that may have settled on the surface of the cell layer or flask.

To measure particle uptake by flow cytometry, cells were seeded at 7.5×10^3 cell/mL and ScreenMAG-labelled for 24 h. Cells were then harvested, centrifugated at 200 g for 5 min, and re-suspended in 200 μ L PBS prior to analysis on a Guava EasyCyte 8HT Flow Cytometer Channel FL2 with InCyte 2.5 Software (Millipore, USA) comparing labelled and unlabelled populations to evaluate percentage uptake based on fluorescent intensity. Analysis was performed using WEASEL (WEHI, Australia) using unlabelled cells as controls to evaluate increased fluorescence. The standard particle concentration used in the study is 10 μ g/mL unless otherwise stated, which was shown to correspond to an intracellular iron load of 20 pg/cell (Markides *et al.*, 2013).

2.3. Fluorescence Imaging of Particle Uptake: Particle uptake was further evaluated visually using an array of fluorescent cell dyes and fluorescent microscopy to evaluate internalisation in relation to cell structure. hMSCs cultured on glass coverslips were labelled with particles and fixed at room temperature for 15 min in 4% (v/v) PFA (VWR, UK). After permeabilisation with 0.1% Triton x-100 for 5 min following two PBS washes, cells were stained for actin filaments using a 1:41 working solution of 6.6 μ M Alexa Fluor® 488 phalloidin in methanol. Slides were incubated in a dark covered container at room temperature for 20 min, and then washed twice with PBS prior to mounting using Vectashield mounting medium

(Vector Laboratories, USA). Imaging was performed using a Leica TCS SP2 confocal laser scanning microscope (CLSM) (Leica Microsystems, Germany).

2.4. Prussian blue staining: hMSC cells were grown in monolayer and labelled with 10 µg/mL MPs for 24 h prior to fixing with 4% PFA for 15 min. Immediately prior to addition to cells, 20% aqueous solution of hydrochloric acid and 10% aqueous solution of potassium ferrocyanide were mixed in equal parts. This staining solution was applied to the fixed monolayer for 5 min and washed three times with PBS. Images were acquired using an Eclipse TS100 inverted microscope (Nikon, Japan).

2.5. Transmission Electron Microscopy (TEM): To confirm the cellular location of the particles, samples were fixed in 3% glutaraldehyde in 0.1 M cacodylate buffer overnight and post-fixed in 1% aqueous osmium tetroxide for 30 min. The samples were then dehydrated in a graded ethanol series and infiltrated with Transmit resin (TAAB, UK) allowed to polymerise overnight at 70°C. Semi-thin sections were cut (0.5 µm) using a Reichert-Jung ultramicrotome and stained with 2% toluidine blue. Ultra-thin sections were cut (70–90 nm) using the same equipment and collected on copper grids. Grids were then contrasted using 50% methanolic uranyl acetate and Reynolds lead citrate (Robards and Wilson 1993). Imaging was performed on a FEI Tecnai 12 Biotwin TEM (FEI, USA) with up to 120kV and x300k magnification.

2.6. Particle Labelling measurement: Flow cytometry was used to measure the level of particle labelling over time. For mitotic arrest, mitomycin C (Sigma Aldrich, UK) treatment was used to halt cell division, using a final concentration of 10 µg/mL for a

2.5 h incubation at 37°C (Nieto *et al.*, 2007). Cells were then washed twice with PBS and harvested for use. Mitotically arrested and control cells were cultured over a 7 day period with cells fixed in 4% PFA for analysis on days 1, 5 and 7. To investigate particle transfer between co-cultured populations, GFP-expressing MSCs (gMSCs) labelled with MPs were cultured with unlabelled hMSCs. Both populations were mitotically arrested prior to co-culture with samples fixed in 4% PFA each day over 7 days, before flow cytometry quantification of particle presence and GFP status.

2.7. Cell Surface Marker Analysis: hMSCs and pMSCs were assessed for expression of multipotent markers (Dominici *et al.*, 2006) performed 24 h after MP labelling (with SiMAG and ScreenMAG respectively), and 14 days after initial labelling with repeated passaging and re-labelling every 3 days to maintain a high MP level throughout. Cells were harvested with trypsin/EDTA and pelleted by centrifugation for 5 min at 200 g, before washing in PBS. Cell pellets were then resuspended in 100 µL PBS supplemented with 5 µL of antibodies against CD29 (Abcam, UK), CD105, CD34 and CD73 (AbdSerotec, UK), CD90 and SSEA4 (eBiosciences, USA) for 30 min at room temperature, before two PBS washes and flow cytometry analysis.

2.8. Cell Viability Assays: The resazurin metabolic assay was performed to determine metabolic changes, using a working solution consisting of 10% (v/v) Presto Blue stock solution prepared according to the manufacturer's instructions. After 45 min of incubation, the fluorescent signal 100 µL samples was measured at

535 nm excitation and 615 nm emission in triplicate using an Infinite 200 PRO plate reader and i-control software (Tecan, Switzerland).

Impact on membrane integrity was assessed using a LIVE/DEAD® Alexa Fluor® 488 fixable viability dye. Cells were harvested with trypsin/EDTA and pelleted by centrifugation for 5 min at 200 g, washed twice with PBS twice, and resuspended in 100 µL amine reactive dye working solution consisting of 1% (v/v) amine reactive DMSO stock in PBS. Following 15 min incubation at room temperature, cells were rinsed twice with PBS and resuspended in 200 µL PBS prior to measurement on a Guava EasyCyte 8HT Flow Cytometer. Unlabelled cells were used as viable controls and DMSO or paraformaldehyde fixative treatments provided toxicity controls.

2.9. Single Cell Gel Electrophoresis (Comet) Assay: Potential damage to the DNA was assessed with the alkaline comet assay (Seedhouse *et al.*, 2006). hMSC were grown in monolayer and either left unlabelled, labelled with 10 µg/mL or 100 µg/mL SiMAG for 24 hours. Following trypsinisation, cells were washed with PBS once and resuspended in low melting point agarose (Trevigen, UK) at 10^5 cells/mL. CometAssay alkaline control cells were used as a positive control for DNA damage (Trevigen). Cell containing agarose was immediately spread on comet slides (Trevigen) and left to harden before complete immersion in cell lysis buffer (Trevigen). Lysis was performed overnight at 4°C in the dark. Following this, lysis buffer was removed and slides immersed in a UV protected electrophoresis tank containing TBE running buffer and allowed to sit for 60 min. Voltage was set at 25V per CM distance between electrodes and running time at 40 min. Following running, slides were removed from the buffer and washed three times in dH₂O before dipping

in ethanol for 1 min and drying overnight. Dry comet slides were stained with 75 μ L of 0.2% SYBR Green in TBE buffer per agarose droplet. Samples were immediately imaged under a rhodamine filter with an Olympus BX40 microscope. Comet tails were analyzed using Comet Assay III image analysis software (Perceptive Instruments, UK). Fifty comet images were obtained from each of the duplicate gel spots and each experimental condition was repeated three times; therefore, 600 images were scored in total for each treatment. The Tail Moment was used in all analysis.

2.10. Mesenchymal Differentiation: For differentiation assays, hMSCs were incubated for a period of 21 days in the relevant differentiation media. For osteogenic assays cells were seeded at 5×10^3 cells/cm² in well plates (Sigma-Aldrich, UK). The medium was then changed (considered as day 0) every 3 days for 21 days with either control medium or osteogenic induction DMEM supplemented with 100 nM dexamethasone, 0.05 mM L-ascorbic acid-2-phosphate and 10 mM β -Glycerophosphate. For adipogenic assays cells were seeded at 1×10^4 cells/cm² in well plates (Sigma-Aldrich, UK). The medium was then changed (considered as day 0) every 3 days for 21 days with either control medium or adipogenic induction high glucose (4500 mg/L) DMEM supplemented with 1 μ M dexamethasone, 500 μ M isobutylmethylxanthine, 10 μ g/mL insulin, and 1 μ M rosiglitazone. For chondrogenic assays cells were seeded at 37.5×10^4 cells/cm² in flasks for labelling duration. Cells were then detached and 200 μ L of 1.25×10^6 cells/mL cell suspensions added to 96 well v-bottom plates (Nalge Nunc International, USA) and spun at 450g for 10 min. Following 24 h attachment duration, medium was then changed every day for 21 days with either control medium or chondrogenic induction high glucose (4500 mg/L)

DMEM supplemented with 2 mM L-glutamine, 0.1 μ M dexamethasone, 50 μ g/mL ascorbic acid phosphate, 1 mM Na-pyruvate, 40 μ g/mL Proline, 10 ng/mL TGF- β and 1x ITS Liquid Media Supplement (Sigma-Aldrich, UK).

2.11. Differentiation Assays: Lipid containing cells were identified using Oil Red O (Sheng *et al.*, 2007). Cells were washed with PBS and fixed at room temperature for 15 min in 4% (v/v) paraformaldehyde (PFA). Cells were then washed twice with dH₂O and incubated Oil Red O working solution added (180 mg/L Oil Red O in 60% isopropanol/40% dH₂O) for 30 min at ambient temperature. Samples were then washed and imaged before extraction of the incorporated stain with isopropanol to measure absorption at 510 nm on an Infinite 200 PRO plate reader and i-control software (Tecan, Switzerland).

Mineralised nodules were identified using Von Kossa staining (Wang *et al.*, 2006). Cells were washed with PBS and fixed at room temperature for 15 min in 4% PFA. Cells were then washed three times with dH₂O and incubated with 1% silver nitrate in dH₂O (Sigma-Aldrich, UK) under a UV lamp for 15 min. Samples were washed three times with dH₂O, incubated for 5 min with 2.5% sodium thiosulfate solution (Sigma-Aldrich, UK), washed again with dH₂O and imaged using an Eclipse TS100 inverted microscope (Nikon, Japan).

Sulfated glycosaminoglycans detected with the dye 1,9-dimethylmethylene blue (DMMB) were used as an indicator of chondrogenesis. Chondrogenic micromasses were freeze-thawed three times to partially disaggregate them followed by papain digestion (sodium phosphate (0.1 M), cysteine hydrochloride (5 mM), EDTA (5 mM) and papain (45.12 μ M) in dH₂O, pH adjusted to 6.5) overnight at 60°C. Aliquots of

digested sample were stained with DMMB dye solution (0.03 M sodium formate, 0.046 mM DMMB, 85.5 mM ethanol and 53 mM formic acid in dH₂O) left for 10 minutes at room temperature and read for absorbance at 540 nm on an Infinite 200 PRO plate reader and i-control software (Tecan, Switzerland). Aliquots of digested sample were also taken for DNA content analysis with CyQUANT® to allow for normalization. CyQUANT® GR dye/cell-lysis buffer was added to samples and incubated for 5 min at room temperature. Samples were analyzed on an Infinite 200 PRO plate reader and i-control software (Tecan, Switzerland).

2.12. Directed migration assays: For the vertical migration model, hMSCs were labelled with concentrations ranging from 2.5µg/mL to 100 µg/mL alongside unlabelled control cells for 24 hours. Cells were harvested and resuspended to a concentration of 1×10^5 cells/mL. 20 µL drops were deposited, in quadruplicate, on the inside of a multiwell plate lid which was carefully placed to form hanging drops suspended above humidified wells. A magnetic array constructed from 10mm x 3mm neodymium magnets (2800 gauss) (Magnet Expert, UK) was placed above each well, and after 24 h the proportion of cells attached to the under surface of the lid was evaluated after toluidine blue staining (0.1% for 10 minutes) and imaging using a 41 Megapixel PureView Zeiss Camera (Nokia, Finland). Quantitative 2D image density analysis was performed using ImageJ (NIH, USA).

For the transmigration assay, SiMAG-labelled pMSCs (0, 1, and 10 10 µg/mL) were seeded at a concentration of 10^4 cells / collagen transwell insert (Corning, UK) and allowed to attach for 24 h. Plates were either placed on a magnetic array mimicking a standard 24-well plate layout or cultured without a magnetic field for 24 hours. The

collagen layer was then gently removed and the transwell completely washed three times with PBS. Migrated cells located on the underside of the transwell were fixed (using 4% formalin for 1 hour), stained with DAPI and imaged with a fluorescent microscope. Five independent areas of the well were imaged (top, bottom, left right and centre) and averaged for each sample.

2.13. MRI imaging: To establish the ex vivo knee model, chondrocytes were isolated from porcine articular knee cartilage (Staffordshire Meat Packers, Stoke-on-Trent, UK) two hours post slaughtering based on a technique adapted from (Hayman *et al.*, 2006). Cartilage was carefully removed from the upper condyles of the knee, finely diced, weighed and rinsed in PBS and 2 % Penicillin/Streptomycin. After overnight incubation in chondrocyte isolation media consisting of DMEM/HAM'S F12 (Lonza Biowhittaker, UK), 2 % Penicillin/Streptomycin, 50 µg/mL sterilised ascorbate (Sigma-Aldrich, UK), 1 mg/mL clostridial collagenase (Sigma-Aldrich, UK) and 0.1 mg/mL DNase (Sigma-Aldrich, UK), the digested cartilage suspension was filtered through 100 µm cell strainer and centrifuged at 600G for 10 min. Chondrocytes were seeded at 2×10^4 cells/cm² and cultured in chondrocytes proliferation media (DMEM/HAM'S F12 supplemented with 10 % FBS, 1 % L-Glutamine and 1 % Penicillin/Streptomycin).

The in vitro MRI visibility threshold of SiMAG-labelled cells populations (0, 1, 5, 10 and 100 µg/mL) was investigated at varying cell densities (5×10^5 , 10^5 and 10^4) in 2 mg/mL rat tail type I collagen gel (BD Biosciences, UK). Samples were then imaged using a 2.3T Bruker animal scanner (NTU, Nottingham, UK) with MSME sequences

using 1000 ms repetition time, 10.25 ms echo time with 8 echoes, and a matrix size of 256x192 with spatial resolution of 0.469x0.625 mm.

Ex vivo imaging was carried out using a cadaveric porcine knee model of articular cartilage damage to assess the visibility threshold of MP-labelled cells in a clinically relevant model of autologous chondrocyte implantation (ACI) to treat cartilage damage (Chiang *et al.*, 2005). Pig legs were processed to remove all surrounding tissue using a surgical scalpel. Once the knee had been isolated, the patella tendon was sliced and the patella pulled back to reveal the articulating ends of the femur and tibia. The knee was then bent to fully expose the upper condyles, and cartilage flaps were created (1.5 cm x 0.5 cm x 1.5 cm) across the upper condyles of the knee. Two defects were created on each condyle (left and right) at least 0.5 cm apart. MP-labelled cells were suspended in a collagen type 1 gel solution (4.5 mg/mL) and injected within the defect while the knee was in the bent upright position, taking care to ensure no bubbles or leakage occurred. After the gels had set (1 hr, 37°C) the leg was straightened, the patella replaced and securely bandaged to prevent excess movement, before storage at -20°C until imaging at the MARIARC centre (Liverpool University) using a Siemens Symphony 1.5 T scanner. One day prior to MR imaging samples were defrosted, placed within a circularly polarised extremity coil, and Double Echo Steady State (DESS) sequences were applied in agreement with MRI scanning conditions implemented in the imaging and diagnosis of human knee pathologies.

2.14. Statistical Analysis: Statistical analysis was in the form of ANOVA performed using GraphPad PRISM (GraphPad Software, USA). Tukey's post hoc analysis was

performed to determine the significance between subgroups of the analysed population. Significance shown as * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ and **** $p < 0.0001$.

3. Results

3.1. Cytocompatibility study

To evaluate the capacity of MSCs to take up MPs, monolayer cultures were incubated overnight with various concentrations of particles. Particle uptake in hMSCs following a 24 h incubation period with MPs was analyzed by fluorescence microscopy and flow cytometry (Figure 1).

Incubation with increasing doses of MPs led to a proportional increase in the fluorescence signal measured for hMSCs (Figure 1A). Time-lapse microscopy (Supplementary File 1) and fluorescence microscopy (Figure 1B) confirmed particle uptake while cells retained morphology post-labelling. Prussian blue staining allowed visualisation of the iron containing particles present within cells (Figure 1C). TEM imaging confirmed the presence of MPs within the cytoplasm, and highlighted their localisation to vesicles found to congregate around the nucleus (Figure 1D). The efficiency of MP uptake was compared under different serum concentrations using flow cytometry, which demonstrated a dose-dependent negative effect of serum on cell labelling (Figure 1E).

Following uptake, particle retention was analyzed over time in culture (Figure 2). In dividing hMSCs, MPs were found to be progressively diluted, until day 7 when they

were not detected (Figure 2A). In mitotically arrested cells however, particles were retained more efficiently and showed a significant retention compared to untreated cells at day 7, suggesting the MP load might be divided between daughter cells. Observation of labelled cells showed the occasional presence of some isolated particles within cell projections (Figure 2B).

To investigate the fate of particles over time, a co-culture experiment was set up to examine whether MPs may be transferred between labelled and unlabelled hMSC populations (Figure 2C). GFP-expressing MSCs (gMSCs) labelled with MPs were mixed with control unlabelled hMSCs, and over 7 days in co-culture cells were analyzed by flow cytometry to evaluate the proportion of MP-containing cells within each MSC population. While a decrease in the percentage of MP-containing gMSCs was seen over time, there was no detectable appearance of MP-containing cells in the unlabelled hMSC population over 7 days.

The effect of MP exposure on cell identity was analyzed through surface marker analysis and cell integrity assays (Figure 3). Using markers associated to MSCs, comparable positive expression of CD90, CD105, CD73, SSEA4 and CD29, with negative expression of CD34, was confirmed between labelled and unlabelled control populations 24 h post labelling (Figure 3A). Cultures exposed to serial MP labelling every three days for 14 days to maintain maximum dose similarly demonstrated retained marker expression, confirming that exposure to MPs did not elicit a significant change in marker identity (Supplementary File 2).

The effect of MP exposure was further investigated through metabolic assays of MSCs labelled with increasing doses of SiMAG MPs using a resazurin based dye Presto Blue. Data gathered demonstrated a slight increase in metabolic activity at

low particle doses and a decreased metabolic activity associated with very high doses 24 h post labelling (Figure 3B). This increased metabolic activity at low MP doses appeared to be lost 48 h post labelling (*data not shown*). Cell membrane integrity assessed using flow cytometry indicated no effect of MP labelling could be detected 24 h (Figure 3C) post labelling for either pMSCs cells or hMSCs.

Since MPs were found to accumulate close to the nucleus, their possible effect on cellular DNA was examined using the comet assay which provides a sensitive measure of DNA damage throughout the population (Figure 3D). No statistically significant increase in DNA damage was observed at 10-100 µg/mL when compared to unlabelled MSC controls ($p>0.05$).

3.2. Application of MSC labelling for regenerative medicine

After establishing the cytocompatibility of particle labelling, the efficiency of the differentiation response obtained under various culture conditions was evaluated in MSCs. hMSCs, either unlabelled or labelled with SiMAG, were treated with osteogenic, adipogenic and chondrogenic media for 7 and 14 days to measure their response with and without MP exposure (Figure 4). After 21 days in culture with relevant differentiation media, histology staining (Fig. 4A-C) showed successful responses as detected through mineral deposition (Von Kossa staining for the osteogenic condition), lipid accumulation (Oil Red O staining for the adipogenic condition) and glycosaminoglycan (GAG) production (Alcian blue staining for the chondrogenic condition). Subsequent quantitative assays revealed no significant difference between unlabelled and MP-labelled cell populations for the osteogenic alkaline phosphatase activity and Alizarin Red O assays (Fig. 4D-E), or for the

adipogenic Oil Red O staining (Figure 4F). Quantitation of GAG formation in response to the chondrogenic treatment (Figure 4G) showed no detrimental effect of MP labelling, which produced a slight detectable increase in signal compared to unlabelled controls. These data demonstrate no reduction in differentiation capacity following particle labelling.

The iron core present in SiMAG particles makes them susceptible to magnetic forces, a feature potentially beneficial for novel tissue engineering approaches. In order to test whether MSC labelling with SiMAG could provide added control over the behaviour of the cells, a migration assay was run to measure the cellular response in vitro (Figure 5). When exposed to a permanent magnet located above the samples for 24 hours (Figure 5A), labelled cells displayed a significant higher vertical migration towards the magnet when compared to unlabelled samples, which failed to migrate and adhere. When observing cells recruited to the lid in response to magnet exposure, cells labelled with higher MP concentrations appeared to aggregate over a smaller more defined area at the centre of the lid, rather than spread over a larger surface area as seen at the lower dose (2.5 µg/mL), possibly due to a stronger cell response at the point of highest field strength, but this 3D aggregation could not be accurately quantified using this 2D adherence assay.

To confirm the magnet-assisted migration response of cells labelled using particle concentrations previously shown to maintain cellular integrity, a further experimental model was used where MSCs were seeded onto a porous collagen scaffold and exposed to a magnetic field (Figure 5B). Cells labelled with 10 µg/mL MPs showed a significantly enhanced migratory capacity compared to unlabelled cells ($p < 0.001$).

SiMAG particles can also act as potential contrast agent which could allow post-delivery of cellular therapies for applications such as cartilage repair. In such approaches, an exogenously-expanded cell population would be delivered to a discrete site, where it needs to be retained in order to promote local tissue repair (El Haj *et al.*, 2014). The ability to image and monitor the implanted cells would allow monitoring of the therapy over time (Markides *et al.*, 2013). In order to identify the variables for cell tracking after labelling, the MRI visibility thresholds required in terms of particle concentration and cell number were established first in vitro and then in a pre-clinical large animal model of cell injection (Figure 6). When monitored in vitro (Figure 6A), SiMAG-labelled MSCs and chondrocytes were clearly detectable by MRI with significant dose-dependent contrast when using doses in the range of 10^4 to 0.5×10^6 cells. T_2^{eff} (Figure 6B) was seen to decrease with increasing cell numbers and particle concentrations corresponding to an increasing Fe content. A minimum visibility threshold of $5 \mu\text{g/mL}$ used with 5×10^5 labelled cells was identified in vitro. The detectability of MSC and chondrocyte cell populations after SiMAG labelling was found to be comparable in this model.

To further evaluate imaging capability in vivo, labelled cell populations were re-suspended in a collagen type I gel, a substrate widely used in cartilage tissue engineering (Deponti *et al.*, 2013), injected into a porcine knee model (Chiang *et al.*, 2005), and MR-imaged using specific T_2 weighted sequences (Figure 6C). In this clinically relevant model, the effect of particle concentration on the MRI detection was analyzed by implanting varying cell doses (10^4 , 10^5 and 5×10^6) of SiMAG-labelled cells to determine the visibility threshold, using 2 particle concentrations ($5 \mu\text{g/mL}$ and $10 \mu\text{g/mL}$). A combination of 10^5 cells labelled with $10 \mu\text{g/mL}$ was found to provide suitable contrast to enable graft detection by MRI within the host tissue.

4. Discussion

4.1. Efficient unassisted labelling of hMSCs:

Magnetic particles offer interesting properties for a multitude of biological and biomedical applications. Superparamagnetic iron oxide nanoparticles already demonstrate clinical efficacy and safety for MRI imaging (Colombo *et al.*, 2012) are now being investigated for more advanced theranostic applications for cell tracking and manipulation (Hu *et al.*, 2014, Corot *et al.*, 2006). MRI agents are developed to be bio-inert in order to minimise interaction with the cells within the body. Conversely, cell labelling agents must interact with the cell of interest to enable labelling without impacting upon its normal function. Thus, characterization of cell-MP interactions needs to be thoroughly assessed for advanced applications in cell-based therapies. In this study, we evaluated the suitability of commercially available 1 μm silica coated particles as a non-toxic labelling agent for cell tracking and manipulation towards both *in vitro* and *in vivo* applications.

24 h-incubation of MSCs with MPs was found to allow efficient labelling of the cell population, with over 95 % of cells labelled at 10 $\mu\text{g/mL}$ as measured by flow cytometry. This is in line with previous publications describing near 100% cell labelling using visual inspection following Prussian Blue staining or iron measurements (Markides *et al.*, 2013, Balakumaran *et al.*, 2010, Kostura *et al.*, 2004, Liu *et al.*, 2011, Pawelczyk *et al.*, 2006). A dose of 10 $\mu\text{g/mL}$ was selected as standard labelling concentration for MSCs, which was comparable to other reports (7 $\mu\text{g/mL}$ (Liu *et al.*, 2011), 25 $\mu\text{g/mL}$ (Kostura *et al.*, 2004)).

Cell labelling experiments demonstrated rapid uptake of MPs into MSCs, resulting in efficient cell labelling without the need for an added chemical carrier. Previous studies have suggested stem cell populations may benefit from assisted MP uptake through cellular targeting (Lewin *et al.*, 2000) or the use of transfection agents including polyethylenimine, protamine sulfate and polylysine (Balakumaran *et al.*, 2010, Kosture *et al.*, 2004, England *et al.*, 2013, Schafer *et al.*, 2010, Jing *et al.*, 2008, Arbab *et al.*, 2004). Interestingly, past reports have mentioned inefficient uptake by rat MSCs (Jing *et al.*, 2008), and undetectable uptake with human MSCs (Kostura *et al.*, 2004) when different particles were used alone. In contrast, our results confirm highly efficient uptake of the SiMAG particles in the absence of any additional facilitator, in line with observations carried out in other stem cell populations (Chen *et al.*, 2013). Particle surface modifications influence the characteristics of size, charge, toxicity and degradability of the particle (Li *et al.*, 2013), and have previously been reported to influence particle-cell interactions (Gupta and Gupta 2005, Zhao *et al.*, 2011, Sakhtianchi *et al.*, 2013). The SiMAG particles used here are silanol-coated, presenting an activated Si-OH surface arrangement. One of the main benefits from the silanol surface is a high colloidal suspension stability even in high volume fractions, through pH changes and electrolyte disturbances (Mulvaney *et al.*, 2000), all of which are likely to occur to some degree during application in a physiological environment. When silanol-coated MPs come into contact with the membrane, their association with the phosphatidyl choline rich regions of the membrane (Zhao *et al.*, 2011) is thought to elicit a membrane wrapping effect as other regions associate with the rigid curvature of the silanol surface. The subsequent entry of the MPs is dependent upon the energy released through the exothermic membrane wrapping effect and the energy required

to bend the membrane around the MP completely. In this situation, the dense nature of these MPs is considered to decrease the energy required for deformation of the membrane, thus facilitating engulfment (Zhao *et al.*, 2011), as suggested by the report that larger MPs are more thermo-dynamically favourable for endocytosis (Slowing *et al.*, 2009). Surface properties of the particles may also influence their interaction with natural proteins from serum (Wiogo *et al.*, 2011). Data presented here further demonstrate that the presence of serum diminishes SiMAG particle labelling in a dose dependant manner, potentially due to diminished accessibility of the surface silanol groups to the membrane, in line with previous reports supporting cell loading in serum-free conditions (Wilhelm and Gazeau 2008).

The efficient uptake of the SiMAG particles allowed labelled stem cell populations to be monitored both through their iron content and fluorescent analysis techniques. Particles appeared to cross the extracellular membrane, possibly through membrane wrapping and engulfment as previously described for silica particles (Zhao *et al.*, 2011), although the exact nature of this process requires further examination. Once inside the cell, particles accumulated at a central location inside endosome-like structures proximal to the nucleus, and no particle was observed inside the nuclear space, likely due to their micron size and contrary to what has been reported for particles below 70nm (Chen and von Mikecz 2005). Such intracellular particle distribution has previously been observed in MSCs (Neuberger *et al.*, 2005, Chang *et al.*, 2012) and other cell types (Wilhelm and Gazeau 2008, Robert *et al.*, 2010, Sun *et al.*, 2012).

4.2. Cellular compatibility:

Whilst previous studies have described the use of different particle types for cell labelling, few have focused on the potential implications of MP labelling for MSC cell health and function. Among these, most reports have investigated the biocompatibility of smaller MPs used with an auxiliary labelling reagent (Balakumaran *et al.*, 2010, Arbab *et al.*, 2004). Here, the suitability of SiMAG labelling for human MSCs was carefully examined through a range of parameters reflecting the integrity and cell health of labelled MSCs. Previously published studies on MP cytocompatibility have largely relied on the assessment of cell morphology combined to MTT/MTS assays, however these have demonstrated questionable reliability for particle and nano-materials studies (Laaksonen *et al.*, 2007). A resazurin-based metabolic measurement was therefore selected here, and indicated a slight increase in metabolic activity after particle labelling at low doses of particle uptake. This mild effect, which has been mentioned in different experimental conditions, could be linked to homeostatic mechanisms increasing lipid membrane synthesis in the cell to compensate for extracellular membrane disturbance associated with particle internalisation (McNeil and Steinhardt 1997, Kowalski *et al.*, 1972). Similarly, MSC surface marker expression analyzed before and after labelling showed that both primary and established MSCs retained their cell identity (Dominici *et al.*, 2006). This matches observations reported for different models and labelling conditions, which reported no significant change in MSCs (Balakumaran *et al.*, 2010), and similar stable marker expression in hematopoietic stem cell populations (Arbab *et al.*, 2004).

Although previous studies have suggested to good MP cytocompatibility for cell cultures (Li *et al.*, 2013, Budde and Frank 2009, Heymer *et al.*, 2008), some observations using small size MPs (60nm) have described changes in MSC

migration, colony-formation efficiency and even differentiation after particle labelling (Schafer *et al.*, 2009). Similar MP concentrations have also been reported to cause significant toxicity on neuronal and glial cells, while they did not appear to affect other cell types such as cardiomyogenic and pancreatic cells (Laurent *et al.*, 2012, Mahmoudi *et al.*, 2011). It is therefore important to evaluate the toxicity of each MP labelling protocol to be used in the target cell model for the application considered. MP-related toxicity may arise from the leaching of ions from metal core and the biodegradation polymer coating, which could cause oxidative stress (Kim *et al.*, 2011) through the leaching of metal ions from the core, or the release of oxidants by enzymatic degradation of the MPs (Mahmoudi *et al.*, 2012). Although iron can be metabolised in the human body (Henning *et al.*, 2009, Bulte *et al.*, 2009, Berry 2005, Kim *et al.*, 2010, Ju *et al.*, 2006), high quantities of Fe can impair viability and normal cell function (Li *et al.*, 2013, He *et al.*, 2007), underlining the need for a suitable balance between high Fe incorporation and safe cell function. Particle concentration ranging from 2.8- 400 µg/mL have been reportedly used for in vivo tracking.(Kim *et al.*, 2010, Jing *et al.*, 2008, He *et al.*, 2007, Kim *et al.*, 2010, Farrell *et al.*, 2009). The particle concentration chosen for this study (10 µg/mL), which was selected within the lower end of this range, showed no significant effect on cell viability or on the level of DNA damage in the MSC population as measured by the comet assay. This was true even for higher concentrations (100 µg/mL), and is in line with other studies which have shown low toxicity of both Fe₃O₄ and Fe₂O₃ –based particles (Karlsson *et al.*, 2009).

In addition to preserving the health of labelled cell populations for future cell therapies, maintaining their functionality is equally critical if they are to deliver a therapeutic effect. Reports published to date provided mixed results for the impact of

MPs on MSC differentiation. While a majority of studies reported no significant change based on histological or molecular assays, some negative effects on chondrogenesis have been observed (Kostura *et al.*, 2004, Bulte *et al.*, 2004). To examine the suitability of SiMAG-labelled MSCs to fulfil a therapeutic role, we examined their ability to differentiate into osteogenic, adipogenic and chondrogenic lineages, and found it to be maintained when examined both qualitatively and quantitatively. Bone nodules and lipid droplets were present in their respective cultures, with no statistically significant difference between unlabelled and labelled cell populations. Chondrogenic differentiation yielded micromass pellets demonstrating positive staining of glycosaminoglycans (GAGs) for both control and MP-labelled cultures. Closer examination revealed an increase in GAGs measured in MP-labelled pellets compared with the unlabelled samples, which could be due to more efficient centrifugal aggregation of the MP-labelled cells, as observed in our culture, since this is an important experimental parameter for the establishment of micromass cultures.

4.3. Control of target cell populations:

The possible dilution of the particle load by either exocytosis or cell division represents an inherent limitation of MPs and MRI based tracking in cell-based therapies, which could be of concern in long term animal studies. MSC labelling was detected here during a 7-day period in the case of dividing cell populations, beyond which the intracellular particle concentrations returned to control levels. This however was not solely dependent upon cell division as previously observed with smaller particles (Wilhelm and Gazeau 2008, Kim *et al.*, 2012), since non-dividing

populations also demonstrated particle loss albeit at a reduced rate. Arrested cells still demonstrated around 30% labelling 7 days post labelling, suggesting the occurrence of particle release or biodegradation in addition to mitotic dilution. Particle loss has been described as size-dependent, with smaller particles reportedly exocytosed at a faster rate than larger particles (Sakhtianchi *et al.*, 2013). Interestingly, this would fit with the observation of MP-labelled mouse MSCs implanted subcutaneously showing halving of the MRI signal over 3 days, and over one third of the initial signal detected by day 7 (Liu *et al.*, 2011). Berman *et al.* suggested particle decrease to be an indicator of viable cells, as non-viable cells may also retain the particles due to an inability to divide or actively exocytose (Berman *et al.*, 2011).

It is unclear whether magnetic labelling of MSCs may be associated with particle loss *in vivo*, and whether this may lead to subsequent unspecific labelling through secondary particle uptake by an unintended population. Results from our co-culture model combining labelled and unlabelled MSCs showed that the gradual loss of particles from a labelled cell population did not result in any significant uptake by neighbouring unlabelled populations. This suggested that transfer of particles either directly or indirectly through release into media is not occurring at a population level. This absence of apparent secondary particle uptake may be due to the presence of protein coronas on released particles obstructing the surface silanol groups from associating with the membrane (Zhao *et al.*, 2011, Foldbjerg *et al.*, 2013), which could decrease subsequent binding and cell internalisation. This may represent a long term experimental and safety benefit ensuring limiting possible leakage of the label from the target cells to unrelated cell populations *in vivo*. Particles released *in vivo* may furthermore be phagocytosed by macrophages, a process typically more

efficient for larger particles such as the ones used here than for smaller ones (Burtea *et al.*, 2008). This would further reduce the amount of released particles available for secondary uptake and limit the putative unspecific labelling of surrounding tissues.

The use of MRI for cell-based therapies bares a dual purpose. Not only can it precisely image the anatomical damage site and track implanted cells, but it can also evaluate the extent of the repair process at the damage site (Beckmann *et al.*, 2003, Henderson *et al.*, 2003). It is therefore important to analyze the extent to which implanted cell populations could be detected within anatomical structures in a realistic clinical model, such as the porcine knee model presented here, which offers dimensions in line with that of human tissue. Implantation of SiMAG-labelled cells generated significant contrast within this system and was clearly detected against anatomical structures. The visibility threshold of SiMAG-labelled cells using a 1.5 T scanner was found to be in agreement with the threshold established *ex vivo* (10^5 cells labelled with 5-10 $\mu\text{g}/\text{mL}$). These values are compatible with published studies varying from single cell detection with 11.7T scanning and micron sized particles (Bulte and Kraitchman 2004, Li *et al.*, 2009), to the detection of 1×10^6 cells labelled with 12 $\mu\text{g}/\text{mL}$ using a 3T machine (Chen *et al.*, 2012). Results presented here thus confirm that SiMAG-based MSC labelling can meet technical criteria outlined for use in preclinical studies (Frank *et al.*, 2004).

5. Conclusions

Beyond imaging, magnetic particles are widely exploited in separation techniques for cell suspensions (Plouffe *et al.*, 2015). *In vitro* experiments carried out in this study confirm their use can be applied to the spatial control of cell populations. Contactless magnetic control of cell movement can further enhance patterning and seeding

procedures both for 2D culture and for 3D tissue engineered scaffolds (Robert *et al.*, 2010, Yanai *et al.*, 2012). Although *ex vivo* models have not reported consistent magnetically-driven migration (Schafer *et al.*, 2010), possibly due to variations in the particles and magnets used, such targeting approaches may open novel therapeutic applications using permanent magnet, electromagnet or MR-assisted cell delivery (El Haj *et al.*, 2012, Robert *et al.*, 2010, Vanecek *et al.*, 2012, Riegler *et al.*, 2010).

Emerging MSC therapies such as Prochymal currently involve the use of high cell doses (in excess of 10^8 cells) (Hare *et al.*, 2009), which may in the future be reduced through improved cell delivery strategies such as magnetically-assisted cell targeting, to reduce the dose needed. Careful prior assessments of the particle uptake, retention profile and biological responses associated with such strategies will be critical to ensure the safe development of enhanced targeting therapies. A recent report introducing the *in vivo* labelling of stem cells prior to their harvest and allogeneic use (Khurana *et al.*, 2013) underlined the requirement to ascertain the cellular innocuousness of MPs for the targeted population. The data presented in our study supports the suitability of $1\mu\text{m}$ SiMAG superparamagnetic iron oxide particles as a possible cell tracking and cell manipulation agent for stem cell-based therapies. Their large size and coating properties facilitating uptake, biocompatibility and visibility for MRI make them favourable candidates for further *in vivo* preclinical research into advanced tissue engineering approaches.

Acknowledgements

We are grateful to Dr Claire Seedhouse and Sahar Aldosari for their advice with the Comet assay, and to Dr David Onion and Denise Christie for their expert help with flow cytometry and TEM respectively. This study was supported by an EPSRC Doctoral Training Centre scholarship (RH and HM), the ARUK Centre for Tissue Engineering and the University of Nottingham.

References

- Arbab AS, Yocum GT, Kalish H, Jordan EK, Anderson SA, Khakoo AY, Read EJ, Frank JA. 2004, Efficient magnetic cell labeling with protamine sulfate complexed to ferumoxides for cellular MRI, *Blood*, **104**(4): 1217-1223.
- Balakumaran A, Pawelczyk E, Ren J, Sworder B, Chaudhry A, Sabatino M, Stroncek D, Frank JA, Robey PG. 2010, Superparamagnetic iron oxide nanoparticles labeling of bone marrow stromal (mesenchymal) cells does not affect their "stemness", *PLoS One*, **5**(7): e11462.
- Beckmann N, Falk R, Zurbrügg S, Dawson J, Engelhardt P. 2003, Macrophage infiltration into the rat knee detected by MRI in a model of antigen-induced arthritis, *Magn Reson Med*, **49**(6): 1047-1055.
- Berman SC, Galpoththawela C, Gilad AA, Bulte JW, Walczak P. 2011, Long-term MR cell tracking of neural stem cells grafted in immunocompetent versus immunodeficient mice reveals distinct differences in contrast between live and dead cells, *Magn Reson Med*, **65**(2): 564-574.
- Berry CC. 2005, Possible exploitation of magnetic nanoparticle–cell interaction for biomedical applications, *J Mater Chem*, **15**: 543-547.
- Budde MD, Frank JA. 2009, Magnetic tagging of therapeutic cells for MRI, *J Nucl Med*, **50**(2): 171-174.
- Bulte JW. 2009, In vivo MRI cell tracking: clinical studies, *AJR*, **193**(2): 314-325.
- Bulte JW, Kraitchman DL. 2004, Iron oxide MR contrast agents for molecular and cellular imaging, *NMR Biomed*, **17**(7): 484-499.
- Bulte JW, Kraitchman DL, Mackay AM, Pittenger MF 2004, Chondrogenic differentiation of mesenchymal stem cells is inhibited after magnetic labeling with ferumoxides, *Blood*, **105**(2): 473.
- Burtea C, Laurent S, Vander Elst L, Muller R. 2008, Contrast agents: magnetic resonance, *Handb Exp Pharmacol*, **185**: 135-165.
- Caplan AI. 2007, Adult mesenchymal stem cells for tissue engineering versus regenerative medicine, *J Cell Physiol*, **213**(2): 341-347.
- Caplan AI, Dennis JE. 2006, Mesenchymal stem cells as trophic mediators, *J Cell Biochem*, **98**(5): 1076-1084.
- Chang YK, Liu YP, Ho JH, Hsu SC, Lee OK. 2012, Amine-surface-modified superparamagnetic iron oxide nanoparticles interfere with differentiation of human mesenchymal stem cells, *J Orthop Res*, **30**(9): 1499-1506.

- Chen CC, Ku MC, D M J, Lai JS, Hueng DY, Chang C. 2013, Simple SPION incubation as an efficient intracellular labeling method for tracking neural progenitor cells using MRI, *PLoS One*, **8**(2): e56125.
- Chen J, Wang F, Zhang Y, Jin X, Zhang L, Feng Y, Lin X, Yang L. 2012, In vivo tracking of superparamagnetic iron oxide nanoparticle labeled chondrocytes in large animal model, *Ann Biomed Eng*, **40**(12): 2568-2578.
- Chen M, von Mikecz A. 2005, Formation of nucleoplasmic protein aggregates impairs nuclear function in response to SiO₂ nanoparticles, *Exp Cell Res*, **305**(1): 51-62.
- Chiang H, Kuo TF, Tsai CC, Lin MC, She BR, Huang YY, Lee HS, Shieh CS, Chen MH, Ramshaw JA, Werkmeister JA, Tuan RS, Jiang CC. 2005, Repair of porcine articular cartilage defect with autologous chondrocyte transplantation, *J Orthop Res*, **23**(3): 584-593.
- Colombo M, Carregal-Romero S, Casula MF, Gutiérrez L, Morales MP, Böhm IB, Heverhagen JT, Prospero D, Parak WJ. 2012, Biological applications of magnetic nanoparticles, *Chem Soc Rev*, **41**(11): 4306-4334.
- Corot C, Robert P, Idée JM, Port M. 2006, Recent advances in iron oxide nanocrystal technology for medical imaging, *Adv Drug Deliv Rev*, **58**(14): 1471-1504.
- Deponti D, Di Giancamillo A, Scotti C, Peretti GM, Martin I. 2013, Animal models for meniscus repair and regeneration, *J Tissue Eng Regen Med*, **9**(5): 512-27.
- Dominici M, Le Blanc K, Mueller I, Slaper-Cortenbach I, Marini F, Krause D, Deans R, Keating A, Prockop Dj, Horwitz E. 2006, Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement, *Cytotherapy*, **8**(4): 315-317.
- El Haj AJ, Glossop JR, Sura HS, Lees MR, Hu B, Wolbank S, van Griensven M, Redl H, Dobson J. 2012, An in vitro model of mesenchymal stem cell targeting using magnetic particle labelling, *J Tissue Eng Regen Med*, **9**(6): 724-733.
- England TJ, Bath PM, Abaei M, Auer D, Jones DR. 2013, Hematopoietic stem cell (CD34+) uptake of superparamagnetic iron oxide is enhanced by but not dependent on a transfection agent, *Cytotherapy*, **15**(3): 384-390.
- Farrell E, Wielopolski P, Pavljasevic P, Kops N, Weinans H, Bernsen MR, van Osch GJ. 2009, Cell labelling with superparamagnetic iron oxide has no effect on chondrocyte behaviour, *Osteoarthritis and Cartilage*, **17**(7): 961-967.
- Fayol D, Frasca G, Le Visage C, Gazeau F, Luciani N, Wilhelm C. 2013, Use of magnetic forces to promote stem cell aggregation during differentiation, and cartilage tissue modeling, *Adv Mater*, **25**(18): 2611-2616.
- Foldbjerg R, Wang J, Beer C, Thorsen K, Sutherland DS, Autrup H. 2013, Biological effects induced by BSA-stabilized silica nanoparticles in mammalian cell lines, *Chem Biol Interact*, **204**(1): 28-38.
- France LA, Scotchford CA, Grant DM, Rashidi H, Popov AA, Sottile V. 2014, Transient serum exposure regimes to support dual differentiation of human mesenchymal stem cells, *J Tissue Eng Regen Med*, **8**(8): 652-663.
- Frank JA, Anderson SA, Kalsih H, Jordan EK, Lewis BK, Yocum GT, Arbab AS. 2004, Methods for magnetically labeling stem and other cells for detection by in vivo magnetic resonance imaging, *Cytotherapy*, **6**(6): 621-625.
- Gilchrist RK, Medal R, Shorey WD, Hanselman RC, Parrott JC, Taylor CB. 1957, Selective inductive heating of lymph nodes. *Ann Surg*. **146**(4): 596-606.
- Gupta AK, Gupta M. 2005, Synthesis and surface engineering of iron oxide nanoparticles for biomedical applications, *Biomaterials*, **26**(18): 3995-4021.

- Hare JM, Traverse JH, Henry TD, Dib N, Strumpf RK, Schulman SP, Gerstenblith G, DeMaria AN, Denktas AE, Gammon RS, Hermiller JB Jr, Reisman MA, Schaer GL, Sherman W. 2009, A randomized, double-blind, placebo-controlled, dose-escalation study of intravenous adult human mesenchymal stem cells (prochymal) after acute myocardial infarction, *J Am Coll Cardiol*, **54**(24): 2277-2286.
- Hayman DM, Blumberg TJ, Scott CC, Athanasiou KA. 2006, The effects of isolation on chondrocyte gene expression, *Tissue Eng*, **12**(9): 2573-2581.
- He G, Zhang H, Wei H, Wang Y, Zhang X, Tang Y, Wei Y, Hu S. 2007, In vivo imaging of bone marrow mesenchymal stem cells transplanted into myocardium using magnetic resonance imaging: a novel method to trace the transplanted cells, *International Journal of Cardiology*, **114**(1): 4-10.
- Henderson IJ, Tuy B, Connell D, Oakes B, Hettwer WH. 2003, Prospective clinical study of autologous chondrocyte implantation and correlation with MRI at three and 12 months, *J Bone Joint Surg Br*, **85**(7): 1060-1066.
- Henning TD, Sutton EJ, Kim A, Golovko D, Horvai A, Ackerman L, Sennino B, McDonald D, Lotz J, Daldrup-Link HE. 2009, The influence of ferucarbotran on the chondrogenesis of human mesenchymal stem cells, *Contrast Media & Molecular Imaging*, **4**(4): 165-173.
- Henstock JR, Rotherham M, Rashidi H, Shakesheff KM, El Haj AJ. 2014, Remotely Activated Mechanotransduction via Magnetic Nanoparticles Promotes Mineralization Synergistically With Bone Morphogenetic Protein 2: Applications for Injectable Cell Therapy, *Stem Cells Transl Med*, **3**(11): 1363-1374.
- Heymer A, Haddad D, Weber M, Gbureck U, Jakob PM, Eulert J, Nöth U. 2008, Iron oxide labelling of human mesenchymal stem cells in collagen hydrogels for articular cartilage repair, *Biomaterials*, **29**(10): 1473-1483.
- Hofmann-Antenbrink M, Hofmann H, Montet X. 2010, Superparamagnetic nanoparticles - a tool for early diagnostics, *Swiss Med Wkly*, **140**: w13081.
- Hu B, Dobson J, El Haj AJ. 2014, Control of smooth muscle α -actin (SMA) up-regulation in HBMSCs using remote magnetic particle mechano-activation, *Nanomedicine*, **10**(1): 45-55.
- Jasmin, Torres AL, Nunes HM, Passipieri JA, Jelicks LA, Gasparetto EL, Spray DC, Campos de Carvalho AC, Mendez-Otero R. 2011, *J Nanobiotechnology*, **9**: 4.
- Jing XH, Yang L, Duan XJ, Xie B, Chen W, Li Z, Tan HB. 2008, In vivo MR imaging tracking of magnetic iron oxide nanoparticle labeled, engineered, autologous bone marrow mesenchymal stem cells following intra-articular injection, *Joint Bone Spine*, **75**(4): 432-438.
- Jing Y, Mal N, Williams PS, Mayorga M, Penn MS, Chalmers JJ, Zborowski M. 2008, Quantitative intracellular magnetic nanoparticle uptake measured by live cell magnetophoresis, *FASEB J*, **22**(12): 4239-4247.
- Ju S, Teng G, Zhang Y, Ma M, Chen F, Ni Y. 2006, In vitro labeling and MRI of mesenchymal stem cells from human umbilical cord blood, *Magnetic Resonance Imaging*, **24**(5): 611-617.
- Karlsson HL, Gustafsson J, Cronholm P, Möller L. 2009, Size-dependent toxicity of metal oxide particles--a comparison between nano- and micrometer size, *Toxicol Lett*, **188**(2): 112-118.
- Khurana A, Chapelin F, Beck G, Lenkov OD, Donig J, Nejadnik H, Messing S, Derugin N, Chan RC, Gaur A, Sennino B, McDonald DM, Kempen PJ, Tikhomirov GA, Rao J, Daldrup-Link HE. 2013, Iron administration before

- stem cell harvest enables MR imaging tracking after transplantation, *Radiology*, **269**(1): 186-197.
- Kim HS, Oh SY, Joo HJ, Son KR, Song IC, Moon WK. 2010, The effects of clinically used MRI contrast agents on the biological properties of human mesenchymal stem cells, *NMR Biomed*, **23**(5): 514-522.
- Kim JA, Åberg C, Salvati A, Dawson KA. 2012, Role of cell cycle on the cellular uptake and dilution of nanoparticles in a cell population, *Nature Nanotechnology*, **7**(1): 62-68.
- Kim T, Momin E, Choi J, Yuan K, Zaidi H, Kim J, Park M, Lee N, McMahon MT, Quinones-Hinojosa A, Bulte JW, Hyeon T, Gilad AA. 2011, Mesoporous silica-coated hollow manganese oxide nanoparticles as positive T1 contrast agents for labeling and MRI tracking of adipose-derived mesenchymal stem cells, *J Am Chem Soc*, **133**(9): 2955-2961.
- Kim TH, Kim JK, Shim W, Kim SY, Park TJ, Jung JY. 2010, Tracking of transplanted mesenchymal stem cells labeled with fluorescent magnetic nanoparticle in liver cirrhosis rat model with 3-T MRI, *Magnetic Resonance Imaging*, **28**(7): 1004-1013.
- Kobayashi T. 2011, Cancer hyperthermia using magnetic nanoparticles, *Biotechnol J*, **6**(11): 1342-1347.
- Kostura L, Kraitchman DL, Mackay AM, Pittenger MF, Bulte JW. 2004, Feridex labeling of mesenchymal stem cells inhibits chondrogenesis but not adipogenesis or osteogenesis, *NMR Biomed*, **17**(7): 513-517.
- Kowalski K, Babiarczyk D, Sato S, Burke G. 1972, Stimulatory effects of induced phagocytosis on the function of isolated thyroid cells, *J Clin Invest*, **51**(11): 2808-2819.
- Kunzmann A, Andersson B, Vogt C, Feliu N, Ye F, Gabrielsson S, Toprak MS, Buerki-Thurnherr T, Laurent S, Vahter M, Krug H, Muhammed M, Scheynius A, Fadeel B. 2011, Efficient internalization of silica-coated iron oxide nanoparticles of different sizes by primary human macrophages and dendritic cells, *Toxicol Appl Pharmacol*, **253**(2): 81-93.
- Laaksonen T, Santos H, Vihola H, Salonen J, Riikonen J, Heikkilä T, Peltonen L, Kumar N, Murzin DY, Lehto VP, Hirvonen J. 2007, Failure of MTT as a toxicity testing agent for mesoporous silicon microparticles, *Chem Res Toxicol*, **20**(12): 1913-1918.
- Laurent S, Burtea C, Thirifays C, Häfeli UO, Mahmoudi M. 2012, Crucial ignored parameters on nanotoxicology: the importance of toxicity assay modifications and "cell vision", *Plos One*, **7**(1): e29997.
- Le Blanc K, Tammik C, Rosendahl K, Zetterberg E, Ringdén O. 2003, HLA expression and immunologic properties of differentiated and undifferentiated mesenchymal stem cells, *Exp Hematol*, **31**(10): 890-896.
- Lewin M, Carlesso N, Tung CH, Tang XW, Cory D, Scadden DT, Weissleder R. 2000, Tat peptide-derivatized magnetic nanoparticles allow in vivo tracking and recovery of progenitor cells, *Nat Biotechnol*, **18**(4): 410-414.
- Li L, Jiang W, Luo K, Song H, Lan F, Wu Y, Gu Z. 2013, Superparamagnetic iron oxide nanoparticles as MRI contrast agents for non-invasive stem cell labeling and tracking, *Theranostics*, **3**(8): 595-615.
- Li Calzi S, Kent DL, Chang KH, Padgett KR, Afzal A, Chandra SB, Caballero S, English D, Garlington W, Hiscott PS, Sheridan CM, Grant MB, Forder JR. 2009, Labeling of stem cells with monocrySTALLINE iron oxide for tracking and localization by magnetic resonance imaging, *Microvasc Res*, **78**(1): 132-139.

- Liu G, Wang Z, Lu J, Xia C, Gao F, Gong Q, Song B, Zhao X, Shuai X, Chen X, Ai H, Gu Z. 2011, Low molecular weight alkyl-polycation wrapped magnetite nanoparticle clusters as MRI probes for stem cell labeling and in vivo imaging, *Biomaterials*, **32**(2): 528-537.
- Mahmoudi M, Laurent S, Shokrgozar MA, Hosseinkhani M. 2011, Toxicity evaluations of superparamagnetic iron oxide nanoparticles: cell "vision" versus physicochemical properties of nanoparticles, *Acs Nano*, **5**(9): 7263-7276.
- Mahmoudi M, Hofmann H, Rothen-Rutishauser B, Petri-Fink A. 2012, Assessing the in vitro and in vivo toxicity of superparamagnetic iron oxide nanoparticles, *Chem Rev*, **112**(4): 2323-2338.
- Mahmoudi M, Sant S, Wang B, Laurent S, Sen T. 2011, Superparamagnetic iron oxide nanoparticles (SPIONs): development, surface modification and applications in chemotherapy, *Adv Drug Deliv Rev*, **63**(1-2): 24-46.
- Markides H, Kehoe O, Morris RH, El Haj AJ. 2013, Whole body tracking of superparamagnetic iron oxide nanoparticle-labelled cells--a rheumatoid arthritis mouse model. *Stem Cell Res Ther*, **4**(5): 126.
- McBride AA, Price DN, Lamoureux LR, Elmaoued AA, Vargas JM, Adolphi NL, Muttill P. 2013, Preparation and characterization of novel magnetic nano-in-microparticles for site-specific pulmonary drug delivery, *Mol Pharm*, **10**(10): 3574-3581.
- McNeil PL, Steinhardt RA. 1997, Loss, restoration, and maintenance of plasma membrane integrity, *J Cell Biol*, **137**(1): 1-4.
- Mulvaney P, Liz-Marzan LM, Giersig M, Ung T. 2000, Silica encapsulation of quantum dots and metal clusters, *J Mater Chem*, **10**: 1259-1270.
- Neuberger T, Schopf B, Hofmann H, Hofmann M, von Rechenberg B. 2005, Superparamagnetic nanoparticles for biomedical applications: Possibilities and limitations of a new drug delivery system, *J Magn Magn Mater*, **293**: 483-496.
- Nieto A, Cabrera CM, Catalina P, Cobo F, Barnie A, Cortés JL, Barroso del Jesus A, Montes R, Concha A. 2007, Effect of mitomycin-C on human foreskin fibroblasts used as feeders in human embryonic stem cells: immunocytochemistry MIB1 score and DNA ploidy and apoptosis evaluated by flow cytometry, *Cell Biol Int*, **31**(3): 269-278.
- Okamoto T, Aoyama T, Nakayama T, Hosaka T, Nishijo K, Nakamura T, Kiyono T, Toguchida JI. 2002, Clonal heterogeneity in differentiation potential of immortalized human mesenchymal stem cells, *Biochem Biophys Res Commun*, **295**: 354-361.
- Pawelczyk E, Arbab AS, Pandit S, Hu E, Frank JA. 2006, Expression of transferrin receptor and ferritin following ferumoxides-protamine sulfate labeling of cells: implications for cellular magnetic resonance imaging, *NMR Biomed*, **19**(5): 581-592.
- Peister A, Mellad JA, Wang M, Tucker HA, Prockop DJ. 2004, Stable transfection of MSCs by electroporation, *Gene Ther*, **11**(2): 224-228.
- Pickard MR, Barraud P, Chari DM. 2011, The transfection of multipotent neural precursor/stem cell transplant populations with magnetic nanoparticles, *Biomaterials*, **32**(9): 2274-2284.
- Plouffe BD, Murthy SK, Lewis LH. 2015, Fundamentals and application of magnetic particles in cell isolation and enrichment: a review, *Rep Prog Phys*, **78**(1): 016601.

- Pooley RA. 2005, Fundamental physics of MR imaging, *Radiographics*, **25**(4): 1087-1099.
- Quarto R, Mastrogiacomo M, Cancedda R, Kutepov SM, Mukhachev V, Lavroukov A, Kon E, Marcacci M. 2001, Repair of large bone defects with the use of autologous bone marrow stromal cells, *N Engl J Med*, **344**(5): 385-386.
- Riegler J, Wells JA, Kyrtatos PG, Price AN, Pankhurst QA, Lythgoe MF. 2010, Targeted magnetic delivery and tracking of cells using a magnetic resonance imaging system, *Biomaterials*, **31**(20): 5366-5371.
- Robert D, Fayol D, Le Visage C, Frasca G, Brulé S, Ménager C, Gazeau F, Letourneur D, Wilhelm C. 2010, Magnetic micro-manipulations to probe the local physical properties of porous scaffolds and to confine stem cells, *Biomaterials*, **31**(7): 1586-1595.
- Robert D, Nguyen TH, Gallet F, Wilhelm C. 2010, In vivo determination of fluctuating forces during endosome trafficking using a combination of active and passive microrheology, *PLoS One*, **5**(4): e10046.
- Robards WA, Wilson AJ. 1993, *Procedures in electron microscopy*, John Wiley and Sons, Chichester, UK
- Sakhtianchi R, Minchin RF, Lee KB, Alkilany AM, Serpooshan V, Mahmoudi M. 2013, Exocytosis of nanoparticles from cells: role in cellular retention and toxicity, *Adv Colloid Interface Sci*, **201**-202: 18-29.
- Schäfer R, Bantleon R, Kehlbach R, Siegel G, Wiskirchen J, Wolburg H, Kluba T, Eibofner F, Northoff H, Claussen CD, Schlemmer HP. 2010, Functional investigations on human mesenchymal stem cells exposed to magnetic fields and labeled with clinically approved iron nanoparticles, *BMC Cell Biol*, **11**: 22.
- Schäfer R, Kehlbach R, Müller M, Bantleon R, Kluba T, Ayturan M, Siegel G, Wolburg H, Northoff H, Dietz K, Claussen CD, Wiskirchen J. 2009, Labeling of human mesenchymal stromal cells with superparamagnetic iron oxide leads to a decrease in migration capacity and colony formation ability, *Cytotherapy*, **11**(1): 68-78.
- Seedhouse CH, Hunter HM, Lloyd-Lewis B, Massip AM, Pallis M, Carter GI, Grundy M, Shang S, Russell NH. 2006, DNA repair contributes to the drug-resistant phenotype of primary acute myeloid leukaemia cells with FLT3 internal tandem duplications and is reversed by the FLT3 inhibitor PKC412, *Leukemia*, **20**(12): 2130-2136.
- Sheng HH, Zhang GG, Cheung WH, Chan CW, Wang YX, Lee KM, Wang HF, Leung KS, Qin LL. 2007, Elevated adipogenesis of marrow mesenchymal stem cells during early steroid-associated osteonecrosis development, *J Orthop Surg Res*, **2**: 15.
- Slowing II, Wu CW, Vivero-Escoto JL, Lin VS. 2009, Mesoporous silica nanoparticles for reducing hemolytic activity towards mammalian red blood cells, *Small*, **5**(1): 57-62.
- Sun JH, Zhang YL, Nie CH, Qian SP, Yu XB, Xie HY, Zhou L, Zheng SS. 2012, In vitro labeling of endothelial progenitor cells isolated from peripheral blood with superparamagnetic iron oxide nanoparticles, *Mol Med Rep*, **6**(2): 282-286.
- Uccelli A. 2008, Adult stem cells for spinal cord injury: what types and how do they work?, *Cytotherapy*, **10**(6): 541-542.
- Vaněček V, Zablotskii V, Forostyak S, Růžička J, Herynek V, Babič M, Jendelová P, Kubinová S, Dejneka A, Syková E. 2012, Highly efficient magnetic targeting of mesenchymal stem cells in spinal cord injury, *Int J Nanomedicine*, **7**: 3719-3730.

- Yanai A, Häfeli UO, Metcalfe AL, Soema P, Addo L, Gregory-Evans CY, Po K, Shan X, Moritz OL, Gregory-Evans K. 2012, Focused magnetic stem cell targeting to the retina using superparamagnetic iron oxide nanoparticles, *Cell Transplant*, **21**(6): 1137-1148.
- Wang YH, Liu Y, Maye P, Rowe DW. 2006, Examination of mineralized nodule formation in living osteoblastic cultures using fluorescent dyes, *Biotechnol Prog*, **22**(6): 1697-1701.
- Wilhelm C, Gazeau F. 2008, Universal cell labelling with anionic magnetic nanoparticles, *Biomaterials*, **29**(22): 3161-3174.
- Wimpenny I, Markides H, El Haj AJ. 2012, Orthopaedic applications of nanoparticle-based stem cell therapies, *Stem Cell Res Ther*, **3**(2): 13.
- Wiogo HT, Lim M, Bulmus V, Yun J, Amal R. 2011, Stabilization of magnetic iron oxide nanoparticles in biological media by fetal bovine serum (FBS), *Langmuir*, **27**(2): 843-850.
- Zhao F, Zhao Y, Liu Y, Chang X, Chen C, Zhao Y. 2011, Cellular uptake, intracellular trafficking, and cytotoxicity of nanomaterials, *Small*, **7**(10): 1322-1337.
- Zhao Y, Sun X, Zhang G, Trewyn BG, Slowing II, Lin VS. 2011, Interaction of mesoporous silica nanoparticles with human red blood cell membranes: size and surface effects, *ACS Nano*, **5**(2): 1366-1375.

Figure Legend:

Figure 1: MSCs labelled with fluorescently labelled MPs analyzed using both flow cytometry and microscopy. (A) Flow cytometry analysis (left) and corresponding quantification (right) showing increased labelling with increasing MP concentrations ($***p < 0.001$, $****p < 0.0001$, $n=3$). (B) Fluorescence imaging of hMSCs labelled with 10 $\mu\text{g/mL}$ particles, showing MPs (yellow), cell outline (Phalloidin, green) and nuclear counterstain (Hoechst 33342, blue); bar: 25 μm . (C) Prussian blue staining highlighting internalised iron-rich MPs within the cell ; bar: 100 μm . (D) TEM imaging of MPs showing that internalised particles are contained within vesicles (arrowhead), which merge into larger vacuoles (arrow) near the nucleus (asterisk). Bar: 5 μm (top) and 10 μm (bottom). (E) Negative effect of serum concentration on the efficiency of MP uptake measured at 24 h after labelling ($*p < 0.05$, $**p < 0.01$, $****p < 0.0001$, $n=3$).

Figure 2: Kinetics of MSC particle retention after labelling with 10 $\mu\text{g/mL}$ particles. (A) Flow cytometry analysis of MPs in labelled cells up to 7 days after labelling showing gradual decrease in dividing cell populations (light grey), while particle dilution is reduced by mitomycin C-mediated inhibition of cell division (dark grey) ($*p < 0.05$, $****p < 0.0001$, $n=3$). (B) Fluorescence microscopy of phalloidin staining (green) with dapi counterstain (blue) showing rare particles (red) detected in cell processes. (C) Distribution of MPs between a labelled (population 1, gMSCs) and unlabelled (population 2, hMSCs) MSC population analyzed by flow cytometry over 7 days of co-culture, showing no evidence of secondary particle uptake. Statistical analysis showing labelling of population 1 between days 0-4 & 6 compared to day 7

but no statistically significant labelling present in population 2 on any of the days (**p < 0.001, n=2).

Figure 3: Cell integrity assessment after particle uptake. (A) MSC marker identity analysed by flow cytometry demonstrating no discernible change in hMSC marker expression following particle labelling (red, 10 µg/mL) compared to unlabelled cells (blue) and the isotype control (grey). (B) Metabolic activity assessed through a resazurin analogue (Presto Blue®) at 24 hours demonstrating no significant negative effect of particle uptake at therapeutic doses (up to 50 µg/mL), when compared to unlabelled controls and DMSO-mediated toxicity (**p < 0.01, ****p < 0.0001, n=3). (C) Cell membrane integrity assay showing stable membrane integrity 24 hours post labelling with MPs (10 µg/mL). Statistical significance calculated compared to DMSO-treated or fixed cells (****p < 0.0001 n=3), no statistically significant difference between treatment groups. (D) DNA integrity analyzed using the comet assay, showing no statistically significant DNA damage in labelled cells at 10 µg/mL and 100 µg/mL. Statistical significance between induced damage (positive control) and other conditions (****p < 0.0001, n=680), no significant difference between unlabelled and MP-labelled conditions.

Figure 4: MSC differentiation in the presence or absence of MPs. (A-C) Differentiation potential under standard culture medium (left panel) or differentiation treatment (right panel) of the hMSC populations towards osteogenic (A), adipogenic (B) and chondrogenic (C) lineages, monitored by von Kossa, Oil Red O and Alcian blue staining respectively. MP-labelled cell populations (10 µg/mL) were compared to unlabelled populations with no detectable decrease in differentiation in vitro. (D-E)

Quantitative assessment of osteogenic response performed at 7 days (alkaline phosphatase activity) and 14 days (Alizarin Red S extraction) showing statistically significant response to induction medium (grey bars) compared to untreated controls (black bars). (F) Adipogenic induction was measured using Oil Red O extraction demonstrating no statistically significant change in lipid accumulation at either concentration. (G) Chondrogenic response assessed using the DMMB assay normalised to DNA content showing increased GAGs in both unlabelled and labelled populations compared to their standard medium treated equivalents (**** $p < 0.0001$, $n = 5$).

Figure 5: Migration of SiMAG-labelled MSCs in vitro. (A) Hanging drops prepared with cells labelled with increasing MP concentrations were incubated in the presence or absence of magnets placed on the upper side of the lid. After 24 hours, surface areas covered by cells recruited to the surface of the lid (inserts) were imaged and measured (**** $p < 0.0001$, $n = 4$). (B) MSCs labelled with MPs (0, 1 and 10 $\mu\text{g/mL}$) over a 24 hour period within a collagen transwell system and exposed to a magnet for 24 hours. Migrated cells counted as the average of 5 fields of view on the underside of each transwell (** $p < 0.001$, $n = 3$).

Figure 6: MRI tracking of SiMAG-labelled hMSCs and chondrocytes. (A-B) Increasing MP concentrations (1, 5 and 10 $\mu\text{g/mL}$) and cell doses (10^5 cells, 5×10^5 cells) showing the MRI visibility threshold of labelled MSCs presented as a T_2^{eff} map (A) and corresponding T_2^{eff} plot (B). (C) Coronal DESS image of labelled chondrocytes implanted in a porcine knee joint (left condyle 10^5 cells, right condyle 5×10^5 cells) analyzed by MRI, using 5 $\mu\text{g/mL}$ (upper panel) and 10 $\mu\text{g/mL}$ (lower

panel) MP concentrations, showing hypointense regions of signal voids (yellow arrows). Red lines highlight the region of interest (ROI).

Supplementary Information:

Supplementary File 1: Time-lapse imaging of hMSC cells incubated in the presence of SiMAG MPs (10 µg/mL), capturing 2 frames/min for 18 hours.

Supplementary File 2: Long term marker expression in MSC cultures analyzed after 14 days of particle labelling. (A) Flow cytometry analysis of hMSCs (A) and pMSCs (B) demonstrating maintenance of MSC surface marker profile following repeated particle labelling (red, 10 µg/mL every three days) compared to unlabelled cells (blue) and the isotype control (grey).