

1 **Genome-wide association and functional studies identify a role for matrix-Gla protein in**
2 **osteoarthritis of the hand**

3
4 WOUTER DEN HOLLANDER^{1¶}, CINDY G. BOER^{2¶}, DEBBIE HART³, MICHELLE S. YAU^{4,5}, YOLANDE F.M. RAMOS¹,
5 SARAH METRUSTRY³, LINDA BROER², JORIS DEELEN^{1,6}, L. ADRIENNE CUPPLES⁷, FERNANDO RIVADENEIRA²,
6 MARGREET KLOPPENBURG⁸, MARJOLEIN PETERS², TIM D. SPECTOR³, ALBERT HOFMAN^{9,10}, P. ELINE SLAGBOOM¹,
7 ROB G.H.H. NELISSEN¹¹, ANDRÉ G. UITTERLINDEN^{2,9}, DAVID T. FELSON^{5,12}, ANNA M. VALDES¹³, INGRID
8 MEULENBELT^{1&*}, JOYCE J.B. VAN MEURS^{2&*}

9
10 ¹*Department of Medical Statistics and Bioinformatics, Section Molecular Epidemiology, Leiden University Medical*
11 *Center, Leiden, The Netherlands*

12 ²*Department of Internal Medicine, Genetic Laboratory, Erasmus Medical Center, Rotterdam, the Netherlands*

13 ³*Department of Twin Research and Genetic Epidemiology, King's College London, London, , UK*

14 ⁴*Institute for Aging Research, Hebrew SeniorLife, Beth Israel Deaconess Medical Center, Harvard Medical School,*
15 *Boston, MA, USA*

16 ⁵*Clinical Epidemiology Research and Training Unit, Boston University School of Medicine, Boston, MA, USA*

17 ⁶*Max Planck Institute for Biology of Ageing, Cologne, Germany*

18 ⁷*Department of Biostatistics, Boston University School of Public Health, Boston, MA, USA; The Framingham Heart*
19 *Study, Framingham, MA, USA*

20 ⁸*Department of Rheumatology, Leiden University Medical Center, Leiden, The Netherlands*

21 ⁹*Department of Epidemiology, Erasmus Medical Center, Rotterdam, The Netherlands*

22 ¹⁰*Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA, USA*

23 ¹¹*Department of Orthopedics, Leiden University Medical Center, Leiden, The Netherlands*

24 ¹²*Arthritis Research UK Epidemiology Unit, University of Manchester, Manchester, UK*

25 ¹³*School of Medicine, University of Nottingham, Nottingham, UK*

26
27 [¶]*These authors contributed equally to this work*

28 [&]*These authors jointly supervised this work*

29 ^{*}*Corresponding authors*

30 j.vanmeurs@erasmusmc.nl (JBJM)

31 i.meulenbelt@lumc.nl (IM)

32

33

34 **Abstract**

35 **Objective** Osteoarthritis (OA) is the most common form of arthritis and the leading cause of disability in
36 the elderly. Of all the joints, genetic predisposition is strongest for OA of the hand, however only few
37 genetic risk loci for hand OA have been identified. Our aim was to identify novel genes associated with
38 hand OA and examine the underlying mechanism.

39 **Methods** We performed a genome-wide association study of a quantitative measure of hand OA in
40 12,784 individuals (discovery:8,743, replication:4,011). Genome-wide significant signals were followed
41 up by analysing gene and allele specific expression in a RNA-sequencing dataset (n=96) of human
42 articular cartilage

43 **Results** We found two significantly associated loci in the discovery set: at chr12 ($P=3.5 \times 10^{-10}$) near the
44 MGP gene and at chr12 ($P=6.1 \times 10^{-9}$) near the CCDC91 gene. The DNA variant near the MGP-gene was
45 validated in three additional studies, which resulted in a highly significant association between the MGP-
46 variant and hand OA (rs4764133, $\text{Beta}_{\text{meta}}=0.83$, $P_{\text{meta}}=1.8 \times 10^{-15}$). This variant is high linkage
47 disequilibrium with a coding variant in *Matrix Gla-Protein (MGP)*, a vitamin K-dependent inhibitor of
48 cartilage calcification. Using RNA-sequencing data from human primary cartilage tissue (n=96), we
49 observed that the hand OA MGP-risk allele was significantly lower expressed compared to the reference
50 allele (40.7%, $P < 5 \times 10^{-16}$).

51 **Conclusions** Our results indicate that the association between the MGP-variant and increased risk for
52 hand OA is caused by a lower expression of *MGP*, which may increase the burden of hand OA by
53 decreased inhibition of cartilage calcification.

54

55 **Keywords:**

56 Genetics, Hand Osteoarthritis, genome-wide association study, functional study, Matrix-Gla protein

57 **Introduction**

58 Osteoarthritis (OA) is the most frequent joint disorder worldwide. An estimated 22% of the adult
59 population has a joint affected by OA and this incidence increases to 49% in individuals over 65 years of
60 age[1]. All synovial joints can be affected by OA, with hand OA as one of the most common forms of OA.
61 Hand OA is characterized by osteophyte formation, bony enlargements of finger joints and cartilage
62 degradation in the joints. One of the factors contributing to cartilage degradation is the increase of
63 calcified cartilage in the joint[2,3]. In addition, hand OA is related to the occurrence of OA at other sites,
64 most notably with knee OA[4,5]. Patients affected by hand OA suffer from pain and disability, impacting
65 their quality of life. OA is a leading cause of chronic disability[6], yet currently no effective therapeutic
66 treatments against osteoarthritis are known. It is therefore imperative to dissect the underlying
67 mechanism of disease aetiology as this may enhance effective and targeted drug development.

68 OA has a strong genetic component. Depending on the joint affected, the heritability of OA is
69 estimated in the range of 40-60%[7,8], with hand OA having the largest heritability, i.e. ~60%[9][10].
70 Therefore, in recent years, several large-scale genetic studies have been performed to identify the
71 underlying genes and pathways leading to OA. Multiple significant associated loci for OA of the hip and
72 knee have been identified through genome-wide association studies (GWAS)[11–18]. However, thus far,
73 only one report has described a robust association with OA of the hand[19]. In this previous report,
74 common variants in the *ALDH1A2* and rare variants in chromosome 1p31 were genome-wide
75 significantly associated with hand OA using a discovery cohort of 837 cases and 77,325 controls.

76 In this study, we aimed to identify novel genes and pathways involved in the aetiology of OA of
77 the hand by performing a large-scale GWAS. We used a semi- quantitative measure for OA of the hand
78 in order to increase statistical power. We gathered a large sample size of 12,754 individuals for analysis.,
79 by combining data from three studies in the discovery phase and an additional three cohorts for

80 replication. Next, we conducted functional follow-up of our top finding to investigate the underlying
81 mechanism.

82

83 **METHODS**

84 **Discovery GWAS, replication and meta-analysis**

85 For a detailed description on the GWAS methods, participating studies, quality control procedures for
86 genotyping and imputation, see supplementary text S1 and Table S1.

87 **Detailed phenotype description of KL sum-score**

88 We have used a semi-quantitative bilateral measure of osteoarthritis of the hand based on the radiographic
89 Kellgren and Lawrence score (KL-score)[20]. Using radiographs of both hands, the KL-score was determined for
90 each joint in the hand. Using these KL-scores we defined the KLsum-score: the total KL-score, the sum, of the
91 following hand joints for both hands (left and right): all Distal Interphalangeal (DIP) joints, all Proximal
92 Interphalangeal (PIP) joints, all Metacarpophalangeal (MCP) joints , the Interphalangeal (IP) joint and the first
93 Carpometacarpal (CMC1) joint. Which gives the sum of 15 joints on each hand, and in total 30 joints for both hands
94 together, resulting in a minimum score of 0 and a maximum score of 120. The Leiden Studies cohort no Kellgren-
95 Lawrence scoring was done of the MCP joints, resulting in a KLsum-score of maximum 88. Individuals lacking KL-
96 grading for both or one hands and individuals with missing age or gender information were excluded from all
97 analyses in all cohorts. As the KL-sum score has a skewed distribution the top finding of the meta-analysis was
98 repeated in the discovery cohorts using a Poisson regression.

99

100 **Visualization of the associated loci and the regulatory landscape**

101 For the top GWAS associated SNP, the LD region ($r^2 > 0.8$) was determined using the 1000G Phase-1
102 population using the HaploReg V3 tool[21]. Using the ROADMAP generated reference epigenomes we
103 determined if any of the variants in high LD were located in potential gene regulatory regions in primary
104 osteoblasts (generated by ENCODE) and bone marrow derived chondrocytes (ROADMAP). [22,23].. The

105 18-state chromatin reference epigenomes were downloaded from the ROADMAP epigenomes data
106 portal[23]. SNPs and regulatory annotations were visualized using the UCSC genome-browser on
107 GRCh37/hg19[24]. For each variant it was also determined if the alternative allele would disrupt a
108 protein binding motif, this was done using the HaploReg V3 tool[21].

109

110 **RNA-sequencing data**

111 Post RNA isolation (Qiagen RNeasy Mini Kit, RIN>7) of 40 knee (15 paired preserved (P) and OA lesioned
112 (OAL), 7 P only, 3 OAL only) and 28 hip (6 paired P and OAL, 14 P only, 2 OAL only) cartilage samples
113 (Supplementary Table S2), paired-end 2x100bp RNA library sequencing (Illumina TruSeq RNA-Library
114 Prep Kit, Illumina HiSeq2000) resulted in an average of 10 million fragments per sample. Reads were
115 aligned using GSNAP against GRCh37/hg19, in which SNPs from the Genome of the Netherlands
116 consortium withMAF>1% were masked to prevent alignment bias. Number of fragments per gene were
117 used to assess quantile-adjusted conditional maximum likelihood (qCML) (edgeR, R-
118 package).Subsequently, differential gene expression analysis was performed pairwise between P and
119 OAL samples for which we had RNA of both (n=21). ASE was assessed using SNVMix2[25] with default
120 settings(min coverage=25, 10 reads per allele). The extent of allele specific expression (ASE) was defined
121 as the fraction of risk allele among all counts at the respective location. Meta-analysis was done only
122 across P samples or OAL when no P counterpart sample was present. P-values were calculated using
123 canonical binominal test (metagen R-package).

124

125 **TaqMan assay**

126 **Conventional TaqMan genotyping was performed on both genomic-DNA (gDNA), articular cartilage**
127 **and Subchondral bone cDNA. An allele-specific custom TaqMan assay for rs1800801 (Thermo Fisher**
128 **Scientific) was used to quantify the allele ratio in cDNA samples and were normalized against the**

129 **gDNA ratio, which was used as an 1:1 allele ratio reference. Each sample has been measured in four**
130 **(cartilage) or eight (subchondral bone) times, while calculations and statistics were performed as**
131 **described previously[19,26]. Cartilage samples which yielded fewer than four measurements (N=2)**
132 **were discarded prior to further analyses. All subchondral bone samples were assessed by eight**
133 **technical replicates.**

134

135

136 **RESULTS**

137 *GWAS of KLsum-score*

138 We conducted a genome-wide association study (GWAS) of a semi-quantitative measure of hand OA, a
139 bilateral summed score of Kellgren- and Lawrence scores[20], that grades radiographic OA severity,
140 across all hand joints (KLsum-score, range of 0 to 120). The discovery set consisted of three Rotterdam
141 Study cohorts (RSI, RSII and, RSIII) and included 8,743 participants with KLsum-scores. Replication was
142 done in another 4,011 individuals from three different cohorts; Leiden studies(LS), Framingham heart
143 study(FHS), and Twins-UK (TUK). General characteristics of the discovery cohorts and replication cohorts
144 can be found in Supplementary Table S3 and in Supplementary Text S1.

145 The discovery analysis yielded two novel independent genome-wide significant loci ($P \leq 5 \times 10^{-8}$)
146 on chromosome 12, an intergenic region between *MGP* and *ERP27* and an intronic region in *CCDC91*. We
147 also identified seven other novel loci with suggestive significance ($P < 5 \times 10^{-6}$) (Figure 1). In total, nine loci
148 were selected for replication in 4,011 individuals from three different cohorts (LS, FHS, and TUK). Using a
149 Bonferroni corrected P -value $< 5.56 \times 10^{-3}$, we significantly replicated one of nine loci, rs4764133
150 ($\text{Beta}_{\text{meta}}=0.83$, $\text{SE}_{\text{meta}}=0.10$, $P\text{-value}_{\text{replication}}=3.4 \times 10^{-07}$, $P\text{-value}_{\text{meta}}=1.8 \times 10^{-15}$) with the same direction of
151 effect as identified in the discovery analysis (Table 1, and Supplementary Figure S1 Forrest plot). This

152 locus maintained genome-wide significance and another locus near *ENPP3* reached near genome-wide
153 significance (chr6:132063842:D, $\text{Beta}_{\text{meta}}=0.58$, $\text{SE}_{\text{meta}}=0.11$, $\text{P-value}_{\text{meta}}=3.8*10^{-7}$) in the combined
154 discovery and replication joint meta-analysis (Table 1). Since the KLsum-score has a skewed distribution
155 the top hit was also re-analysed in the discovery set using a Poisson regression (rs4764133,
156 $\text{Beta}_{\text{poisson}}=0.12$, $\text{SE}_{\text{poisson}}=0.02$, $\text{P-value}_{\text{poisson}}=1.98*10^{-11}$).

157 Our top replicated and genome-wide significant finding, rs4764133 [T] ($\text{P}_{\text{meta}}=1.80*10^{-15}$,
158 $\text{Beta}=0.83$, $\text{MAF}=0.39$) is located in a non-coding intergenic region between *MGP* (Matrix Gla-protein)
159 and *ERP27* (Endoplasmic Reticulum Protein 27). However, variants in high linkage disequilibrium (LD)
160 with rs4764133 ($r^2 \geq 0.8$) span a ~80Kb region encompassing multiple genes, including *MGP* and an open-
161 reading frame *C12orf60* (Figure 2A). Moreover, several of these variants are located in an mRNA
162 transcript, including a nonsynonymous variant in *MGP*, and variants in 3' and 5'UTR of *MGP* and
163 *C12orf60* (Table 2, Figure 2B). The nonsynonymous variant in *MGP*, rs4236, is predicted to be non-
164 damaging (STIFT=1, tolerated; PolyPhen=0, benign) causing a threonine to alanine amino acid
165 substitution. Two variants are located in predicted active promoter region of *MGP* (rs1800801) and
166 *C12orf60* (rs9668569) in chondrogenic cells and primary osteoblasts (Table 2).

167 Next, we investigated the association of rs4764133 with bilateral severe hand OA and bilateral
168 finger OA using the discovery set (RS-I, RS-II and RS-III). We found a strong association with finger OA (P-
169 $\text{value}=3.09*10^{-8}$, $\text{OR}=1.25$) and nominal significant association with severe hand OA ($\text{p-value}=2.80*10^{-}$
170 2 , $\text{OR}=1.36$), which has a low frequency in the population (Table S4), we also see a nominal significant
171 association with cartilage thickness in the hip joint (minimal joint space width. mJSW). To see if
172 rs4764133 also confers risk for other forms of osteoarthritis, i.e. osteoarthritis of the hip and knee, we
173 used the GWAS summary data of the treat the treat OA consortium[27] and the recently published
174 mJSW meta-analysis[18]. No association was found between rs4764133 and hip or knee OA (Table S4).

175 However, we did find a nominal significant association between rs rs1049897 ($r^2= 0.98$ with
176 rs4764133)($P\text{-value}=1.28*10^{-2}$, $\text{Beta}=-0.398$).

177

178 *Gene expression analyses*

179 In order to identify potential causal genes located in the LD block surrounding rs4764133, we assessed
180 gene expression of *MGP*, *ERP27*, *ART4*, *SMOC3* (*C12orf69*) and *C12orf60* in articular cartilage, the
181 primary OA affected tissue. RNA sequencing was obtained on articular cartilage from primary OA
182 patients who had total joint replacement surgeries of either the knee ($n=25$) or hip ($n=22$) joint.
183 Expression levels of *ERP27*, *C12orf60*, *ART4* and *SMOC3* were substantially lower than *MGP* expression
184 levels in articular cartilage (Figure. S2A). Nonetheless, neither *MGP*, *ERP27*, *ART4* nor *SMOC3* and
185 *C12orf60* showed significant difference in gene expression between paired preserved(P) and OA
186 lesioned(OAL) articular cartilage. However, while these genes are not differentially expressed in OA
187 affected cartilage, it is possible that the identified GWAS SNPs affect gene transcription. When we
188 analysed the relationship between the top SNP and expression analysis in a classical eQTL (expression
189 Quantitative Trait Loci) analysis, we did not to detect significant correlations between rs1049897, rs4236
190 or rs1800801 and absolute *MGP*, *ERP27*, *ART4*, *SMOC3* or *C12orf60* expression levels (Figure S2B).
191 However, we did observe several variants in high LD located in the mRNA transcript of *MGP* and
192 *C12orf60*, allowing us to assess allele specific expression (ASE) for these genes. We were unable to study
193 ASE for *ART4*, *SMOC3* and *ERP27*, since no SNP in high LD with rs4764133 is present in the coding region.
194 In ASE the influence of exonic alleles on gene expression *in-cis* is measured within heterozygote
195 subjects, circumventing strong effects from environmental or trans-acting influences. This property
196 results in ASE analysis to be a more statistically powerful approach, when compared to classical eQTL
197 analysis [28]. Subsequently, we found that the OA risk alleles for three coding variants in high LD with
198 the lead variant, rs4236 (Figure. 3SA, 39.6% C allele, $P<5*10^{-16}$), rs1049897 (Figure. S3B, 44.4% A allele,

199 $P < 5 \times 10^{-10}$), and rs1800801 (Figure. 3A, 40.7% T allele, $P < 5 \times 10^{-16}$), were significantly correlated with
200 lower expression of *MGP*, marking imbalanced expression among heterozygotes, independent of the
201 disease status of the articular cartilage. No allele specific expression was observed between SNPs
202 rs11276, rs3088189 and rs1861698 (residing in *C12orf60* and in high LD with the lead SNP, $r^2 > 0.8$, Table
203 2). Technical and biological replication was performed using a custom allele specific TaqMan assay for
204 rs1800801 in eight additional heterozygous individuals for which we isolated RNA from P cartilage (n=2),
205 OAL (n=2) or both (n=4) from primary knee OA patients and confirmed the observed imbalance in
206 preserved articular cartilage (Figure 3B, relative allelic difference=0.92, $P < 1 \times 10^{-6}$), as well as in 8 knee
207 subchondral bone samples (Figure 3C, relative allelic difference=0.78, $P < 1 \times 10^{-4}$).

208

209 **Discussion**

210 Here, we show for the first time, that there is a robust genome-wide significant association between
211 rs4764133, located near *MGP*, and hand OA. Furthermore we performed functional validation showing
212 that *MGP* coding variants in LD with rs4764133 are associated with allele-specific expression of *MGP*
213 which may increase risk of hand OA by lowering inhibition of articular cartilage calcification, since *MGP*
214 is an essential inhibitor of cartilage calcification[29,30]. These findings suggest that *MGP* could be
215 considered a prioritized drug target for hand OA, since genetically supported drug targets double the
216 success rate of therapeutics in clinical development[31].

217 *MGP* is an essential inhibitor of cartilage calcification, and genetic deficiencies of *MGP* in
218 humans and mice have been linked to abnormal mineralization of soft tissues, including cartilaginous
219 tissue[29,32]. Furthermore *MGP* has been previously implicated in relation to OA. A small candidate
220 study reported marginally significant association between hand OA and genetic variants in *MGP*
221 (rs1800802 and rs4236)[33]. This is consistent with our findings that the minor allele for rs4764133 and
222 related coding variants in high LD ($r^2 > 0.8$), rs1800802 and rs4236, increase the risk of hand OA and that

223 we found high expression of *MGP* in both preserved and OA lesioned articular cartilage. In contrast,
224 another study showed that an MGP protein complex is excreted by healthy articular chondrocytes, but
225 not by OA affected chondrocytes[34], although we only assed *MGP* expression and not MGP protein
226 complex excretion.

227 Although the loci with allele specific expression (ASE) are known to be enriched for eQTLs[35],
228 we were unable to detect an association between the MGP-genotype and *MGP* RNA-expression levels in
229 cartilage. This could have been due to our modest sample size (knee joint, n=25 and or hip joint, n=22) in
230 combination with large heterogeneity of the tissue. Notably, the available cartilage samples originated
231 from different joint sites (knee, hip) and different disease stage (preserved versus affected), and had
232 large age range of the individuals. Also, It is known that ASE is a more powerful technique then classical
233 eQTL analysis to identify functional SNPs influencing expression of genes[28]. While the extent of
234 imbalance could be considered relatively modest, an increasing number of OA associated SNP alleles
235 appear to mark ASE by comparable amount[19,36–38]. From a more biological perspective, one could
236 consider a prolonged, albeit slight, deviation from homeostasis due to modest ASE of cartilage relevant
237 genes to be of substantial influence over time. This latter hypothesis could contain the molecular basis
238 for increased risk towards developing OA among the ageing population. Additionally, we observed that
239 the rs1800801 alleles also affected expression of MGP in subchondral bone samples. This could imply
240 that, in parallel to an effect in cartilage, he presumed disturbed cartilage homeostasis is further affected
241 by the underlying bone. Further enabling the view that OA is a pathology of the entire joint.

242 Our findings may give an explanation for the known vitamin K association with OA: MGP
243 mediated calcification inhibition is dependent on γ -carboxylation by vitamin K[39]. It has been shown
244 that low vitamin K intake is correlated with OA[40]. Thus vitamin K intake may be a potential therapeutic
245 treatment in OA. Recently, a first randomized control trial testing the effects of vitamin K on OA was
246 published, which reported no overall effect of vitamin K on hand OA[41]. Despite the low power of the

247 trial, there was a significant beneficial effect on joint space narrowing (cartilage degradation) among
248 those individuals that were VitK deficient at the start of the trial[41]. Thus, an adequately powered study
249 of vitamin K may be justified based on the found MGP association. Furthermore, genetic predisposition
250 for hand OA, was not taken into account in the trial. Perhaps, genetic predisposition for hand OA (*MGP*-
251 risk variants) in combination with insufficient vitamin K intake might potentiate cartilage calcification
252 and subsequent risk for developing hand OA. Therefore, future OA trails, therapeutic and preventive
253 treatments might benefit from taking a personalized medicine approach since genetically supported
254 drug targets double the success rate of therapeutics in clinical development[31].

255 Styrkarsdottir *et al.*(2014) reported on common genetic variants that associate with severe hand
256 OA, among the replication cohorts were the Leiden and Rotterdam cohorts[19]. Although we observe
257 suggestive signals at the reported locus (*ALDH1A2* gene, 1p31) the respective variants did not meet the
258 genome-wide significance threshold in our analyses (Supplementary Table S5). This difference is likely
259 caused by the markedly different phenotypes that were used for either analyses. Where Styrkarsdottir
260 *et al.* studied a dichotomous severe hand OA phenotype, our phenotype was semi-quantitatively
261 phenotype.

262 To conclude, we here present coding variants in *MGP*, that are associated with radiographic
263 hand OA, and the hand OA risk allele marks lower expression of *MGP* in articular cartilage. Our findings
264 suggest that *MGP* might play an important role in hand OA pathogenesis through pathways related to
265 articular cartilage calcification and vitamin K. Better understanding of *MGP* gene and protein regulation
266 and its relation to vitamin K intake and OA, may reveal novel therapeutic drug targets for hand OA.

267

268

269 **ACKNOWLEDGMENTS**

270 This study was funded by The Netherlands Society for Scientific Research (NWO) VIDI Grant 917103521.
271 The Rotterdam Study is funded by Erasmus Medical Center and Erasmus University, Rotterdam,
272 Netherlands Organization for the Health Research and Development (ZonMw), the Research Institute for
273 Diseases in the Elderly (RIDE), the Ministry of Education, Culture and Science, the Ministry for Health,
274 Welfare and Sports, the European Commission (DG XII), and the Municipality of Rotterdam. The authors
275 are grateful to the study participants, the staff from the Rotterdam Study and the participating general
276 practitioners and pharmacists.

277 The generation and management of GWAS genotype data for the Rotterdam Study (RS I, RS II,
278 RS III) was executed by the Human Genotyping Facility of the Genetic Laboratory of the Department of
279 Internal Medicine, Erasmus MC, Rotterdam, The Netherlands. The GWAS datasets are supported by the
280 Netherlands Organisation of Scientific Research NWO Investments (nr. 175.010.2005.011, 911-03-012),
281 the Genetic Laboratory of the Department of Internal Medicine, Erasmus MC, the Research Institute for
282 Diseases in the Elderly (014-93-015; RIDE2), the Netherlands Genomics Initiative (NGI)/Netherlands
283 Organisation for Scientific Research (NWO) Netherlands Consortium for Healthy Aging (NCHA), project
284 nr. 050-060-810. We thank Pascal Arp, Mila Jhamai, Marijn Verkerk, Lizbeth Herrera and Marjolein
285 Peters, MSc, and Carolina Medina-Gomez, MSc, for their help in creating the GWAS database, and Karol
286 Estrada, PhD, Yurii Aulchenko, PhD, and Carolina Medina-Gomez, MSc, for the creation and analysis of
287 imputed data

288 The Leiden University Medical Centre, the Dutch Arthritis Association and Pfizer Inc., Groton, CT,
289 USA support the GARP study, whilst the LLS was supported by the Netherlands Organization of Scientific
290 Research (MW 904-61-095, 911-03-016, 917-66-344 and 911-03-012), Leiden University Medical Centre,
291 and by the “Centre of Medical System Biology” and the “Netherlands Consortium of Healthy Aging” in
292 the framework of the Netherlands Genomics Initiative (NGI). Furthermore, the research leading to the

293 RAAK biobank and the current results has received funding from the Dutch Arthritis Association (DAA
294 2010_017) and the European Union's Seventh Framework Programme (FP7/2007-2011) under grant
295 agreement no. 259679. We thank Nico Lakenberg, Ruud van der Breggen and Eka Suchiman, for their
296 help in preparing DNA and RNA samples.

297 TwinsUK is funded by the Wellcome Trust, Medical Research Council, European Union, the
298 National Institute for Health Research (NIHR)-funded BioResource, Clinical Research Facility and
299 Biomedical Research Centre based at Guy's and St Thomas' NHS Foundation Trust in partnership with
300 King's College London.

301 The Framingham Heart Study of the National Heart, Lung, and Blood Institute of the National
302 Institutes of Health and Boston University School of Medicine was supported by the National Institutes
303 of Health (Contract No. HHSN268201500001I, N01-HC-25195, AG18393, AR47785) and its contract with
304 Affymetrix, Inc. for genotyping services (N02-HL-6-4278). Analyses reflect intellectual input and resource
305 development from the Framingham Heart Study investigators participating in the SNP Health
306 Association Resource (SHARe) project. M.S.Y. is supported by the National Institutes of Aging
307 (T32AG023480).

308

309 **AUTHOR CONTRIBUTIONS**

310 W.H. and C.G.B. contributed equally to this work. D.H., M.S.Y., Y.F.M.R. and S.M. performed replication
311 analysis for this work, L.B. provided analysis help. L.S.C. and F.R. provided data. M.K. provided
312 phenotypic contribution to the GARP study. M.P. provided data and analyses. T.D.S. contributed data for
313 replication. A.H. contributed data of the RS cohorts. J.D., M.B. and P.E.S contributed to genotyping data
314 and analyses of LLS cohort. R.G.H.H.N. provided contribution to the RAAK study. A.G.U. contributed
315 genotype data of RS cohorts , D.T.F. and A.V. contributed replication data for this work. I.M., and J.B.J.M.
316 jointly supervised this work.

317

318 **Competing Interests**

319 The Authors declare no competing interests.

320

321 **Materials & Correspondence**

322 GWAS summary statistics can be requested from J.B.J.M. All correspondence should be addressed to
323 J.B.J.M. and I.M.

324

325

326

327 **References**

- 328 1 CDC. Prevalence of Doctor-Diagnosed Arthritis and Arthritis-Attributable Activity Limitation — United
329 States, 2010–2012. *MMWR Morb Mortal Wkly Rep* 2013;**62**:869–73. doi:10.1001/jama.296.22.2671
- 330 2 Patel N, Buckland-Wright C. Advancement in the zone of calcified cartilage in osteoarthritic hands of
331 patients detected by high definition macroradiography. *Osteoarthr Cartil* 1999;**7**:520–
332 5.http://www.idealibrary.com (accessed 2 Nov 2016).
- 333 3 Fuerst M, Bertrand J, Lammers L, *et al.* Calcification of articular cartilage in human osteoarthritis. *Arthritis*
334 *Rheum* 2009;**60**:2694–703. doi:10.1002/art.24774
- 335 4 Nelson AE, DeVellis RF, Renner JB, *et al.* Quantification of the whole-body burden of radiographic
336 osteoarthritis using factor analysis. *Arthritis Res Ther* 2011;**13**:R176. doi:10.1186/ar3501
- 337 5 Jonsson H, Helgadóttir GP, Aspelund T, *et al.* Hand Osteoarthritis Severity is Associated with Total Knee
338 Joint Replacements Independently of BMI. The Ages-Reykjavik Study. *Open Rheumatol J* 2011;**5**:7–12.
339 doi:10.2174/1874312901105010007
- 340 6 CDC. Prevalence of Disabilities and Associated Health Conditions Among Adults --- United States, 1999.
341 *MMWR Morb Mortal Wkly Rep* 2009;**58**:421–6.
- 342 7 Spector TD, Cicuttini F, Baker J, *et al.* Genetic influences on osteoarthritis in women: a twin study. *BMJ*
343 1996;**312**:940–3. doi:10.1136/bmj.312.7036.940
- 344 8 Felson DT. Risk factors for osteoarthritis: understanding joint vulnerability. *Clin Orthop Relat Res*
345 2004;:S16–21. doi:10.1097/01.blo.0000144971.12731.a2
- 346 9 Bijkerk C, Houwing-Duistermaat JJ, Valkenburg H a., *et al.* Heritabilities of radiologic osteoarthritis in
347 peripheral joints and of disc degeneration of the spine. *Arthritis Rheum* 1999;**42**:1729–35.
348 doi:10.1002/1529-0131(199908)42:8<1729::AID-ANR23>3.0.CO;2-H
- 349 10 Ishimori ML, Altman RD, Cohen MJ, *et al.* Heritability patterns in hand osteoarthritis: the role of
350 osteophytes. *Arthritis Res Ther* 2010;**12**:R180. doi:10.1186/ar3144
- 351 11 Kerkhof HJM, Lories RJ, Meulenbelt I, *et al.* A genome-wide association study identifies an osteoarthritis
352 susceptibility locus on chromosome 7q22. *Arthritis Rheum* 2010;**62**:499–510. doi:10.1002/art.27184
- 353 12 Nakajima M, Takahashi A, Kou I, *et al.* New sequence variants in HLA class II/III region associated with
354 susceptibility to knee osteoarthritis identified by genome-wide association study. *PLoS One* 2010;**5**:e9723.
355 doi:10.1371/journal.pone.0009723
- 356 13 Castaño Betancourt MC, Cailotto F, Kerkhof HJ, *et al.* Genome-wide association and functional studies
357 identify the DOT1L gene to be involved in cartilage thickness and hip osteoarthritis. *Proc Natl Acad Sci U S*
358 *A* 2012;**109**:8218–23. doi:10.1073/pnas.1119899109
- 359 14 Consortium A, Collaborators A. Identification of new susceptibility loci for osteoarthritis (arcOGEN): a
360 genome-wide association study. *Lancet* 2012;**380**:815–23. doi:10.1016/S0140-6736(12)60681-3
- 361 15 Day-Williams AG, Southam L, Panoutsopoulou K, *et al.* A variant in MCF2L is associated with osteoarthritis.
362 *Am J Hum Genet* 2011;**89**:446–50. doi:10.1016/j.ajhg.2011.08.001
- 363 16 Evangelou E, Kerkhof HJ, Styrkarsdóttir U, *et al.* A meta-analysis of genome-wide association studies
364 identifies novel variants associated with osteoarthritis of the hip. *Ann Rheum Dis* 2013;:1–7.

- 365 doi:10.1136/annrheumdis-2012-203114
- 366 17 Miyamoto Y, Shi D, Nakajima M, *et al.* Common variants in DVWA on chromosome 3p24.3 are associated
367 with susceptibility to knee osteoarthritis. *Nat Genet* 2008;**40**:994–8. doi:10.1038/ng.176
- 368 18 Castaño-Betancourt MC, Evans DS, Ramos YFM, *et al.* Novel Genetic Variants for Cartilage Thickness and
369 Hip Osteoarthritis. *PLoS Genet* 2016;**12**:e1006260. doi:10.1371/journal.pgen.1006260
- 370 19 Styrkarsdottir U, Thorleifsson G, Helgadottir HT, *et al.* Severe osteoarthritis of the hand associates with
371 common variants within the ALDH1A2 gene and with rare variants at 1p31. *Nat Genet* 2014;**46**:498–502.
372 doi:10.1038/ng.2957
- 373 20 Kellgren JH, Lawrence JS. Radiological Assessment of Osteo-Arthrosis. *Ann Rheum Dis* 1957;**16**:494–502.
374 doi:10.1136/ard.16.4.494
- 375 21 Ward LD, Kellis M. HaploReg: A resource for exploring chromatin states, conservation, and regulatory motif
376 alterations within sets of genetically linked variants. *Nucleic Acids Res* 2012;**40**:930–4.
377 doi:10.1093/nar/gkr917
- 378 22 Chadwick LH. The NIH Roadmap Epigenomics Program data resource. *Epigenomics* 2012;**4**:317–24.
379 doi:10.2217/epi.12.18
- 380 23 Kundaje A, Meuleman W, Ernst J, *et al.* Integrative analysis of 111 reference human epigenomes. *Nature*
381 2015;**518**:317–30. doi:10.1038/nature14248
- 382 24 Kent WJ, Sugnet CW, Furey TS, *et al.* The Human Genome Browser at UCSC. *Genome Res* 2002;**12**:996–
383 1006. doi:10.1101/gr.229102
- 384 25 Goya R, Sun MGF, Morin RD, *et al.* SNVMix: predicting single nucleotide variants from next-generation
385 sequencing of tumors. *Bioinformatics* 2010;**26**:730–6. doi:10.1093/bioinformatics/btq040
- 386 26 Bos SD, Bovée JVMG, Duijnisveld BJ, *et al.* Increased type II deiodinase protein in OA-affected cartilage and
387 allelic imbalance of OA risk polymorphism rs225014 at DIO2 in human OA joint tissues. *Ann Rheum Dis*
388 2012;**71**:1254–8. doi:10.1136/annrheumdis-2011-200981
- 389 27 Kerkhof HJM, Doherty M, Arden NK, *et al.* Large-scale meta-analysis of interleukin-1 beta and interleukin-1
390 receptor antagonist polymorphisms on risk of radiographic hip and knee osteoarthritis and severity of knee
391 osteoarthritis. *Osteoarthr Cartil* 2011;**19**:265–71. doi:10.1016/j.joca.2010.12.003
- 392 28 Pastinen T. Genome-wide allele-specific analysis: insights into regulatory variation. *Nat Rev Genet*
393 2010;**11**:533–8. doi:10.1038/nrg2815
- 394 29 Luo G, Ducy P, McKee MD, *et al.* Spontaneous calcification of arteries and cartilage in mice lacking matrix
395 GLA protein. *Nature* 1997;**386**:78–81. doi:10.1038/386078a0
- 396 30 Yagami K, Suh JY, Enomoto-Iwamoto M, *et al.* Matrix GLA protein is a developmental regulator of
397 chondrocyte mineralization and, when constitutively expressed, blocks endochondral and
398 intramembranous ossification in the limb. *J Cell Biol* 1999;**147**:1097–
399 108. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2169349&tool=pmcentrez&rendertype=abstract>
400
- 401 31 Nelson MR, Tipney H, Painter JL, *et al.* The support of human genetic evidence for approved drug
402 indications. *Nat Genet* 2015;**47**:856–60. doi:10.1038/ng.3314
- 403 32 Munroe PB, Olgunturk RO, Fryns JP, *et al.* Mutations in the gene encoding the human matrix Gla protein

404 cause Keutel syndrome. *Nat Genet* 1999;**21**:142–4. doi:10.1038/5102

405 33 Misra D, Booth SL, Crosier MD, *et al.* Matrix Gla protein polymorphism, but not concentrations, is
406 associated with radiographic hand osteoarthritis. *J Rheumatol* 2011;**38**:1960–5.
407 doi:10.3899/jrheum.100985

408 34 Wallin R, Schurgers LJ, Loeser RF. Biosynthesis of the vitamin K-dependent matrix Gla protein (MGP) in
409 chondrocytes: a fetuin-MGP protein complex is assembled in vesicles shed from normal but not from
410 osteoarthritic chondrocytes. *Osteoarthritis Cartilage* 2010;**18**:1096–103. doi:10.1016/j.joca.2010.05.013

411 35 Hango G, Williams DE, Falkenburg JH, *et al.* Influence of IL-1 alpha and -1 beta on the survival of human
412 bone marrow cells responding to hematopoietic colony-stimulating factors. *J Immunol* 1989;**142**:4329–
413 34. <http://www.ncbi.nlm.nih.gov/pubmed/2656864> (accessed 22 Aug 2016).

414 36 Gee F, Clubbs CF, Raine EVA, *et al.* Allelic expression analysis of the osteoarthritis susceptibility locus that
415 maps to chromosome 3p21 reveals cis-acting eQTLs at GNL3 and SPCS1. *BMC Med Genet* 2014;**15**:53.
416 doi:10.1186/1471-2350-15-53

417 37 Raine EVA, Dodd AW, Reynard LN, *et al.* Allelic expression analysis of the osteoarthritis susceptibility gene
418 COL11A1 in human joint tissues. *BMC Musculoskelet Disord* 2013;**14**:85. doi:10.1186/1471-2474-14-85

419 38 Bos SD, Bovée JVMG, Duijnisveld BJ, *et al.* Increased type II deiodinase protein in OA-affected cartilage and
420 allelic imbalance of OA risk polymorphism rs225014 at DIO2 in human OA joint tissues. *Ann Rheum Dis*
421 2012;**71**:1254–8. doi:10.1136/annrheumdis-2011-200981

422 39 Schurgers LJ, Uitto J, Reutelingsperger CP. Vitamin K-dependent carboxylation of matrix Gla-protein: a
423 crucial switch to control ectopic mineralization. *Trends Mol Med* 2013;**19**:217–26.
424 doi:10.1016/j.molmed.2012.12.008

425 40 Shea MK, Kritchevsky SB, Hsu F-C, *et al.* The association between vitamin K status and knee osteoarthritis
426 features in older adults: the Health, Aging and Body Composition Study. *Osteoarthritis Cartilage*
427 2015;**23**:370–8. doi:10.1016/j.joca.2014.12.008

428 41 Neogi T, Felson DT, Sarno R, *et al.* Vitamin K in hand osteoarthritis: results from a randomised clinical trial.
429 *Ann Rheum Dis* 2008;**67**:1570–3. doi:10.1136/ard.2008.094771

430

431

432 **Figures**

433 **Figure 1.** GWAS results for association with the KLsum score in the discovery phase. Manhattan plot for association
434 with the KLsum-score, adjusted for age and sex, in the discovery cohorts of RSI, RSII and RSIII. The $-\log_{10}$ P-values,
435 for each of the ~11 million SNPs analyzed (remaining after EASYQC quality control) as part of the genome wide
436 association with the KLsum-score, plotted against their position per chromosome. The red dotted horizontal line
437 corresponds to the genome-wide significant threshold ($P = 5 \times 10^{-8}$). The dotted grey line corresponds to the
438 selection for replication threshold ($P = 5 \times 10^{-6}$). SNP location represented by [], if the SNP is localized intergenic the
439 dashes denotes the distance, $-\leq 10$ kb, $-\leq 100$ kb, $-\leq 1000$ kb, $-\leq 1$ Mkb, ≥ 1 Mkb.

440
441 **Figure 2.** Locus zoom plot for rs4764133. locus zoom plot for rs4764133, 150 kb upstream and downstream of
442 rs4764133 has been taken as plotted region (A). Zoom in on *MGP* and three SNPs in high LD with top SNP that are
443 located in the *MGP* mRNA transcript (B). Also represented is ROADMAP chromatin 18-state data of two tissue
444 types: human Mesenchymal Stem Cell (hMSC) derived cultured chondrogenic cells and primary osteoblasts. In
445 both these cell types the chromatin contains active marks surrounding the *MGP* promoter.

446
447 **Figure 3.** Allelic imbalanced expression of *MGP* marked by the alleles among heterozygotes of rs1800801 (A), in
448 the assessed cartilage RNA sequencing dataset. Validation of selected rs1800801 using a custom TaqMan assay
449 confirmed the imbalance (B). Allelic imbalance was also assed in with a custom TagMan assay in subchondral bone
450 samples (C). Preserved (P) and OA lesioned (OAL) samples are shown respectively in blue and red, and genomic
451 DNA (TaqMan control) in black (G). For ASE results for rs4236 and rs1049879, see Supplementary Figure S3 and for
452 information on the samples see Supplementary Table S2.

453

454

455 **Table 1.** Results GWAS quantitative bilateral phenotype of osteoarthritis of the hand (KLsum-score), discovery, replication and meta-analysis

SNP	Chr	Position (hg19)	Effect Allele	Other Allele	EAF†	Discovery*			Replication**			Combined			Genomic location††
						Beta	SE	P-value	Beta	SE	P-value	Beta	SE	Pval	
rs1494593	5	23574856	T	C	0.88	-0.83	0.18	4.46E-06	-0.15	0.32	6.31E-01	-0.67	0.16	2.28E-05	<i>PRDM9</i> --[]--- <i>C5orf17</i>
rs114370021	5	167398535	A	G	0.27	-0.84	0.18	2.12E-06	0.48	0.32	1.28E-01	-0.53	0.16	6.96E-04	[<i>TENM2</i>]
rs7770034	6	44447004	A	G	0.48	-0.54	0.12	3.42E-06	-0.36	0.21	8.71E-02	-0.50	0.10	1.01E-06	<i>CDC5L</i> --[]--- <i>SUPT3H</i>
6:132063842:D‡	6	132063842	D	I	0.27	0.64	0.13	1.28E-06	0.41	0.23	7.82E-02	0.58	0.11	3.79E-07	[<i>ENPP3</i>]
11:90657297:D	11	90657297	D	I	0.11	0.93	0.19	1.35E-06	0.33	0.33	3.11E-01	0.78	0.17	2.91E-06	<i>DISC1FP1</i> -[]---- <i>FAT3</i>
rs4764133	12	15064363	T	C	0.39	0.75	0.12	3.45E-10	1.11	0.22	3.34E-07	0.83	0.10	1.80E-15	<i>MGP</i>--[]- <i>ERP27</i>
rs7139060	12	28693144	A	G	0.67	-0.73	0.12	6.12E-09	0.11	0.22	6.16E-01	-0.52	0.11	1.47E-06	[<i>CCDC91</i>]
rs1950427	14	25955502	T	C	0.12	0.86	0.18	1.09E-06	-0.28	0.30	3.56E-01	0.57	0.15	1.80E-04	<i>STXBP6</i> ---[]--- <i>NOVA1</i>
rs6108226	20	8960884	T	C	0.77	0.70	0.15	3.94E-06	0.20	0.27	4.43E-01	0.58	0.13	1.14E-05	<i>PLCB1</i> --[]-- <i>PLCB4</i>

* Discovery: RS-I, RS-II, RS-III, $n = 8,743$

** Replication : GARP, LSS, TwinsUK & FHS, $n = 4,011$

† EAF: Effect Allele Frequency

††SNP location represented by [], if the SNP is localized intergenic the dashes denotes the distance, -≤10 kb, --≤100kb, ---≤1000kb, ----≤1Mkb, -----≥1Mkb

‡For TwinsUK a proxy SNP was used: rs3850251 $r^2=1$ $D'=1$ (as calculated in the RSI, RSII and RSIII cohorts)

456 **Table 2.** rs4764133 LD block ($r^2 > 0.8$) annotation of potential functional elements in osteoblasts and
 457 chondrogenic cells, X marks no potential functional annotation i.e. enhancer region, promoter region or
 458 altered protein binding motifs.

SNP	P-value Discovery	r^2	Annotation*	Regulatory Chromatin Marks**		Altered Protein Binding Motifs (Haploreg V3)
				Chondrogenic cells	Osteoblasts	
rs1049897	3.48E-09	0.88	MGP 3'-UTR	Transcription	X	X
rs4236	4.16E-09	0.86	MGP non-synonymous	Enhancer region	X	HNF4, PLAG1
rs1800801	1.12E-09	0.95	MGP 5'UTR	Promoter region	Promoter region	Zfp410
rs7310951	4.04E-09	0.86	C12orf60	Enhancer region	X	DMRT7, Gfi1, Pax-5
rs12320004	4.04E-09	0.86	C12orf60	Enhancer region	X	BHLHE40, P300, HEN1, LBP-1, RAD21, TATA, Zfx
rs10772814	3.76E-09	0.88	C12orf60	Enhancer region	X	HNF4
rs10492151	1.21E-09	0.95	C12orf60	Enhancer region	X	AIRE, Hoxa13
rs725445	3.58E-08	0.82	C12orf60	Enhancer region	X	Hand1
rs725444	3.92E-09	0.87	C12orf60	Enhancer region	X	Foxf1, Foxi1, Foxo, Foxq1, Mef2
rs4764131	6.31E-10	0.97	C12orf60	Enhancer region	Enhancer region	Myc
rs9668569	5.91E-10	0.97	C12orf60	Promoter region	Promoter region	X
rs2430687	2.44E-09	0.89	C12orf60	Enhancer region	Enhancer region	BHLHE40
rs12311463	6.91E-10	0.97	C12orf60	Enhancer region	Enhancer region	Pou1f1, Pou2f2, TATA
rs67482087	4.61E-10	0.95	C12orf60	Enhancer region	Enhancer region	Foxp1, Irx, Pou1f1, Pou2f2, Pou3f3, TATA
rs67436073	6.76E-10	0.97	C12orf60	Enhancer region	Enhancer region	Foxj2, Foxk1, Foxo, GATA, Mef2, Pou2f2, Pou3f2, Pou6f1, TATA, Zfp
rs11276	8.05E-09	0.96	C12orf60 non-synonymous	X	X	SPIB, NF-AT
rs3088189	9.46E-09	0.96	C12orf60 synonymous	X	X	SPIB
rs1861698	3.56E-09	0.96	C12orf60 synonymous	X	X	Bbx, Pou1f1, TATA

*Gene annotation based on the hg19 release of the UCSC Genome Browser

** Regulatory chromatin marks taken from the ROADMAP Epigenomes project chromatin state learning core 18-state model