1	Examining evidence for behavioural mimicry of parental eating by
2	adolescent females: an observational study
3	Maxine Sharps <sup>1</sup> , Suzanne Higgs <sup>2</sup> , Jackie Blissett <sup>2</sup> , Arie Nouwen <sup>3</sup> , Magdalena Chechlacz <sup>4</sup> ,
4	Harriet A Allen <sup>5</sup> , Eric Robinson <sup>1</sup>
5	
6	<sup>1</sup> University of Liverpool
7	<sup>2</sup> University of Birmingham
8	<sup>3</sup> Middlesex University
9	<sup>4</sup> University of Oxford
10	<sup>5</sup> University of Nottingham
11	
12	Corresponding Author:
13	Maxine Sharps, Psychological Sciences,
14	Eleanor Rathbone Building,
15	University of Liverpool,
16	Liverpool, L69 7ZA, UK
17	Email – Maxine.sharps@liv.ac.uk
18	Funding statement: The current study was funded in part, by the European Foundation for the
19	Study of Diabetes (EFSD)/ Novo Nordisk European Clinical Research Programme in
20	Adolescents with Type 2 Diabetes.
21	The authors report no conflicts of interest
22	Word count: 4405
22	Kay words: mimicry: social modelling: social eating

#### **ABSTRACT**

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

Behavioural mimicry is a potential mechanism explaining why adolescents appear to be influenced by their parents' eating behaviour. In the current study we examined whether there is evidence that adolescent females mimic their parents when eating. Videos of thirty-eight parent and female adolescent dyads eating a lunchtime meal together were examined. We tested whether a parent placing a food item into their mouth was associated with an increased likelihood that their adolescent child would place any food item (non-specific mimicry) or the same item (specific mimicry) in their mouth at three different time frames, namely during the same second or within the next fifteen seconds (+15), five seconds (+5) or two second (+2)period. Parents and adolescents' overall food intake was positively correlated, whereby a parent eating a larger amount of food was associated with the adolescent eating a larger meal. Across all of the three time frames adolescents were more likely to place a food item in their mouth if their parent had recently placed that same food item in their mouth (specific food item mimicry), however there was no evidence of non-specific mimicry. This observational study suggests that when eating in a social context there is evidence that adolescent females may mimic their parental eating behaviour, selecting and eating more of a food item if their parent has just started to eat that food.

Social context has been shown to have a strong influence on eating behaviour (Herman, Roth 42 & Polivy., 2003; Goldman et al., 1991). Social modelling research has shown that the eating 43 behaviour of adults and children can be influenced by the amount of food other diners are 44 eating; eating more when others are eating more and less when they are eating less 45 (Bevelander et al., 2012; Hermans et al., 2009). A variety of potential explanations of these 46 effects have been suggested. For example, modelling may occur because the behaviour of 47 one's peers sets a norm of what constitutes a socially appropriate amount to eat (Herman et 48 al., 2003; Vartanian et al., 2013) or because it acts as an informational cue to guide behaviour 49 50 (Robinson et al., 2013). 51 Parents are thought to be one of the most important social influences on child and adolescent 52 53 eating behaviour (Salvy et al., 2011), influencing health beliefs, behaviours and dietary intake (Oliveria et al., 1992; Lau et al., 1990). Moreover, parental and child food consumption tend 54 to be correlated in terms of the type and amounts of food that both eat (McGowan et al., 55

2012; Wroten et al., 2012; Sweetman et al., 2011). Likewise, research has shown that
children are more likely to try a food if they observe their parent eating that same food
(Harper et al., 1975). More recent research has also shown in an experimental setting that the
presence of a parent shapes the amount and types of food adolescents eat (Salvy et al., 2011).

However, the mechanisms underlying the processes by which adolescents adapt their eating

to match parental behaviour when eating has received less attention.

60

61

62

63

64

65

66

One possibility is that adolescents mimic or synchronise to their parents' eating behaviour when dining together. Behavioural mimicry refers to the process whereby a person imitates the behaviour of another person without conscious awareness and is thought to occur due to a tight neural link between perception and action (Chartrand & Bargh., 1999; Chartrand et al.,

2009), such that observing another person's movements may trigger one's own motor system to perform that same movement (Lakin & Chartrand., 2003; Iacoboni., 2009), e.g. taking a bite of food. Mimicry has been suggested to occur for a number of behaviours (Larsen et al, 2009; Neumann & Strack., 2000; Bernieri., 1988) and more recently the role of behavioural mimicry in social eating contexts has been examined. Hermans et al. (2012) found that when two female adults ate the same meal together, participants were more likely to pick up and eat the food if their eating partner had done so in the proceeding five seconds. Similarly, Bevelander et al. (2013) found that when a young child (aged 6-11) picked up and ate a chocolate covered peanut, this was associated with an increased likelihood that their eating partner would subsequently pick up and eat that food. Thus, previous studies have only investigated behavioural mimicry in child only or adult only groupings (Hermans et al., 2012, Bevelander et al., 2013) and as research supports that adolescents' eating behaviour may be affected by the eating behaviour of a present parent (Salvy et al., 2011), it will be important to understand whether mimicry of eating behaviour may occur between a parent and an adolescent. It may be the case that mimicry of parental eating is a mechanism explaining parental influence on adolescent eating behaviour.

In studies to date examining behavioural mimicry during social eating, participants have only been provided with a single food item to eat (Hermans et al., 2012; Bevelander et al., 2013). From these studies it is therefore not possible to infer whether participants were mimicking eating of a specific food type (if you take food x, I then take food x) or whether participants were simply synchronising the rate of their food intake in a more general/non-specific manner. For example, it may be that watching another person pick up a food item triggers an automatic reaction to reach for any food item (non-specific food item mimicry) or only the same food item (specific food item mimicry). Differentiating between these two possibilities

is of importance because it may signal mechanisms that underlie mimicry. If automatic synchrony of gestures is of importance (Hermans et al., 2012; Iacoboni et al., 1999) then we may expect to see evidence for non-specific mimicry, because mimicry of the action of eating is key. Conversely, if mimicry occurs because an eating partner sets a norm about which foods are and are not appropriate to eat (Vartanian et al., 2013; Herman et al., 2003), then only mimicry of congruent food items may be observed. These questions are also of importance because in naturalistic social eating contexts such as family meal times, a variety of food items are likely to be available.

In the present study we aimed to examine whether there is evidence that adolescents mimic the eating behaviour of their parents when eating together. In order to assess mimicry, videos of parent-adolescent dyads eating a multi-item lunchtime meal were examined. We examined whether there was evidence of both 'non-specific food item mimicry' and 'specific food item mimicry'. Based on previous studies of eating mimicry (Bevelander et al., 2013; Hermans et al., 2012), it was hypothesised that a parent placing a food item in their mouth would be associated with an increased likelihood that their adolescent child would also place a food item in their mouth. However, we reasoned that if evidence of mimicry was observed, it may only be food item specific, as parental behaviour during a meal may primarily signal which foods are appropriate to eat and when.

#### **METHOD**

Background

The videos analyzed were of adolescents and parents eating a multi-item lunchtime meal together, which were recorded as part of a test day for a larger study examining brain

activations and responsiveness to food cues. In the larger study, participants arrived at the laboratory on the morning of their test day where they underwent an MRI scanning session, which was followed by a multi-item lunch. Participants were aware that their lunch time meal would be video recorded. However, participants were not explicitly told that their food intake would be measured or that mimicry would be later examined. Three groups of participants were recruited as part of the larger study; adolescents with type 2 diabetes, overweight and obese adolescents (without type 2 diabetes) and healthy weight adolescents (without type 2 diabetes). See supplemental material for more detailed information about the selection criteria for the larger study.

## **Participants**

From the original data collected we were unable to use ten videos due to equipment failure or error and one video was excluded because the participant did not eat anything. In addition, we opted to focus on female adolescents only, due to the consistency of which social influence effects have been replicated amongst females (Hermans et al., 2012; Pliner and Mann., 2004; Roth et al., 2001) and there only being a small number of videos of adolescent males available. Therefore, nine videos of adolescent males were not coded or analyzed. Thus, the total sample for the present research consisted of 38 dyads containing female adolescents eating with a parent. See Table 1 for sample ethnicity and socio-economic status. There were 33 female parents and 5 male parents. The adolescents were aged 12.0 - 18.8 years, with a mean age of 15.4 years, SD = 1.9. Adolescent weight categories were classified according to the defined International Obesity Task Force age specific cut offs (Cole et al, 2000, Cole et al, 2007). Eleven of the adolescents were classed as being in the healthy weight range (BMI 18.5-24.9), fourteen were classed as overweight and obese (BMI 18.5-24.9), fourteen were classed as overweight mean adolescent BMI 18.5-24.90, fourteen were classed as overweight mean adolescent BMI 18.5-24.90, fourteen were classed as overweight mean adolescent BMI 18.5-24.90, fourteen were classed as overweight mean adolescent BMI 18.5-24.90, fourteen were classed as overweight mean adolescent BMI 18.5-24.90, fourteen were classed as overweight mean adolescent BMI 18.5-24.90, fourteen were classed as overweight mean adolescent BMI 18.5-24.90, fourteen were classed as overweight mean adolescent BMI

30.6, SD = 9.7. Mean parental BMI = 30.1, SD = 5.8. See Table 2 for adolescent and parental BMI information for the healthy weight, overweight and obese, and diabetic groups separately.

For our planned analyses we did not have any hypotheses relating to whether the weight or diabetes status of adolescent participants would moderate or influence any tendency to mimic parental eating, because social influence on food intake has been shown to be a relatively consistent effect and observed to a similar degree in both healthy weight and overweight individuals (Conger et al., 1980, Herman et al., 2003, Robinson et al., 2014). We did however check if this was the case by conducting our planned analyses (see later section) and included adolescent group (healthy weight, overweight and obese, diabetic) as an additional factor.

There was no evidence that adolescent group significantly moderated any mimicry effects (p > 0.05). Thus, as the number of adolescents in each group was relatively small and we did not have strong a-priori hypotheses, the results we report throughout are for all adolescent participants combined.

#### Lunch time meal

All sessions took place in an eating laboratory at the University of Birmingham. The room was furnished with a table and two chairs. Adolescents and parents were served a standardized multi-item meal each on separate trays. Each lunch item was on a separate plate and the meal consisted of a cheese sandwich (369 kcals), an individual Chicago Town cheese pizza (453 kcal), small bowl of cherry tomatoes (18kcal), an Activia strawberry yoghurt (123 kcal), an apple (45kcal), a Satsuma (18kcal), 25g Walkers ready salted crisps (131 kcal) and two Maryland double chocolate cookies (112kcal). A jug of water and 2 glasses was also provided. They were asked not to share food from each other's trays and told that they were

not expected to eat all the food, but to eat until they were full. The lunchtime meals were recorded using video cameras and participants were made aware of this prior to participating in the study.

#### **ANALYSIS**

- Strategy of analysis for overall food consumption
- Our first aim was to test whether there was evidence that adolescents' overall consumption may have been influenced by their parents' consumption. We did this by correlating the total amount of food adolescents ate (in kcals) with the amount of food their parent ate (kcals)
- using a Spearman's correlation.

Coding of video data

The first step in order to investigate whether there was evidence that the adolescents may have mimicked the eating of their parents was to code the video data by recording every time an adult or adolescent placed a food item into their mouth, the name of that food item (e.g. pizza) and the time that the food entered the mouth. All occurrences of eating were recorded by the first author. A random sample constituting 10% of these codings were independently checked by one of the other authors and there were no disagreements. The first author then coded each time an adolescent placed food into their mouth during the sensitive and nonsensitive time periods of the meal (see next section 'Defining sensitive and non-sensitive periods'). All of this coding was then cross-checked by an independent research assistant blind to the study hypotheses. Only a small number of discrepancies were noted (7 instances of mimicry were coded incorrectly, which constituted less than 1% of total coding) and they were resolved after discussion between the research assistant and lead author.

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

Defining sensitive and non-sensitive periods

Previous studies have examined if participants are more likely to eat a food item in the 5 or 15 seconds after a dining partner has placed food in their mouth (known as a 'sensitive period'), in comparison to the other periods of the meal when a partner had not recently placed food into their mouth (known as a 'non-sensitive period') (Hermans et al., 2012; Bevelander et al., 2013; Larsen et al., 2010). In the present study we examined three sensitive time period cut off points (+2, +5, +15 seconds), because we reasoned that mimicry may also occur in a shorter time frame (i.e. within + 2 seconds of a person eating) than previous studies have tested, as mimicry has been suggested to be automatic (Iacoboni et al., 1999). The three timeframe cut off points (+2, +5, +15) were treated as *separate* timeframes. Each meal was split into sensitive (the times during the meal in which a parent had recently placed food into their mouth) and non-sensitive time periods (all other times during the meal; i.e. the times during the meal in which a parent had not recently placed food in their mouth) for each of the three separate time frames (+2, +5, +15). This approach would allow us to test whether the rate at which adolescents placed food into their mouth differed between sensitive vs. nonsensitive periods, for the three time frames individually. See 1 for a detailed example. We presumed that if adolescents ate at a quicker rate during sensitive vs. non-sensitive periods, this would constitute evidence of mimicry. We calculated the rate of placing food into the mouth (defined as a consumption ratio, see next section) as opposed to just the number of times food was placed in the mouth, in order to account for there being differences in total sensitive vs. non-sensitive time during each meal.

214

215

Strategy of analysis for mimicry

As noted, we coded how frequently adolescents placed food items into their mouth during the sensitive periods (times when the parent **had** recently placed food in their mouth) and during the non-sensitive periods (times when the parent **had not** recently placed food in their mouth) of the lunchtime meal, for the three time frames separately. We then quantified this formally by computing 'consumption ratios'; the number of times a food item was placed into an adolescents' mouth per second<sup>2</sup>. Following this we compared the consumption ratio observed for the sensitive periods vs. non-sensitive periods of the meal using a Wilcoxon signed ranks test<sup>3</sup> for the three different time frames individually (+2, +5, +15). We adjusted the analyses using a Bonferroni correction to account for multiple comparisons. This allowed us to compare the consumption ratios (the number of times a food item was placed into an adolescents' mouth per second) for the periods of the meal in which a parent had recently placed into their mouth vs. periods of the meal in which the parent had not recently placed food into their mouth. Importantly, we computed these consumption ratios for both *non-specific* food item mimicry and *specific food* item mimicry.

#### Non-specific food item mimicry

In order to compute consumption ratios for **non-specific** food item mimicry we used the aforementioned analysis strategy and examined the rate at which adolescents placed **any** food item into their mouth during the sensitive periods vs. the rate at which adolescents placed **any** food into their mouth during the non-sensitive periods. This analysis allowed us to examine whether adolescents more frequently placed **any** food item in their mouth in periods when their parent had recently placed **any** food item in their mouth, as opposed to periods of the meal when a parent had not recently placed **any** food in their mouth.

## Specific food item mimicry

In order to compute consumption ratios for **specific** food item mimicry here we instead examined the rate at which adolescents placed the **same** food item into their mouth which their parent had placed in their mouth in the proceeding 2, 5, or 15 seconds (sensitive period) vs. times when the parent **had not** placed a food item into their mouth in the proceeding 2, 5, or 15 seconds (non-sensitive periods). This analysis allowed us to examine whether adolescents more frequently placed a food item in their mouth in the periods of the meal in which their parent had recently placed the **same** food item in their mouth, as opposed to all other time periods of the meal.

Thus, we were able to examine whether there was evidence of *specific* food item and *non-specific* food item mimicry using +2, +5 and +15 time frames individually.

## **RESULTS**

254 Total food intake

Parents ate a mean of 816.1 ( $\pm$ 204.8) calories during the lunchtime meal and adolescents ate a mean of 697.6 ( $\pm$ 238.3) calories during the meal. A Spearman's correlation showed that the amount eaten by the parents and children was significantly correlated [r (38) = .49, p < .001], whereby a parent eating a larger number of calories was associated with their adolescent child also eating a larger number of calories.

Meal length and frequency of food being placed into the mouth

Mean meal length was 18 minutes and 13 seconds (SD = 6.37). The mean number of times that parents placed any food item into their mouth was 59.50 (SD = 19.07). The mean number of times that adolescents placed any food item into their mouth was 77.84 (SD = 24.19). On average, parents placed food into their mouth every 19.88 seconds (SD = 8.98), which

constitutes a mean consumption ratio = 0.06 bites per second during the meal, while, adolescents placed food into their mouth every 14.53 seconds (SD = 4.93) on average, which constitutes a mean consumption ratio = 0.08 bites per second during the meal.

- Non-specific mimicry
- There was little evidence of non-specific food item mimicry during the meal. The consumption ratios for each of the three sensitive time periods were not significantly higher than the consumption ratios observed during the equivalent non-sensitive periods; +2 (z =-.17, p = .26, r = .03) + 5 (z = -1.47, p = .42, r = .24), and +15 (z = -2.27, p = .06, r = .37). See Table 3 for consumption ratio values. This indicates that the rate at which adolescents placed any food into their mouth (the consumption ratios) were similar in the periods of the meal in which their parent had recently placed any food into their mouth (sensitive periods) and all other periods of the meal in which their parent had not recently placed any food into their mouth (non-sensitive periods), regardless of whether 'sensitive' was defined as being within +2, +5 or +15 seconds after a parent had placed food into their mouth. Thus, it was not the case that adolescents were significantly more likely to place any food item into their mouth if their parent had recently placed a food item into their mouth.

- Specific mimicry
- For specific food items, there was evidence of mimicry for the +2 (z = -3.42, p < .001, r = .55), +5 (z = -3.90, p < .001, r = .63) and +15 (z = -3.73, p < .001, r = .60) second timeframes, as consumption ratios during these sensitive time periods were higher than the consumption ratios observed during the equivalent non-sensitive periods. See Table 3 for consumption ratio values. This indicates that the rate at which adolescents placed a food into their mouth was greater in the periods of the meal in which their parent had recently eaten that same food

item (sensitive periods) compared to the other remaining periods of the meal in which their parent had not recently eaten that same food item (non-sensitive periods), regardless of whether 'sensitive' was defined as being within +2, +5 or +15 seconds after a parent had placed food into their mouth. Thus, there was evidence that adolescents were significantly more likely to place a food item in their mouth if their parent had recently placed that same food item into their mouth.

# 299 DISCUSSION

The present study examined whether there is evidence that female adolescents may mimic their parents when eating together during a lunchtime meal. In line with previous work (Story et al., 2002), there was evidence of a positive correlation between parent and adolescent food consumption; adolescents consumed more calories during their lunch when their parent consumed more calories. We also examined if behavioural mimicry may underlie the influence that parents can have on their adolescents' eating behaviour. Results indicated that a parent placing a food item into their mouth was associated with an increased likelihood that their adolescent child subsequently picked up and ate the *same* food item during the following two, five and fifteen second periods. However, we did not find evidence that a parent placing a food item into their mouth was associated with an increased likelihood of their child placing *any* food item into their mouth in these time periods. Thus, adolescents appeared to mimic eating of specific food items only.

As in previous eating behaviour studies in adults and children (Hermans et al., 2012;

Bevelander et al., 2013) this observational data appears to support behavioural mimicry of eating. However, the current study expands on these previous studies as we found evidence of

behavioural mimicry in a different dyad than has previously been examined (adolescents and parents) and we were able to test whether adolescents mimicked the *specific* type of foods their parents were eating, or whether this process of mimicry was not food item specific; i.e. whether the parent placing a food into their mouth would simply increase the likelihood that the adolescent would place any food in their mouth. The findings of the present study suggest that adolescents were not simply synchronising their gestures or eating speed to match their parents (due to a lack of evidence for non-specific mimicry), as has been previously suggested as a potential explanation for social influence on eating (Hermans et al., 2012). Instead adolescents may have been using their parents as a reference point about which food items to eat and when, which could be interpreted through either a normative or informational account of social influence on eating (Robinson et al., 2013; Herman et al., 2003), although further studies will need to address this proposition more directly. The main novel finding of the present work was that we found evidence of specific food item mimicry during a shorter time frame (during the same or subsequent two seconds after a parent had placed food into their mouth), and within a different relationship than has been previously tested (Hermans et al., 2012; Bevelander, 2013), which suggests that there may be evidence for mimicry of eating behaviour in a shorter time frame than has been previously assumed.

333

334

335

336

337

338

339

340

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

One possibility is that we did not find evidence for non-specific mimicry (i.e. a parent placing food into their mouth was not associated with an increased likelihood that the adolescent subsequently placed *any* food into their mouth) because the rate of adolescent eating was relatively high during the meal. It could be argued that a high eating pace across all periods of the meal would make it difficult to observe differences between periods of the meal in which a parent had vs. had not recently eaten, possibly due to a form of ceiling effect. Further research examining food-item specific vs. non-food item specific mimicry in other meal

settings which promote a slower pace of eating would now be valuable. It is also possible that the influence parents appeared to have on adolescent eating may be in part explained by a form of visual attentional bias (Laibson, 2001, Wardle, 2007, Hardman et al, 2014), whereby adolescents visually followed parental gaze or hand movement to food choices, and parents visually attending to a specific food increased the likelihood that the adolescent then followed that cue and ate the same food.

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

341

342

343

344

345

346

A strength of the present study was that we examined parent-adolescent child dyads eating in a semi-naturalistic environment, rather than examining behavioural mimicry when a member of the dyad had been instructed on how much to eat (i.e. the confederate) (Hermans et al., 2012; Bevelander et al., 2013). Moreover, we examined mimicry during a multi-item lunch time meal and this allowed us to examine the extent to which adolescents mimicked specific food choices. It is not clear whether this finding of specific mimicry is unique to this dyad or whether it may occur in other relationships, therefore, further research is now needed. Due to the cross-sectional nature of the present study one possibility we cannot rule out is that some of the specific mimicry we observed may have been explained by the adolescents and parents already sharing similar meal/food item order preferences, thus, further work could build on the findings reported here by examining the effect of experimentally manipulating a parent's behaviour during a meal on the extent to which their adolescent child mimics this behaviour. One limitation that could also be addressed in further work is to investigate evidence of mimicry between adolescent males and their parents. Here our sample was female. However, recently Bevelander et al (2013) found that both male and female children (6-11 years old) were more likely to eat after witnessing a peer reaching for snack food than without such a cue. Therefore, it is possible that adolescent males may model the eating behaviour of their parents, and that mimicry may underlie this modelling. Finally, we did not examine whether

state (e.g. hunger) or trait (e.g. the quality of the relationship between the parent and adolescent) factors may have moderated the likelihood of mimicry, so further work designed to specifically explore the factors which may make mimicry more or less likely would now be valuable.

## Conclusions

This observational study suggests that when eating in a social context there is evidence that adolescent females may mimic their parental eating behaviour, selecting and eating more of a food item if their parent has just started to eat that food.

## Notes

Taking the +2 time frame as an example, the 'sensitive periods' of the meal were all seconds of the meal which occurred within the same or next 2 seconds after a parent had placed food into their mouth. The 'non-sensitive' periods of the meal were all other seconds during the meal. Likewise, for the +5 time frame, the 'sensitive periods' of the meal were all seconds of the meal which occurred within the same or next 5 seconds after a parent had placed food into their mouth. The 'non-sensitive' periods of the meal were all other seconds during the meal. Thus, for each participant the meal was split into 'sensitive' and 'non sensitive' time using three different sensitive period cut-off points (+2, +5, +15 seconds).

Consumption ratios were calculated by counting the number of times that the adolescent placed food into their mouth within a period and dividing this by the total amount of seconds in that period.

In the Wilcoxon signed ranks test the sensitive periods were deducted from the non-

<sup>3</sup> In the Wilcoxon signed ranks test the sensitive periods were deducted from the non-sensitive periods. The negative ranks indicate the sensitive periods while the positive ranks indicate the non-sensitive periods. No ties were observed in the analysis.

## 392 **REFERENCES**

194.

415

2	$\sim$	1
٦.	ч	۲.

Addessi, E., Galloway. A,T., Visalberghi, E., Birch LL. (2005) Specific social influences on 394 the acceptance of novel foods in 2-5 year old children. Appetite, 45, 264-71. 395 Bernieri, F.J. (1988) Coordinated movement and rapport in teacher-student interactions. 396 Journal of Nonverbal Behaviour, 12, 120-138. 397 Bevelander, K. E., Anschutz., D.J., Engels, R.C.M.E (2012) Social norms in food intake 398 among normal weight and overweight children. Appetite, 58, 864-872. 399 Bevelander, K.E., Lichtwarck-Aschoff, A., Anschutz, D.J., Hermans, R.C.J., Engels, 400 R.C.M.E. (2013) Imitation of snack food intake among normal-weight and overweight 401 children. Frontiers in Psychology, 4, 949. 402 403 Chartrand, T.L., & Bargh, J.A. (1999) The chameleon effect: The perception-behaviour link and social interaction, Journal of Personality and Social Psychology, 76, 893-910. 404 Chartrand, T. L., Maddux, W. W., and Lakin, J. L. (2009). "Beyond the perception-behavior 405 link: the ubiquitous utility and motivational moderators of nonconscious mimicry 406 2005," in The New Unconscious, Oxford Series in Social Cognition and Social 407 Neuroscience, eds R. R. Hassin, J. S. Uleman, and J. A. Bargh (New York, NY: 408 Oxford University Press), 334–361. 409 Cole, T. J., Bellizzi, M. C., Flegal, K. M., & Dietz, W. H. (2000). Establishing a standard 410 411 definition for child overweight and obesity worldwide: international survey. BMJ, *320*(7244), 1240-1243. 412 Cole, T. J., Flegal, K. M., Nicholls, D., & Jackson, A. A. (2007). Body mass index cut offs to 413 define thinness in children and adolescents: international survey. BMJ, 335(7612), 414

416 Conger, J.C., Conger, A.J., Philip, R., K.L., Matter, J.A. (1980). The effect of social cues on the eating behaviour of obese and normal subjects, Journal of Personality, 48, 258-417 271. 418 Goldman, S.J., Herman, C.P., Polivy, J. (1991) Is the effect of a social model attenuated by 419 hunger? Appetite, 17, 129-140. 420 Hardman, C.A., Scott, J., Field, M., & Jones, A. (2014) To eat or not to eat; the effects of 421 expectancy on reactivity to food cues. Appetite, 76, 153-160. 422 Harper, L.V., Sanders, K.M (1975). The effect of adults' eating on young children's 423 acceptance of unfamiliar foods. Journal of Experimental Child Psychology, 20, 206-424 2014 425 Herman, C.P., Roth, D.A., Polivy, J. (2003). Effects of the presence of others on food intake: 426 427 A normative interpretation. *Psychological Bulletin*, 129, 873-886. Hermans, R.C.J., Larsen, J.K., Herman, C.P., Engels, R.C.M.E. (2009) Effects of social 428 modelling on young women's nutrient dense food intake. Appetite, 53, 135-138 429 Hermans, R.C.J., Lichtwarck-Aschoff, A., Bevelander, K.E, Herman, C.P, Larsen, J.K, 430 Engels, R.C.M.E. (2012) Mimicry of food intake: The dynamic interplay between 431 eating companions. PLoS ONE, 7:e31027. Doi:10.1371/journal.pone.0031027 432 Iacoboni, M., Woods, R.P., Brass, M., Bekkering, H., Mazzoitta, J.C. et al (1999) Cortical 433 mechanisms of human imitation, Science, 286, 2526-2528. 434 435 Laibson, D. (2001) A cue-theory of consumption, The Quarterly Journal of Economics, 116, 81-119. 436 Lakin, J.L. & Chartrand, T.L. (2003) Using Nonconscious Behavioural Mimicry to Create 437 Affiliation and Rapport, *Psychological Science*, 14, 334-339. 438

Larsen, H., Engels, R.C.M.E., Souren, P.M., Overbeek, G.J., Granic, I. (2010) Peer influence 439 in the micro-perspective: imitation of alcoholic and non-alcoholic beverages, 440 Addictive behaviours, 35, 49-52. 441 Lau, R.R., Quadrel, M.J., Hartman, K.A (1990) Development and change of young adults' 442 preventive health beliefs and behaviour: influence from parents and peers. Journal of 443 Health and Social Behaviour, 31, 240-59 444 McGowan, L., Croker, H., Wardle, J., Cooke, L.J (2012) Environmental and individual 445 determinants of core and non-core food and drink intake in preschool-aged children in 446 447 the United Kingdom. European Journal of Clinical Nutrition, 66, 322-328. Neumann, R., & Strack, F. (2000) "Mood contagion": The automatic transfer of mood 448 between persons, Journal of Personality and Social Psychology, 79, 211-223. 449 450 Oliveria, S., Ellison, R., Moore, L., Gillman, M., Garrahie, E., Singer, M. (1992) Parent-child relationships in nutrient intake: The Framingham children's study. *American Journal* 451 of Clinical Nutrition, 56, 593-8 452 Pliner, P. & Mann N (2004) Influence of social norms and palatability on amount consumed 453 and food choice. Appetite, 42, 227-237 454 Robinson, E., Tobias, T., Shaw, L., Freeman, E., Higgs, S. (2011) Social matching of food 455 intake and the need for social acceptance. Appetite, 56, 747-752. 456 Robinson, E., Benwell, H., Higgs, S. (2013) Food intake norms increase and decrease snack 457 458 food intake in a remote confederate study. Appetite, 65, 20-24 Robinson, E., Blissett, J., Higgs, S. (2013) Social influences on eating: implications for 459 nutritional interventions, Nutritional Research Reviews, 26, 166-176. 460 Robinson, E., Sharps, M., Price, N., Dallas, R. (2014) Eating like you are overweight: The 461 effect of overweight models on food intake in a remote confederate study, Appetite, 462 *82*, 119-123. 463

464	Roth, D.A., Herman, C.P., Polivy, J., Pliner, P. (2001) self-presentational conflict in social
465	eating situations: A normative perspective. Appetite, 36, 165-171
466	Salvy, S.J., Elmo, A., Nitecki, L.A., Kluczynski, A., Roemmich, J.N. (2011) Influence of
467	parents and friends on children's and adolescents' food intake and food selection.
468	American Journal of Clinical Nutrition, 93, 87-92.
469	Salvy, S-J., de la Haye, K., Bowker, J.C., Hermans, R.C.J. (2012) Influence of peers and
470	friends on children's and adolescents' eating and activity behaviours. Physiology and
471	Behavior, doi:10.1016/j.physbeh.2012.03.022
472	Story, M., Neumark-Sztainer, D., French, S. (2002) Individual and environmental influences
473	on adolescent eating behaviours, Journal of the American Dietetic Association,
474	<i>102</i> ,40-51.
475	Sweetman, C., McGowan, L., Croker, H., Cooke, L. (2011) Characteristics of family
476	mealtimes affecting children's vegetable consumption and liking. Journal of the
477	American dietetic association, 111, 269-273
478	Vartanian LR, Sokol N, Herman CP, Polivy J (2013) Social Models Provide a Norm of
479	Appropriate Food Intake for Young Women. PLoS ONE 8(11): e79268.
480	doi:10.1371/journal.pone.0079268
481	Wardle, J. (2007) Eating Behaviour and Obesity, Obesity Reviews, 8, 73-75.
482	Wroten, K.C., O'Neil, C.E., Stuff, J.E., Liu, Y., Nicklas, T.A. (2012) Resemblance of dietary
483	intakes of snacks, sweets, fruit and vegetables among mother-child dyads from low
484	income families. Appetite, 59, 316-323.

 Table 1. Demographic information of sample

4	8	7

		Parent	Adolescent
Demographics		n = 38	n = 38
Ethnicity	White	50%	55.3%
	Asian	39.5%	36.8%
	Black	5.3%	2.6%
	Chinese	2.6%	2.6%
	Other/ Mixed	2.6%	2.6%
Income*	<£15,000	41.7%	n/a
	£15,000-60,000	44.4%	n/a
	212,000 00,000	11.170	11/ 6
	>£60,000	13.9%	n/a
	,		
Education level	Secondary school	21 100/	n/a
Education level	Secondary School	21.10/0	11/ a
	GCSE	28.90%	n/a
	A-level/ College	26.30%	n/a
	**		
	University	<b>-</b> 000 /	,
	Graduate	7.90%	n/a
	Post-graduate	15.80%	n/a

\*n=36 for income, information not available for 2 parents.

**Table 2.** Mean BMI (SD) for healthy weight, overweight and obese, and diabetic adolescent groups

	Healthy weight adolescents	Overweight and obese	Type 2 diabetic
	(n=11)	Adolescents (n=14)	(n=13)
Adolescent BMI	21.8 (1.7)	33.3 (6.9)	34.7 (11.6)
Parental BMI	26.1 (4.7)	32.1 (5.0)	31.3 (6.0)

**Table 3.** Consumption ratios for food item specific and non-food item specific mimicry during sensitive and non-sensitive periods (n=38)

	Food item specific mimicry		Non-food item specific mimicry	
	Sensitive	Non-sensitive	Sensitive	Non-sensitive
		+2 seconds		
Mean (SD)	0.02 (0.02)	0.01 (0.03)	0.08 (0.03)	0.08 (0.04)
Median	0.02*	0.01	0.07	0.07
		+5 seconds		
Mean (SD)	0.02 (0.02)	0.01 (0.01)	0.08 (0.03)	0.08 (0.05)
Median	0.02*	0.01	0.07	0.07
		+15 seconds		
Mean (SD)	0.02 (0.02)	0.01 (0.01)	0.07 (0.03)	0.10 (0.11)
Median	0.02*	0.01	0.07	0.07

Consumption ratios indicate the number of times per second adolescents placed a food item into their mouth within sensitive and non-sensitive periods. A higher ratio indicates a greater rate of placing food items into the mouth.

<sup>\*</sup>indicates a significant difference between the sensitive and non-sensitive consumption ratios at p < 0.01.