

# Cities are hotspots for threatened species.

Christopher D. Ives<sup>1\*+</sup>, Pia E. Lentini<sup>2\*</sup>, Caragh G. Threlfall<sup>3</sup>, Karen Ikin<sup>4</sup>, Danielle F. Shanahan<sup>5</sup>,  
Georgia E. Garrard<sup>1</sup>, Sarah A. Bekessy<sup>1</sup>, Richard A. Fuller<sup>5</sup>, Laura Mumaw<sup>1</sup>, Laura Rayner<sup>4</sup>, Ross  
Rowe<sup>4,6,7</sup>, Leonie E. Valentine<sup>8</sup>, Dave Kendal<sup>9</sup>

<sup>1</sup> School of Global, Urban and Social Studies, RMIT University.

<sup>2</sup> School of BioSciences, University of Melbourne

<sup>3</sup> School of Ecosystem and Forest Sciences, University of Melbourne

<sup>4</sup> Fenner School of Environment and Society, Australian National University

<sup>5</sup> School of Biological Sciences, University of Queensland

<sup>6</sup> Australian Government Department of the Environment

<sup>7</sup> National Environmental Research Program, Landscapes and Policy Research Hub, University of  
Tasmania

<sup>8</sup> School of Plant Biology, The University of Western Australia.

<sup>9</sup> Australian Research Centre for Urban Ecology

\* These authors contributed equally to this work.

+ Current address: Faculty of Sustainability, Leuphana University, Lüneburg Germany.

chris.ives@leuphana.de

**Keywords:** Australia, biodiversity, conservation policy, threatened species, urbanization, species  
distributions

**Running title:** The importance of cities for threatened species.

**Corresponding author:** Christopher Ives

**Word counts:** Abstract 296; Main body 4,118; References 1,158

26 **Abstract**

27

28 **Aim**

29 Although urbanisation impacts many species, there is little information on the patterns of threatened  
30 species occurrences in urban relative to non-urban areas. By assessing the extent of threatened  
31 species distributions across all Australian cities, we aim to investigate the currently under-utilised  
32 opportunity cities present to national biodiversity conservation.

33

34 **Location**

35 Australian mainland, Tasmania and offshore islands.

36

37 **Methods**

38 We assessed the distributions of Australia's 1,643 terrestrial threatened species and the extent to  
39 which they overlapped with 99 cities (of > 10,000 people), with all non-urban areas, and with  
40 simulated 'dummy' cities which covered the same area and bioregion as the true cities but were  
41 non-urban. We analysed differences between animals and plants, and examined variability within  
42 these groups using species accumulation modelling. Threatened species richness of true versus  
43 dummy cities was analysed using generalised linear mixed-effects models.

44

45 **Results**

46 Australian cities support substantially more nationally threatened animal and plant species than all  
47 other non-urban areas on a unit-area basis. Thirty percent of threatened species were found to occur  
48 in cities. Distribution patterns differed between plants and animals: threatened animals were  
49 generally distributed across multiple cities, while more individual plant species were found in each  
50 city with a greater proportion of their distributions occurring in urban areas. Individual cities tended  
51 to comprise unique suites of threatened species, and especially plants. The analysis of true versus

52 dummy cities demonstrated that, even after accounting for factors such as net primary productivity  
53 and distance to the coast, cities still consistently supported a greater number of threatened species.

54

55 Main conclusions

56 This research highlights that Australian cities are important for threatened species conservation, and  
57 that the species assemblages of individual cities are relatively distinct. National conservation policy  
58 should recognise that cities play an integral role when planning for and managing threatened  
59 species.

## 60 **1. Introduction**

61 Threatened species can be found in cities all over the world. Twenty-two percent of the known  
62 occurrences of endangered plants in the USA fall within the 40 largest cities (Schwartz *et al.*, 2002),  
63 and in an analysis of 54 cities Aronson *et al.* (2014) found that nearly a third are known to contain  
64 globally threatened birds. Indeed, the probability of a species being listed on the IUCN Red List  
65 increases with the percentage of its range that is urbanised (McDonald *et al.*, 2008). The reasons for  
66 this are becoming well understood: cities are often located in areas of high biological diversity  
67 (Luck, 2007), and urbanisation is a significant and expanding land use change that leads to habitat  
68 loss and fragmentation (Seto *et al.*, 2012). While the impacts of urbanisation on biodiversity are  
69 undeniable, this may also make cities especially important for achieving conservation outcomes.  
70 However, little is known about the relative importance of cities for conserving different kinds of  
71 organisms.

72

73 Urban areas occupy < 0.5% of the Earth's total land area (Schneider *et al.*, 2009), yet some  
74 threatened species are highly reliant on urban environments. For example, in the United Kingdom,  
75 the song thrush *Turdus philomelos*, a declining species of national conservation concern, occurs at  
76 densities more than three times higher in urban habitats than in the surrounding rural environment  
77 (Mason, 2000). The endangered Nielsen Park She-oak (*Allocasuarina portuensis*) also occurs  
78 exclusively within the metropolitan area of greater Sydney. Despite examples such as these, the  
79 designation of protected areas remote from human disturbance remains the dominant conservation  
80 paradigm worldwide (Miller & Hobbs, 2002). We have known for a long time that such wilderness  
81 thinking does not reflect ecological reality (Williams, 1980; Cronon, 1995). Yet conservation  
82 decision-making continues to implicitly, and sometimes explicitly, exclude urban environments  
83 from conservation investment (e.g. Sanderson *et al.*, 2002; Mittermeier *et al.*, 2003), as the negative  
84 pressures associated with urban development are seen to render urban habitats as 'lost causes' from

85 a biodiversity perspective (Cavin, 2013). By ignoring urban areas, important conservation  
86 opportunities are potentially missed.

87

88 On the Australian continent more than 1,600 species are considered threatened with extinction  
89 (Walsh *et al.*, 2013). Australian environmental policies and legislation are similar to those of other  
90 jurisdictions in that they tend to prioritise existing natural environments over disturbed or human-  
91 modified areas for biodiversity conservation or investment. Indeed, the second principle  
92 underpinning Australia's Biodiversity Conservation Strategy is that "*biodiversity is best conserved*  
93 *by protecting existing natural environments*" (Natural Resource Management Ministerial Council,  
94 2010, p16). Under the *Environment Protection and Biodiversity Conservation Act 1999* (EPBC  
95 Act), threats to listed species of conservation concern occurring in areas of highly modified or  
96 degraded habitat within city boundaries may be less likely to be deemed significant. This is because  
97 decision makers need to consider, among other factors, the "*sensitivity of the environment which*  
98 *will be impacted*", as well as whether the action will lead to a long-term decrease in the size of a  
99 population (Department of the Environment, 2013, p5). Consequently, certain projects within cities  
100 may not trigger impact assessment and approval requirements because the long-term viability of the  
101 population or habitat is assessed as having already been compromised. This set of circumstances,  
102 particularly in the case of small scale urban expansion, has the potential to lead to death by a  
103 thousand cuts, whereby incremental habitat destruction can lead to significant landscape-scale  
104 biodiversity loss (Dales, 2011; McCauley *et al.*, 2013).

105

106 The aim of this study is to assess the extent to which threatened species are reliant on conservation  
107 within cities. To explore this, we use the continent of Australia, which has very high endemic  
108 biodiversity (Chapman, 2009), as a case example, and investigate how the geographic distributions  
109 of species of national conservation concern overlap with urban areas. Specifically we measure how  
110 restricted threatened species' geographic ranges are to cities, and whether this is different for plants

111 versus animals. Finally, we explore the potential contribution that individual cities can make to  
112 biodiversity conservation by examining how the composition of threatened species varies in  
113 different cities across the continent.

114

## 115 **2. Methods**

### 116 *2.1 Threatened species and city data*

117 All 1,643 species (1,215 plants and 428 animals) that are considered to be of ‘national  
118 environmental significance’ under Australia’s EPBC Act were included in our analyses. This  
119 includes nationally-listed threatened species, native migratory species listed under international  
120 conventions or agreements, and marine species that use terrestrial areas for nesting (Commonwealth  
121 of Australia, 2014a). We hereafter refer to all of these species as ‘threatened species’. The listing  
122 criteria and categories used under the EPBC Act are adapted from those used to list species under  
123 the IUCN Red List of Threatened Species (Walsh *et al.*, 2013), with the main difference being the  
124 absence of a ‘near threatened’ category from the EPBC Act making the list more conservative  
125 (Commonwealth of Australia, 2014a). The majority of these species were from the flowering plant  
126 class Magnoliopsida (857 species) followed by lilies (Liliopsida, 289 species), birds (181 species),  
127 mammals (84 species), and reptiles (50 species).

128

129 Polygons representing the modelled distribution of each species were sourced from the Australian  
130 Department of the Environment’s ‘Environment Resources Information Network’ (Commonwealth  
131 of Australia, 2014b). The Australian Government uses these data to inform management and policy  
132 decisions and to undertake preliminary assessments of whether proposed developments or land use  
133 changes trigger targeted assessment and approval under the EPBC Act. The polygons were  
134 modelled from observation records, ecological data and research information provided from a range  
135 of Australian government, industry and non-government organisations, in addition to national-scale  
136 environmental data. For migratory species, distributions refer only to breeding sites, sites of

137 significance, or known locations rather than the entire range of the species. The polygons are not  
138 intended to be definitive maps of species occurrence, and generalisations made in the modelling  
139 process preclude detailed analyses of species distributions at fine scales. However, a reasonable  
140 level of spatial certainty is possible through classification of the polygons by the likelihood of  
141 species occurrence. For our analyses, only polygons where species are ‘known to occur’ (restricted  
142 to preferred habitat near observation records) and ‘likely to occur’ (preferred habitat within species  
143 range) were used. Polygons indicating where species ‘may occur’ (areas within environmental  
144 envelope or geographic region) were excluded. Polygons were projected to Geocentric Datum of  
145 Australia 1994 Australian Albers, and clipped to a shapefile representing terrestrial areas (the  
146 Australian mainland, Tasmania, and offshore territorial islands).

147

148 A layer representing the urban areas of Australia was derived from Australian Bureau of Statistics  
149 data (Section of State Ranges classification based on Statistical Area 1 polygons; Australian Bureau  
150 of Statistics, 2011a). This is a standard categorisation of land in Australia, used by government and  
151 non-government agencies. According to the dataset, land was classified as of “urban character” if:  
152 (i) the urban ‘Mesh Block’ (the smallest census unit) population is  $\geq 45\%$  of the total population of  
153 the Statistical Area 1 polygon and dwelling density  $\geq 45$  dwellings per sq km; or (ii) the population  
154 density is  $\geq 100$  persons per sq km and dwelling density  $\geq 50$  dwellings per sq km; or (iii) the  
155 population density is  $\geq 200$  persons per sq km (Australian Bureau of Statistics, 2011b, p19). Only  
156 urban polygons with populations  $> 10,000$  people were selected (hereafter referred to as ‘cities’ for  
157 simplicity), thereby excluding the smallest settlements. Following our criteria, the 99 cities in  
158 Australia cover  $17,420 \text{ km}^2$  (0.23% of terrestrial land mass), and range in size from  $10.5 \text{ km}^2$  for  
159 Nelson Bay, New South Wales, to  $2597.4 \text{ km}^2$  for Melbourne, Victoria (mean =  $175.3 \text{ km}^2$ , median  
160 =  $50.0 \text{ km}^2$ , SD =  $420.2 \text{ km}^2$ ). Although designated as ‘urban’ in character, the scale at which these  
161 areas were classified meant that they contained a range of land covers including built and natural  
162 lands.

163

164 *2.2 The importance of cities for threatened species*

165 Using ArcMap (v10.2, ESRI Redlands CA USA), we identified areas where the city polygons  
166 intersected with threatened species distribution polygons. From this, we calculated the proportion of  
167 each species' distribution that was urban and created a threatened species list for each city. To  
168 analyse the unique contribution of each city to the total assemblage of species located in urban  
169 areas, presence/absence species accumulation curves were generated using the 'specaccum'  
170 function in the 'vegan' package in R (R Core Team 2014, vers 3.1.0). We also generated a pairwise  
171 Jaccard dissimilarity matrix for the presence and absence of plant and animal species per city and  
172 carried out a hierarchical cluster analysis (using the 'average' linkage method and the 'hclust'  
173 function) to assess differences in community composition between cities. We then mapped mean  
174 dissimilarity values for each of the cities to help visualise patterns of beta diversity across the  
175 continent.

176

177 We converted the polygons representing threatened species to 1 km<sup>2</sup>-resolution binary rasters using  
178 the 'rasterize' function in R's 'raster' package (vers 2.2-31). Raster cells were given a value of 1 if  
179 the centre of the cell overlapped with the associated polygon, or 0 if there was no overlap. We  
180 calculated the number of threatened species that were known or likely to occur in each cell by  
181 summing the values across all of the threatened species rasters.

182

183 As a conservative comparative analysis, we repeated the processes outlined above using only those  
184 polygons that represented where species were 'known' to occur. As the difference between these  
185 analyses was minimal (see Appendix S1) we consequently present only the results from the  
186 combined 'known' and 'likely' distributions here, as this includes the larger complement of species.

187

188 *2.3 Mixed-effects models to account for potentially confounding factors*



189 To account for potentially confounding environmental variables that might influence the threatened  
190 species richness of a city irrespective of urbanisation, for each of our 99 ‘true’ cities we generated a  
191 paired ‘dummy’ city of equivalent area which was randomly positioned within the same bioregion  
192 (of which there are 89 across Australia). We then calculated both total threatened species richness  
193 of each true and dummy city, and the mean richness of the raster cells that comprised them. Both  
194 total and mean threatened species richness were analysed using mixed-effects regression models in  
195 the ‘lme4’ package in R. Total threatened species richness was fitted as a generalised linear mixed-  
196 effects model against a Poisson distribution using a log link with the ‘glmer’ function, and mean  
197 threatened species richness as a linear mixed-effects model with the ‘lmer’ function. The models  
198 were fitted with five fixed predictor variables; (i) categorical city type (i.e. true v dummy), (ii) mean  
199 net primary productivity (NPP, calculated as the mean across the months of 2014 and downloaded  
200 as a 0.1 degree raster from NASA Earth Observations 2015), (iii) city area, (iv) distance from the  
201 coast (measured from the nearest city edge), and (v) latitude. Continuous variables were centred and  
202 scaled prior to the analysis. The bioregion in which the true or dummy city occurred was fitted as a  
203 random effect in both models. We also noted that protected areas made up a substantially smaller  
204 proportion of the landmass in the true cities (mean =  $0.03 \pm 0.17$  SD) than the dummy cities (mean  
205 =  $0.12 \pm 0.33$  SD), but because this was strongly correlated with city type it was not included in the  
206 models.

207

### 208 **3. Results**

#### 209 *3.1 The distribution of threatened species in cities versus non-urban areas*

210 Of the 1,643 threatened species in our analysis, 503 (30%) had distributions that intersected with  
211 cities. This proportion differed for plants and animals, with 25% of listed plants and 46% of listed  
212 animals having at least part of their distributions located in cities. Species distribution size varied  
213 considerably (many species had relatively small distributions and only a small number had very  
214 large distributions) but distribution size was not strongly correlated with the proportion of a species’

215 distribution located in cities (Spearman's  $\rho = 0.33$ ). The distributions of animals (mean = 4.5  
216 million ha, median = 63,743 ha) tended to be much larger than those of plants (mean = 240,000 ha,  
217 median = 13,463 ha). Threatened species richness was higher in coastal areas and around the edges  
218 of cities (Fig. 1).

219

220 < Figure 1 >

221

222 There was substantial variation in the degree to which the distributions of threatened species  
223 included cities; species that were at least partially urban were found in an average of six cities  
224 ( $\pm 11.8$  SD). While some species were found in many cities (e.g. the eastern great egret *Ardea*  
225 *modesta* was found in 90 urban settlements), 258 threatened species (51%) occurred in one urban  
226 settlement only (Fig 2a). The distributions of eight threatened species (all plants) entirely  
227 overlapped with cities, while 51 (10%) of the 503 threatened species found in cities had >30% of  
228 their distribution in urban areas (Fig. 2b). Patterns were quite different for threatened plants and  
229 animals; plants tended to be found in fewer cities (mean =  $1.95 \pm 2.34$  SD) than animals (mean =  
230  $12.57 \pm 16.63$  SD) and were thus more spatially restricted, but had a larger proportion of their  
231 distribution in cities (plant mean =  $0.16 \pm 0.26$  SD, animal mean =  $0.04 \pm 0.08$  SD, Fig. 2).

232

233 < Figure 2 >

234

### 235 3.2 The importance of cities for threatened species

236 All 99 cities were known or likely to contain threatened animal species, and 88 cities (89%)  
237 contained threatened plant species or appropriate habitat (see Appendix S2 for city-specific details).  
238 Cities coincided with the distributions of substantially more threatened species than all other non-  
239 urban areas on a per-unit-area basis (Fig. 3). This was true for both animals and plants, with a very  
240 high proportion of non-urban cells containing no threatened plant species. The mean threatened

241 species richness for 1 km<sup>2</sup> city cells was 10.04 ( $\pm$  3.79 SD), and 2.72 ( $\pm$  2.88 SD) for non-urban  
242 cells.

243

244 < Figure 3 >

245

246 On average, cities contained 32 threatened species ( $\pm$ 25.5 SD). Sydney contained the most  
247 threatened species (124 species), but only a few (large) cities contained a high diversity of  
248 threatened species (Fig. 4a). This was especially pronounced for plants, with only 12% of cities  
249 containing >10 threatened plant species (see Fig. 4a).

250

251 Individual cities contained distinct sets of threatened species, and contributed unique species to the  
252 total urban assemblage with no evidence of an asymptote in the threatened species accumulation  
253 curves (Figure 4b). This differentiation among cities was driven primarily by threatened plants.  
254 Hierarchical cluster analysis supported this result, demonstrating that few cities had a similar  
255 threatened species composition (Appendix S3, Fig S3.1 and S3.2). The mean Jaccard dissimilarity  
256 score between cities for animals was 26.94 ( $\pm$  3.63 SD), with Kalgoorlie-Boulder supporting the  
257 most unique animal assemblage and Port Macquarie the least (Fig. S3.3). Plant communities were  
258 even more dissimilar between cities, with a mean Jaccard dissimilarity score of 26.76 ( $\pm$  3.76 SD);  
259 Kempsey supported the most unique plant assemblage while Taree's assemblage was most similar  
260 to other cities (Fig. S3.4).

261

262 < Figure 4 >

263

264 Our comparison of true versus non-urban dummy cities reinforced the findings of our broader  
265 analysis. As noted above, total threatened species richness ranged from 2-124 for true cities (mean  
266 = 31.49,  $\pm$  25.39 SD), and for dummies this range was 1-61 (mean = 12.12,  $\pm$  11.07 SD). The mean

267 threatened species richness of cells was 0.19-18.36 for true cities (mean = 9.04,  $\pm$  3.78 SD), and  
268 0.02-14.07 for dummies (mean = 7.26,  $\pm$  3.88 SD).

269

270 Regression modelling demonstrated that non-urban dummy cities had consistently lower total  
271 threatened species richness (coefficient estimate -0.84,  $\pm$  0.05 SE) and mean 1 km<sup>2</sup> cell threatened  
272 species richness (-1.67,  $\pm$  0.42 SE) than the true cities, even once potentially confounding factors  
273 had been accounted for (Fig. 5, see Appendix S4 for all coefficient estimates). Other factors which  
274 appeared to have strong effects on threatened species richness included net primary productivity,  
275 which was positively associated with mean cell richness (1.15,  $\pm$  0.34 SE), and distance from the  
276 coast, which had a negative effect on both mean cell richness (-1.21,  $\pm$  0.38 SE), and total richness  
277 (-0.72,  $\pm$  0.09 SE, Fig. 5).

278

279 < Figure 5 >

280

## 281 **4. Discussion**

### 282 *4.1 The importance of cities for conservation*

283 This is the first study to demonstrate at a continental scale that cities contain more threatened  
284 species per unit area than non-urban areas. Our analyses have shown that all Australian cities  
285 harbour or are likely to harbour threatened species, and 30% of Australia's threatened species  
286 occur, or are likely to occur, in cities that cover only 0.23% of the total land area. The elevated  
287 importance of cities for threatened species richness remained evident even when accounting for  
288 other biogeographic factors that may affect species richness such as primary productivity, distance  
289 from the coast, and latitude. This extends on the findings of Schwartz *et al.* (2002), who revealed  
290 that 22% of the occurrences of US endangered plant populations were located in the 40 largest  
291 metropolitan areas (comprising 8.4% of the land area). We note, however, that these findings may  
292 be influenced by the fact that both Australian and US cities are relatively young on a global scale,

293 and may be carrying extinction debts (Hahs *et al.*, 2009). Further, it is likely that the regions defined  
294 as ‘urban’ in this study contain a more heterogeneous composition of land covers than other studies  
295 in the literature. We therefore reaffirm the need for clear definitions of urbanisation to be reported  
296 in urban biodiversity studies, as has been called for by other scholars (McDonnell & Hahs, 2013).

297

298 The greater richness of threatened species in cities compared with equivalent non-urban dummy  
299 cities was more pronounced for total threatened species richness than for mean cell threatened  
300 species richness (Fig. 5). This suggests that the assemblages of threatened species in cities vary  
301 more greatly across their area than equivalent non-urban areas. Cities are known to have high levels  
302 of landscape heterogeneity (Alberti, 2005), with patches of remnant habitat commonly interspersed  
303 with highly disturbed areas. This landscape configuration may favour a wider variety of threatened  
304 species, thus increasing beta diversity and contributing to the higher total threatened species  
305 richness observed in cities. This is plausible in Australia where native ecosystems commonly  
306 remain within and around cities and adjacent to other land uses (Bekessy *et al.*, 2012; Newton *et al.*,  
307 2001).

308

#### 309 *4.2 Spatial patterning of species distributions*

310 The composition of threatened species varies among cities (Fig. 4b, Appendix S3). This suggests  
311 that the pattern identified by Aronson *et al.*, (2014), whereby city biotas reflect regional species  
312 pools, extends to threatened species. This trend may be especially pronounced in Australia given  
313 that the cities included in our study cover a vast spatial area with huge variation in environmental  
314 conditions. Patterns were different for plants and animals. Unique sets of threatened plants were  
315 found in individual cities, while threatened animals tended to be found in multiple cities (Fig. 4b).  
316 These results strongly suggest that all cities ought to be considered carefully in threatened species  
317 conservation and management.

318

319 We found that a small subset of threatened species were highly restricted to cities, and that this  
320 pattern was more pronounced for plants than it was for animals. Individual plant species were  
321 usually found within few cities, however a large proportion of their distribution was contained  
322 within those cities. In contrast, few animal species had a substantial share of their distributions  
323 located in cities (Fig. 2b). Most threatened plants in our dataset have relatively small distributions,  
324 and would be considered local endemics that are unique to certain bioclimatic regions of Australia.  
325 For example, the fringed spider-orchid *Caladenia thysanochila* is an endangered species with a  
326 small distribution, found entirely within a rapidly urbanizing region of Melbourne, Victoria  
327 (Department of the Environment, 2014). In contrast, some animals had very large distributions,  
328 occurring in 30 or more cities (Fig. 2a). This pattern of distribution for plants likely contributes to  
329 our finding of higher total threatened species richness per city than mean cell threatened species  
330 richness. Our finding that some threatened plants are found exclusively in urban environments is  
331 similar to that for North American floras (Schwartz *et al.*, 2002) and highlights that cities can be  
332 important for the conservation of rare and unique plants.

333

#### 334 *4.3 Implications for conservation policy and practice*

335 The disproportionate representation of threatened species in Australian cities identified in this study  
336 suggests that practitioners should seek to identify and act upon conservation opportunities in urban  
337 environments. It is important to note, though, that cities contain both threats and opportunities for  
338 biodiversity conservation. The animals in our dataset included several nationally migrant and  
339 nomadic species, such as the grey-headed flying-fox, *Pteropus poliocephalus* (Eby & Collins, 1999)  
340 and swift parrot, *Lathamus discolor* (Swift Parrot Recovery Team, 2001), that move across large  
341 areas as food resources (e.g. nectar, fruit or blossoms) become seasonally available. Often these  
342 resources are found in non-remnant, human-modified habitats. Indeed, Carnaby's black cockatoo,  
343 *Calyptorhynchus latirostris*, relies on an introduced pine plantation within the city of Perth for food,  
344 despite the fact that this represents a comparatively small proportion of their range (Valentine *et al.*,

345 2014). Cities may be especially valuable to these kinds of species, as they can provide more stable  
346 resources throughout the year as a result of human planting selection and supplementary watering  
347 (Parris & Hazell, 2005; Williams *et al.*, 2006). In contrast, other species rely on remnant patches of  
348 vegetation for their survival, many of which are under threat or in a degraded condition. The fringed  
349 spider-orchid, for example, is unlikely to persist if its remaining historical habitat is developed for  
350 housing, and its occurrence may even represent an extinction debt given the amount of habitat  
351 remaining. Irrespective of whether threatened species are threatened by urbanisation or supported  
352 by urban conditions, this study highlights the need for conservation action in cities. Depending on  
353 the nature of conservation threats and opportunities, a suite of conservation tools should be  
354 employed, such as spatial planning of urban development (e.g. Bekessy *et al.*, 2012), focussed  
355 recovery planning, and active management, restoration, and improvement of habitats (Hahs *et al.*,  
356 2009; Standish *et al.*, 2012).

357

#### 358 *4.4 Caveats and future research opportunities*

359 As with any spatial data compiled from multiple sources over a period of time, our species data may  
360 contain mapping errors. The most pertinent errors are those of commission and omission as a result  
361 of incomplete and unequal sampling effort. Few systematic biodiversity surveys have been  
362 conducted in Australia, yet those that have been done have often excluded urban areas (e.g. the  
363 regional forest agreement process; Slee, 2001). On the other hand, it is possible that ad-hoc  
364 databases may have an over-representation of urban records, as survey effort will arguably be  
365 greater in more populous areas. Ultimately, despite any inaccuracies, the results presented here are  
366 noteworthy since the datasets are those used by decision makers when assessing development  
367 applications and generating species recovery plans. Nevertheless, while our conservative analysis  
368 indicated that modelling assumptions did not have a large impact on our inference relating to the  
369 distribution of threatened species in cities, future research could explore the role of possible  
370 sampling biases further.

371

372 Finally, we note that while presence of a population in a location does not indicate its fitness or  
373 long-term viability in that location, it signals a potential conservation opportunity. In their  
374 multidisciplinary review of 787 urban biodiversity conservation studies, Shwartz *et al.* (2014) found  
375 only eight papers reported similar or improved levels of population viability of species of  
376 conservation significance in urban areas compared to nearby greener environments. Yet they also  
377 note that only three studies specifically set out to test this condition of viability, all of which  
378 reported in the affirmative. From these results Shwartz *et al.* (2014) concluded that “the importance  
379 of urban areas for general conservation is not convincingly supported by scientific research” (p. 43).  
380 Nevertheless, we argue that even if threatened species experience lower levels of population  
381 viability in urban environments, their overrepresentation in these areas makes cities even more  
382 important for conservation management and planning, noting too that doing nothing may reduce  
383 viability even further. We echo Shwartz *et al.*’s call for further research into the population  
384 dynamics of significant species in cities as a way of shedding light on ecological mechanisms that  
385 influence species persistence, as it can help determine which specific conservation actions are  
386 required.

387

## 388 **5. Conclusion**

389 Using Australia as a case example, this study is the first to demonstrate at a continental scale that  
390 cities contain disproportionately more threatened species than equivalent non-urban areas. Some  
391 species (particularly plants) have a much greater proportion of their distribution within urban areas  
392 than others, and all Australian cities are home to different suites of threatened species. These  
393 findings highlight and reinforce the global importance of planning and managing urban landscapes  
394 to conserve biodiversity (Secretariat of the Convention on Biological Diversity, 2012). We  
395 recommend that practitioners seriously consider the contribution that urban environments could



396 make to national biodiversity conservation, and incorporate this information into species recovery  
397 planning.

398

### 399 **Acknowledgements**

400 This research arose from a workshop held in Bungendore, NSW, in 2013, supported by the  
401 Australian Research Council (ARC) Centre of Excellence for Environmental Decisions (CEED).  
402 CDI, PEL, KI, DFS, SAB, RAF and LEV are supported by the Australian Government National  
403 Environmental Research Program Environmental Decisions Hub (NERP ED). CDI, KI, GEG, SAB,  
404 LM, LR and LEV are supported by the ARC CEED. GEG is supported by The Myer Foundation.  
405 SAB and RAF are ARC Future Fellows. DFS is supported through ARC Discovery Grant  
406 DP120102857. DK is supported by the Baker Foundation. CGT is supported through ARC Linkage  
407 Grant LP110100686. We thank three anonymous reviewers and two editors whose comments  
408 substantially improved this manuscript.

409 **References**

- 410 Alberti, M. (2005) The Effects of Urban Patterns on Ecosystem Function. *International Regional*  
411 *Science Review*, 28, 168–192.
- 412 Aronson, M. F. J., La Sorte, F. A., Nilon, C. H., Katti, M., Goddard, M. A., Lepczyk, C. A.,  
413 Warren, P. S., Williams, N. S. G., Cilliers, S., Clarkson, B., Dobbs, C., Dolan, R., Hedblom,  
414 M., Klotz, S., Kooijmans, J. L., Kuhn, I., MacGregor-Fors, I., McDonnell, M., Mortberg, U.,  
415 Pysek, P., Siebert, S., Sushinsky, J., Werner, P. & Winter, M. (2014). A global analysis of the  
416 impacts of urbanization on bird and plant diversity reveals key anthropogenic drivers.  
417 *Proceedings of the Royal Society B*, 281(20133330).
- 418 Australian Bureau of Statistics (2011a). Australian Statistical Geography Standard (ASGS).  
419 Available online: <http://www.abs.gov.au/ausstats/abs@.nsf/Lookup/2901.0Chapter23102011>  
420 (Accessed 6 December 2014).
- 421 Australian Bureau of Statistics (2011b). Australian Statistical Geography Standard (ASGS):  
422 Volume 4 - Significant Urban Areas, Urban Centres and Localities, Section of State. Available  
423 online:  
424 [http://www.ausstats.abs.gov.au/ausstats/subscriber.nsf/0/1080B7CB374FC771CA257A980013](http://www.ausstats.abs.gov.au/ausstats/subscriber.nsf/0/1080B7CB374FC771CA257A980013D404/$File/1270055004_july%202011.pdf)  
425 [D404/\\$File/1270055004\\_july%202011.pdf](http://www.ausstats.abs.gov.au/ausstats/subscriber.nsf/0/1080B7CB374FC771CA257A980013D404/$File/1270055004_july%202011.pdf) (Accessed 6 December 2014).
- 426 Bekessy, S. A., White, M., Gordon, A., Moilanen, A., Mccarthy, M. A., & Wintle, B. A. (2012).  
427 Transparent planning for biodiversity and development in the urban fringe. *Landscape and*  
428 *Urban Planning*, 108, 140–149.
- 429 Cavin, J. S. (2013). Beyond prejudice: Conservation in the City. A case study from Switzerland.  
430 *Biological Conservation*, 166, 84–89.
- 431 Chapman, A. D. (2009). *Numbers of Living Species in Australia and the World*. A Report for the  
432 Australian Biological Resources Study September 2009. Australian Biodiversity Information  
433 Services, Toowoomba, Australia. URL <http://www.environment.gov.au/node/13866> (Accessed  
434 2 July 2015).

435 Commonwealth of Australia, (2014a). Environment Protection and Biodiversity Conservation Act  
436 1999. Available online: <http://www.environment.gov.au/epbc> (Accessed 6 December 2014).

437 Commonwealth of Australia, (2014b). Species of National Environmental Significance. Available  
438 online: <http://www.environment.gov.au/science/erin/databases-maps/snes> (Accessed 6  
439 December 2014).

440 Cronon, W. J. (1995). The Trouble with Wilderness: Or, Getting Back to the Wrong Nature. In  
441 *Uncommon Ground: Rethinking the human place in nature* (Vol. 1, p. 7).

442 Dales, J. (2011). Death by a thousand cuts: Incorporating cumulative effects in Australia's  
443 Environment Protection and Biodiversity Conservation Act. *Pac. Rim L. & Policy J.*, 20(1),  
444 149–178.

445 Department of the Environment, (2014). Policy statement for Melbourne urban development  
446 proposals needing consideration under Parts 7,8 and 9 of the EPBC Act. Department of the  
447 Environment, Australian Government, Canberra. Available online:  
448 [http://www.environment.gov.au/resource/melbourne-urban-development-%C2%96-policy-  
449 statement-environment-protection-and-biodiversity](http://www.environment.gov.au/resource/melbourne-urban-development-%C2%96-policy-statement-environment-protection-and-biodiversity) (Accessed 6 December 2014).

450 Department of Environment, (2013). Matters of national environmental significance, Significant  
451 impact guidelines 1.1, Environment Protection and Biodiversity Conservation Act 1999,  
452 Department of the Environment, Australian Government, Canberra. Available online:  
453 [http://www.environment.gov.au/epbc/publications/significant-impact-guidelines-11-matters-  
454 national-environmental-significance](http://www.environment.gov.au/epbc/publications/significant-impact-guidelines-11-matters-national-environmental-significance) (Accessed 6 December 2014).

455 Eby, P., & Collins, L. (1999). The distribution, abundance and vulnerability to population reduction  
456 of a nomadic nectarivore, the Grey-headed Flying-fox *Pteropus poliocephalus* in New South  
457 Wales, during a period of resource concentration. *Australian Zoologist*, 31(1), 240–253.

458 Hahs, A. K., McDonnell, M. J., McCarthy, M. A., Vesk, P. A., Corlett, R. T., Norton, B. A.,  
459 Clemants, S. E., Duncan, R. P., Thompson, K., Schwartz, M. W. & Williams, N. S. G. (2009).  
460 A global synthesis of plant extinction rates in urban areas. *Ecology Letters*, 12(11), 1165–73.

461 Luck, G. W. (2007). A review of the relationships between human population density and  
462 biodiversity. *Biological Reviews of the Cambridge Philosophical Society*, 82(4), 607–45.

463 McDonnell, M.J. & Hahs, A.K. (2013) The future of urban biodiversity research: Moving beyond  
464 the “low-hanging fruit.” *Urban Ecosystems*, 16, 397–409.

465 Mason, C. F. (2000). Thrushes now largely restricted to the built environment in eastern England.  
466 *Diversity and Distributions*, 6, 189–194.

467 McCauley, L. A., Jenkins, D. G., & Quintana-Ascencio, P. F. (2013). Isolated Wetland Loss and  
468 Degradation Over Two Decades in an Increasingly Urbanized Landscape. *Wetlands*, 33(1),  
469 117–127.

470 McDonald, R. I., Kareiva, P., & Forman, R. T. T. (2008). The implications of current and future  
471 urbanization for global protected areas and biodiversity conservation. *Biological Conservation*,  
472 141(6), 1695–1703.

473 Miller, J. R., & Hobbs, R. J. (2002). Conservation Where People Live and Work. *Conservation*  
474 *Biology*, 16(2), 330–337.

475 Mittermeier, R. a, Mittermeier, C. G., Brooks, T. M., Pilgrim, J. D., Konstant, W. R., da Fonseca,  
476 G. a B., & Kormos, C. (2003). Wilderness and biodiversity conservation. *Proceedings of the*  
477 *National Academy of Sciences of the United States of America*, 100(18), 10309–10313.

478 NASA Earth Observations (2015). Net primary productivity (1 month – terra/modis),  
479 [http://neo.sci.gsfc.nasa.gov/view.php?datasetId=MOD17A2\\_M\\_PSN](http://neo.sci.gsfc.nasa.gov/view.php?datasetId=MOD17A2_M_PSN) (Accessed 2 July 2015).

480 Natural Resource Management Ministerial Council (2010). Australia’s Biodiversity Conservation  
481 Strategy 2010-2030, Australian Government, Department of Sustainability, Environment,  
482 Water, Population and Communities, Canberra

483 Newton PW, Baum S, Bhatia K et al (2001) Human settlements theme, Australia state of the  
484 environment report 2001. CSIRO Publishing, Canberra, Australia

485 Parris, K. M., & Hazell, D. L. (2005). Biotic effects of climate change in urban environments: The  
486 case of the grey-headed flying-fox (*Pteropus poliocephalus*) in Melbourne, Australia.  
487 *Biological Conservation*, 124(2), 267–276.

488 R Core Team (2014). R: A language and environment for statistical computing. R Foundation for  
489 Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.

490 Sanderson, E. W., Jaiteh, M., Levy, M. a., Redford, K. H., Wannebo, A. V., & Woolmer, G. (2002).  
491 The Human Footprint and the Last of the Wild. *BioScience*, 52(10), 891.

492 Schneider, A., Friedl, M.A. & Potere, D. (2009) A new map of global urban extent from MODIS  
493 satellite data. *Environmental Research Letters*, 4, 044003.

494 Schwartz, M. W., Jurjavcic, N. L., & Brien, J. M. O. (2002). Conservation’s Disenfranchised Urban  
495 Poor. *BioScience*, 52(7), 601–606.

496 Secretariat of the Convention on Biological Diversity. (2012). *Cities and Biodiversity Outlook:  
497 Action and Policy*. Montreal. URL: <https://www.cbd.int/doc/health/cbo-action-policy-en.pdf>  
498 (Accessed 2 July 2015).

499 Seto, K.C., Güneralp, B. & Hutyrá, L.R. (2012) Global forecasts of urban expansion to 2030 and  
500 direct impacts on biodiversity and carbon pools. *Proceedings of the National Academy of  
501 Sciences of the United States of America*, 109, 16083–8.

502 Shwartz, A., Turbé, A., Julliard, R., Simon, L., & Prévot, A.-C. (2014). Outstanding challenges for  
503 urban conservation research and action. *Global Environmental Change*, 28, 39–49.

504 Slee, B., (2001). Resolving production-environment conflicts: the case of the regional forest  
505 agreement process in Australia. *Forest Policy and Economics* 3, 17–30.

506 Standish, R. J., Hobbs, R. J., & Miller, J. R. (2012). Improving city life: options for ecological  
507 restoration in urban landscapes and how these might influence interactions between people and  
508 nature. *Landscape Ecology*. 28(6): 1213–1221.

509 Swift Parrot Recovery Team (2001) Swift Parrot (*Lathamus discolor*) Recovery Plan 2001-2005.  
510 Tasmanian Department of Primary Industries, Water and Environment. URL

- 511 <http://www.environment.gov.au/resource/swift-parrot-lathamus-discolor-recovery-plan-2001->  
512 [2005](http://www.environment.gov.au/resource/swift-parrot-lathamus-discolor-recovery-plan-2001-) (Accessed 2 July 2015).
- 513 Valentine, L.E., Fisher, R., Wilson, B.A., Sonneman, T., Stock, W.D., Fleming, P.A. and Hobbs,  
514 R.J. (2014) Time since fire influences food resources for an endangered species, Carnaby's  
515 cockatoo, in a fire-prone landscape. *Biological Conservation*, 175: 1–9.
- 516 Walsh, J.C., Watson, J.E.M., Bottrill, M.C., Joseph, L.N. & Possingham, H.P. (2013) Trends and  
517 biases in the listing and recovery planning for threatened species: an Australian case study.  
518 *Oryx*, 47, 1–10.
- 519 Williams, N. S. G., Mcdonnell, M. J., Phelan, G. K., Keim, L. D., & Van Der Ree, R. (2006). Range  
520 expansion due to urbanization: Increased food resources attract Grey-headed Flying-foxes  
521 (*Pteropus poliocephalus*) to Melbourne. *Austral Ecology*, 31(2), 190–198.
- 522 Williams, R. (1980). Ideas of Nature. In *Problems in Materialism and Culture*. London: Verso.  
523

524 **Biosketch**

525 The authors of this study are Australian researchers with interests in urban ecological systems and  
526 biodiversity conservation. Together, a wide range of disciplines is represented including ecology,  
527 social science and environmental policy. This article is an output from a workshop funded by the  
528 Australian Research Council Centre of Excellence for Environmental Decisions. Many of the  
529 authors are affiliated with the Environmental Decisions Group (EDG): a network of conservation  
530 researchers working on the science of effective decision making to better conserve biodiversity.  
531 More details about EDG can be found at <http://www.edg.org.au/>

532

533 **Figure legends**

534 Figure 1. Threatened species richness across Australia, with darker colours representing greater  
535 richness. Urban areas are outlined in black. Cities shown in greater detail in boxes are (a) Perth, (b)  
536 Brisbane and (c) Melbourne.

537

538 Figure 2. Plots of (a) species ranked according to the number of cities in which they occur and (b)  
539 the proportion of their distributions that fall in cities. Species are ordered on the x-axes by their  
540 rank, with the species occurring in the most cities, or with the greatest proportion of their  
541 distribution as urban, assigned the rank of 1.

542

543 Figure 3. The proportion of 1 km<sup>2</sup> cells in Australia, classified as either urban (white) or non-urban  
544 (grey) which support different numbers of threatened species. Data are presented for (a) all  
545 threatened species, (b) animals and (c) plants. Bars being skewed to the left of the plots indicates  
546 that a greater proportion of cells support fewer threatened species. Across Australia a small number  
547 of cells contained from 19 up to 32 threatened species, but the plot has been truncated at 18 along  
548 the x-axis because bars were not visible when the proportion was <0.005.


549

550 Figure 4. Plots of (a) ranked and (b) cumulative richness of threatened species in cities. The lack of  
551 asymptote in the species accumulation curves (b) suggests that each city contributes different  
552 species to the overall pool of threatened species found in urban areas.



553

554 Figure 5. Model curves comparing cities and equivalent ‘dummy cities’ within bioregions for (a)  
555 total threatened species richness, and (b, c) mean 1 km<sup>2</sup> richness of threatened species. Higher  
556 richness is consistently observed for cities, even once distance from the coast and net primary  
557 productivity are accounted for.






560 **Figure 2.**




561

562 **Figure 3.**




563  
564

565 **Figure 4.**




566



567 **Figure 5.**

568



569

570 **List of Supplementary Materials**

571

572 Appendix S1: Comparative analysis between known and known and/or likely to occur distributions

573

574 Appendix S2: List of Australian cities, with human population size and total, animal, and plant  
575 threatened species richness.

576

577 Appendix S3: Analysis of differences in threatened species composition between cities including  
578 hierarchical cluster analysis of (i) animals and (ii) plants, and maps of mean threatened species  
579 community similarity across Australia for (iii) animals and (iv) plants.

580

581 Appendix S4: Models of (i) total city threatened species richness, and (ii) mean 1km<sup>2</sup> cell  
582 threatened species richness for true cities versus dummy cities (non-urban controls).