

1 **Title:** Activity profiles of elite wheelchair rugby players during competition

2 **Submission Type:** Original investigation

3 **Authors:**

4 James M Rhodes<sup>1</sup>, Barry S Mason<sup>1</sup>, Bertrand Perrat<sup>2</sup>, Martin J Smith<sup>2</sup>, Laurie A Malone<sup>3</sup>,  
5 Victoria L Goosey-Tolfrey<sup>1</sup>

6 **Affiliations:**

7 <sup>1</sup>Peter Harrison Centre for Disability Sport, School of Sport, Exercise and Health Sciences,  
8 Loughborough University, UK.

9 <sup>2</sup>Faculty of Engineering, University of Nottingham, UK

10 <sup>3</sup>Research Department, Lakeshore Foundation, UAB/Lakeshore Foundation Research  
11 Collaborative, Birmingham, Alabama, USA

12 **Corresponding Author:**

13 Victoria L Goosey-Tolfrey

14 Peter Harrison Centre for Disability Sport,  
15 School of Sport, Exercise and Health Sciences,  
16 Loughborough University, UK.

17 Tel: +44 (0)1509 226386.

18 Email: [V.L.Tolfrey@lboro.ac.uk](mailto:V.L.Tolfrey@lboro.ac.uk)

19 **Running Head:** Activity profiles of wheelchair rugby

20 **Abstract Word Count:** 248

21 **Text-only Word Count:** 3215

22 **Number of figures:** 2

23 **Number of tables:** 4

## Abstract

**Purpose:** To quantify the activity profiles of elite wheelchair rugby and establish classification-specific arbitrary speed zones. Additionally, indicators of fatigue during full matches were explored. **Methods:** Seventy-five elite wheelchair rugby players from eleven national teams were monitored using a radio-frequency based, indoor tracking system across two international tournaments. Players who participated in complete quarters ( $n = 75$ ) and full matches ( $n = 25$ ) were included and grouped by their International Wheelchair Rugby Federation functional classification: group I (0-0.5), II (1.0-1.5), III (2.0-2.5) and IV (3.0-3.5). **Results:** During a typical quarter, significant increases in total distance (m), relative distance ( $\text{m}\cdot\text{min}^{-1}$ ), and mean speed ( $\text{m}\cdot\text{s}^{-1}$ ) were associated with an increase in classification group ( $P < 0.001$ ), with the exception of group III and IV. However, group IV players achieved significantly higher peak speeds ( $3.82 \pm 0.31 \text{ m}\cdot\text{s}^{-1}$ ) than groups I ( $2.99 \pm 0.28 \text{ m}\cdot\text{s}^{-1}$ ), II ( $3.44 \pm 0.26 \text{ m}\cdot\text{s}^{-1}$ ) and III ( $3.67 \pm 0.32 \text{ m}\cdot\text{s}^{-1}$ ). Groups I and II differed significantly in match intensity during very low/low speed zones and the number of high-intensity activities in comparison with groups III and IV ( $P < 0.001$ ). Full match analysis revealed that activity profiles did not differ significantly between quarters. **Conclusions:** Notable differences in the volume of activity were displayed across the functional classification groups. However, the specific on-court requirements of defensive (I and II) and offensive (III and IV) match roles appeared to influence the intensity of match activities and consequently training prescription should be structured accordingly.

**Keywords:** movement demands, performance analysis, fatigue, classification, Paralympic

## Introduction

Quantifying the activity profiles of elite athletes during competition facilitates the prescription of training programmes specific to the demands of the sport, which can optimise performance and minimise injury risk for individuals.<sup>1</sup> Typically, automatic video tracking techniques and global positioning systems (GPS) have been used to identify activity profiles within able-bodied (AB) team sports.<sup>2-6</sup> Unfortunately, owing largely to technological limitations, an accurate quantification of the activity profiles during indoor sports such as wheelchair rugby (WCR) remains relatively unknown.

A limited number of studies have previously investigated the activity profiles of WCR.<sup>7,8</sup> Spörner et al.<sup>7</sup> revealed that WCR players typically covered  $2364 \pm 956$  m at a mean speed of  $1.33 \pm 0.25 \text{ m}\cdot\text{s}^{-1}$  during match-play. Unfortunately, this information was derived

57 using a wheel-mounted data logger, which has been associated with inaccuracies during high  
58 speed ( $> 2.5 \text{ m}\cdot\text{s}^{-1}$ ) movements.<sup>9</sup> Moreover, the analysis was confined to recreational players  
59 and was therefore not representative of an elite population. Through the use of image-based  
60 processing techniques, Sarro et al.<sup>8</sup> reported that elite WCR players covered greater distances  
61 ( $4540 \pm 817 \text{ m}$ ) at a mean speed of  $1.14 \pm 0.21 \text{ m}\cdot\text{s}^{-1}$ . However, as a result of the time  
62 consuming analysis procedures involved using this method, the results were restricted to a  
63 small sample size ( $n = 8$ ).

64 Whilst only limited information regarding the volume of activity performed has been  
65 addressed in WCR, little is also known about the impact of functional classification on  
66 activity profiles. At present, WCR players are classified into one of eight classification  
67 groups based on their functional ability, ranging from 0 (least function) to 3.5 (most function).  
68 Previous research has shown classification-dependant trends in performance, with higher  
69 game-efficiency patterns,<sup>10</sup> and greater total distance and mean speed values (Sarro et al.,  
70 2010) associated with higher functional classifications. Moreover, Sarro et al.<sup>8</sup> also suggested  
71 that fatigue was more prominent in players with reduced function, due to a greater decrease in  
72 distance and mean speed values across match-halves. Despite this, previous research has  
73 demonstrated total distance to be a weak indicator of fatigue across competitive match-play  
74 in AB sports such as soccer.<sup>11,12</sup> Alternatively, high-intensity activities,<sup>4</sup> relative distance,<sup>13</sup>  
75 and peak speeds<sup>5</sup> have been advocated as better indicators of fatigue over time.

76 To further quantify the intensity of exercise during competition and training, activities  
77 have commonly been categorised into pre-determined arbitrary speed zones.<sup>2,11</sup> Arbitrary  
78 speed zones facilitate the longitudinal assessment of an athlete's performance over time.  
79 However, given that sprint performance has been shown to be dependent on functional  
80 classification in WCR,<sup>14,15</sup> the use of arbitrary speed zones for all classification groups is  
81 likely to misrepresent match-play intensity. Subsequently, recent studies have improved the  
82 specificity by relativizing speed zone design through the use of an individual's peak  
83 speed.<sup>16,17</sup> Whilst technological limitations have previously prevented the analysis of such  
84 variables in WCR, the recent development and validation of a radio-frequency based indoor  
85 tracking system (ITS) has enabled a broader assessment of elite WCR match-play to be  
86 possible.<sup>18</sup> Therefore, through the use of the ITS the aims of the current study were to: (1)  
87 quantify the demands of WCR between classification groups and to establish arbitrary speed  
88 zones specific to each classification; and (2) to explore any changes in activity profiles across  
89 full matches to establish indicators of fatigue in WCR.

90

## Methods

91

### 92 **Participants**

93 A total of 11 national WCR teams participated in the study with data collected across 21  
94 competitive matches over two international tournaments (2013 European and Americas Zonal  
95 Championships). Approval for the study was obtained from the International Wheelchair  
96 Rugby Federation (IWRF) and the organising committee of each tournament in addition to  
97 the University's local ethical advisory committee. Written informed consent was provided by  
98 each player prior to data collection. Data was collected from all consenting teams and players  
99 (age =  $32 \pm 7$  years), however data was only presented for players who completed complete  
100 quarters ( $n = 75$ ) or full matches ( $n = 25$ ). Players were categorised into four groups  
101 according to their IWRF classification, based on previous guidelines.<sup>14,19</sup> The breakdown of  
102 data collected from each group is presented in Table 1.

103 \*\*\*INSERT TABLE 1 HERE\*\*\*

### 104 **Design**

105 Data was collected during WCR matches using a radio-frequency based ITS (Ubisense,  
106 Cambridge, UK). Sensors were located in each of the four corners of the court, with two  
107 additional sensors parallel to the halfway line, to maximise court coverage. Each sensor was  
108 secured to ceiling beams above the court, at a height of 5-7 m (depending on venue). A  
109 calibration procedure outlined by Rhodes et al.<sup>18</sup> was performed at the beginning of each day.

110 Each player was equipped with a small, lightweight tag (size = 40 x 40 x 10 mm;  
111 mass = 25 g) which was fixed to the foot-strap of the wheelchair. Where a foot-strap was not  
112 feasible (e.g. lower limb amputees), the tag was securely mounted onto the wheelchair frame  
113 as close to the foot-strap region as possible. All players were familiarised with the tags during  
114 training sessions prior to the start of the competitions. Tags sampled at 8 Hz, which has  
115 previously been confirmed as an acceptable sampling frequency for the collection of  
116 movement parameters specific to WCR.<sup>18</sup> Data collection commenced at the beginning of  
117 each quarter and terminated at the end of the quarter. Collection was only paused during any  
118 periods of extended stoppages (time-outs, equipment calls) throughout each quarter since  
119 WCR players also remain active during the stopped game clock.<sup>8</sup> This resulted in a mean  
120 collection time of 15.1 ( $\pm 1.4$ ) minutes per quarter. Raw data files were exported using  
121 software developed specifically for WCR (Nottingham, UK).

### 122 **Measures**

123 Total distance (m) and relative distance covered ( $\text{m}\cdot\text{min}^{-1}$ ; relative to time spent on court),  
124 mean and peak speed ( $\text{m}\cdot\text{s}^{-1}$ ) was determined for each player during complete quarters of

125 WCR. Using an approach similar to Venter et al.<sup>16</sup> and Cahill et al.,<sup>17</sup> five arbitrary speed  
126 zones were established specific to each classification. Using the ‘mean’ peak speed ( $V_{max}$ )  
127 of each classification group the following five speed zones, relative to  $V_{max}$  were calculated:  
128 very low ( $\leq 20\% V_{max}$ ), low (21-50%  $V_{max}$ ), moderate (51-80%  $V_{max}$ ), high (81-95%  
129  $V_{max}$ ), and very high ( $> 95\% V_{max}$ ). The time spent in each of the arbitrary speed zones  
130 was calculated for each classification. Analyses of high-intensity (HI) activities (high and  
131 very high speed zones) were extended to include the total number of HI activities performed  
132 and both the mean and maximum duration and distance of these activities.

133 To assess the influence of fatigue on activity profiles across full matches of WCR,  
134 total distance (m), relative distance ( $m \cdot min^{-1}$ ), mean speed, peak speed ( $m \cdot s^{-1}$ ), and HI  
135 activities were explored. Only full match datasets (all 4 quarters completed by an individual)  
136 were analysed, with movement variables compared between quarters and halves.

### 137 **Statistical Analyses**

138 Data analysis was performed using the Statistical Package for the Social Sciences (SPSS  
139 version 21, Chicago, IL). Descriptive statistics (mean  $\pm$  standard deviation [SD]) were  
140 calculated for each participant for all movement variables. Normality and homogeneity of  
141 variance was confirmed by Shapiro-Wilk’s and Levene’s tests respectively. Since players  
142 differed in the number of repeated quarters they participated in and the varying sample sizes  
143 between classification groups, mixed linear modelling was applied to account for the  
144 unbalanced design.<sup>20</sup> Interactions between classification and quarter were also analysed using  
145 the full match datasets. Main effects and interactions were accepted as statistically significant  
146 whereby  $P \leq 0.05$ . Pairwise comparisons were utilised to explore any significant main effects  
147 between classification groups (I, II, III and IV), with a Bonferroni-corrected alpha level used  
148 to account for multiple contrasts ( $P = 0.008$ ). Effect sizes (ES), estimated from the ratio of  
149 the mean difference to the pooled standard deviation were also calculated. The magnitude of  
150 the effect size was classed as trivial ( $< 0.2$ ), small ( $\geq 0.2-0.6$ ), moderate ( $\geq 0.6-1.2$ ), large ( $\geq$   
151  $1.2-2.0$ ), and very large ( $\geq 2.0$ ) based on previous guidelines.<sup>21</sup>

## 152 **Results**

### 153 **Activity profiles during complete quarters of wheelchair rugby**


154 Functional classification significantly influenced the total distance, relative distance, mean  
155 speed and peak speed achieved during complete quarters of WCR ( $P < 0.001$ ). As  
156 demonstrated in Table 2, significant increases in total distance, relative distance and mean

157 speed were revealed with an increase in functional classification, except for groups III and IV  
158 ( $P \geq 0.704$ ;  $ES \leq 0.1$ ). Alternatively, peak speed was significantly higher as classification  
159 increased across all groups (Table 2).

160 \*\*\*INSERT TABLE 2 HERE\*\*\*

161 **Arbitrary speed zones.** The ‘mean’ peak speed values displayed (Table 2) established  
162 arbitrary speed zones specific to each classification group (Table 3). In general, WCR players  
163 spent 31% of a typical quarter in the very low speed zone, with the majority of time spent in  
164 the low speed zone (47%). The moderate speed zone accounted for 20% of the quarter  
165 duration, with 1.5% and 0.5% spent in the high and very high zones respectively. As  
166 illustrated in Figure 1, classification had no significant effect on the times spent in the  
167 moderate ( $P = 0.099$ ), high ( $P = 0.081$ ) and very high ( $P = 0.636$ ) speed zones. However the  
168 time spent in the very low and low speed zones was influenced by classification ( $P < 0.001$ ).  
169 Groups I and II spent a significantly greater time in the very low speed zone than groups III  
170 and IV ( $P < 0.001$ ;  $ES = 0.7 - 1.1$ ). Alternatively, groups III and IV spent a significantly  
171 greater time in the low speed zone, compared to groups I and II ( $P < 0.001$ ;  $ES = 0.8 - 1.4$ ).

172 \*\*\*INSERT TABLE 3 HERE\*\*\*



173  
174 **Figure 1** – Time spent (min) within five arbitrary speed zones between classification groups  
175 during a typical WCR quarter. #Significantly different to group III. †Significantly different to  
176 group IV. Data presented as means  $\pm$  SD.


177

178 **High-intensity activities.** The number of HI activities differed between classifications ( $P =$   
 179 0.005). As highlighted in Table 4, group I performed more HI activities than groups III ( $P =$   
 180 0.005; ES = 0.6) and IV ( $P = 0.004$ ; ES = 0.6). Classification had no significant effect on the  
 181 mean ( $P = 0.347$ ) and maximum ( $P = 0.629$ ) duration of HI activities. However a significant  
 182 main effect for the mean ( $P < 0.001$ ) and maximum ( $P = 0.031$ ) distance of each HI activity  
 183 was revealed. The mean distance of each HI activity was significantly greater for groups III  
 184 and IV compared to I and II (Table 4). Despite this, pairwise comparisons failed to reach  
 185 statistical significance between all classification groups for maximum distance ( $P \geq 0.009$ ;  
 186 ES  $\leq 0.5$ ).

187 \*\*\*INSERT TABLE 4 HERE\*\*\*

188 **Activity profiles during full matches of wheelchair rugby**

189 Total distance ( $P \geq 0.827$ ), relative distance ( $P \geq 0.963$ ), mean speed ( $P \geq 0.946$ ) and peak  
 190 speed ( $P \geq 0.944$ ) did not differ across quarters or halves (Figure 2). No significant changes  
 191 in the number ( $P \geq 0.964$ ), mean duration ( $P \geq 0.990$ ) maximum duration ( $P \geq 0.641$ ), mean  
 192 distance ( $P \geq 0.998$ ) or maximum distance ( $P \geq 0.592$ ) of HI activities performed were  
 193 identified across quarters and halves. Moreover, no interactions existed for any movement  
 194 parameter between classification group and quarters and match-half ( $P \geq 0.545$ ).



195

196 **Figure 2** – Total distance (a), relative distance (b), mean speed (c) and peak speed (d) values  
197 of each classification group during each of the four quarters in full matches of WCR. Data  
198 presented as means  $\pm$  SD.

199

200

## Discussion

201 The results of the current study revealed that functional classification is closely associated  
202 with the volume of activity elicited over typical quarters of WCR match-play. In addition, the  
203 ability to perform greater peak speeds increased with functional classification. Whilst the  
204 current study was the first to establish arbitrary speed zones for WCR, results revealed that  
205 match-play intensity was also influenced by functional classification, particularly during low  
206 speeds, which has practical implications on classification-specific training prescription.  
207 Furthermore, comparison of activity profiles across full WCR matches indicated no  
208 deterioration of physical performance was evident, regardless of functional classification.

209 The present study demonstrated that total distance, relative distance and mean speed  
210 values increased in association with higher functional classification across a typical quarter,  
211 yet no significant difference between classification groups III (2.0-2.5) and IV (3.0-3.5) was  
212 observed. Such findings are consistent with previous WCR match-play research, in which  
213 game efficiency patterns did not significantly differ between these classification groups.<sup>19</sup>  
214 Practical implications of these findings may impact upon team selection, in which group III  
215 players (2.0-2.5) do not seemingly restrict the functional ability of the team, whilst  
216 subsequently reducing the total on-court classification points (8.0 points permitted at any one  
217 time). This could partially explain why the present study observed a wider number of  
218 participants within group III ( $n = 28$ ) than in group IV ( $n = 13$ ). Despite this, sprint  
219 performance differed across all classification groups, with group IV capable of reaching  
220 significantly higher peak speeds ( $3.82 \pm 0.31 \text{ m}\cdot\text{s}^{-1}$ ) than groups I ( $2.99 \pm 0.28 \text{ m}\cdot\text{s}^{-1}$ ), II ( $3.44$   
221  $\pm 0.26 \text{ m}\cdot\text{s}^{-1}$ ), and III ( $3.67 \pm 0.32 \text{ m}\cdot\text{s}^{-1}$ ). This could, however, be attributed to the superior  
222 trunk function associated with higher classification groups.<sup>22</sup> While the ability to apply force  
223 to the hand-rim is a prerequisite for successful sprint performance, trunk function has  
224 previously been established as an important determinant of hand-rim force.<sup>22,23</sup> Subsequently,  
225 improved trunk function was likely to attribute to an increase in applied hand-rim force and,  
226 as such, greater peak speeds can be expected in higher functional players.<sup>23</sup> Nevertheless, the  
227 volume of activity along with the peak speeds performed during WCR match-play advocates  
228 the need for classification-specific training drills.



229           Given that peak speeds are influenced by functional classification, the use of arbitrary  
230 speed zones for all classification groups was likely to misrepresent individual intensity  
231 profiles. Thus, the creation of arbitrary zones specific to each classification group was an  
232 important outcome of the current study. Accordingly, the data suggested that elite WCR  
233 match-play is typically played at low speeds, with at least 75% of a typical quarter spent  
234 within the very low and low speed zones ( $\leq 50\%$  Vmax) regardless of functional  
235 classification. Specifically, groups I and II spent a significantly greater amount of time within  
236 the very low zone compared to groups III and IV. Such a finding may be attributed to the  
237 varying on-court roles, in which groups I and II (0-1.5) have previously been identified as  
238 low point players who predominantly occupy defensive roles, whereas groups III and IV (2.0-  
239 3.5) have been identified as high point players occupying offensive roles.<sup>24,25</sup> These on-court  
240 roles require low point players to ‘pick’ the opposition (block and trap opponents), which  
241 may account for the longer durations of static/very low speed activity. Alternatively, groups  
242 III and IV (2.0-3.5) spent significantly more time within the low speed zone, equating to 54%  
243 and 52% of the total quarter duration respectively, as opposed to groups I (39%) and II (41%).  
244 These findings indicate the contrasting intermittent match intensities between low and high  
245 point players, suggesting the need for role specific training drills.

246           The present data also indicated that HI activities were influenced by on-court roles  
247 during a typical WCR quarter. The significantly greater number of HI activities exhibited by  
248 low point (I & II) compared to high point players (III & IV) indicate that this is a key  
249 requirement for the defensive on-court role. The rationale for such a finding may be  
250 attributed to the fact that low point players do not possess the physical function of high point  
251 players,<sup>26</sup> and therefore must perform high intensity activities more frequently to compete  
252 with more functionally able opponents. Furthermore, typical HI durations of 1.7 to 1.9  
253 seconds were observed, with no significant differences across classification groups. This  
254 could be partly attributed to opposing players and court dimensions, preventing the capacity  
255 to generate prolonged durations of HI activities. Despite this, the higher speeds attained by  
256 high point players are likely to have attributed to the significant differences found in the  
257 mean distance of HI activities. Nevertheless, these findings further emphasize that on-court  
258 roles seem to dictate the intensity of activity profiles in WCR, highlighting the necessity for  
259 role specific training drills, in addition to classification-specific drills required for the volume  
260 of activity.

261           As part of the largest study to monitor activity profiles across full WCR matches, our  
262 results revealed elite WCR players covered approximately  $4213 \pm 626$  m at a mean speed of

263  $1.17 \pm 0.14 \text{ m}\cdot\text{s}^{-1}$ . These results were in accordance with the total distance ( $4540 \pm 817 \text{ m}$ )  
264 and mean speed values ( $1.14 \pm 0.21 \text{ m}\cdot\text{s}^{-1}$ ) previously reported by Sarro et al.<sup>8</sup> However, in  
265 contrast to Sarro et al.,<sup>8</sup> the present study revealed that activity profiles did not seem to  
266 deviate significantly across full WCR matches, suggesting match-play activity was not  
267 influenced by fatigue. Sarro et al.<sup>8</sup> further suggested that this decline was greater within low  
268 point players (distance - 9.9%; mean speed - 19.1%) than high point players (distance - 4.2%;  
269 mean speed - 10.1%). This would appear to suggest that WCR players now display far  
270 superior physical capabilities than the previous data from match-play collected in 2008.<sup>8</sup>  
271 Indeed, the advancement of sport science support and the development of conditioning  
272 strategies may partly explain these contrasting findings. The continuous roll-on substitutions  
273 in WCR may also attribute to these results, whereby if activity was perceived to be  
274 deteriorating then the likelihood is they would be substituted. Despite this, future analysis of  
275 game efficiency (e.g. ball-handling skills) across full matches may further contribute to the  
276 current understandings of fatigue during WCR match-play.

### 277 **Practical Applications**

278 In order to facilitate the development of WCR training programmes a better understanding of  
279 the match-play demands are required to improve the key training principles; specificity and  
280 individualisation of training. The current findings suggest that training programmes should be  
281 classification specific when related to activity volume, and designed to elicit the levels of  
282 aerobic demands sufficient to cope with match distances of up to ~4,600 m, combined with  
283 the anaerobic demands required for ~38 high-intensity bouts per match. Such programmes  
284 should also be extended to accommodate the various intensities attributed to the specific on-  
285 court roles of low (0-1.5) and high point players (2.0-3.5).

286 As identified in previous work, the ability to accelerate from a standstill is a key  
287 indicator of performance in WCR.<sup>27,28</sup> Yet owing to the sensitivity of the ITS when sampling  
288 at 8 Hz, a limitation of the current study was the inability to accurately measure acceleration  
289 values. Further work utilising the ITS alongside accelerometry technology may provide a  
290 more in-depth insight into the activity profiles during WCR match-play. However, as  
291 acceleration values over the first two pushes have previously been shown to range between  
292  $1.69$  and  $1.81 \text{ m}\cdot\text{s}^{-2}$  in elite WCR athletes,<sup>24</sup> such values would not have registered as HI  
293 activities within the present study, and as a consequent the true HI activities seen during  
294 match-play may be underestimated. Whilst the dynamic nature of WCR match-play has been  
295 explored in the present study, it is recommended that future research investigates the effect of  
296 situational variables (e.g. team rank, match outcome) on WCR activity profiles as seen in AB

297 sports<sup>2,3,29</sup> to establish which measures of performance are associated with successful  
298 performance.

299

300

### **Conclusions**

301 The present investigation demonstrated notable differences in the volume of activity profiles  
302 across functional classification during elite WCR match-play. Additionally, the use of  
303 individualised peak speeds in determining arbitrary speed zones provided new insights into  
304 the classification-specific differences in match-play intensity. However, these differences  
305 were exacerbated between groups I and II (0-1.5) compared with groups III and IV (2.0-3.5).  
306 Such differences can be attributed to the varying on-court roles of defensive (I and II) and  
307 offensive (III and IV) players. Furthermore, as opposed to previous reports, the match-play  
308 activities monitored in the current study were not shown to be associated with a physical  
309 decline across full WCR matches. The current results highlight the importance of both  
310 classification and role-specific training drills in WCR.

311

### **Acknowledgements**

312 The authors would like to acknowledge the financial support from UK Sport and the Peter  
313 Harrison Centre for Disability Sport. We would also like to further acknowledge the  
314 assistance of John Bishop at the IWRF, along with all the tournament organisers and the  
315 Lakeshore Foundation facilities team for their assistance and support. Appreciation is also  
316 extended to Jen Bryant for her assistance with data collection and to all the coaches and  
317 athletes who agreed to participate in this study.

- 319 1. Gabbett TJ, King T, Jenkins D. Applied physiology of rugby league. *Sports Med.*  
320 2008; 38:119-138.  
321
- 322 2. Rampinini E, Coutts AJ, Castagna C, Sassi R, Impellizzeri FM. Variation in top level  
323 soccer match performance. *Int J Sports Med.* 2007; 28:1018-1024.  
324
- 325 3. Di Salvo V, Gregson W, Atkinson G, Tordoff P, Drust B. Analysis of high intensity  
326 activity in premier league soccer. *Int J Sports Med.* 2009; 30:205-212.  
327
- 328 4. Bradley PS, Di Mascio M, Peart D, Olsen P, Sheldon B. High intensity activity  
329 profiles of elite soccer players at different performance levels. *J Strength Cond Res.*  
330 2010; 24:2343-2351.  
331
- 332 5. McLellan C, Lovell D, Gass G. Performance analysis of elite rugby league match play  
333 using global positioning systems. *J Strength Cond Res.* 2011; 25:1703-1710.  
334
- 335 6. Waldron M, Twist C, Highton J, Worsfold P, Daniels M. Movement and  
336 physiological match demands of elite rugby league using portable global positioning  
337 systems. *J Sports Sci.* 2011; 29:1223-1230.  
338
- 339 7. Spomer ML, Grindle GG, Kelleher A, Teodorski EE, Cooper R, Cooper RA.  
340 Quantification of activity during wheelchair basketball and rugby at the National  
341 Veterans Wheelchair Games: a pilot study. *Prosthet Orthot Int.* 2009; 33:210-217.  
342
- 343 8. Sarro KJ, Misuta MS, Burkett B, Malone LA, Barros RML. Tracking of wheelchair  
344 rugby players in the 2008 demolition derby final. *J Sports Sci.* 2010; 28:193-200.  
345
- 346 9. Sindall P, Whytock K, Lenton JP, et al. Criterion validity and accuracy of global  
347 positioning satellite and data logging devices for wheelchair tennis court movement. *J*  
348 *Spinal Cord Med.* 2013; 36:383-93.  
349
- 350 10. Molik B, Lubelska E, Kosmol A, et al. An examination of the international  
351 wheelchair rugby federation classification system utilizing parameters of offensive  
352 game efficiency. *Adapt Phys Act Quart.* 2008; 25: 335-351.  
353
- 354 11. Bangsbo J. The physiology of soccer-with special reference to intense intermittent  
355 exercise. *Acta Physiol Scand Suppl.* 1994; 619:1-155.  
356
- 357 12. Mohr M, Krstrup P, Bangsbo J. Match performance of high-standard soccer players  
358 with special reference to development of fatigue. *J Sports Sci.* 2003; 21:519-528.  
359
- 360 13. Austin DJ, Kelly SJ. Professional rugby league positional match-play analysis through  
361 the use of global positioning system. *J Strength Cond Res.* 2014; 28:187-193.  
362
- 363 14. Morgulec-Adamowicz N, Kosmol A, Molik B, Yilla AB, Laskin JJ. Aerobic,  
364 anaerobic, and skill performance with regard to classification in wheelchair rugby  
365 athletes. *Res Q Exerc Sport.* 2011; 82:61-69.

- 366 15. Goosey-Tolfrey VL, Leicht CA. Field-based physiological testing of wheelchair  
367 athletes. *Sports Med.* 2013; 43:77-91.  
368
- 369 16. Venter RE, Opperman E, Opperman S. The use of global positioning system (GPS)  
370 tracking devices to assess movement demands and impacts in under-19 rugby union  
371 match play. *Afr J Phys, Health Ed, Recr Dance.* 2011; 17:1-8.  
372
- 373 17. Cahill N, Lamb K, Worsfold P, Headey R, Murray S. The movement characteristics  
374 of English Premiership rugby union players. *J Sports Sci.* 2013; 31:229-237.  
375
- 376 18. Rhodes JM, Mason BS, Perrat B, Smith M, Goosey-Tolfrey VL. The validity and  
377 reliability of a novel indoor player tracking system for use within wheelchair court  
378 sports. *J Sports Sci.* 2014; doi: 10.1080/02640414.2014.910608.  
379
- 380 19. Morgulec-Adamowicz N, Kosmol A, Bogdan M, et al. Game efficiency of wheelchair  
381 rugby athletes at the 2008 Paralympic Games with regard to player classification.  
382 *Hum Mov.* 2010; 11:29-36.  
383
- 384 20. Cnaan A, Laird NM, Slasor P. Using the general linear mixed model to analyse  
385 unbalanced repeated measures and longitudinal data. *Stat Med.* 1997; 16:2349-2380.  
386
- 387 21. Batterham AM, Hopkin WG. Making meaningful inferences about magnitudes. *Int J*  
388 *Sports Physiol Perform.* 2006; 1:50-57.  
389
- 390 22. Vanlandewijck YC, Verellen J, Tweedy S. Towards evidence-based classification in  
391 wheelchair sports: impact of seating position on wheelchair acceleration. *J Sports Sci.*  
392 2011; 29:1089-1096.  
393
- 394 23. Vanlandewijck YC, Spaepen AJ, Lysens RJ. Relationship between the level of  
395 physical impairment and sports performance in elite wheelchair basketball athletes.  
396 *Adapt Phys Act Quart.* 1995; 12:139-150.  
397
- 398 24. Mason BS, van der Woude LH, Goosey-Tolfrey VL. Influence of glove type on  
399 mobility performance for wheelchair rugby players. *Am J Phys Med Rehabil.* 2009;  
400 88:559-570.  
401
- 402 25. Orr K, Malone LA. Wheelchair rugby. In: Goosey-Tolfrey VL, ed. *Wheelchair sport:*  
403 *A complete guide for athletes, coaches, and teachers.* Champaign, IL: Human  
404 Kinetics; 2010:151-166.  
405
- 406 26. van der Woude LH, Bouten C, Veegar HEJ, et al. Aerobic work capacity in elite  
407 wheelchair athletes. A cross sectional analysis. *Am J Phys Med Rehabil.* 2002;  
408 81:261-271.  
409
- 410 27. Vanlandewijck YC, Theisen D, Daly D. Wheelchair propulsion biomechanics:  
411 implications for wheelchair sports. *Sports Med.* 2001; 31:339-367.  
412

- 413 28. Mason BS, Porcellato L, van der Woude LH, Goosey-Tolfrey VL. A qualitative  
414 examination of wheelchair configuration for optimal mobility performance in  
415 wheelchair sports. *J Rehabil Med.* 2010; 42:141-149.  
416
- 417 29. Taylor JB, Mellalieu SD, James N, Shearer DA. The influence of match location,  
418 quality of opposition, and match status on technical performance in professional  
419 association football. *J Sports Sci.* 2008; 26:885-895.  
420

**Table 1.** A breakdown of classification profiles, datasets and sample size.

Group	IWRF Classification	Full Quarters	Participants ( <i>n</i> )	Full Matches	Participants ( <i>n</i> )
I	0-0.5	38	12	2	2
II	1.0-1.5	138	22	12	9
III	2.0-2.5	122	28	9	8
IV	3.0-3.5	108	13	12	6
	<b>Total</b>	<b>406</b>	<b>75</b>	<b>35</b>	<b>25</b>

**Table 2.** Descriptive statistics (mean  $\pm$  SD) for movement variables during a typical WCR quarter

Variables	<b>I</b> ( <i>n</i> = 38)		<b>II</b> ( <i>n</i> = 138)		<b>III</b> ( <i>n</i> = 122)		<b>IV</b> ( <i>n</i> = 108)	
	<i>Mean</i>	<i>SD</i>	<i>Mean</i>	<i>SD</i>	<i>Mean</i>	<i>SD</i>	<i>Mean</i>	<i>SD</i>
Total distance (m)	881 <sup>*#†</sup>	137	1011 <sup>#†</sup>	142	1155	196	1153	172
Relative distance (m·min <sup>-1</sup> )	59.9 <sup>*#†</sup>	6.5	69.7 <sup>#†</sup>	8.4	77.1	7.4	78.4	10.1
Mean speed (m·s <sup>-1</sup> )	1.01 <sup>*#†</sup>	0.11	1.15 <sup>#†</sup>	0.13	1.27	0.13	1.29	0.16
Peak speed (m·s <sup>-1</sup> )	2.99 <sup>*#†</sup>	0.28	3.44 <sup>#†</sup>	0.26	3.67 <sup>†</sup>	0.32	3.82	0.31

Note: *n* = number of datasets per classification group. \*Significant to group II ( $P < 0.05$ ); #Significant to group III ( $P < 0.05$ ); †Significant to group IV ( $P < 0.05$ ).

**Table 3.** Arbitrary speed zones ( $\text{m}\cdot\text{s}^{-1}$ ) as proposed for use within WCR

Zones		<b>I</b> ( $n = 38$ )	<b>II</b> ( $n = 138$ )	<b>III</b> ( $n = 122$ )	<b>IV</b> ( $n = 108$ )
Very low	$\leq 20\% V_{max}$	$\leq 0.60$	$\leq 0.69$	$\leq 0.73$	$\leq 0.76$
Low	$21-50\% V_{max}$	0.61-1.50	0.70-1.72	0.74-1.84	0.77-1.91
Moderate	$51-80\% V_{max}$	1.51-2.39	1.73-2.75	1.85-2.94	1.92-3.06
High	$81-95\% V_{max}$	2.40-2.84	2.76-3.27	2.95-3.49	3.07-3.63
Very High	$> 95\% V_{max}$	$> 2.84$	$> 3.27$	$> 3.49$	$> 3.63$

**Table 4.** Descriptive statistics (mean  $\pm$  SD) for HI activities performed during a typical WCR quarter

HI activities	<b>I</b> ( $n = 38$ )		<b>II</b> ( $n = 138$ )		<b>III</b> ( $n = 122$ )		<b>IV</b> ( $n = 108$ )	
	<i>Mean</i>	<i>SD</i>	<i>Mean</i>	<i>SD</i>	<i>Mean</i>	<i>SD</i>	<i>Mean</i>	<i>SD</i>
Number	13 <sup>#†</sup>	7	11	6	9	5	9	6
Mean duration (s)	1.7	0.8	1.7	0.7	1.8	0.8	1.9	0.8
Max duration (s)	4.3	1.9	4.2	1.9	4.4	2.3	4.0	1.8
Mean distance (m)	4.7 <sup>#†</sup>	2.3	5.4 <sup>#†</sup>	2.1	6.3	2.6	6.4	2.8
Max distance (m)	11.7	5.2	13.5	6.2	15.4	8.4	14.8	6.6

Note:  $n$  = number of datasets per classification group. <sup>#</sup>Significant to group III ( $P < 0.05$ ); <sup>†</sup>Significant to group IV ( $P < 0.05$ ).