

# One-Pot Cannizzaro Cascade Synthesis of *ortho*-Fused Cycloocta-2,5-dien-1-ones from 2-Bromo(hetero)aryl Aldehydes

Laurence Burroughs, Lee Eccleshare, John Ritchie, Omkar Kulkarni, Barry Lygo, Simon Woodward,\* and William Lewis


**Abstract:** An intramolecular Cannizzaro-type hydride transfer to an *in situ* prepared allene enables the synthesis of *ortho*-fused 4-substituted cycloocta-2,5-dien-1-ones with unprecedented technical ease for an eight-ring carboannulation. Various derivatives could be obtained from commercially available (hetero)aryl aldehydes, trimethylsilylacetylene, and simple propargyl chlorides in good yields.

Readily (often commercially) available 2-bromo(hetero)aryl aldehydes (**1**, Scheme 1) constitute powerful and popular units for the construction of *ortho*-fused carbocyclic rings.<sup>[1]</sup> Two strategies commonly employed in the multitudinous syntheses described in the literature involve 1) the derivatization of **1** to intermediates of type **A**, with a  $\pi$ -system attacking an electrophile that is ultimately derived from the aldehyde by functional-group manipulations (e.g., an alcohol, acetal, or even the aldehyde itself); or 2) formyl group homologation to an anion-stabilizing unit that attacks a suitably activated  $\pi$ -bond (**B**). Whereas many elegant catalytic systems that use intermediates **A** or **B** have already been described,<sup>[2,3]</sup> the preparation of the suitable precursors is often step-intensive, raising questions on the overall process

efficiency in the “age of sustainability”.<sup>[4]</sup> We speculated that a new annulation strategy, based on a Cannizzaro-type reaction via intermediate **C**, might be possible. As the hydride transfer simultaneously exposes a powerful Michael acceptor (ynone or equivalent) and a carbanion in close proximity, efficient annulation might be expected to occur (especially if **C** is directly attained *in situ*). Herein, we describe the use of such a strategy for the formation of eight-membered rings. The paucity of direct single-pot/step procedures to such medium rings attests to the known issues associated with their synthesis;<sup>[5]</sup> they therefore provide a stringent test for strategy **C**. Although redox-based Cannizzaro reactions, including asymmetric and triggered C–C bond-forming versions, have recently been developed,<sup>[6]</sup> to the best of our knowledge, they have not been employed for carboannulation reactions to medium rings. Genuine single-pot/step procedures to eight-membered rings are very limited—Repe’s historical catalytic access to cyclooctatetraene<sup>[7]</sup> and Murakami’s [4+2+2] approach<sup>[8]</sup> being notable exceptions. Modern alternatives frequently describe catalytic cyclizations that proceed with exquisitely high yields (for the final step). Unfortunately, their overall synthetic efficiency is often compromised by the multistep syntheses required to obtain the cyclization precursors.<sup>[9]</sup>

Initial investigation of our own proposal (**C**) centered on the combination of readily available 2-bromobenzaldehyde (**1a**), trimethylsilylacetylene (**2a**), and propargyl chloride **3a**. Encouragingly, even an initial run returned significant amounts of benzo[8]annulene derivative **4aaa** (Table 1, entry 1, ca. 7%) whose identity and regiochemistry was confirmed by X-ray crystallography (see the Supporting Information) as initial NMR data was not conclusive.<sup>[10]</sup> Unfortunately, under the conditions of entry 1, multiple side products were also produced. Upon changing from an organocopper to an organocuprate formulation (entry 2), the reaction became much cleaner, but the yield of **4aaa** remained low. The recovered side products of entry 2 included significant amounts of the alcohols **5a** and **6a** together with unreacted **3a**. We could confirm that formation of **D** was almost quantitative in < 20 min under all conditions tried (Et<sub>2</sub>O or THF, –78 to 0°C, based on recovered **5a** in independent reactions). We therefore suspected issues associated with both *n*BuLi–halogen exchange from **D** and the copper-promoted S<sub>N</sub>2’ reaction leading to the progenitor for eight-ring formation. Further experiments were designed to probe these hypotheses (Table 1).

Organometallic compounds derived from **D** were poorly soluble in Et<sub>2</sub>O so the reaction solvent was switched to THF (entry 3). This led to an improved reaction that enabled us to




**Scheme 1.** Proposed use of 2-bromo(hetero)aryl aldehydes in Cannizzaro-triggered annulation cascades compared to traditional approaches. EWG = electron-withdrawing group, LA = Lewis acid, R<sup>1</sup>–R<sup>3</sup> are generic groups, *n* and *m* are typically 0–1.

[\*] Dr. L. Burroughs, L. Eccleshare, J. Ritchie, O. Kulkarni, Prof. Dr. B. Lygo, Prof. Dr. S. Woodward, Dr. W. Lewis  
School of Chemistry, University of Nottingham  
University Park, Nottingham NG7 2RD (UK)  
E-mail: simon.woodward@nottingham.ac.uk

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201505347>.

© 2015 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.


**Table 1:** Optimization of the Cannizzaro-triggered cascade reaction to eight-ring compound **4aaa**.<sup>[a]</sup>


Entry	Solvent	Halogen exchange of intermediate <b>D</b>	Transmetalation with CuBr-SMe <sub>2</sub>	Addition of <b>3a</b> and cyclization	Yield <b>4aaa</b> [%]
1	Et <sub>2</sub> O	<i>t</i> BuLi (2 equiv), -50°C, 20 min	1 equiv, -50°C, 20 min	1 equiv, -50°C to RT over 5 h	< 10
2	Et <sub>2</sub> O	<i>n</i> BuLi (1 equiv), 0°C, 10 min	0.5 equiv, -50°C, 5 min	0.5 equiv, -50°C to +10°C over 1.2 h	14 <sup>[b]</sup>
3	THF	<i>n</i> BuLi (1 equiv), -50°C, 15 min	0.5 equiv, -50°C, 10 min	0.5 equiv, -50°C to 10°C over 1.2 h	53
4	THF	<i>n</i> BuLi (1 equiv), -50°C, 15 min	0.5 equiv, -50°C, 1 h	0.5 equiv, -50°C to -10°C over 1.5 h; kept at -10°C for 1 h	70 (69) <sup>[c]</sup>
5	THF	<i>n</i> BuLi (1 equiv), -50°C, 15 min	0.5 equiv, -50°C, 1 h	0.5 equiv, -50°C to -10°C over 1.5 h; kept at -10°C for 2 h	35

[a] Reaction conditions: **1a** (0.71 mmol), **2a** (0.77 mmol)/*n*BuLi (0.72 mmol), CuBr-SMe<sub>2</sub> (0.36–0.71 mmol), **3a** (0.35–0.70 mmol) in Et<sub>2</sub>O or THF (1.0 mL). The yields of **4aaa** were determined by GC analysis against genuine samples and an internal standard (1-methylnaphthalene, 50 μL). [b] The mass balance of the reaction included **3a**, **5a**, and **6a** in a 2.4:1.0:1.8 ratio. [c] The yield of isolated product for a reaction performed with 1.39 mmol of **3a** is shown in parentheses.

identify that the onset of the Cannizzaro-type cascade reaction occurs at -20 to -10°C for this substrate. Independent treatment of THF solutions containing **D** with *n*BuLi followed by quenching with D<sub>2</sub>O confirmed that complete Br/Li exchange had occurred within 15 min at -40 to -60°C, and that the resulting organolithium species was rather stable at these temperatures. However, above approximately -15°C, the same species began to decompose rapidly. Thus, -40 to -50°C was selected as the temperature range for optimizing the transmetalation to CuBr-SMe<sub>2</sub>. When the nominal diorganocuprate was prepared under the conditions of entry 3 (-50°C, 10 min, from the aryl lithium intermediate with CuBr-SMe<sub>2</sub>) and then treated with methyl vinyl ketone, very poor conversions into the 1,4-addition product were attained. This strongly suggested that transmetalation to Cu<sup>I</sup> is rather slow for this system. Gratifyingly, extending the time for Li/Cu exchange to one hour (entry 4) significantly increased the quantity of **4aaa** formed, and the GC yields were comparable to those attained through isolation (70% vs. 69%). Further extending the transmetalation time to two hours at -10°C had a detrimental effect on the yield of **4aaa** (entry 5).

With an optimized procedure for the preparation of parent compound **4aaa** in hand, we studied the scope of the reaction with respect to the 2-bromo(hetero)aryl aldehyde component **1**. Using the conditions of entry 4 (Table 1) and aldehydes **1b–1j**, the eight-ring compounds that are shown in Scheme 2 were attained. Substitution at the 3-position was better tolerated than at the 1- and 4-positions (products **4baa/4caa** vs. **4daa/4faa**). Monosubstitution at the 2-position was also tolerated. A comparison of the yields of **4baa–4gaa** indicates that steric factors are more important than electronic ones; for example, intramolecular OMe steric contacts increase the *ortho* steric profile of **4eaa** versus **4jaa**. The yields of compounds **4haa–4jaa** also support this trend. Whereas the yields of **4aaa–4jaa** are modest, it should be borne in mind that they represent one-pot processes to analytically pure crystalline products on up to multigram scales, using only inexpensive commercial starting materials. The new process compares very favorably to the consolidated



**Scheme 2.** Variation of the 2-bromo(hetero)aryl aldehyde; yields starting from the commercial aldehydes are given.<sup>[9]</sup> [a] The precursor aldehyde was obtained from isatin (3 steps).

yields and time efficiencies of previously reported eight-ring syntheses.<sup>[9]</sup>

By varying the acetylene component using the readily attained starting materials **2b–2f**, five further examples of eight-ring compounds **4** were synthesized (Scheme 3). The terminal alkyne substituent could encompass alkyl, vinyl, or aryl units, and the corresponding products were all obtained in acceptable yields. Finally, variation of the propargyl chloride was investigated using simple **3b–3i** (Scheme 3). The aryl moiety could accommodate both electron-donating and -withdrawing substituents and some functional groups (**4aae** and **4aaf**).

Whereas the full mechanistic features of our Cannizzaro-triggered cascade are still under investigation, a working hypothesis is shown in Scheme 4. Sequential 1,2-acetylide addition, halogen exchange, and transmetalation to copper affords intermediate **E**. This copper(I) species undergoes a  $\gamma$ -selective S<sub>N</sub>2' addition to propargyl chloride **3** affording the non-isolable allene **C'**, which undergoes the Cannizzaro-type hydride shift proposed in Scheme 1. The resulting allyl anion



development and demonstration (308768). J.R. is grateful to European Thermodynamics Ltd and the Engineering Physical Sciences Research Council (EPSRC) for provision of a studentship. L.B. thanks the EPSRC's National Service for Computational Chemistry (NSCCS) for providing computational facilities, and O.K. acknowledges a Kishore Vaigyanik Protsahan Yojana (KVPJ) Fellowship.

**Keywords:** aldehydes · annulation · carbocycles · medium-ring compounds · synthetic methods

**How to cite:** *Angew. Chem. Int. Ed.* **2015**, *54*, 10648–10651  
*Angew. Chem.* **2015**, *127*, 10794–10797

- [1] a) *Science of Synthesis—Compounds with two carbon-heteroatom bonds, aldehydes*, Vol. 25 (Ed.: R. Brückner), Georg Thieme, Stuttgart, **2007**, p. 901, also Sections 25.4–25.6, pp. 463–710; b) P. Merino in *Science of Synthesis—Monocyclic Arenes, Quasiarenes, and Annulenes*, Vol. 45a (Eds.: J. S. Siegel, Y. Tobe), Georg Thieme, Stuttgart, **2009**, pp. 253–327, also Sections 45.6.3–45.6.4.
- [2] For selected recent examples of strategy A, see: a) A. R. Jagdale, J. H. Park, S. W. Youn, *J. Org. Chem.* **2011**, *76*, 7204–7215; b) D. Das, S. Pratihar, S. Roy, *Org. Lett.* **2012**, *14*, 4870–4873; c) W. Zi, F. D. Toste, *J. Am. Chem. Soc.* **2013**, *135*, 12600–12603; d) P. Sarnpitak, K. Trongchit, Y. Kostenko, S. Sathalalai, M. P. Gleeson, S. Ruchirawat, P. Ploypradith, *J. Org. Chem.* **2013**, *78*, 8281–8296; e) M. Chen, N. Sun, Y. Liu, *Org. Lett.* **2013**, *15*, 5574–5577; f) A. S. K. Hashmi, J. Hofmann, S. Shi, A. Schütz, M. Rudolph, C. Lotzschütz, M. Wietek, M. Bührle, M. Wölfle, F. Rominger, *Chem. Eur. J.* **2013**, *19*, 382–389 (also for reactivity of type B).
- [3] For selected recent examples of strategy B, see: a) H. Zhang, R. C. Larock, *J. Org. Chem.* **2002**, *67*, 7048–7056; b) M. Tian, P. Belmont, *J. Org. Chem.* **2008**, *73*, 4101–4109; c) S. P. Shukla, R. Tiwari, A. K. Verma, *Tetrahedron* **2012**, *68*, 9035–9044; d) X.-C. Wang, M.-J. Zhong, Y.-M. Liang, *Org. Biomol. Chem.* **2012**, *10*, 3636–3641; e) C. H. Oh, J. H. Kim, B. K. Oh, J. R. Park, J. H. Lee, *Chem. Eur. J.* **2013**, *19*, 2592–2596; f) S. Manojveer, R. Balamurugan, *Org. Lett.* **2014**, *16*, 1712–1715.
- [4] For relevant opinions on reaction efficiency, see: a) R. A. Sheldon, *Chem. Soc. Rev.* **2012**, *41*, 1437–1451; b) *Green Chemistry in the Pharmaceutical Industry* (Eds.: P. J. Dunn, A. S. Wells, M. T. Williams), Wiley-VCH, Weinheim, **2010**, pp. 370; c) P. A. Clarke, A. V. Zaytzev, A. C. Whitwood, *Tetrahedron Lett.* **2007**, *48*, 5209–5212.
- [5] For reviews on the synthesis of medium-ring carbocycles, see: a) L. Yet, *Tetrahedron* **1999**, *55*, 9349–9403 (via radicals); b) *Topics in Organometallic Chemistry—Alkene Metathesis in Organic Synthesis*, Vol. 1 (Ed.: A. Fürstner), Springer, New York, USA, **1998**, p. 231 (by metathesis); c) C. Galli, L. Mandolini, *Eur. J. Org. Chem.* **2000**, 3117–3125 (ring-strain effects); d) K. C. Majumdar, B. Chattopadhyay, *Curr. Org. Chem.* **2009**, *13*, 731–757 (by palladium catalysis).
- [6] For selected “modern era” Cannizzaro reactions, see: a) A. E. Russell, S. P. Miller, J. P. Morken, *J. Org. Chem.* **2000**, *65*, 8381–8383; b) M. S. Abaee, R. Sharifi, M. M. Mojtahedi, *Org. Lett.* **2005**, *7*, 5893–5895; c) K. Ishihara, T. Yano, M. Fushimi, *J. Fluorine Chem.* **2008**, *129*, 994–997; d) C. Wuensch, H. Lechner, S. M. Glueck, K. Zangger, M. Hall, K. Faber, *ChemCatChem* **2013**, *5*, 1744–1748; e) P. Wang, W.-J. Tao, X.-L. Sun, S. Liao, Y. Tang, *J. Am. Chem. Soc.* **2013**, *135*, 16849–16852; f) Q. Xu, J. Chen, H. Tian, X. Yuan, S. Li, C. Zhou, J. Liu, *Angew. Chem. Int. Ed.* **2014**, *53*, 225–229; *Angew. Chem.* **2014**, *126*, 229–233.
- [7] A. S. Travis, W. J. Hornix, R. Bud, P. J. T. Morris, *Br. J. Hist. Sci.* **2009**, *25*, 145–167, and references therein.
- [8] M. Murakami, S. Ashida, T. Matsuda, *J. Am. Chem. Soc.* **2006**, *128*, 2166–2167.
- [9] For a selection of representative recent examples (the mode of cyclization, the number of steps required to prepare the cyclization precursor from inexpensive starting materials, and the overall yield for the total synthesis of the eight-ring product (synthesis of the precursors and subsequent cyclization) are given in parentheses); see: a) M. Utsugi, Y. Kamada, M. Nakada, *Tetrahedron Lett.* **2008**, *49*, 4754–4757 (Pd-cat. Heck reaction, 20, 36%); b) Y. Oonishi, A. Hosotani, Y. Sato, *J. Am. Chem. Soc.* **2011**, *133*, 10386–10389 (Rh-cat. [6+2], 7, 21–23%); c) C. S. Penkett, J. A. Woolford, *Org. Lett.* **2012**, *14*, 5704–5707 (photochemistry, 1, <5%); d) Y. Oonishi, A. Hosotani, Y. Sato, *Angew. Chem. Int. Ed.* **2012**, *51*, 11548–11551; *Angew. Chem.* **2012**, *124*, 11716–11719 (Rh-cat. [6+2], 4–7, 8–25%); e) T. Iwai, H. Okochi, H. Ito, M. Sawamura, *Angew. Chem. Int. Ed.* **2013**, *52*, 4239–4242; *Angew. Chem.* **2013**, *125*, 4333–4336 (Au-cat. alkyne cyclization, 5–7 and 3 for the catalyst synthesis, 14–15%); f) C. B. Nielsen, T. Brock-Nannestad, P. Hammershøj, T. K. Reenberg, M. Schau-Magnussen, D. Trpevski, T. Hensel, R. Salcedo, G. V. Baryshnikov, B. F. Minaev, M. Pittelkow, *Chem. Eur. J.* **2013**, *19*, 3898–3904 (Friedel–Crafts, 5, 34–39%); g) T. Nishinaga, T. Ohmae, K. Aita, M. Takase, M. Iyoda, T. Araib, Y. Kunugi, *Chem. Commun.* **2013**, *49*, 5354–5356 (Cu-cat. oxidation, 3, 16%); h) K. Aita, T. Ohmae, M. Takase, K. Nomura, H. Kimura, T. Nishinaga, *Org. Lett.* **2013**, *15*, 3522–3525 (Ru-cat. metathesis, 7, <5%); i) Y. Wang, J. Song, L. Xu, Y. Kann, J. Shi, H. Wang, *J. Org. Chem.* **2014**, *79*, 2255–2262 (Cu-cat. oxidation, 3–5, 12–21%); j) Y. Wang, D. Gao, J. Shi, Y. Kan, J. Song, C. Li, H. Wang, *Tetrahedron* **2014**, *70*, 631–636 (Cu-cat. oxidation, 1, 44%); k) S. Hirai, M. Utsugi, M. Iwamoto, M. Nakada, *Chem. Eur. J.* **2015**, *21*, 355–359 (Pd-cat. alkenylation, >10, <40%); l) C. Hulot, S. Amiri, G. Blond, P. R. Schreiner, J. Suffert, *J. Am. Chem. Soc.* **2009**, *131*, 13387–13398 (Ni-cat. electrocyclization, 8, 28–57%); m) M. Charpenay, A. Boudhar, C. Hulot, G. Blond, J. Suffert, *Tetrahedron* **2013**, *69*, 7568–7591 (Ni-cat. electrocyclization, 6, 39–75%). According to this analysis (excluding **4iaa**), the performance of our own system is: Cannizzaro-triggered, 2–3, 16–62%, average: 42%, 19 examples with >40%. For the preparation of our own starting materials, costs were in the range of €0.3–8 mmol<sup>-1</sup>. Compound **4iaa** is included to demonstrate the need to avoid rather long starting-material syntheses.
- [10] For these derivatives, the proton attached to C4 gives rise to a characteristic low-field resonance at approximately 5.0 ppm, presumably owing to its proximity to the aryl ring current. Compounds **4** are often highly crystalline, and in a number of cases, they could be isolated by a simple recrystallization of the crude reaction material on gram scale. The structures of **4haa** and **4jaa** were confirmed by X-ray crystallography (see the Supporting Information).
- [11] Y. Zhao, D. G. Truhlar, *Theor. Chem. Acc.* **2008**, *120*, 215–241; J. A. Montgomery, M. J. Frisch, J. W. Ochterski, G. A. Petersson, *J. Chem. Phys.* **1999**, *110*, 2822–2827; J. A. Montgomery, Jr., M. J. Frisch, J. W. Ochterski, G. A. Petersson, *J. Chem. Phys.* **2000**, *112*, 6532–6542.
- [12] M. Alajarin, B. Bonillo, M. Marin-Luna, P. Sanchez-Andrada, A. Vidal, *Chem. Eur. J.* **2013**, *19*, 16093–16103.
- [13] Y. Zhang, F. Yang, L. Zheng, Q. Dang, X. Bai, *Org. Lett.* **2014**, *16*, 6041–6043.
- [14] S. L. Schreiber, S. E. Kelly, *Tetrahedron Lett.* **1984**, *25*, 1757–1760. We thank a referee for this suggestion.

Received: June 11, 2015

Published online: July 29, 2015