

A GENERAL METHOD FOR CONSTRUCTING ESSENTIAL UNIFORM ALGEBRAS

J. F. FEINSTEIN AND ALEXANDER J. IZZO

ABSTRACT. A general method for constructing essential uniform algebras with prescribed properties is presented. Using the method, the following examples are constructed: an essential, natural, regular uniform algebra on the closed unit disc; an essential, natural counterexample to the peak point conjecture on each manifold-with-boundary of dimension at least three; and an essential, natural uniform algebra on the unit sphere in \mathbb{C}^3 containing the ball algebra and invariant under the action of the 3-torus. These examples show that a smoothness hypothesis in some results of Anderson and Izzo cannot be omitted.

1. INTRODUCTION

Let X be a compact space, and let $C(X)$ be the algebra of all continuous complex-valued functions on X with the supremum norm $\|f\|_X = \sup\{|f(x)| : x \in X\}$. A *uniform algebra* on X is a closed subalgebra of $C(X)$ that contains the constant functions and separates the points of X . One can often obtain a uniform algebra A with particular properties on a specific space X by finding a uniform algebra B with the desired properties on a subspace E of X and then taking A to consist of those continuous functions on X whose restrictions to E belong to B . However, sometimes it is of interest to know whether there are examples on the space X that do not arise from algebras on a subspace in this trivial manner. This issue is made precise using the notion of essential set and essential uniform algebra, as introduced by Bear in [7] (see also [8, pp. 144–147]). The *essential set* E for a uniform algebra A on a space X is the smallest closed subset of X such that A contains every continuous function on X that vanishes on E . Note that A contains every continuous function whose restriction to E lies in the restriction of A to E . The uniform algebra A is said to be *essential* if $E = X$. The reader may also wish to consult [31], where Tomiyama determined the connection between the essential set of A and the antisymmetric decomposition of the maximal ideal space of A .

2010 *Mathematics Subject Classification.* Primary 46J10.

Key words and phrases. essential uniform algebra, regular algebra, normal algebra.

In this paper, we present a general method for constructing essential uniform algebras. We then use the method to obtain three particular examples of essential uniform algebras with special properties. These examples demonstrate contrasts between uniform algebras generated by smooth functions and those not generated by such functions.

In [11], de Paepe gave a very different method for constructing essential uniform algebras with specified properties. However, de Paepe's algebras and the compact spaces they are defined on do not have the properties we require.

We say that a uniform algebra A on X is *nontrivial* if $A \neq C(X)$, and is *natural* (on X) if X is the maximal ideal space of A (under the usual identification of points of X with evaluation functionals). For the definitions of other terms used in this paper, see the next section.

Our main result is as follows.

Theorem 1.1. *Let A be a nontrivial uniform algebra on a compact space K . Let X be a compact metric space every non-empty open subset of which contains a nowhere dense subspace homeomorphic to K . Then there exists a sequence $\{K_n\}_{n=1}^{\infty}$ of pairwise disjoint, nowhere dense subspaces of X each homeomorphic to K such that $\bigcup_{n=1}^{\infty} K_n$ is dense in X and $\text{diam}(K_n) \rightarrow 0$. If homeomorphisms $h_n : K_n \rightarrow K$ are chosen and we set $A_n = \{f \circ h_n : f \in A\}$, then the collection of functions $\tilde{A} = \{f \in C(X) : f|_{K_n} \in A_n \text{ for all } n\}$ is an essential uniform algebra on X . The equality $\tilde{A}|_{K_n} = A_n$ holds for all n .*

Furthermore, the following relations hold between the properties of A and the properties of \tilde{A} :

- (i) \tilde{A} is natural if and only if A is natural;
- (ii) \tilde{A} is regular on X if and only if A is regular on K ;
- (iii) \tilde{A} is normal if and only if A is normal;
- (iv) every point of X is a peak point for \tilde{A} if and only if every point of K is a peak point for A ;
- (v) \tilde{A} has bounded relative units if and only if A has bounded relative units;
- (vi) \tilde{A} is strongly regular if and only if A is strongly regular;
- (vii) \tilde{A} is a Ditkin algebra if and only if A is a Ditkin algebra.

The uniform algebra \tilde{A} has bounded relative units, with bound 1, at every point of $X \setminus \bigcup_{n=1}^{\infty} K_n$, and (hence) each of these points is a peak point for \tilde{A} , at each of these points \tilde{A} satisfies Ditkin's condition, and at each of these points \tilde{A} is strongly regular.

One could also consider a more general situation in which, rather than starting with one space K and uniform algebra A , one deals with a sequence

of spaces K_n , possibly not all homeomorphic, with dense union in X , and algebras A_n on the K_n . We leave this to the interested reader.

Note that every compact metric space is homeomorphic to a nowhere dense subset of the Hilbert cube [21, Theorem V 4]. This allows us to apply Theorem 1.1 whenever A is a nontrivial uniform algebra on a metrizable compact space K in order to construct an essential uniform algebra on the Hilbert cube sharing many properties with A . However we are mostly interested in examples on spaces whose topological dimension is finite.

Our original motivation came from the following question: Does there exist an essential, natural, regular uniform algebra on the closed unit disc? As an application of our main theorem, we prove that the answer is affirmative. To our knowledge, the first example of an essential, natural, regular uniform algebra on a locally connected compact metric space was given in [16, Example 2.9], using the algebra $R(K)$ for a suitable compact plane set K . However the set K obtained was a ‘classical’ Swiss cheese set, and so (although both connected and locally connected), was infinitely connected.

Theorem 1.2. *There exists an essential, natural, regular uniform algebra on the closed unit disc $\overline{\mathbb{D}} = \{z \in \mathbb{C} : |z| \leq 1\}$.*

However, applying a theorem of Michael Freeman [19, Theorem 4.1], we shall see that no such uniform algebra can be generated by smooth functions. (To say that a collection of functions F generates A means that the collection of all polynomials in the functions in F is a dense subset of A .)

Theorem 1.3. *There does not exist an essential, natural, regular uniform algebra generated by C^1 -smooth functions on a compact two-dimensional C^1 -manifold-with-boundary.*

A theorem of John Anderson and the Alexander Izzo [2, Theorem 4.2] classifies all the natural uniform algebras containing the identity function z and generated by C^1 -smooth functions on the closed disc. The example constructed below in the proof of Theorem 1.2 is easily seen to contain the function z and thus shows that this classification does not continue to hold without the smoothness hypothesis.

Our other two examples are related to the so called peak point conjecture. This conjecture asserted that if a uniform algebra A is natural on X and every point of X is a peak point for A , then $A = C(X)$. (A point $x \in X$ is said to be a *peak point* for A if there exists $f \in A$ with $f(x) = 1$ and $|f(y)| < 1$ for all $y \in X \setminus \{x\}$.) A counterexample to this peak-point

conjecture was produced by Brian Cole in 1968 [9] (or see [8, Appendix], or [29, Chapter 3, Section 19]), and other counterexamples have been given since then with a variety of additional properties, and including examples which are generated by smooth functions on manifolds (see, for example, [6, 12, 13, 14, 22, 23, 30]). Nevertheless, Anderson and Izzo [1, 2, 3] and Anderson, Izzo, and Wermer [4, 5] have established peak point theorems under certain smoothness hypotheses. One of those results, which we state here, asserts that certain uniform algebras can never be essential.

Theorem 1.4 ([2], Theorem 1.2). *Let A be a uniform algebra on a compact C^1 -manifold-with-boundary M . Assume that A is generated by C^1 -smooth functions, that A is natural on M , and that every point of M is a peak point for A . Then the uniform algebra A is not essential. In fact, the essential set for A has empty interior in M .*

Our next example shows that this theorem becomes false if the hypothesis that the algebra is generated by C^1 -smooth functions is dropped.

Theorem 1.5. *On every compact manifold-with-boundary X of dimension greater than or equal to 3 there exists an essential, natural uniform algebra such that every point of X is a peak point. In addition, we may arrange for this uniform algebra to have bounded relative units.*

Another theorem of Anderson and Izzo asserts that the peak point conjecture holds for uniform algebras generated by C^1 -smooth functions on a compact manifold-with-boundary of dimension *two*. It remains an open question whether the smoothness hypothesis can be dropped from that theorem.

For our final example, we modify so as to make essential, an example constructed by the second author [23, Theorem 2.4] in response to a question raised by Ronald Douglas in connection with his work on a conjecture of William Arveson in operator theory. Before stating the result we recall some notions used in the statement.

The ball algebra $A(S)$ on the unit sphere $S \subset \mathbb{C}^n$ consists of the restrictions to the sphere of the functions that are continuous on the closed unit ball $\overline{B}_n \subset \mathbb{C}^n$ and holomorphic on the open unit ball B_n . The action of the n -torus T^n on S is the map $T^n \times S \rightarrow S$ given by $((e^{i\theta_1}, \dots, e^{i\theta_n}), (z_1, \dots, z_n)) \mapsto (e^{i\theta_1} z_1, \dots, e^{i\theta_n} z_n)$. To say that an algebra A on S is invariant under the action of the n -torus means that the function $(z_1, \dots, z_n) \mapsto f(e^{i\theta_1} z_1, \dots, e^{i\theta_n} z_n)$ is in A for each point $(e^{i\theta_1}, \dots, e^{i\theta_n}) \in T^n$ whenever f is in A .

Theorem 1.6. *There exists an essential, natural uniform algebra on the unit sphere S in \mathbb{C}^3 that contains the ball algebra and is invariant under the action of the 3-torus on S .*

The example in [23] is generated by C^∞ -smooth functions and satisfies the conditions above except that it is not essential. Note that since every point of the sphere is a peak point for the ball algebra, Theorem 1.4 shows that it is not possible for the algebra to be simultaneously both essential and generated by smooth functions.

Although the uniform algebras we construct are essential, they do include many non-constant, real-valued functions, and so they are not antisymmetric. This leaves open the question of whether or not there are antisymmetric uniform algebras with these properties. For example, we do not know whether there exists an antisymmetric, natural, regular uniform algebra on the closed unit disc.

Theorem 1.1 is proved in Section 3. In Section 4, we prove Theorems 1.2, 1.5, and 1.6. Theorem 1.3 is proved in Section 5.

2. PRELIMINARIES

Throughout the paper, by a *compact space* we shall mean a non-empty, compact, Hausdorff topological space; by a *compact plane set* we shall mean a non-empty, compact subset of the complex plane.

We assume that the reader has some familiarity with uniform algebras. For further background we refer the reader to [8, 20, 29]. The reader may also consult [10, 25] for the general theory of commutative Banach algebras, and more details concerning regularity conditions and their applications.

Definition 2.1. Let A be a uniform algebra on a compact space X . Recall that we say that A is natural (on X) if the only non-zero multiplicative linear functionals on A are given by evaluations at the points of X , in which case X may be identified with the maximal ideal space of A . We say that A is *regular on X* if, for each closed subset $E \subseteq X$ and each $x \in X \setminus E$, there is a function $f \in A$ with $f(x) = 1$ and $f(E) \subseteq \{0\}$; the algebra A is *normal on X* if, for each pair of disjoint closed subsets E and F of X , there is a function $f \in A$ with $f(E) \subseteq \{0\}$ and $f(F) \subseteq \{1\}$. We say that A is *regular* if it is natural and regular on X ; A is *normal* if it is natural and normal on X .

It is standard that the uniform algebra A on X is normal if and only if it is natural and regular on X [29, Theorem 27.2], and that this holds if and only if A is normal on X [29, Theorem 27.3].

The following elementary result was implicitly assumed throughout [18] (in the more general setting of Banach function algebras). The proof involves an elementary compactness argument: the details are given in [17, Proposition 2.1].

Lemma 2.2. *A uniform algebra A on X is regular on X if and only if for every pair of distinct points x_0 and x_1 in X there is a function f in A such that f is zero on a neighborhood of x_0 and $f(x_1) = 1$.*

The next lemma is proved using a similar elementary compactness argument: we leave the details to the reader.

Lemma 2.3. *A uniform algebra A on X is normal (on X) if and only if for every pair of distinct points x_0 and x_1 in X there is a function f in A such that f is zero on a neighborhood of x_0 and one on a neighborhood of x_1 .*

We now introduce some important ideals, and recall some stronger regularity conditions.

Definition 2.4. Let A be a uniform algebra on a compact space X . Let $x \in X$. We define the ideals M_x, J_x in A as follows:

$$\begin{aligned} M_x &= \{f \in A : f(x) = 0\}; \\ J_x &= \{f \in A : f^{-1}(\{0\}) \text{ is a neighborhood of } x\}. \end{aligned}$$

We say that A is *strongly regular at x* if J_x is dense in M_x ; A satisfies *Ditkin's condition at x* if, for every $f \in M_x$ and every $\varepsilon > 0$, there is $g \in J_x$ with $\|gf - f\|_X < \varepsilon$; for $C \geq 1$, A has *bounded relative units at x with bound C* if, for every compact set $F \subseteq X \setminus \{x\}$, there is $f \in J_x$ with $\|f\|_X \leq C$, such that $f(F) \subseteq \{1\}$. We say that A *has bounded relative units at x* if there exists $C \geq 1$ such that A has bounded relative units at x with bound C . The algebra A is *strongly regular* if it is strongly regular at every point of X ; A is a *Ditkin algebra* if it satisfies Ditkin's condition at each $x \in X$; A is a *strong Ditkin algebra* if it is strongly regular and, for every $x \in X$, the ideal M_x has a bounded approximate identity; A has *bounded relative units* if it has bounded relative units at each $x \in X$.

The reader may find a short survey of the relationships between these regularity conditions for uniform algebras in [15]. In particular we note the following. Let A be a uniform algebra on a compact space X , and let $x \in X$. If A has bounded relative units at x , then A satisfies Ditkin's condition at x , A is strongly regular at x , M_x has a bounded approximate identity and,

for all $C > 1$, A has bounded relative units at x with bound C . Thus if A has bounded relative units, then every $C > 1$ serves as a global bound. (For general Banach function algebras the bound may genuinely depend on the point.) The uniform algebra A has bounded relative units if and only if it is a strong Ditkin algebra and this implies that A is a Ditkin algebra; if A is a Ditkin algebra then A is strongly regular; if A is strongly regular then A is natural and regular on X and hence normal.

Now suppose that X is metrizable. It is then standard that M_x has a bounded approximate identity if and only if x is a peak point for A . (See, for example, [8, p. 101], or [10, Theorem 4.3.5], and note that strong boundary points coincide with peak points when X is metrizable.) Thus, if A has bounded relative units at x , then x must be a peak point for A . In this setting, A has bounded relative units if and only if A is strongly regular and every point of X is a peak point.

Examples of non-trivial, strongly regular uniform algebras on compact metrizable spaces were given in [12], including some examples with bounded relative units.

We shall need some results about metrizability. The first of these is essentially [24, Corollary 26.16]. We include a short proof for the reader's convenience.

Lemma 2.5. *Let X be a compact metrizable space and Y be a quotient space of X . If Y is Hausdorff, then Y is metrizable.*

Proof. It is well known that a compact Hausdorff space Z is metrizable if and only if $C(Z)$ is separable. Let $q : X \rightarrow Y$ be a quotient map of X onto Y . The map $q^* : C(Y) \rightarrow C(X)$ given by $q^*(f) = f \circ q$ embeds $C(Y)$ isometrically into $C(X)$. Thus the hypotheses of the lemma give that $C(Y)$ is isometric to a subspace of a separable metric space and hence is separable. Consequently, Y is metrizable. \square

Let (X, d) be a metric space, let $x \in X$, and let $r > 0$. We denote by $B_r(x)$ the set $\{y \in X : d(y, x) < r\}$, which is called *the open ball of radius r about x* .

Lemma 2.6. *Let (X, d) be a compact metric space, let (K_n) be a sequence of pairwise disjoint, closed subsets of X whose diameters go to zero, and let Y be the quotient space obtained from X by identifying each K_n to a point. Then Y is metrizable.*

Proof. By Lemma 2.5, it suffices to show that Y is Hausdorff. Denote the quotient map of X onto Y by q . Let a and b be distinct points of Y . Then

$q^{-1}(\{a\})$ and $q^{-1}(\{b\})$ are disjoint closed sets in X . Choose disjoint neighborhoods U_a and U_b of $q^{-1}(\{a\})$ and $q^{-1}(\{b\})$, respectively. Define \mathcal{K}_a to be the collection of those K_n that intersect U_a but are not contained in U_a and define \mathcal{K}_b in the same way but with U_a replaced by U_b . Set

$$V_a = U_a \setminus \left(\bigcup_{K \in \mathcal{K}_a} K \right) \quad \text{and} \quad V_b = U_b \setminus \left(\bigcup_{K \in \mathcal{K}_b} K \right).$$

Then V_a and V_b are disjoint sets containing $q^{-1}(\{a\})$ and $q^{-1}(\{b\})$, and each of V_a and V_b is saturated (i.e., $q^{-1}(q(V_a)) = V_a$ and $q^{-1}(q(V_b)) = V_b$). Thus to complete the proof, it suffices to show that V_a and V_b are open. Let $x \in V_a$ be arbitrary. Since U_a is open, there is an $r > 0$ such that the open ball $B_r(a)$ of radius r about a is contained in U_a . Since the diameters of the K_n go to zero, there are at most finitely many K_n , say K_{n_1}, \dots, K_{n_k} , that intersect $B_{r/2}(a)$ and are not contained in U_a . Then $B_{r/2}(a) \setminus (K_{n_1} \cup \dots \cup K_{n_k})$ is an open set about a contained in V_a . Thus V_a is open. The same argument applies to V_b . \square

3. PROOF OF THEOREM 1.1

Proof of Theorem 1.1. Since X is a compact metric space, there is a countable dense subset $\{a_n\}_{n=1}^\infty$ in X . Set $n_1 = 1$ and $r_1 = 1$. By hypothesis there is a nowhere dense subspace K_1 of the ball $B_1(a_1)$ homeomorphic to K . Let n_2 be the smallest positive integer such that $n_1 < n_2$ and a_{n_2} is not in K_1 , choose r_2 such that $0 < r_2 < 1/2$ and the ball $B_{r_2}(a_{n_2})$ is disjoint from K_1 , and then choose a nowhere dense subspace K_2 of $B_{r_2}(a_{n_2})$ homeomorphic to K . Now suppose we have chosen positive integers $n_1 < n_2 < \dots < n_m$, positive numbers r_1, \dots, r_m , and nowhere dense subspaces K_1, \dots, K_m of X . By the Baire category theorem, the union $K_1 \cup \dots \cup K_m$ is nowhere dense in X . Thus infinitely many points of $\{a_n\}_{n=1}^\infty$ lie in the complement of $K_1 \cup \dots \cup K_m$. Let n_{m+1} be the smallest positive integer such that $n_m < n_{m+1}$ and $a_{n_{m+1}}$ is not in $K_1 \cup \dots \cup K_m$, choose r_{m+1} such that $0 < r_{m+1} < 1/(m+1)$ and such that the ball $B_{r_{m+1}}(a_{n_{m+1}})$ is disjoint from $K_1 \cup \dots \cup K_m$, and then choose a nowhere dense subspace K_{m+1} of $B_{r_{m+1}}(a_{n_{m+1}})$ homeomorphic to K . By induction, we then obtain a sequence $\{K_n\}_{n=1}^\infty$ as in the third sentence of the theorem.

Now suppose that homeomorphisms $h_n : K_n \rightarrow K$ are given, and define A_n and \tilde{A} as in the statement of the theorem.

We shall frequently use certain quotient spaces of X in the remainder of the proof. Let Y be the quotient space obtained from X by identifying each K_n to a point, and let q be the quotient map of X onto Y . Also, for

each positive integer m , let Y_m be the quotient space obtained from X by identifying each K_n with $n \neq m$ to a point, and let q_m be the quotient map of X onto Y_m . By Lemma 2.6, Y and all of the Y_m are metrizable.

We now establish the equality $\tilde{A}|_{K_m} = A_m$ for all m . Fix a positive integer m . We may identify K_m with the subset $q_m(K_m)$ of Y_m . Given an arbitrary function g that belongs to A_m , apply the Tietze extension theorem to extend g to a continuous complex-valued function f on Y_m . Then the function $f \circ q_m$ is obviously in \tilde{A} and $(f \circ q_m)|_{K_m} = g$.

We next show that \tilde{A} is a uniform algebra. All conditions are obvious except that \tilde{A} separates points. For that let a and b be two distinct points of X . If there is a positive integer m such that both a and b belong to K_m , then the equality $\tilde{A}|_{K_m} = A_m$ gives at once a function in \tilde{A} that separates a and b . Otherwise we may use the quotient space Y defined above: since in this case $q(a) \neq q(b)$, there is a continuous real-valued function f on Y that separates $q(a)$ and $q(b)$. The function $f \circ q$ is then obviously in \tilde{A} and separates a from b .

To show that \tilde{A} is essential, first note that by the Baire category theorem, the union of the K_n has empty interior in X . Thus letting U be an arbitrary nonempty open subset of X , we know that there is a point a in U lying in none of the K_n . Now choose $r > 0$ such that the ball $B_{2r}(a)$ is contained in U . Note that there must be infinitely many K_n that intersect the ball $B_r(a)$. Because $\text{diam}(K_n) \rightarrow 0$, it must then be that for some positive integer m , the set K_m is contained in U . Because the restriction of each function in \tilde{A} to K_m lies in A_m and A is a nontrivial uniform algebra, it follows that U must intersect the essential set for \tilde{A} . Consequently, \tilde{A} is essential.

At this point it is convenient to make an observation concerning the separation of certain pairs of closed sets, as this will help at a number of points in the remainder of the proof. Let Y and q be as above. We claim that, for each pair of closed subsets E and F of X such that $q(E) \cap q(F) = \emptyset$, there is a real-valued function $f \in \tilde{A}$ with $\|f\|_X = 1$ such that $f \equiv 0$ on some neighborhood of E and $f \equiv 1$ on some neighborhood of F . We shall refer to this below as *our observation about closed sets*. Note here that $q(E) \cap q(F) = \emptyset$ if and only if $E \cap F = \emptyset$ and there is no positive integer m such that E and F both meet K_m .

To prove our claim, let E and F be such a pair of closed subsets of X . Since $q(E)$ and $q(F)$ are disjoint closed subsets of Y , and Y is normal, we may choose, in Y , a pair of disjoint closed neighborhoods N_E, N_F of

$q(E)$, $q(F)$ respectively. By Urysohn's lemma, there is a continuous real-valued function $g \in C(Y)$ with $\|g\|_Y = 1$ and such that $g(N_E) \subseteq \{0\}$ and $g(N_F) \subseteq \{1\}$. Set $f = g \circ q$. Then $f \in \tilde{A}$, $\|f\|_X = 1$, $f \equiv 0$ on the neighborhood $q^{-1}(N_E)$ of E , and $f \equiv 1$ on the neighborhood $q^{-1}(N_F)$ of F . Thus f has the desired properties.

We now establish the properties of the points of $X \setminus \bigcup_{n=1}^{\infty} K_n$ from the end of the statement of the theorem. Let $x \in X \setminus \bigcup_{n=1}^{\infty} K_n$. The fact that \tilde{A} has bounded relative units at x with bound 1 is immediate from our observation about closed sets, taking $E = \{x\}$ and considering an arbitrary compact subset F of $X \setminus \{x\}$. It follows that \tilde{A} satisfies Ditkin's condition at x , \tilde{A} is strongly regular at x , and x is a peak point for \tilde{A} . (The fact that x is a peak point may also be proved directly.)

We now turn to establishing the relations between the properties of \tilde{A} and those of A .

(i) It is well known that given a uniform algebra B on a compact space Σ and a closed subspace E of Σ , the maximal ideal space of the uniform algebra $\overline{B|_E}$ can be identified with a subspace of the maximal ideal space of B . (This subspace is the *B-convex hull* of E .) Furthermore, under this identification, the maximal ideal space of B is the disjoint union of the maximal ideal spaces of its restrictions to its maximal sets of antisymmetry. (See the remark after [20, Theorem II.13.2].) It follows that a uniform algebra is natural if and only if each of its restrictions to a maximal set of antisymmetry is natural.

Now let Y be the quotient space defined above. The continuous real-valued functions separate points on Y , and each of these functions induces a function on X belonging to \tilde{A} . It follows that each set of antisymmetry for \tilde{A} is either contained in some K_n or is a singleton. Furthermore, it then follows from the equality $\tilde{A}|_{K_n} = A_n$ that the sets of antisymmetry for \tilde{A} contained in K_n are precisely the sets of antisymmetry for A_n . Consequently, the collection of maximal sets of antisymmetry for \tilde{A} consists precisely of the maximal sets of antisymmetry for the A_n and the singletons lying in no K_n .

That \tilde{A} is natural if and only if A is natural follows at once from the conclusions of the preceding two paragraphs.

(ii) The proof of this is similar to the proof of (iii) so we leave it as an exercise for the reader.

(iii) Suppose A is normal. Let x_0 and x_1 be distinct points of X . We seek a function in \tilde{A} that is zero on a neighborhood of x_0 and one on a

neighborhood of x_1 . The case when x_0 and x_1 do not lie in a common K_n is easy, as the sets $E = \{x_0\}$ and $F = \{x_1\}$ then satisfy the conditions for our observation about closed sets.

Assume now that for some m , both x_0 and x_1 belong to K_m . Let Y_m and q_m be the quotient space and quotient map defined above. Since by hypothesis A is normal, there is a function h in A_m such that h is zero on a neighborhood of x_0 in K_m and one on a neighborhood of x_1 in K_m . Now regard K_m as a subspace of Y_m , and choose disjoint neighborhoods V_0 and V_1 of x_0 and x_1 , respectively, in Y_m , such that $h = 0$ on $V_0 \cap K_m$ and $h = 1$ on $V_1 \cap K_m$. Then choose neighborhoods W_0 and W_1 of x_0 and x_1 in Y_m with $\overline{W_0} \subset V_0$ and $\overline{W_1} \subset V_1$. Now define g on $K_m \cup \overline{W_0} \cup \overline{W_1}$ by

$$g(x) = \begin{cases} h(x) & \text{for } x \in K_m \\ 0 & \text{for } x \in \overline{W_0} \\ 1 & \text{for } x \in \overline{W_1} \end{cases}$$

Then g is well-defined and continuous. Apply the Tietze extension theorem to extend g to a continuous function h on Y_m . Then the function $f = h \circ q_m$ is the function we seek.

That normality of \tilde{A} implies normality of A is trivial.

(iv) Suppose every point of K is a peak point for A . We are to show that an arbitrary point x in X is a peak point for \tilde{A} . When x lies in no K_n , we already know that x is a peak point for \tilde{A} , as noted earlier.

Consider now the case when x belongs to K_m . Let Y_m and q_m be as above. By hypothesis there is a function g in A_m that peaks at x . By the Tietze extension theorem, g extends to a continuous complex-valued function h on Y_m with supremum norm 1. Let ρ be a nonnegative continuous function on Y_m that is identically equal to 1 on K_m and strictly less than 1 everywhere else on Y_m . Then the function $(\rho h) \circ q_m$ belongs to \tilde{A} and peaks at x .

That every point of K is a peak point for A if every point of X is a peak point for \tilde{A} is trivial.

(v) As noted in Section 2, a uniform algebra on a compact metric space has bounded relative units if and only if it is strongly regular and every point is a peak point. Thus (v) may be deduced from (iv) and (vi). However, we include a direct proof.

As the property of having bounded relative units passes to restriction algebras, it follows that if \tilde{A} has bounded relative units, then so does A . Now suppose that A has bounded relative units. We show that \tilde{A} has bounded relative units. We already know that \tilde{A} has bounded relative units at every point of $X \setminus \bigcup_{n=1}^{\infty} K_n$. Now let $m \in \mathbb{N}$, and let $x \in K_m$. Then A_m has

bounded relative units at x , with bound C , say. We show that \tilde{A} also has bounded relative units at x with bound C .

Let E be a compact subset of $X \setminus \{x\}$, and set $F = E \cap K_m$ (which may be empty). Then there is a $g \in A_m$ with $\|g\|_{K_m} \leq C$ such that g is constantly 0 on some relatively open neighborhood U of x in K_m , and g is constantly 1 on F . Let Y_m and q_m be as above. As before, we may identify K_m with the subset $q_m(K_m)$ of Y_m , and then we have $x \in Y_m \setminus q_m(E)$. Let N be a compact neighborhood of x in $Y_m \setminus q_m(E)$ such that $N \cap K_m \subseteq U$. Then there is an obvious continuous extension h of g to $K_m \cup N \cup q_m(E)$ such that $h \equiv 0$ on N and $h \equiv 1$ on $q_m(E)$, and this extension h still has uniform norm at most C . We may then apply the Tietze extension theorem to extend h to all of Y_m with the same norm: we also call this extension h . Set $f = h \circ q_m$. Then $f \in \tilde{A}$, $\|f\|_X \leq C$, f is constantly 0 on the neighborhood $q^{-1}(N)$ of x , and f is constantly 1 on E . This shows that \tilde{A} has bounded relative units at x with bound C .

(vi) As in (v), only one implication requires a proof, as the other is trivial.

Suppose that A is strongly regular. We already know that \tilde{A} is strongly regular at every point of $X \setminus \bigcup_{n=1}^{\infty} K_n$. Now let $m \in \mathbb{N}$, and let $x \in K_m$. Then A_m is strongly regular at x . We show that \tilde{A} is also strongly regular at x . Let $f \in \tilde{A}$ with $f(x) = 0$, and let $\varepsilon > 0$. Then there exists $g \in A_m$ with g constantly 0 on a relatively open neighborhood U of x in K_m and such that $\|f|_{K_m} - g\|_{K_m} < \varepsilon$. Now let Y_m and q_m be as above. Regarding K_m as a subset of Y_m , we may use the Tietze extension theorem as before to extend g to a function $h \in C(Y_m)$ such that h vanishes on a neighborhood N of x in Y_m . Set $\tilde{g} = h \circ q_m$. Then $\tilde{g} \in J_x$ in \tilde{A} , and $\tilde{g}|_{K_m} = g$, so $\|f - \tilde{g}\|_{K_m} < \varepsilon$.

Set $F = K_m$ and set $E = \{x \in X : |f(x) - \tilde{g}(x)| \geq \varepsilon\}$. Then E and F satisfy the conditions for our observation about closed sets, so there is a function $a \in \tilde{A}$ with $\|a\|_X = 1$ such that $a \equiv 0$ on a neighborhood of E and $a \equiv 1$ on a neighborhood of F . We have $\|a(f - \tilde{g})\|_X < \varepsilon$. Set $b = (1-a)(f - \tilde{g})$. Then $b \equiv 0$ on a neighborhood of K_m and $\|b - (f - \tilde{g})\|_X = \|a(f - \tilde{g})\|_X < \varepsilon$. Thus $b + \tilde{g} \in J_x$, and $\|(b + \tilde{g}) - f\|_X < \varepsilon$. This shows that (in \tilde{A}) J_x is dense in M_x , as required.

(vii) This is similar to (vi), and we leave the details to the reader. We note only that the function g from (vi) should now have the form $g_1 f|_{K_m}$ for some g_1 in A_m which vanishes on a relatively open neighborhood of x in K_m . This function g_1 should then be extended to a function $\tilde{g}_1 \in J_x$ in \tilde{A} , at which point we set $\tilde{g} = f\tilde{g}_1$. This ensures that the function $b + \tilde{g}$ from (vi) is in fJ_x . \square

4. CONSTRUCTION OF SPECIFIC EXAMPLES

Recall that, for a compact plane set K , the uniform algebra $R(K)$ is the uniform closure in $C(K)$ of the set of restrictions to K of rational functions with no poles in K .

Proof of Theorem 1.2. Let K be a compact set in the plane such that $R(K)$ is regular but nontrivial. (An example of such a set was constructed by McKissick [27] and is commonly referred to as McKissick's Swiss cheese. McKissick's example is also presented in [29, pp. 344–355], and a substantial simplification of part of the argument involved is given in [26].) Set $A = R(K)$ and $X = \mathbb{D}$. The set K is nowhere dense in the plane, and the uniform algebra $R(K)$ is natural. Thus the uniform algebra \tilde{A} furnished by Theorem 1.1 is essential, natural, and regular. \square

Proof of Theorem 1.5. Let E be McKissick's Swiss cheese [27] mentioned in the proof of Theorem 1.2 above. Recall that E is nowhere dense in the plane. Let K be the Cartesian product of E and a circle. Richard Basener [6] (or see [29, Example 19.8]) produced a nontrivial, natural uniform algebra A on K with every point of K a peak point. Since a solid torus embeds in \mathbb{R}^3 , the same is true of K , and because E is nowhere dense in the plane, K embeds in \mathbb{R}^3 so as to be nowhere dense. Thus every open subset of the manifold X contains a nowhere dense subspace homeomorphic to K . Now the uniform algebra \tilde{A} furnished by Theorem 1.1 is essential and natural, and every point of X is a peak point for \tilde{A} .

If we wish to ensure, in addition, that \tilde{A} has bounded relative units, then instead of Basener's example, we may use the nontrivial uniform algebra with bounded relative units on a compact metric space L constructed in [12, Theorem 3.6], as long as we ensure that L has topological dimension 1. This can be done by starting the construction from McKissick's example, as McKissick's Swiss cheese has topological dimension 1, and the construction preserves the topological dimension of the maximal ideal space. The well-known embedding theorem for spaces of finite topological dimension due to Menger and Nöbeling ([21, Theorem V 2] or [28, Theorem 50.5]) then gives that L embeds in \mathbb{R}^3 . Consequently every open subset of the manifold X contains a nowhere dense subspace homeomorphic to L , and applying Theorem 1.1 with L in place of K yields the desired example. \square

Proof of Theorem 1.6. This result does not follow directly from Theorem 1.1 because the requirement that the algebra be invariant under the action of the 3-torus precludes defining the algebra in terms of the behavior of

functions on arbitrarily small sets. Instead we combine ideas from the proof of Theorem 1.1 with the proof of [23, Theorem 2.4].

Let K be a compact set in the plane such that $R(K)$ is nontrivial but is such that the only Jensen measures for $R(K)$ are the point masses (for instance McKissick's Swiss cheese). Then K is nowhere dense in the plane. Consider the open quarter disc

$$Q = \{a + bi \in \mathbb{C} : a > 0, b > 0, \text{ and } |a|^2 + |b|^2 < 1\}.$$

The argument in the first paragraph of the proof of Theorem 1.1 can be repeated to obtain a sequence $\{K_n\}_{n=1}^\infty$ of disjoint, nowhere dense subspaces of Q each homeomorphic to K such that $\bigcup_{n=1}^\infty K_n$ is dense in Q and $\text{diam}(K_n) \rightarrow 0$. For each n , define X_n by

$$X_n = \{(z_1, z_2, z_3) \in S : |z_1| + |z_2|i \in K_n\}.$$

Then clearly each X_n is a compact subset of S that is invariant under the action of the 3-torus. Note that the functions z_1, z_2 , and z_3 have no zeros on X_n . Let $\pi_n : X_n \rightarrow K_n$ be defined by $\pi_n(z_1, z_2, z_3) = |z_1| + |z_2|i$. Now let B_n be the uniform algebra on X_n generated by the functions $z_1, z_2, z_3, z_1^{-1}, z_2^{-1}, z_3^{-1}$ and all functions of the form $g \circ \pi_n$ with $g \in R(K_n)$. Then B_n is clearly invariant under the action of the 3-torus. Finally let $A = \{f \in C(S) : f|_{X_n} \in B_n \text{ for all } n\}$. Then A is invariant under the action of the 3-torus. Clearly A contains the ball algebra $A(S)$. Thus the proof will be complete once we show that A is natural and essential.

The quotient space obtained from \overline{Q} by identifying each K_n to a point is metrizable by Lemma 2.6, so the real-valued continuous functions on this quotient space separate points. It follows that each set of antisymmetry for A is either contained in some X_n or is a singleton. It is shown in the proof of [23, Theorem 2.4] that the algebras B_n are natural. That A is also natural now follows from the fact, used above to prove Theorem 1.1(i), that the maximal ideal space of a uniform algebra is the union of the maximal ideal spaces of its restrictions to its maximal sets of antisymmetry.

The proof that A is essential is similar to the proof that \tilde{A} is essential in Theorem 1.1. Let $\pi : S \rightarrow \overline{Q}$ be defined by $\pi(z_1, z_2, z_3) = |z_1| + |z_2|i$. Because A is invariant under the action of the 3-torus, the essential set for A is also invariant under that action. Consequently, to show that A is essential, it suffices to show that for every nonempty open set U of Q , the set $\pi^{-1}(U)$ intersects the essential set for A . Let U be an arbitrary nonempty open set of Q . By the Baire category theorem we know that there is a point p in U lying in none of the K_n . Now choose $r > 0$ such that the ball $B_{2r}(p)$ is contained in

U . Note that there must be infinitely many K_n that intersect the ball $B_r(p)$. Because $\text{diam}(K_n) \rightarrow 0$, it must then be that for some positive integer m , the set K_m is contained in U . Then $X_m = \pi^{-1}(K_m)$ is contained in $\pi^{-1}(U)$. The restriction of each function in A to X_m lies in B_m , and it is shown in the proof of [23, Theorem 2.4] that B_m is a nontrivial uniform algebra. It follows that $\pi^{-1}(U)$ intersects the essential set for A . \square

5. PROOF OF THEOREM 1.3

Proof of Theorem 1.3. We suppose that there is such a uniform algebra A on a compact two-dimensional C^1 -manifold-with-boundary M and derive a contradiction. Let F be a collection of C^1 -smooth functions that generate A . Following Freeman [19], we define the *exceptional set* E of F by

$$E = \{p \in M : df(p) \wedge dg(p) = 0 \text{ for all } f, g \text{ in } F\}.$$

Note that E is a closed set in M . By [19, Theorem 4.1], the essential set for A is contained in the exceptional set. Thus we must have $E = M$. Note that this says that at each point p of M , the vector space of complex differentials spanned by the set $\{df(p) : f \in F\}$ has dimension at most one. Consequently, if there is a function f in F whose real and imaginary parts form a local coordinate system for M on some open set U , then every function in A is holomorphic on U in the complex coordinate system given by f . (See [19, p. 43].) This contradicts the regularity of A , so there is no such function f in F . We conclude that for each point p in M , the vector space of *real* differentials spanned by the set

$$\{du(p) : u \text{ is the real or imaginary part of a function } f \in F\}$$

has dimension at most one. But then the functions in A fail to separate points on M , contrary to the definition of a uniform algebra. \square

Acknowledgements. The research in this paper was begun while the second author was visiting the United Kingdom supported by the London Mathematical Society (LMS Scheme 2, grant ref 21212). Progress was also made on the paper while the second author was a visitor at Indiana University. He thanks the Department of Mathematics for its hospitality.

REFERENCES

- [1] J. T. Anderson and A. J. Izzo, *A peak point theorem for uniform algebras generated by smooth functions on a two-manifold*, Bull. London Math. Soc. 33 (2001), 187–193.

- [2] J. T. Anderson and A. J. Izzo, *Peak point theorems for uniform algebras on manifolds*, Math. Zeit. 261 (2009), 65–71.
- [3] J. T. Anderson and A. J. Izzo, *A peak point theorem for uniform algebras on real-analytic varieties*, Math. Ann. 364 (2016), 657–665.
- [4] J. Anderson, A. Izzo and J. Wermer, *Polynomial approximation on three-dimensional real-analytic submanifolds of \mathbb{C}^n* , Proc. Amer. Math. Soc. 129 (2001) no. 8, 2395–2402.
- [5] J. T. Anderson, A. J. Izzo and J. Wermer, *Polynomial approximation on real-analytic varieties in \mathbb{C}^n* , Proc. Amer. Math. Soc. 132 (2003) no. 5, 1495–1500.
- [6] R. F. Basener, *On rationally convex hulls*, Trans. Amer. Math. Soc. 182 (1973), pp. 353–381.
- [7] Bear, H. S., *Complex function algebras*, Trans. Amer. Math. Soc. 90 (1959), 383–393.
- [8] A. Browder, *Introduction to Function Algebras*, Benjamin, New York (1969).
- [9] B. J. Cole, *One-Point Parts and the Peak Point Conjecture*, Ph.D. dissertation, Yale Univ., 1968.
- [10] H. G. Dales, *Banach Algebras and Automatic Continuity*, London Mathematical Society Monographs, New Series, vol. 24, The Clarendon Press Oxford University Press, New York, 2000.
- [11] P. J. de Paepe, *Essential function algebras with large Šilov boundary*, Math. Scand. 43 (1978), 325–328.
- [12] J. F. Feinstein, *A nontrivial, strongly regular uniform algebra*, J. London Math. Soc. 45 (1992), 288–300.
- [13] J. F. Feinstein, *Trivial Jensen measures without regularity*, Studia Math. 148 (2001), 67–74.
- [14] J. F. Feinstein, *A counterexample to a conjecture of S. E. Morris*, Proc. Amer. Math. Soc. 132 (2004), 2389–2397.
- [15] J. F. Feinstein and M. J. Heath, *Regularity and amenability conditions for uniform algebras*, Function Spaces, 159–169, Contemp. Math., 435, Amer. Math. Soc., Providence, RI, 2007
- [16] J. F. Feinstein and M. J. Heath, *Swiss cheeses, rational approximation and universal plane curves*, Studia Math., 196 (2010), 289–306.
- [17] J. F. Feinstein and R. Mortini, *Partial regularity and t -analytic sets for Banach function algebras*, Math. Z., 271 (2012), 139–155.
- [18] J. F. Feinstein and D. W. B. Somerset, *Non-regularity for Banach function algebras*, Studia Math., 141 (2000), 53–68.

- [19] M. Freeman, *Some conditions for uniform approximation on a manifold*, in: *Function Algebras*, F. Birtel (ed.), Scott, Foresman and Co., Chicago, 1966, pp. 42–60.
- [20] T. W. Gamelin, *Uniform Algebras*, 2nd ed., Chelsea Publishing Company, New York, NY, 1984.
- [21] W. Hurewicz and H. Wallman, *Dimension Theory*, Princeton University Press, Princeton, 1948.
- [22] A. J. Izzo, *Failure of polynomial approximation on polynomially convex subsets of the sphere*, *Bull. London Math. Soc.* 28 (1996), 393–397.
- [23] A. J. Izzo, *Uniform algebras on the sphere invariant under group actions*, *Math. Ann.* 344 (2009), 989–995.
- [24] G. J. O. Jameson, *Topology and normed spaces*, Chapman and Hall, London, 1974.
- [25] E. Kaniuth, *A Course in Commutative Banach Algebras*, Graduate Texts in Mathematics 246, Springer, New York, 2009.
- [26] T. W. Körner, *A cheaper Swiss cheese*, *Studia Math.* 83 (1986), 33–36.
- [27] R. McKissick, *A nontrivial normal sup norm algebra*, *Bull. Amer. Math. Soc.* 69 (1963), 391–395.
- [28] J. R. Munkres, *Topology*, 2nd ed., Prentice-Hall, Upper Saddle River, New Jersey, 2000.
- [29] E. L. Stout, *The Theory of Uniform Algebras*, Bogden & Quigley, New York, 1971.
- [30] E.L. Stout, *Polynomial Convexity*, Birkhäuser Boston Inc., Boston, MA, 2007.
- [31] J. Tomiyama, *Some remarks on antisymmetric decompositions of function algebras*, *Tohoku Math. J.*, 16 (1964), 340–344.

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF NOTTINGHAM, UNIVERSITY PARK, NOTTINGHAM NG7 2RD, UK

E-mail address: Joel.Feinstein@nottingham.ac.uk

DEPARTMENT OF MATHEMATICS AND STATISTICS, BOWLING GREEN STATE UNIVERSITY, BOWLING GREEN, OH 43403, USA

E-mail address: aizzo@math.bgsu.edu