

Laboratory strains of *Escherichia coli* K-12: things are seldom what they seem

Douglas F. Browning^{1,*}, Jon L. Hobman² and Stephen J. W. Busby³

Abstract

Escherichia coli K-12 was originally isolated 100 years ago and since then it has become an invaluable model organism and a cornerstone of molecular biology research. However, despite its pedigree, since its initial isolation *E. coli* K-12 has been repeatedly cultured, passaged and mutagenized, resulting in an organism that carries many genetic changes. To understand more about this important model organism, we have sequenced the genomes of two ancestral K-12 strains, WG1 and EMG2, considered to be the progenitors of many key laboratory strains. Our analysis confirms that these strains still carry genetic elements such as bacteriophage lambda (λ) and the F plasmid, but also indicates that they have undergone extensive laboratory-based evolution. Thus, scrutinizing the genomes of ancestral *E. coli* K-12 strains leads us to examine whether *E. coli* K-12 is a sufficiently robust model organism for 21st century microbiology.

DATA SUMMARY

Figs S1–S14 and File S1 are available with the online version of this article. All genome sequence data and assemblies have been deposited in National Center for Biotechnology Information (NCBI) GenBank under BioProject ID PRJNA848777. The assembled and annotated genomes of WG1 and EMG2 have been deposited with the accession numbers CP099590 and CP099591 (WG1) and CP099588 and CP099589 (EMG2).

INTRODUCTION

Escherichia coli K-12 was originally isolated in 1922 from a convalescent diphtheria patient and, later in the 1940s, adopted by Charles Clifton and Edward Tatum as a model organism [1–3]. Since then, *E. coli* K-12 has become the ‘workhorse’ of molecular biology, becoming arguably the premier model organism in bioscience today. MG1655 was the first *E. coli* K-12 strain to have its genome sequence published, followed by W3110, resulting in an explosion of genomic research and comparative genomics [4, 5]. However, despite its prestige, *E. coli* K-12 was stored on agar plates, stabs or slopes before cryopreservation became established, and has been repeatedly subcultured and mutagenized (Fig. 1), resulting in an organism that carries various genetic changes and has lost the ability to produce many surface-associated structures [3]. For example, *E. coli* K-12 laboratory strains are unable to synthesize O antigen on their lipopolysaccharide and no longer carry the F plasmid nor bacteriophage λ [3, 6–9]. One major strength of using *E. coli* K-12 strains for cloning and heterologous gene expression is that K-12 strains cannot establish in the human gut [10, 11] and, thus, even so-called ‘wild-type’ *E. coli* K-12 strains, like MG1655 and W3110, are very different from commensal or environmental isolates [3, 4, 12, 13].

To understand more about this important model organism, we have sequenced the genomes of two *E. coli* K-12 strains, WG1 and EMG2, the proposed ancestors of key laboratory strains (Fig. 1) [1, 2]. Isolates EMG2 (source: Clowes and Hayes) [2] and WG1 (source: J. Lederberg) are both held in the Coli Genetic Stock Center (CGSC) at Yale University (USA), labelled as the

Received 11 July 2022; Accepted 02 November 2022; Published 06 February 2023

Author affiliations: ¹School of Biosciences, College of Health and Life Sciences, Aston University, Aston Triangle, Birmingham B4 7ET, UK; ²School of Biosciences, University of Nottingham, Sutton Bonington Campus, Sutton Bonington, Loughborough LE12 5RD, UK; ³Institute of Microbiology and Infection, School of Biosciences, University of Birmingham, Birmingham B15 2TT, UK.

*Correspondence: Douglas F. Browning, d.browning@aston.ac.uk

Keywords: bacteriophage lambda; *Escherichia coli* K-12; F plasmid; genomic analysis; laboratory-based evolution.

Abbreviations: CDS, coding sequence; CGSC, Coli Genetic Stock Center; ONT, Oxford Nanopore Technologies; SNV, single nucleotide variant.

Data statement: All supporting data, code and protocols have been provided within the article or through supplementary data files. 14 supplementary figures and a supplementary file are available with the online version of this article.

000922 © 2023 The Authors



This is an open-access article distributed under the terms of the Creative Commons Attribution License. This article was made open access via a Publish and Read agreement between the Microbiology Society and the corresponding author's institution.

Impact Statement

Since its isolation in 1922, *Escherichia coli* K-12 has become arguably the premier model organism for contemporary science. Adoption of *E. coli* K-12 by many microbiologists across the globe has meant that it has a complex pedigree and, although many *E. coli* K-12 strains have been sequenced, little is known about the early versions of K-12, which still carry the F plasmid and bacteriophage λ . To understand more about this important model organism, we have sequenced two ancestral K-12 strains, WG1 and EMG2, which are considered to be the progenitors of many of the laboratory strains used today.

E. coli K-12 wild-type strain [1, 2]. However, it is unclear when each strain was deposited. For EMG2, it is specifically stated in the CGSC data entry that it does not contain any known mutations, picked up during laboratory culture, which is likely based on phenotypic analysis. WG1 is reported in the CGSC entry to be *rfb51* [8] and to potentially have a truncated F plasmid. However, it is still classified as wild-type *E. coli* K-12 in the CGSC database. The standard laboratory strains MG1655 and W3110 are both derived from strain W1485, which itself was derived directly from ‘wild-type *E. coli* K-12’ (EMG2/WG1) (Fig. 1) [1, 12]. After comparing the genomes of these ancestral strains, our analysis confirms that these strains carry genetic elements such as phage λ and the F plasmid, but indicates that they have also undergone mutational alteration during their time in laboratories.

METHODS**Bacterial strains and whole-genome sequencing**

E. coli K-12 strains WG1 and EMG2 were obtained for the CGSC, strain numbers CGSC 5073 and CGSC 4401, respectively [1, 2]. Each strain was sequenced using the enhanced sequencing option from MicrobesNG (<https://microbesng.com/>), which uses a combination of Illumina and Oxford Nanopore Technologies (ONT). Cell cultures were grown in LB medium and the cell pellet was isolated by centrifugation and resuspended in cryo-preserved in a Microbank tube (Pro-Lab Diagnostics). Approximately 2×10^9 cells were used for high molecular mass DNA extraction using a Nanobind CCB Big DNA kit (Circulomics). DNA was quantified with the Qubit dsDNA HS assay in a Qubit 3.0 device (Invitrogen). Long-read genomic DNA libraries were prepared with the SQK-LSK109 kit (ONT) with Native Barcoding EXP-NBD104/114 (ONT), using 400–500 ng high molecular mass DNA. Twelve to twenty-four barcoded samples were pooled in a single sequencing library and loaded on a FLO-MIN106 (R.9.4 or R.9.4.1) flow cell in a GridION device (ONT). Illumina reads were adapter trimmed using Trimmomatic 0.30 with a sliding window quality cut-off of Q15 [14]. Unicycler v0.4.0 was used for genome assembly [15] and Prokka 1.11 to annotate contigs [16]. Sequence data have been deposited at GenBank/ENA/DDBJ with the accession numbers CP099590 and CP099591 for WG1, and CP099588 and CP099589 for EMG2.

Bioinformatic analysis of genome sequences

For single nucleotide variant (SNV) calling, reads from EMG2 were aligned to the WG1 reference genome using BWA-MEM and processed using SAMtools 1.2. Variants were called using VarScan with two thresholds, sensitive and specific, where the variant allele frequency is greater than 90 and 10%, respectively. The effects of variants were predicted and annotated using SnpEff. Draft genomes were visualized using Artemis [17], and comparisons between *E. coli* K-12 genomes were made using the Basic Local Alignment Search Tool (BLAST) from the National Center for Biotechnology Information (NCBI) (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>), the Artemis Comparison Tool (ACT) [18] and the Proksee server (<https://proksee.ca/>) [19]. Genome representations were drawn using the Proksee server [19] and ACT [18]. Plasmid replicons were detected in draft genomes with PlasmidFinder 2.1 [5], using software at the Center for Genomic Epidemiology (CGE) (<http://www.genomicepidemiology.org/>). Insertion sequences were located using ISfinder (<https://isfinder.biotoul.fr/blast.php>) [20].

RESULTS**Comparison of the WG1 and EMG2 genomes**

Whole-genome sequencing of WG1 and EMG2 resulted in draft genome sequences, each comprising two contigs; the larger contig, contig 1, is the chromosomal sequence, and the smaller, contig 2, is the F plasmid (Figs 2, 3 and S1, Tables 1 and 2). Since both strains carry bacteriophage λ and the F plasmid, their genomes are slightly bigger than other sequenced *E. coli* K-12 strains, such as MG1655 and W3110 (Table 1) [4, 12]. Comparison of the genomes of both WG1 and EMG2 with those of MG1655 and W3110 indicated that, unlike W3110, no major chromosomal rearrangements had occurred in these strains (Fig. S2) [19, 21]. However, we identified a number of obvious regions of difference (Figs 2, S1 and S3). For example, both EMG2 and W3110 have lost the cryptic prophage CPZ-55, and EMG2 has lost the *gatYZABDR* locus, which is involved in galactitol metabolism [22] (Figs 2 and S4). Interestingly, the *gatYZABDR* genes appear to have been a hotspot for insertion sequence element-mediated disruption in both MG1655 and W3110, which affects expression of this region (Fig. S5) [22]. Similarly, the region upstream of

Table 1. Comparison of the genomes of different *E. coli* K-12 laboratory strains

| | WG1 | EMG2 | MG1655 | W3110 | NCM3722 | LS5218 |
|--------------------|----------------------|----------------------|-------------|-------------|-----------------------|----------------|
| Accession no. | CP099590 CP099591 | CP099588 CP099589 | NC_000913.3 | NC_007779.1 | CP011495.1 CP011496.1 | MVJG00000000.1 |
| Genome size* | 4735495 bp | 4774480 bp | 4641652 bp | 4646332 bp | 4745591 bp | 4699198 bp |
| Plasmid | F plasmid | F plasmid | None | None | F plasmid | F plasmid |
| Total no. of CDSs† | 4431 | 4457 | 4285 | 4213 | 4539 | 4368 |
| G+C content | 50.75mol% | 50.73mol% | 50.79mol% | 50.8mol% | 50.76mol% | 50.72mol% |

*Genome size includes the F plasmid for WG1, EMG2, NCM3722 and LS5218.

†Number of coding sequences (CDSs) is as predicted by each genome annotation.

flhDC locus, which controls flagella production, also seems to have been targeted by different transposable elements (Figs 2 and S6) [23, 24]. Note that strains that have been stored in agar stabs for many years accumulate deleterious mutations due to wholesale transposition of insertion sequences [25–27]. As insertion of different elements into this region influences motility in other *E. coli* K-12 strains, it is likely that the sequence heterogeneity found in this region produces a spectrum of effects [23, 24]. For WG1, we detected the loss of cryptic prophage CP4-6 and a large deletion of the lipopolysaccharide O-antigen biosynthetic cluster, previously termed *rfb-51* (Figs S1, S3 and S4) [8]. Note that EMG2, MG1655 and W3110 carry the alternative *rfb-50* mutation (an IS5 disruption of the rhamnose transferase encoding gene *wbbL*), which appears to be common to most *E. coli* K-12 strains [28]. Thus, neither WG1 nor EMG2 produce O-antigen (Fig. S4) [8, 9]. Loss of O-antigen production seems to be an adaptation to laboratory life, with both the first *E. coli* strain NCTC 86 (isolated in 1885) and commonly used B strains [e.g. BL21(DE3)] all being rough in nature [13, 29, 30]. As well as these differences, WG1 also carries a block of additional genes, encoding an LPS export ABC transporter permease (*lptG*), an acyl-carrier protein (*acpP*) and a NAD-dependent epimerase/dehydratase (*oleD*), which are flanked by IS5 elements (Figs 2 and S7).

The F plasmid

Comparison with MG1655 confirmed that WG1 and EMG2 both carry the F plasmid; however, the two versions of F differ markedly in size, with that from EMG2 (99158 bp) similar in size to the previously sequenced F plasmid (AP001918.199159 bp), whilst F from WG1 is considerable smaller (67408 bp) (Table 2, Figs 3 and S8). This can be attributed to the loss of a large section of F in WG1, carrying the AIDA-I like autotransporter adhesin genes *ycbB* and *ychA*, the *ompP* omptin and the IncFIB replicon (Table 2, Figs 3 and S9) [7, 31]. Surprisingly, F from WG1 carries additional DNA, which includes an IncFII RepA protein (Figs S8 and S9). As in the previously sequenced F plasmid (AP001918.1), EMG2 F carries an IS3 insertion in the *finO* gene, which leads to constitutive F transfer [7, 32, 33]. However, this insertion sequence is absent from the WG1 F (Figs 3, S8 and S9), suggesting that conjugative transfer is regulated in this plasmid and that the insertion of IS3 must have occurred in the immediate ancestor of EMG2. Thus, it is clear that F plasmids from both EMG2 and WG1 have undergone significant laboratory-based evolution, leading to two very different plasmids.

Bacteriophage λ

Comparison with MG1655 indicated that, as expected, both WG1 and EMG2 carry the bacteriophage λ prophage integrated between the *bioA* and *ybhC* genes (Figs 2, S1 and S10). However, comparison with the previous sequenced λ genome (NC_001416) identified some differences in λ from WG1 and EMG2, in particular with the genes encoding tail fibres J, Stf and Tfa (Fig. S11). Of

Table 2. Comparison of the F plasmid from different *E. coli* K-12 laboratory strains

| | WG1 | EMG2 | F plasmid | NCM3722 | LS5218 |
|--------------------|------------------------|-----------------------------|-----------------------------|------------------------|------------------------|
| Accession no. | CP099591 | CP099589 | AP001918.1 | CP011496.1 | CM007715.1 |
| F plasmid size | 67408 bp | 99158 bp | 99159 bp | 67545 bp | 67502 bp |
| Total no. of CDSs* | 73 | 98 | 105 | 79 | 83 |
| G+C content | 51.66mol% | 48.17mol% | 48.17mol% | 51.67mol% | 51.67mol% |
| Plasmid replicons† | IncFIA, IncFIC(FII) | IncFIA, IncFIB, IncFIC(FII) | IncFIA, IncFIB, IncFIC(FII) | IncFIA, IncFIC(FII) | IncFIA, IncFIC(FII) |

*Number of coding sequences (CDSs) is as specified by genome annotation.

†Plasmid replicons were detected using PlasmidFinder 2.1 using software at the Center for Genomic Epidemiology (CGE) [5].




Fig. 1. The pedigree of *E. coli* K-12 strains. The figure details the pathway of *E. coli* K-12 evolution from its isolation in 1922 to the generation of MG1655 and W3100 strains [1, 3, 4, 12]. Blood agar indicates selection on blood agar plates; UV indicates irradiation with ultraviolet light; EMB-gal indicates selection for utilization of galactose on eosin methylene blue indicator plates. Dotted lines represent uncertain evolutionary lineage events.

note is *stf* (side tail fibre), which in λ (NC_001416) carries a frameshift disrupting the gene into two ORFs (*orf-401* and *orf-314*) [34, 35]. Bacteriophage λ carrying this lesion (λ PaPa) forms larger λ plaques [6, 35]. Thus, as *stf* remains intact in WG1 and EMG2, it is likely that both strains would produce a small plaque phenotype [6, 35].

Similarities and differences between WG1 and EMG2

SNV calling showed that *E. coli* K-12 strains WG1 and EMG2 also differ in a number of key genes involved in important cellular functions (File S1). For example, in EMG2, the gene encoding the major sigma factor σ^{70} (*rpoD*) carries a substitution, which results in Tyr at position 571 (Fig. S12a). This is also found in MG1655 and W3110, whilst most *E. coli* strains carry His at this position. Substitutions at σ^{70} residue 571 have been shown to affect transcription at the *lac*, *araBAD*, *merT*, *merR* and the P22 phage *ant* promoters, as well as interfering with σ^{70} binding to core RNA polymerase and its ability to compete with alternative sigma factors [36–40]. Conversely, in WG1, the gene encoding the α subunit of RNA polymerase carries a mutation that results in a Gly to Arg substitution at position 311 (Fig. S12b). This alteration affects expression from both the *merT* and *merR* promoters and the anaerobically activated *pepT* promoter in *Salmonella enterica* serovar Typhimurium [38, 41]. (Note that α in *E. coli* and *S. enterica* serovar Typhimurium are identical.) As for many K-12 strains, both WG1 and EMG2 carry a truncation in *rpoS*, which encodes the stress and stationary phase sigma factor σ^S (Fig. S12c). (Note that the *rpoS* gene in MG1655 is the pseudo revertant *rpoS* 33Q allele [4, 12].) Additionally, *E. coli* K-12 strains also carry changes in genes that influence translation. Like MG1655 and W3110, EMG2 carries a mutation in the gene encoding release factor RF2 (*prfB*) (Thr at position 246) and a mutation in *rpsG* (30S ribosomal protein S7), which results in C-terminal extension of the S7 protein product (Fig. S12d, e). Both substitutions have been shown to affect translation, with the mutation in RF2 resulting in poor termination at UGA stop codons and the trans-translational tagging of S7 with the SsrA peptide [42–46]. Thus, it is clear that, for both EMG2 and WG1,




Fig. 3. Comparison of the F plasmids from different *E. coli* K-12 strains. The figure shows the comparison of the F plasmid (AP001918.1) with that from EMG2 and WG1 using Proksee [19]. The outer two rings display the genes and features of the F plasmid, with selected genes labelled. The green and brown rings illustrate the BLAST results when the F plasmid sequences from EMG2 and WG1, respectively, are compared to the original F plasmid sequence.

rfb-50, *rpoA*, *mdtF* and *nfi*), suggesting that it is distinct from these strains (Fig. 1). It is worth noting that WG1 is similar to *E. coli* strains NCM3722 [54] and LS5218 [55]. Strain NCM3722 (CGSC 12355) was first detailed by Sydney Kustu [48] and LS5218 is an industrial strain used for the production of fatty acid derived products [55]. Both strains carry bacteriophage λ , a smaller version of the F plasmid (Tables 1 and 2, Figs S13 and S14) and contain many of the mutations carried by WG1 [54, 55].

In addition to lineage-specific mutations, it is clear that WG1 and EMG2 have undergone their own laboratory-based evolution events, such as loss of cryptic prophages and gene disruption. The suggestion is that the selection of particular traits by microbiologists has driven laboratory-based evolution. Hence, insertion sequence inactivation of *finO* in F made plasmid transfer easier to study, larger plaques enabled the intricacies of λ lysogeny to be examined and lack of O-antigen enhances plasmid transformation [6, 7, 30, 35]. Thus, our interpretation of *E. coli* biology has been inadvertently biased. Moreover, many other laboratory strains, handed down for generations, are as yet unsequenced, so it is unclear what other changes lie within those strains.

Heterogeneity in bacterial laboratory strains and plasmids has been observed many times and we are at a stage when even the same *E. coli* K-12 stock strains can produce different outcomes, calling reproducibility into question [27, 48, 56–59]. It is clear that there are significant major differences between K-12 and other commensal *E. coli* strains, and these differences became fixed in the ancestors of the very widely used MG1655 and W3110 strains. Given the different mutations seen in WG1 compared to EMG2, it seems likely that identical or similar mutations will be present in other K-12 lineages. However, due to the extensive genetic systems that have been developed, demonstration of safe use, and lack of ability to colonize humans, *E. coli* K-12 strains will justifiably continue to be widely used [10, 11, 52]. We think it is important that there is an awareness of the mutations present in K-12 strains, and the effects of these mutations on the physiology and metabolism of these strains. An understanding of the conditions that might select for mutants in laboratories, and the use of cost-effective and accurate sequencing of laboratory stocks should help to prevent further undetected mutations arising in K-12 strains, which could compromise our understanding of fundamental biological processes. Thus, it is hoped that the next century will continue to provide more insight into the complex

biology and evolution of this versatile organism. Indeed, appreciation of various K-12 strains, as well differences between various bacterial families, is sure to enhance our understanding of life.

Funding information

This work was generously supported by Biotechnology and Biological Sciences Research Council (BBSRC) research grants BB/R017689/1 and BB/W00285X/1 to D.F.B. and S.J.W.B., and BBSRC grant BB/E01044X/1 to J.L.H.

Acknowledgements

We thank MicrobesNG for genome sequencing, in particular Andrew Holmes and Emily Jane Richardson for bioinformatics and SNV calling.

Author contributions

D.F.B., J.L.H. and S.J.W.B. conceived the study, selected samples, carried out bioinformatic analyses and wrote the manuscript. All authors read and approved the final version of the manuscript.

Conflicts of interest

The authors declare that there are no conflicts of interest.

Ethical statement

No ethical clearance was required for this study.

References

- Bachmann BJ. Pedigrees of some mutant strains of *Escherichia coli* K-12. *Bacteriol Rev* 1972;36:525–557.
- Clowes RC, Hayes W. *Experiments in Microbial Genetics*. Oxford, Edinburgh: Blackwell Scientific Publications; 1968.
- Hobman JL, Penn CW, Pallen MJ. Laboratory strains of *Escherichia coli*: model citizens or deceitful delinquents growing old disgracefully? *Mol Microbiol* 2007;64:881–885.
- Blattner FR, Plunkett G, Bloch CA, Perna NT, Burland V, *et al.* The complete genome sequence of *Escherichia coli* K-12. *Science* 1997;277:1453–1462.
- Carattoli A, Zankari E, García-Fernández A, Voldby Larsen M, Lund O, *et al.* In silico detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. *Antimicrob Agents Chemother* 2014;58:3895–3903.
- Casjens SR, Hendrix RW. Bacteriophage lambda: early pioneer and still relevant. *Virology* 2015;479–480:310–330.
- Koraimann G. Spread and persistence of virulence and antibiotic resistance genes: a ride on the F plasmid conjugation module. *EcoSal Plus* 2018;8:ESP-0003–2018.
- Liu D, Reeves PR. *Escherichia coli* K12 regains its O antigen. *Microbiology* 1994;140:49–57.
- Stevenson G, Neal B, Liu D, Hobbs M, Packer NH, *et al.* Structure of the O antigen of *Escherichia coli* K-12 and the sequence of its rfb gene cluster. *J Bacteriol* 1994;176:4144–4156.
- Smith HW. Survival of orally administered *E. coli* K 12 in alimentary tract of man. *Nature* 1975;255:500–502.
- Anderson ES. Viability of, and transfer of a plasmid from, *E. coli* K12 in human intestine. *Nature* 1975;255:502–504.
- Hayashi K, Morooka N, Yamamoto Y, Fujita K, Isono K, *et al.* Highly accurate genome sequences of *Escherichia coli* K-12 strains MG1655 and W3110. *Mol Syst Biol* 2006;2:2006.0007.
- Dunne KA, Chaudhuri RR, Rossiter AE, Beriottio I, Browning DF, *et al.* Sequencing a piece of history: complete genome sequence of the original *Escherichia coli* strain. *Microb Genom* 2017;3:000106.
- Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for illumina sequence data. *Bioinformatics* 2014;30:2114–2120.
- Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 2017;13:e1005595.
- Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 2014;30:2068–2069.
- Rutherford K, Parkhill J, Crook J, Horsnell T, Rice P, *et al.* Artemis: sequence visualization and annotation. *Bioinformatics* 2000;16:944–945.
- Carver TJ, Rutherford KM, Berriman M, Rajandream M-A, Barrell BG, *et al.* ACT: the Artemis Comparison Tool. *Bioinformatics* 2005;21:3422–3423.
- Grant JR, Stothard P. The CGView server: a comparative genomics tool for circular genomes. *Nucleic Acids Res* 2008;36:W181–W184.
- Zhang Z, Schwartz S, Wagner L, Miller W. A greedy algorithm for aligning DNA sequences. *J Comput Biol* 2000;7:203–214.
- Hill CW, Harnish BW. Inversions between ribosomal RNA genes of *Escherichia coli*. *Proc Natl Acad Sci USA* 1981;78:7069–7072.
- Nobelmann B, Lengeler JW. Molecular analysis of the gat genes from *Escherichia coli* and of their roles in galactitol transport and metabolism. *J Bacteriol* 1996;178:6790–6795.
- Zhang Z, Kukita C, Humayun MZ, Saier MH. Environment-directed activation of the *Escherichia coli* flhDC operon by transposons. *Microbiology* 2017;163:554–569.
- Barker CS, Prüss BM, Matsumura P. Increased motility of *Escherichia coli* by insertion sequence element integration into the regulatory region of the flhD operon. *J Bacteriol* 2004;186:7529–7537.
- Naas T, Blot M, Fitch WM, Arber W. Insertion sequence-related genetic variation in resting *Escherichia coli* K-12. *Genetics* 1994;136:721–730.
- Naas T, Blot M, Fitch WM, Arber W. Dynamics of IS-related genetic rearrangements in resting *Escherichia coli* K-12. *Mol Biol Evol* 1995;12:198–207.
- Freddolino PL, Amini S, Tavazoie S. Newly identified genetic variations in common *Escherichia coli* MG1655 stock cultures. *J Bacteriol* 2012;194:303–306.
- Kuhnert P, Nicolet J, Frey J. Rapid and accurate identification of *Escherichia coli* K-12 strains. *Appl Environ Microbiol* 1995;61:4135–4139.
- Jeong H, Barbe V, Lee CH, Vallenet D, Yu DS, *et al.* Genome sequences of *Escherichia coli* B strains REL606 and BL21(DE3). *J Mol Biol* 2009;394:644–652.
- Browning DF, Wells TJ, França FLS, Morris FC, Sevastyanovich YR, *et al.* Laboratory adapted *Escherichia coli* K-12 becomes a pathogen of *Caenorhabditis elegans* upon restoration of O antigen biosynthesis. *Mol Microbiol* 2013;87:939–950.
- Yen MR, Peabody CR, Partovi SM, Zhai Y, Tseng YH, *et al.* Protein-translocating outer membrane porins of Gram-negative bacteria. *Biochim Biophys Acta* 2002;1562:6–31.
- Cheah KC, Skurray R. The F plasmid carries an IS3 insertion within finO. *J Gen Microbiol* 1986;132:3269–3275.
- Yoshioka Y, Ohtsubo H, Ohtsubo E. Repressor gene finO in plasmids R100 and F: constitutive transfer of plasmid F is caused by insertion of IS3 into F finO. *J Bacteriol* 1987;169:619–623.

34. Sanger F, Coulson AR, Hong GF, Hill DF, Petersen GB. Nucleotide sequence of bacteriophage lambda DNA. *J Mol Biol* 1982;162:729–773.
35. Hendrix RW, Duda RL. Bacteriophage lambda PaPa: not the mother of all lambda phages. *Science* 1992;258:1145–1148.
36. Siegele DA, Hu JC, Gross CA. Mutations in rpoD, the gene encoding the sigma 70 subunit of *Escherichia coli* RNA polymerase, that increase expression of the lac operon in the absence of CAP-cAMP. *J Mol Biol* 1988;203:29–37.
37. Siegele DA, Hu JC, Walter WA, Gross CA. Altered promoter recognition by mutant forms of the sigma 70 subunit of *Escherichia coli* RNA polymerase. *J Mol Biol* 1989;206:591–603.
38. Caslake LF, Ashraf SI, Summers AO. Mutations in the alpha and sigma-70 subunits of RNA polymerase affect expression of the mer operon. *J Bacteriol* 1997;179:1787–1795.
39. Jishage M, Kvint K, Shingler V, Nyström T. Regulation of sigma factor competition by the alarmone ppGpp. *Genes Dev* 2002;16:1260–1270.
40. Cicero MP, Sharp MM, Gross CA, Kreuzer KN. Substitutions in bacteriophage T4 AsiA and *Escherichia coli* sigma(70) that suppress T4 motA activation mutations. *J Bacteriol* 2001;183:2289–2297.
41. Lombardo MJ, Bagga D, Miller CG. Mutations in rpoA affect expression of anaerobically regulated genes in *Salmonella typhimurium*. *J Bacteriol* 1991;173:7511–7518.
42. Schaub RE, Hayes CS. Deletion of the RluD pseudouridine synthase promotes SsrA peptide tagging of ribosomal protein S7. *Mol Microbiol* 2011;79:331–341.
43. Pavlov MY, Freistroffer DV, Dincbas V, MacDougall J, Buckingham RH, *et al.* A direct estimation of the context effect on the efficiency of termination. *J Mol Biol* 1998;284:579–590.
44. Collier J, Binet E, Bouloc P. Competition between SsrA tagging and translational termination at weak stop codons in *Escherichia coli*. *Mol Microbiol* 2002;45:745–754.
45. Dincbas-Renqvist V, Engström A, Mora L, Heurgué-Hamard V, Buckingham R, *et al.* A post-translational modification in the GGQ motif of RF2 from *Escherichia coli* stimulates termination of translation. *EMBO J* 2000;19:6900–6907.
46. Uno M, Ito K, Nakamura Y. Functional specificity of amino acid at position 246 in the tRNA mimicry domain of bacterial release factor 2. *Biochimie* 1996;78:935–943.
47. Jensen KF. The *Escherichia coli* K-12 “wild types” W3110 and MG1655 have an rph frameshift mutation that leads to pyrimidine starvation due to low pyrE expression levels. *J Bacteriol* 1993;175:3401–3407.
48. Soupene E, van Heeswijk WC, Plumbridge J, Stewart V, Bertenthal D, *et al.* Physiological studies of *Escherichia coli* strain MG1655: growth defects and apparent cross-regulation of gene expression. *J Bacteriol* 2003;185:5611–5626.
49. Lawther RP, Calhoun DH, Gray J, Adams CW, Hauser CA, *et al.* DNA sequence fine-structure analysis of ilvG (IlvG+) mutations of *Escherichia coli* K-12. *J Bacteriol* 1982;149:294–298.
50. Schuster S, Vavra M, Greim L, Kern WV. Exploring the contribution of the AcrB homolog MdtF to drug resistance and dye efflux in a multidrug resistant *E. coli* isolate. *Antibiotics* 2021;10:503.
51. Weiss B. Endonuclease V of *Escherichia coli* prevents mutations from nitrosative deamination during nitrate/nitrite respiration. *Mutat Res* 2001;461:301–309.
52. Zimmer C. *Microcosm: E. coli and the New Science of Life*. New York: Pantheon Books; 2008.
53. Neidhardt FC, Curtiss R. *Escherichia coli and Salmonella: Cellular and Molecular Biology*. Washington, DC: American Society for Microbiology; 1996.
54. Brown SD, Jun S. Complete genome sequence of *Escherichia coli* NCM3722. *Genome Announc* 2015;3:e00879-15.
55. Rand JM, Gordon GC, Mehrer CR, Pflieger BF. Genome sequence and analysis of *Escherichia coli* production strain LS5218. *Metab Eng Commun* 2017;5:78–83.
56. Nahku R, Peebo K, Valgepea K, Barrick JE, Adamberg K, *et al.* Stock culture heterogeneity rather than new mutational variation complicates short-term cell physiology studies of *Escherichia coli* K-12 MG1655 in continuous culture. *Microbiology* 2011;157:2604–2610.
57. Spira B, de Almeida Toledo R, Maharjan RP, Ferenci T. The uncertain consequences of transferring bacterial strains between laboratories – rpoS instability as an example. *BMC Microbiol* 2011;11:248.
58. Hooton SPT, Pritchard ACW, Asiani K, Gray-Hammerton CJ, Stekel DJ, *et al.* Laboratory stock variants of the archetype silver resistance plasmid pMG101 demonstrate plasmid fusion, loss of transmissibility, and transposition of Tn7/*pco/sil* into the host chromosome. *Front Microbiol* 2021;12:723322.
59. Jishage M, Ishihama A. Variation in RNA polymerase sigma subunit composition within different stocks of *Escherichia coli* W3110. *J Bacteriol* 1997;179:959–963.

Five reasons to publish your next article with a Microbiology Society journal

1. When you submit to our journals, you are supporting Society activities for your community.
2. Experience a fair, transparent process and critical, constructive review.
3. If you are at a Publish and Read institution, you'll enjoy the benefits of Open Access across our journal portfolio.
4. Author feedback says our Editors are 'thorough and fair' and 'patient and caring'.
5. Increase your reach and impact and share your research more widely.

Find out more and submit your article at microbiologyresearch.org.