

1 **Mitochondrial DNA copy number associates with insulin sensitivity and aerobic capacity,**
2 **and differs between sedentary, overweight middle-aged males with and without type 2**
3 **diabetes**

4
5 Dumitru Constantin-Teodosiu¹, Despina Constantin¹, Maurice M. Pelters², Lex B. Verdijk², Luc
6 van Loon², and Paul. L. Greenhaff¹

7
8 ¹MRC/ARUK Centre for Musculoskeletal Ageing Research, National Institute for Health
9 Research Nottingham Biomedical Research Centre, School of Life Sciences, Nottingham
10 University Medical School, Nottingham, NG7 2UH, United Kingdom, ²NUTRIM School for
11 Nutrition and Translational Research in Metabolism, Department of Human Biology and
12 Movement Sciences, Maastricht University Medical Centre, Maastricht, Netherlands.

13
14 Short running title: insulin resistance and mitochondria

15
16 Corresponding author: Dumitru Constantin-Teodosiu, PhD

17 Email: tim.constantin@nottingham.ac.uk

18 Tel no +44-115-8230111

19 Abstract word counts:

20 Main body text word counts

21 Reference number:

22 Table number: 3

23 Figure number: 2

24

Abstract

Background/objectives: Increased risk of type 2 diabetes mellitus (T2DM) is linked to impaired muscle mitochondrial function and reduced mitochondrial DNA copy number (mtDNA^{num}). However, studies have failed to control for habitual physical activity levels, which directly influences both mtDNA copy number and insulin sensitivity. We, therefore, examined whether physical conditioning status (maximal oxygen uptake, $\dot{V}O_{2max}$) was associated with skeletal muscle mitochondrial volume and mtDNA^{num}, and was predictive of T2DM in overweight, middle-aged men.

Methods: Whole-body physiological (ISI-insulin sensitivity index, HOMA-IR, $\dot{V}O_{2max}$) and muscle biochemical/molecular (vastus lateralis; mtDNA^{num}, mitochondrial and glycolytic enzymes activity, lipid content and markers of lipid peroxidation) measurements were performed in 3 groups of overweight, middle-aged male volunteers ($n=10$ per group): sedentary T2DM (ST2DM); sedentary control (SC) and non-sedentary control (NSC), who differed in aerobic capacity (ST2DM<SC<NSC).

Results: mtDNA^{num} was greater in NSC vs SC and ST2DM ($P<0.001$; $P<0.001$), and less in ST2DM vs SC ($P<0.01$). Across all groups, mtDNA^{num} positively correlated with ISI ($P<0.001$; $r=0.688$) and $\dot{V}O_{2max}$ (normalised to free fat mass; $r=0.684$, $P<0.001$), and negatively correlated to HOMA-IR ($r=-0.544$, $P<0.01$). The activity of mitochondrial enzymes (GluDH, CS and β -HAD) was greater in NSC than ST2DM ($P<0.01$, $P<0.001$ and $P<0.05$) and SC ($P<0.05$, $P<0.01$ and $P<0.05$), but similar between ST2DM and SC. Intramuscular free fatty acids, triglycerides and malondialdehyde contents were similar between ST2DM and SC.

Conclusions: Body composition and indices of muscle mitochondrial volume/function were similar between SC and ST2DM. However, mtDNA^{num} differed and was positively associated with ISI, HOMA-IR and $\dot{V}O_{2max}$ across all groups. Collectively, the findings support the

49 contention that habitual physical activity is a key component of T2DM development, possibly
50 by influencing mtDNA^{num}.

51

52 **Introduction**

53 Excess caloric intake and lack of physical activity are primary causes of increasing obesity
54 prevalence worldwide. Being overweight or obese is commonly associated with elevated
55 circulating free fatty acid concentrations and increased risk of metabolic inflexibility (defined
56 as the inability of skeletal muscle to switch from fat to carbohydrate oxidation in response to
57 increased circulating glucose and insulin concentrations), a central feature of insulin resistance
58 (IR) and type 2 diabetes (T2DM). However, not all obese people develop T2DM, and not all
59 T2DM patients are obese. Alternatively, increased risk of T2DM development has been linked
60 to reduced muscle mitochondrial function (defects in intrinsic mitochondrial ATP production)
61 and reduced mitochondrial DNA copy number (mtDNA^{num})¹⁻⁵. However, these observations
62 have not been consistent across studies, and some would argue that declines in mitochondrial
63 function are normalised when differences in physical activity levels ($\dot{V}O_2$), mitochondrial
64 content and insulin action are considered⁶⁻⁸. It is also noteworthy that mitochondrial respiration
65 (with or without normalisation for mitochondrial content) does not change following gastric-
66 bypass induced improvements in insulin sensitivity⁹. Therefore, whether reduced intrinsic
67 mitochondrial function is causative in the induction of insulin resistance/T2DM, or contributes
68 to increased susceptibility to T2DM, or arises as a consequence of existing insulin resistance
69 remains an openly debated topic.

70

71 Mitochondria are the site of cellular oxidative phosphorylation processes in which fat,
72 carbohydrates, and amino acids are oxidatively decarboxylated to produce reducing equivalents,
73 which are subsequently used to generate ATP. Also, mitochondrial function is an integral part
74 of glucose-stimulated insulin secretion in pancreatic beta-cells¹⁰. The dynamic equilibrium
75 between mtDNA synthesis and degradation determines the mtDNA^{num}, which is relatively stable
76 under normal physiological conditions. However, changes in mtDNA^{num} are associated with

77 pathological changes in tissues and organs. Human mtDNA resides in hundreds to thousands of
78 copies in each cell and encodes for 13 structural proteins, which are subunits of the oxidative
79 phosphorylation electron transport chain, in addition to 2 ribosomal RNAs (rRNA) and 22
80 transfer RNAs (tRNA). However, mtDNA replication, transcription, translation and repair is
81 controlled by proteins encoded by nuclear DNA (nDNA) ¹¹. Qualitative changes in the mtDNA
82 sequence induced by the mitochondrial reactive oxygen species (ROS), such as mutations and
83 deletions, have been implicated in the pathogenesis of T2DM ¹². However, this can only account
84 for a small proportion of patients with T2DM ¹³.

85 Nevertheless, it was demonstrated that the content of mtDNA decreased in patients with T2DM
86 ¹⁴⁻¹⁷ and that reduced mtDNA levels precede the development of diabetes ¹⁴, although not
87 consistently ^{7, 8}. A confounding factor that may have contributed to these conflicting
88 observations is age since an age-related decline in mtDNA^{num} was previously identified in
89 isolated human islets ^{15, 16} and rodent skeletal muscle ¹⁸. Similarly, regular physical activity by
90 amplifying the signal for mitochondrial biogenesis can increase mitochondrial content and
91 function in young and older volunteers ¹⁹, and also increases insulin sensitivity (IS) ²⁰, while
92 deconditioning has the opposite effects ²¹. Overall, therefore, it would be useful if one could
93 control for those factors known to contribute to variation in the mtDNA copy number, such as
94 age and aerobic training status (maximal oxygen uptake), to provide more informative insight
95 of the role of mtDNA^{num} in T2DM risk.

96
97 The present study therefore aimed to identify whether indices of whole-body insulin sensitivity
98 (ISI) and insulin resistance (IR) were associated with (1) maximal oxygen uptake, (2) indices of
99 skeletal muscle intrinsic mitochondrial function, (3) measures of skeletal muscle mitochondrial
100 volume or (4) skeletal muscle mtDNA^{num} in a cohort of middle-aged male volunteers clustered

101 into sedentary T2DM (ST2DM), sedentary control (SC), and non-sedentary control (NSC) sub-
102 cohorts.

103 **Materials and Methods**

104 *Study participants*

105 This study was part of a previous project from which skeletal muscle fatty acid transporter
106 protein expression has been reported ²². A total of 10 male ST2DM patients, 10 normoglycaemic
107 SC male volunteers, and 10 normoglycaemic NSC male volunteers provided informed consent
108 to participate in this study, which was approved by the Maastricht University Medical Ethics
109 Committee. All volunteers were overweight (BMI >25, Table 1). The ST2DM and SC
110 volunteers were matched for age, BMI and whole-body fat mass (Table 1), and none was or had
111 been engaged in a physical activity training programme.

112 In contrast, the NSC volunteers reported cycling 3-4 times each week for more than 45 min. The
113 inclusion of ST2DM patients was based entirely on their medical condition, and confirmation
114 was verified with an oral glucose tolerance test (OGTT). Patient medication included
115 Metformin, Amaryl, Lipitor, Glucophage, Avandia, Tolbutamide, Daonil and Statins. The
116 average duration of clinical T2DM up to the start of the study was 7.5 ± 1.1 yrs. Anthropometric,
117 physiological and biochemical parameters for each group of volunteers are shown in Table 1.

118

119 *Study protocol*

120 Subjects reported to the laboratory after an overnight fast. Following 30 min of supine rest, a
121 vastus lateralis muscle biopsy sample was obtained from each volunteer under local anaesthesia
122 (lignocaine 2%) using a Bergstrom biopsy needle. The muscle biopsy specimens were snap-
123 frozen in liquid nitrogen and stored at -80°C until analyses were undertaken at a later date.

124 Following a further 30 min of supine rest, a catheter (Baxter BV, Utrecht, the Netherlands) was
125 inserted into an antecubital vein, and a blood sample was drawn ($t = 0$ min), after which 75 g of
126 glucose (dissolved in 250 mL water) was ingested, and a further blood sample was collected at
127 $t = 120$ min. Plasma glucose concentrations (Table 1) were measured (Yellow Spring glucose

128 analyser) to assess glucose intolerance and type 2 diabetes according to the American Diabetes
 129 Association guidelines (www.diabetes.org) while serum, which was stored for 2 yrs at -80°C,
 130 was used to assess insulin concentration (ELISA kit; Mercodia, Uppsala, Sweden). A small
 131 blood specimen collected at 0 times was used to measure the glycosylated haemoglobin HbA1c
 132 using an A1CNow⁺ device (Medisave, UK).

133
 134 Maximal power output (W_{max}) and maximal oxygen uptake ($\dot{V}O_{2max}$) were determined on an
 135 electronically braked cycle ergometer (Lode Excalibur, Groningen, the Netherlands) during an
 136 incremental exhaustive exercise test undertaken one week before muscle biopsy sampling
 137 (Table 1). Oxygen uptake ($\dot{V}O_2$) and carbon dioxide production ($\dot{V}CO_2$) were measured
 138 continuously (Oxycon; Mijnhart, Breda, the Netherlands). Body composition was assessed
 139 using the hydrostatic weighing method in the morning after an overnight fast. Simultaneously,
 140 residual lung volume was measured by the helium-dilution technique using a spirometer
 141 (Volugraph 2000; Mijnhart, Bunnik, the Netherlands). Body weight was measured with a digital
 142 balance. Body fat percentage was calculated using Siri's equation²³. Fat-free mass (FFM) was
 143 calculated by subtracting fat mass (FM) from total body weight (Table 1).

144
 145 *Insulin sensitivity index (ISI_{0, 120})*: was calculated using serum insulin and plasma glucose
 146 concentrations in a fasted state (0 min) and 120 min post-oral glucose ingestion according to
 147 Gutt *et al.*²⁴. The ISI_{0,120} index (ml²/kg/μIU/min⁻¹) was defined as:

$$148 \quad ISI_{0, 120} = \frac{75,000 + (G_0 - G_{120}) \times 0.19 \times m}{120 \times G_{mean} \times \log(I_{mean})}$$

149 where 75,000 represents the oral glucose load in mg, G₀ represents fasting plasma glucose
 150 concentration (mg/dL), G₁₂₀ - represents plasma glucose concentration at 120 min (mg/dL), 0.19
 151 represents glucose space in L/kg body weight, m represents body mass (kg), 120 represents

152 duration of the test (min), I_{mean} represents mean serum insulin concentration during the test
153 (mIU/L) and G_{mean} represents mean plasma glucose concentration during the test (mmol/L).

154
155 *HOMA-IR*: The homeostatic model assessment (HOMA), which is a method used to quantify
156 insulin resistance, was calculated as:

$$157 \frac{\text{Plasma glucose concentration (mmol/L)} \times \text{serum insulin concentration (mIU/L)}}{22.5}$$

159 Where 22.5 is a normalising factor representing the product of normal fasting plasma glucose
160 concentration of 4.5 mmol/L and normal fasting plasma insulin concentration of 5 μ IU/mL.

161
162 *Muscle lipid peroxidation*: Muscle malondialdehyde (MDA) content was determined as an
163 indicator of lipid peroxidation products based on the method of Erdelmeier *et al.*²⁵. Briefly,
164 frozen muscle tissue was homogenised in 5 mmol/L butylated hydroxytoluene in 20 mmol/L
165 phosphate buffer pH 7.4, followed by centrifugation at 3,000 g at 4°C. Clear muscle lysate was
166 acidic hydrolysed at 60°C for 80 min followed by mixing with N-methyl-2-phenylindol in 3:1
167 (v/v) acetonitrile: methanol, incubation at 45°C for 60 min and finally centrifuged at 15,000 g
168 for 10 min to clarify. Absorbance was measured spectrophotometrically at 586 nm. The
169 concentration of malondialdehyde (μ mol/L/mg protein) was calculated using 1,1,3,3-
170 tetramethoxypropane as a standard.

171
172 *Muscle free fatty acid and triglyceride content*:
173 Frozen muscle aliquots were homogenized in a Potter glass homogenizer for 3 min with 200 μ l
174 buffer (10 mmol/L Tris/HCl, pH 7.0 containing 154 mmol/L NaCl, 1 mmol/L EDTA, and 1%
175 Triton X-100)/mg wet weight. The muscle lysates were then centrifuged at 24,000 g for 10 min.
176 The pellets were discarded, and each supernatant was split into two aliquots. In the first aliquot,

177 levels of unbound free fatty acids (NEFA) only were measured using a WAKO NEFA assay kit,
178 while in the second aliquot, through alkaline hydrolysis, the pools of free NEFA and NEFA
179 released from triglycerides hydrolysis were determined. A 10 point standard curve generated
180 from 1 mmol/L stock oleic acid solution was run in parallel. The triglycerides content was
181 calculated by subtracting the free NEFA from the pooled NEFA values.

182
183 *Muscle enzymes activities:* The activities of muscle mitochondrial glutamate dehydrogenase
184 (GluDH), citrate synthase (CS), and β -hydroacyl-CoA dehydrogenase (HAD) and cytosolic
185 Gly3P dehydrogenase (Gly3PDH) were measured as described previously ²¹. Briefly, following
186 the lysis of frozen muscle pieces (~5 mg wet weight) in buffer containing K₂HPO₄ and Triton
187 X-100 using a Potter Elvehjem homogenizer the enzyme activities were measured
188 spectrophotometrically in the presence of suitable cofactors, activators, and buffers specific for
189 each enzyme.

190
191 *Relative mtDNA copy number:* The extraction of nuclear (nDNA) and mitochondrial DNA
192 (mtDNA) from skeletal muscle was accomplished according to the manufacturer's
193 recommendations using Qiagen QIAamp® DNA Mini kit. Briefly, the procedure involved
194 initial tissue lysis in a buffer containing proteinase K, incubation for 3 hrs at 56°C to digest the
195 myofibril proteins followed by the spinning of the lysates on silica-membrane-based nucleic
196 acid purification columns and elution of the mtDNA and nDNA with appropriate buffers. Before
197 the addition of buffer AL (Qiagen), 4 μ l of free DNase activity RNase A stock solution 7,000
198 U/ml was added to each sample lysate. The quality and quantity of DNAs were assessed by
199 measurements at 260, 280 and 230 nm. The expression level of selected markers of nDNA and
200 mtDNA used to evaluate their abundance was accomplished by using TaqMan probe real-time
201 PCR. The TaqMan probe design for the detection of nDNA levels was based on interrogation

202 of the intron sequence spanning between exons 3-4 of the genomic hydroxymethylbilane
203 synthase (HMBS) gene. The probe design for detection of mtDNA levels was based on
204 interrogation of a stable fragment of the mtDNA loop, namely the mitochondrially encoded
205 NADH:ubiquinone oxidoreductase core subunit 1 (ND1). The $2^{-\Delta C_t}$ formula, where $\Delta = C_{tND1} -$
206 C_{tHMBS} , was used to express the relative number of mtDNA copies to nDNA.

207
208 *Statistical analysis*

209 Data in text, tables, and figures are expressed as mean \pm SEM, with $n=10$ in each experimental
210 group. Between-group differences were determined using one-factor analysis of variance
211 (ANOVA). A Scheffe's post-hoc test was applied in the case of a significant F-ratio to locate
212 between-group differences. Significance was set at the 0.05 level of confidence. The strength of
213 the linear correlations between the investigated variables (r) was obtained using Pearson's
214 correlation. Correlations were deemed to be significant at 0.05 and 0.01 levels (2-tailed). The
215 contribution of the variation in mitochondrial enzyme activity, mtDNA, FFM and $\dot{V}O_{2max}/FFM$
216 to overall variation in ISI and HOMA-IR were determined using a linear regression model (IBM
217 SPSS Statistics 24 package). Sample size calculation was calculated using G-Power software
218 (version 3.1.9.2, Dusseldorf University, Germany) for ANOVA one-way fixed effects given α
219 = 0.05, number of groups = 3, power = 0.8 and effect size = 0.6.

220

221 **Results**

222 *Participant anthropometric, physiological and biochemical characteristics*

223 The anthropometric, physiological and biochemical characteristics of the ST2DM, SC and NSC
224 groups are shown in Table 1. Subjects did not differ in age and whole-body fat-free mass. All
225 volunteers were overweight (BMI >25), and body mass index in ST2DM was significantly
226 greater than in NSC. Whole-body fat (%) in NSC was significantly less than in ST2DM and SC
227 and similar between ST2DM and SC groups.

228
229 Fasting plasma glucose concentration in ST2DM was significantly greater than NSC and SC.
230 Post-feeding (120 min) plasma glucose concentration in ST2DM was significantly greater than
231 SC and NSC. Fasting serum insulin concentration in ST2DM was significantly greater than
232 NSC. Post-feeding (120 min) serum insulin concentration was not different across groups,
233 although it tended to be less in NSC than in SC and ST2DM. The insulin sensitivity index (ISI)
234 in SC and NSC was significantly greater than in ST2DM, and HOMA-IR in SC and NSC was
235 significantly less than in ST2DM. Percentage glycated haemoglobin (HbA1c) in ST2DM was
236 significantly greater than in NSC and SC. Maximal oxygen uptake in ST2DM was no different
237 from SC, but when normalised to free fat mass (ml/min/kg FFA), $\dot{V}O_{2\max}$ in SC was significantly
238 greater than in ST2DM. Irrespective of the reference base, $\dot{V}O_{2\max}$ in NSC was significantly
239 greater than in ST2DM and SC (Table 1). Maximal power output in NSC was significantly
240 greater than SC and ST2DM.

241
242 *Muscle biopsy analyses*

243 The muscle content of free fatty acids, triglycerides, determined as indices of muscle lipid
244 availability and the muscle content of malondialdehyde (MDA), determined as an index of lipid
245 peroxidation are presented in Table 2. Due to a scarcity of muscle tissue in some of the subjects

246 in the NSC group, muscle metabolites could not be determined in all volunteers, and therefore,
247 this group was omitted. Nevertheless, no difference was observed between ST2DM and SC for
248 any parameter.

249 The individual and the mean relative mtDNA copy number displayed in Fig. 1 illustrates that
250 on average NSC had a significantly greater number of mtDNA copies than SC ($P<0.001$) and
251 ST2DM ($P<0.001$; $1,461\pm52$, 749 ± 34 , 454 ± 58 , respectively). The SC mtDNA copy number
252 was also greater than ST2DM ($P<0.01$).

253 The maximal activity of 3 mitochondrial enzymes (GluDH, CS and HAD) was determined as
254 indices of mitochondrial volume and function (CS). The activity of all was greater in NSC
255 compared with ST2DM and SC, but no differences were seen between ST2DM and SC (Fig.
256 2A). The maximal activity of the Gly3PDH enzyme was used as a marker of capacity for
257 glycolytic energy production. There was no significant difference between SC and ST2DM
258 volunteers, but Gly3PDH activity was less in NSC compared with both groups (both $P<0.001$;
259 Fig. 2B).

260 Pearson correlations between the relative mtDNA copy number and ISI, HOMA-IR, $\dot{V}O_{2max}$
261 normalised to FFM and mitochondrial enzyme activities across groups are presented in Table 3.
262 Across all individuals, mtDNA^{num} highly associated with ISI, HOMA-IR, mitochondrial enzyme
263 activities (GluDH, and CS) and $\dot{V}O_{2max}/FFM$.

264

265 **Discussion**

266 The present study demonstrates that in a cohort of overweight, middle-aged male volunteers,
267 mtDNA^{num} is ordered as NSC > SC > ST2DM. Furthermore, across all individuals, mtDNA^{num}
268 was highly correlated with ISI, HOMA-IR, mitochondrial volume markers (GlutDH and CS
269 activities) and $\dot{V}O_{2max}$ normalised to FFM, while the associations between indices of
270 mitochondrial volume and ISI and HOMA-IR were less robust. These observations, together
271 with the knowledge that there was no difference in body composition or muscle lipids between
272 SC and ST2DM volunteers, leads to the conclusion that mtDNA^{num} is a sensitive index of insulin
273 sensitivity, which most likely reflects mitochondrial mass and supports the notion that regular
274 exercise exerts a protective role against the development of IR and T2DM.

275

276 In the present study, a significant difference in mtDNA^{num} was observed between SC and
277 ST2DM groups, despite no between-group differences in muscle CS activity. Frequently, CS
278 activity shows concordance with proteins entirely coded by mtDNA, such as complex I, II and
279 IV activity in both young and older subjects^{26, 27}, and it is often used as an index of muscle
280 ‘mitochondrial content/volume’. However, it has to be recognised that CS protein is coded by
281 nDNA, rather than by mtDNA, indicating that the discordance between CS activity and mtDNA
282 in the present study may not be entirely unexpected. In line with the CS findings, other
283 mitochondrial enzyme activity measurements (GlutDH and HAD), body composition
284 measurements (% whole body fat and FFM), muscle levels of free fatty acids, triglycerides and
285 malondialdehyde (a marker of lipid peroxidation; all of which have been reported to be causative
286 of muscle insulin resistance), were similar in the SC and ST2DM groups. However, the $\dot{V}O_{2max}$
287 normalised to FFM in SC was significantly greater than in the T2DM group, presumably
288 reflecting greater habitual physical activity levels in SC compared with ST2DM. In line with
289 this contention, Table 3 illustrates that mtDNA^{num} was found to associate strongly with indices

290 of insulin sensitivity (ISI and HOMA-IR) and $\dot{V}O_{2max}$ normalised to FFM across all volunteers,
291 and far better than the other muscle level measurements made.

292 It is worth commenting that the whole-body ISI did not associate with $\dot{V}O_{2max}$ across all study
293 volunteers. This finding may be accounted for by the presence of additional factors that
294 contribute to the biological variation of ISI and $\dot{V}O_{2max}$. Indeed, our linear regression model
295 showed that 68% of the variation of ISI was accounted for by the variation of the mtDNA,
296 GlutDH and FFM (all muscle related), while the variation of $\dot{V}O_{2max}$ was more most likely to
297 be accounted for by adaptations of the cardiovascular and pulmonary systems in addition to
298 those adaptations occurring at the muscle level.

299
300 Declines in mtDNA^{num} and mitochondrial function have been linked to human ageing²⁸ and
301 thereby age-related reductions in physical function²⁹. This age-related decline in mtDNA^{num}
302 does not appear to be gender-related as mtDNA^{num} in women and men was found to be almost
303 the same³⁰. In contrast, mtDNA^{num} appears to be preserved in heart muscle with ageing,
304 presumably due to its continuous contraction state¹⁸. In keeping with these observations, human
305 mtDNA^{num} also appears to be tissue-specific. Thus, values are reported to be greatest in muscle
306 tissue, followed by blood vessels, and lowest in leucocytes (in both T2DM and control subjects)
307³¹. Although oxidative stress stimulates mitochondrial biogenesis, it also induces a greater
308 degree of apoptosis in T2DM, resulting in a decrease in muscle tissue mtDNA^{num}³².

309
310 In conclusion, we report here the existence of a significant difference in ISI, HOMA-IR and
311 mtDNA^{num} across SC and T2DM volunteers, despite indices in mitochondrial volume and
312 function, and body composition and muscle free fatty acids, triglycerides and malondialdehyde
313 being similar between sedentary and T2DM volunteers. Moreover, we found that mtDNA^{num}
314 strongly correlates with indices of insulin sensitivity and $\dot{V}O_{2max}$, which most likely reflect

315 mitochondrial mass and supports the evidence that non-sedentary behaviour in the form of
316 regular exercise exerts a protective role against the development of IR and T2DM.

317 **Acknowledgements**

318 This work was supported by the Medical Research Council [grant number MR/P021220/1].

319 The authors would like to thank the volunteers who participated in the present study.

320 **Conflict of interest**

321 There are no competing or conflicting interests

322 **Author contributions**

323 DTC - conducting experiments, collection, analysis, and interpretation of data, figures, literature
324 search, writing of the manuscript, and final approval.

325 DC - data collection, analysis and interpretation, writing of the manuscript, and final approval.

326 MMP - study design, conducting experiments, and manuscript final approval.

327 LBV - study design, conducting experiments, and manuscript final approval.

328 LVL - study design, data interpretation, and manuscript final approval.

329 PLG - data interpretation, writing of the manuscript, and final approval.

330 **Patients and healthy volunteers consent:** obtained

331

References

- 332
333
334 1. Kelley DE, He J, Menshikova EV, Ritov VB. Dysfunction of mitochondria in human
335 skeletal muscle in type 2 diabetes. *Diabetes* 2002; **51**(10): 2944-50.
- 336 2. Befroy DE, Petersen KF, Dufour S, Mason GF, de Graaf RA, Rothman DL *et al.*
337 Impaired mitochondrial substrate oxidation in muscle of insulin-resistant offspring of
338 type 2 diabetic patients. *Diabetes* 2007; **56**(5): 1376-81.
- 339 3. Petersen KF, Dufour S, Befroy D, Garcia R, Shulman GI. Impaired mitochondrial
340 activity in the insulin-resistant offspring of patients with type 2 diabetes. *N Engl J Med*
341 2004; **350**(7): 664-71.
- 342 4. Schrauwen-Hinderling VB, Kooi ME, Hesselink MK, Jeneson JA, Backes WH, van
343 Echteld CJ *et al.* Impaired in vivo mitochondrial function but similar intramyocellular
344 lipid content in patients with type 2 diabetes mellitus and BMI-matched control subjects.
345 *Diabetologia* 2007; **50**(1): 113-20.
- 346 5. Sleigh A, Raymond-Barker P, Thackray K, Porter D, Hatunic M, Vottero A *et al.*
347 Mitochondrial dysfunction in patients with primary congenital insulin resistance. *J Clin*
348 *Invest* 2011; **121**(6): 2457-61.
- 349 6. Hey-Mogensen M, Hojlund K, Vind BF, Wang L, Dela F, Beck-Nielsen H *et al.* Effect
350 of physical training on mitochondrial respiration and reactive oxygen species release in
351 skeletal muscle in patients with obesity and type 2 diabetes. *Diabetologia* 2010; **53**(9):
352 1976-85.
- 353 7. Boushel R, Gnaiger E, Schjerling P, Skovbro M, Kraunsoe R, Dela F. Patients with type
354 2 diabetes have normal mitochondrial function in skeletal muscle. *Diabetologia* 2007;
355 **50**(4): 790-6.
- 356 8. Asmann YW, Stump CS, Short KR, Coenen-Schimke JM, Guo Z, Bigelow ML *et al.*
357 Skeletal muscle mitochondrial functions, mitochondrial DNA copy numbers, and gene

- 358 transcript profiles in type 2 diabetic and nondiabetic subjects at equal levels of low or
359 high insulin and euglycemia. *Diabetes* 2006; **55**(12): 3309-19.
- 360 9. Lund MT, Larsen S, Hansen M, Courraud J, Floyd AK, Stockel M *et al.* Mitochondrial
361 respiratory capacity remains stable despite a comprehensive and sustained increase in
362 insulin sensitivity in obese patients undergoing gastric bypass surgery. *Acta Physiol*
363 (*Oxf*) 2018; **223**(1): e13032.
- 364 10. Kwak SH, Park KS, Lee KU, Lee HK. Mitochondrial metabolism and diabetes. *J*
365 *Diabetes Investig* 2010; **1**(5): 161-9.
- 366 11. Koopman WJ, Willems PH, Smeitink JA. Monogenic mitochondrial disorders. *N Engl J*
367 *Med* 2012; **366**(12): 1132-41.
- 368 12. Kadowaki T, Kadowaki H, Mori Y, Tobe K, Sakuta R, Suzuki Y *et al.* A subtype of
369 diabetes mellitus associated with a mutation of mitochondrial DNA. *N Engl J Med* 1994;
370 **330**(14): 962-8.
- 371 13. Alcolado JC, Thomas AW. Maternally inherited diabetes mellitus: the role of
372 mitochondrial DNA defects. *Diabet Med* 1995; **12**(2): 102-8.
- 373 14. Lee HK, Song JH, Shin CS, Park DJ, Park KS, Lee KU *et al.* Decreased mitochondrial
374 DNA content in peripheral blood precedes the development of non-insulin-dependent
375 diabetes mellitus. *Diabetes Res Clin Pract* 1998; **42**(3): 161-7.
- 376 15. Nile DL, Brown AE, Kumaheri MA, Blair HR, Heggie A, Miwa S *et al.* Age-related
377 mitochondrial DNA depletion and the impact on pancreatic Beta cell function. *PLoS One*
378 2014; **9**(12): e115433.
- 379 16. Cree LM, Patel SK, Pyle A, Lynn S, Turnbull DM, Chinnery PF *et al.* Age-related
380 decline in mitochondrial DNA copy number in isolated human pancreatic islets.
381 *Diabetologia* 2008; **51**(8): 1440-3.

- 382 17. Czajka A, Ajaz S, Gnudi L, Parsade CK, Jones P, Reid F *et al.* Altered Mitochondrial
383 Function, Mitochondrial DNA and Reduced Metabolic Flexibility in Patients With
384 Diabetic Nephropathy. *EBioMedicine* 2015; **2**(6): 499-512.
- 385 18. Barazzoni R, Short KR, Nair KS. Effects of aging on mitochondrial DNA copy number
386 and cytochrome c oxidase gene expression in rat skeletal muscle, liver, and heart. *J Biol*
387 *Chem* 2000; **275**(5): 3343-7.
- 388 19. Menshikova EV, Ritov VB, Fairfull L, Ferrell RE, Kelley DE, Goodpaster BH. Effects
389 of exercise on mitochondrial content and function in aging human skeletal muscle. *J*
390 *Gerontol A Biol Sci Med Sci* 2006; **61**(6): 534-40.
- 391 20. Borghouts LB, Keizer HA. Exercise and insulin sensitivity: a review. *Int J Sports Med*
392 2000; **21**(1): 1-12.
- 393 21. Wibom R, Hultman E, Johansson M, Matherei K, Constantin-Teodosiu D, Schantz PG.
394 Adaptation of mitochondrial ATP production in human skeletal muscle to endurance
395 training and detraining. *J Appl Physiol (1985)* 1992; **73**(5): 2004-10.
- 396 22. Pelsers MM, Tsintzas K, Boon H, Jewell K, Norton L, Luiken JJ *et al.* Skeletal muscle
397 fatty acid transporter protein expression in type 2 diabetes patients compared with
398 overweight, sedentary men and age-matched, endurance-trained cyclists. *Acta Physiol*
399 *(Oxf)* 2007; **190**(3): 209-19.
- 400 23. Siri WE. The gross composition of the body. *Adv Biol Med Phys* 1956; **4**: 239-80.
- 401 24. Gutt M, Davis CL, Spitzer SB, Llabre MM, Kumar M, Czarnecki EM *et al.* Validation
402 of the insulin sensitivity index (ISI(0,120)): comparison with other measures. *Diabetes*
403 *Res Clin Pract* 2000; **47**(3): 177-84.
- 404 25. Erdelmeier I, Gerard-Monnier D, Yadan JC, Chaudiere J. Reactions of N-methyl-2-
405 phenylindole with malondialdehyde and 4-hydroxyalkenals. Mechanistic aspects of the
406 colorimetric assay of lipid peroxidation. *Chem Res Toxicol* 1998; **11**(10): 1184-94.

- 407 26. Larsen S, Nielsen J, Hansen CN, Nielsen LB, Wibrand F, Stride N *et al.* Biomarkers of
408 mitochondrial content in skeletal muscle of healthy young human subjects. *J Physiol*
409 2012; **590**(14): 3349-60.
- 410 27. Crane JD, Devries MC, Safdar A, Hamadeh MJ, Tarnopolsky MA. The effect of aging
411 on human skeletal muscle mitochondrial and intramyocellular lipid ultrastructure. *J*
412 *Gerontol A Biol Sci Med Sci* 2010; **65**(2): 119-28.
- 413 28. Short KR, Bigelow ML, Kahl J, Singh R, Coenen-Schimke J, Raghavakaimal S *et al.*
414 Decline in skeletal muscle mitochondrial function with aging in humans. *P Natl Acad*
415 *Sci USA* 2005; **102**(15): 5618-5623.
- 416 29. Hebert SL, Marquet-de Rouge P, Lanza IR, McCrady-Spitzer SK, Levine JA, Middha S
417 *et al.* Mitochondrial Aging and Physical Decline: Insights From Three Generations of
418 Women. *J Gerontol A Biol Sci Med Sci* 2015; **70**(11): 1409-17.
- 419 30. Kaaman M, Sparks LM, van Harmelen V, Smith SR, Sjolín E, Dahlman I *et al.* Strong
420 association between mitochondrial DNA copy number and lipogenesis in human white
421 adipose tissue. *Diabetologia* 2007; **50**(12): 2526-33.
- 422 31. Hsieh CJ, Weng SW, Liou CW, Lin TK, Chen JB, Tiao MM *et al.* Tissue-specific
423 differences in mitochondrial DNA content in type 2 diabetes. *Diabetes Res Clin Pract*
424 2011; **92**(1): 106-10.
- 425 32. Lee HC, Wei YH. Oxidative stress, mitochondrial DNA mutation, and apoptosis in
426 aging. *Exp Biol Med (Maywood)* 2007; **232**(5): 592-606.
- 427

428 **Table 1.** Anthropometric, physiological and biochemical characteristics of participants

429

| | Type 2 diabetes <i>n</i> = 10 | Sedentary control <i>n</i> = 10 | Non-sedentary control <i>n</i> = 10 |
|--|---|---|---|
| Age (yrs) | 58.9±1.7 | 60.0±2.1 | 57.4±0.9 |
| BMI (kg/m ²) | 28.9±1.2 | 27.5±0.5 | 25.5±0.7* |
| Whole-body fat (%) | 28.8±1.8 | 29.2±1.3 | 17.2±1.2*# |
| Whole-body FFM (kg) | 64.0±1.9 | 61.3±1.5 | 63.6±1.0 |
| Fasting plasma glucose (mmol/L) | 9.0±0.4 | 5.5±0.2* | 5.7±0.1* |
| Plasma glucose _{120 min} (mmol/L) | 16.8±1.0 | 5.3±0.49* | 5.3±0.4* |
| Fasting serum insulin _{0 min} (mIU/L) | 8.8±0.9 | 7.9±1.6 | 5.1±0.6* |
| Serum insulin _{120 min} (mIU/L) | 45.2±7.8 | 48.4±8.0 | 29.4±6.3 |
| Insulin sensitivity index (ISI) ^a | 46.8±1.8 | 83.6±5.8* | 103.0±8.6** |
| HOMA-IR | 3.63±0.35 | 1.82±0.37*** | 1.30±0.15*** |
| HbA1c (%) | 7.30±0.3 | 5.83±0.2* | 5.78±0.1* |
| $\dot{V}O_2$ max (L/min) | 2.90±0.20 | 3.19±0.19 | 3.80±0.12**# |
| $\dot{V}O_2$ max (mL/min/kg FFM) | 45.0±2.3 | 52.0±2.7* | 59.8±1.6***# |
| Wmax | 205±16 | 206±18 | 300±9***# |

430 BMI - body mass index, FFM - free fat mass, ^aml²/kg/μIU/min. *, **, *** Significantly431 different from Type 2 diabetes group (*P*<0.05, *P*<0.01 and *P*<0.001, respectively).432 #Significantly different from the sedentary control group (*P*<0.05).

433

434 **Table 2.** Muscle content of lipids and their oxidation product (malondialdehyde) in sedentary
435 control volunteers ($n=10$) and sedentary T2DM patients ($n=10$).

436

| Muscle metabolites | Sedentary control | Sedentary T2DM patients |
|---|--------------------------|--------------------------------|
| Malondialdehyde* | 30.1±4.2 | 32.4±5.4 |
| Intramuscular free fatty acids [#] | 11.66±0.68 | 15.21±2.92 |
| Intramuscular triglycerides [#] | 1.10±0.04 | 0.85±0.13 |

437 * $\mu\text{mol/mg}$ protein, [#] mmol/kg dry matter

Table 3. Pearson correlations between muscle mtDNA^{num}, whole body insulin sensitivity index (ISI) and insulin resistance (HOMA-IR), and several muscle mitochondrial capacity indices (glutamate dehydrogenase - GlutDH, citrate synthase - CS, and β -hydroxy acyl-CoA dehydrogenase activity - HAD) and $\dot{V}O_{2\max}$ normalised to free fat mass (FFM), in three groups of late middle-aged males clustered according to aerobic capacity and the presence of T2DM.

| | | mtDNA | ISI | HOMA- IR | GlutDH | CS | HAD | $\dot{V}O_{2\max}/\text{FFM}$ |
|-------------------------------|---------------------|--------|--------|-------------|--------|--------|--------|-------------------------------|
| mtDNA | Pearson correlation | - | 0.688 | -0.542 | 0.603 | 0.604 | 0.382 | 0.684 |
| | Sig (2-tailed) | - | 0.001 | 0.002 | 0.001 | 0.001 | 0.037 | 0.001 |
| ISI | Pearson correlation | 0.688 | - | -0.662 | 0.488 | 0.328 | 0.187 | 0.178 |
| | Sig (2-tailed) | 0.001 | - | -0.001 | 0.006 | 0.077 | 0.321 | 0.348 |
| HOMA-IR | Pearson correlation | -0.542 | -0.662 | - | -0.424 | -0.325 | -0.112 | -0.229 |
| | Sig (2-tailed) | 0.002 | 0.001 | - | 0.020 | 0.080 | 0.554 | 0.224 |
| GlutDH | Pearson correlation | 0.603 | 0.488 | -0.424 | - | 0.715 | 0.605 | 0.217 |
| | Sig (2-tailed) | 0.001 | 0.006 | 0.020 | - | 0.001 | 0.001 | 0.249 |
| CS | Pearson correlation | 0.604 | 0.328 | -0.325 | 0.715 | - | 0.654 | 0.161 |
| | Sig (2-tailed) | 0.001 | 0.077 | 0.080 | 0.001 | - | 0.001 | 0.394 |
| HAD | Pearson correlation | 0.382 | 0.187 | -0.112 | 0.605 | 0.654 | - | 0.350 |
| | Sig (2-tailed) | 0.037 | 0.321 | 0.554 | 0.001 | 0.001 | - | 0.105 |
| $\dot{V}O_{2\max}/\text{FFM}$ | Pearson correlation | 0.684 | 0.278 | -0.229 | 0.217 | 0.161 | 0.350 | - |
| | Sig (2-tailed) | 0.001 | 0.240 | 0.224 | 0.249 | 0.394 | 0.105 | - |

Figure Legends

Figure 1 Relative mtDNA copy number in vastus lateralis muscle of sedentary Type 2 diabetes mellitus (ST2DM), sedentary control (SC) and non-sedentary control (NSC) volunteers. Data represent mean \pm SEM and individual values. Significant difference between groups depicted as: ** ($P<0.01$); *** ($P<0.001$).

Figure 2. Mitochondrial volume markers (glutamate dehydrogenase, citrate synthase and β -hydroxyacyl-CoA dehydrogenase activity; Fig. 2A) and glycolytic capacity index (glyceraldehyde-3P-dehydrogenase; Fig. 2B) in vastus lateralis muscle of sedentary Type 2 diabetes mellitus (ST2DM), sedentary control (SC) and non-sedentary control (NSC) volunteers. Data represent mean \pm SEM and individual values. *, **, *** Significantly different from ST2DM; $P<0.05$, $P<0.01$, and $P<0.010$, respectively. †, ††, ††† Significantly different from SC; $P<0.05$, $P<0.01$, $P<0.001$, respectively.




Figure 1




Figure 2