

Article

Not peer-reviewed version

Conofolidine, a Natural Plant Alkaloid Causes Apoptosis and Senescence in Cancer Cells

[Mohammed Zuhair Al-Hayali](#) , Choy-Eng Nge , Kuan Hon Lim , Hilary M Collins , Toh Seok Kam ,
[Tracey Dawn Bradshaw](#) *

Posted Date: 8 January 2024

doi: 10.20944/preprints202401.0602.v1

Keywords: conofolidine; bisindole alkaloids; carcinoma cell lines; cytotoxicity; apoptosis; senescence; ROS generation; γ -H2AX



Preprints.org is a free multidiscipline platform providing preprint service that is dedicated to making early versions of research outputs permanently available and citable. Preprints posted at Preprints.org appear in Web of Science, Crossref, Google Scholar, Scilit, Europe PMC.

Copyright: This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

Article

Conofolidine, a Natural Plant Alkaloid Causes Apoptosis and Senescence in Cancer Cells

Mohammed Al-Hayali ^{1,2,*}, Choy-Eng Nge ³, Kuan Hon Lim ⁴, Hilary Collins ², Toh-Seok Kam ³ and Tracey D. Bradshaw ^{2,*}

¹ Al-Kitab University, College of Pharmacy, P.O. Box: 1068, Altun Kupri, 36015, Iraq

² School of Pharmacy, Biodiscovery Institute, University of Nottingham, University Park, Nottingham NG7 2RD, UK; hilary.collins@nottingham.ac.uk (H.C.)

³ Department of Chemistry, Faculty of Science, University of Malaya, 50603 Kuala Lumpur, Malaysia; choyeng86@gmail.com (C.E.N.); tskam@um.edu.my (T.S.K.)

⁴ School of Pharmacy, University of Nottingham Malaysia, Jalan Broga, 43500 Semenyih, Selangor Darul Ehsan, Malaysia; kuanhon.lim@nottingham.edu.my (K.H.L.)

* Correspondence: mohammed.zuhair@uoalkitab.edu.iq (M.A.-H.); tracey.bradshaw@nottingham.ac.uk (T.D.B.)

Simple Summary: Cancer is the second leading global cause of death. For decades, scientists exploited plants as the main source of numerous chemotherapeutic agents. Our research aim is to identify a new potential anticancer agent from a library of plant-derived compounds and explore its mode(s) of action. Conofolidine, isolated from the Malayan plant *Tabernaemontana corymbosa*, elicited the most potent anticancer activity (among the four alkaloids tested) against human cancer cells derived from the colon, breast, lung and pancreas. Conofolidine triggered two death pathways: apoptosis and senescence. At the molecular level, conofolidine increased oxidative stress and interacted with essential regulatory proteins responsible for cancer cell proliferation and DNA repair. To our knowledge, this is the first work investigating conofolidine's anticancer action. Further studies of this promising natural product are justified.

Abstract: Natural products contribute substantially to anticancer therapy; the plant kingdom provides an important source of molecules. Conofolidine is a novel *Aspidosperma-Aspidosperma* bisindole alkaloid isolated from the Malayan plant *Tabernaemontana corymbosa*. Herein, we report conofolidine's anticancer activity together with that of three other bisindoles - conophylline, leucophyllidine and bipleiophylline against human-derived carcinoma cell lines. Remarkably, conofolidine was able to induce apoptosis (as observed in MDA-MB-468 breast cancer cells) or senescence (as detected in HT-29 colorectal carcinoma cells). Annexin V-FITC/PI, caspase activation and PARP cleavage confirmed the former while positive β -gal staining corroborated the latter. Evident cell cycle perturbations were observed comprising S-phase depletion, accompanied by downregulated CDK2, and cyclins (A2, D1) with p21 upregulation. Confocal imaging of HCT-116 cells revealed induction of aberrant mitotic phenotypes - multi-nucleation, membrane blebbing and DNA-fragmentation. The DNA integrity assessment of HCT-116 and MDA-MB-468 showed irreparable damage identified by increased fluorescent γ -H2AX during the G1 cell cycle phase. Furthermore, γ -H2AX foci were visually validated in HCT-116 and MDA-MB-468 cells using confocal microscopy. Conofolidine increased oxidative stress, preceding apoptosis- and senescence-induction in most carcinoma cell lines as seen by enhanced ROS levels accompanied by NQO1 expression. Collectively, we present conofolidine as a potential anticancer candidate capable of inducing heterogeneous modes of cancer cell death *in vitro*, encouraging further preclinical evaluation of this natural product.

Keywords: conofolidine; bisindole alkaloids; carcinoma cell lines; cytotoxicity; apoptosis; senescence; ROS generation; γ -H2AX

1. Introduction

The impact of natural products on new medicines' discoveries has been indispensable. Over the past 30 years, molecules inspired by nature constituted the mainstay of chemotherapy [1]. Moreover, natural products continue to provide novel lead structures which serve as templates for the synthesis of more potent and safer compounds [2], highlighting the importance of continued research for new anticancer agents from nature.

Apoptosis-based chemotherapy constitutes a core focus for anticancer drug development as achieving this goal is expected to profoundly improve clinical response [3,4]. However, apoptosis evasion is a cancer hallmark [5], and non-apoptotic death mechanisms e.g. senescence have also been shown to improve the clinical outcome of anticancer chemotherapy [6]. While apoptosis can induce early killing of rapidly dividing cancer cells by high doses of anticancer agents; tumour recurrence, drug resistance and normal tissue damage are common complications [6]. Alternatively, senescence may exert its tumour-suppressive effect at sub-apoptotic doses *via* induction of permanent cell cycle arrest offering an effective strategy against cancer with fewer adverse effects [3,7].

Tabernaemontana corymbosa (Apocynaceae) is a Malaysian rainforest species and a rich source of promising anticancer agents. We report herein for the first time *in vitro* dual apoptosis- and senescence-inducing abilities of a novel *Aspidosperma-Aspidosperma* bisindole, conofolidine from this plant. Conofolidine's indole moieties closely resemble that of jerantinines A and B also isolated from *Tabernaemontana corymbosa*; jerantinines A and B displayed potent antiproliferative activities *via* suppression of tubulin polymerization [8,9]. In this study, we demonstrate conofolidine's anticancer activity, together with that of three other bisindoles - conophylline, leucophyllidine and bipleiophylline, Figure 1 [10–13] - against human-carcinoma cell lines derived from breast, colorectal, lung and pancreatic carcinomas and B-cell lymphomas. In 2016, Nge *et al.* reported the isolation and structure elucidation of conofolidine (as a new member of the conophylline-group; Figure 1) and preliminary growth inhibitory effects [14]. Herein, further anticancer activities, cytotoxicity, cell cycle and death pathways triggered (apoptosis and senescence) have been investigated after treatment of carcinoma cells with conofolidine. Preliminary data showed a remarkable potency against carcinoma cells, e.g., $GI_{50}=0.054 \mu\text{M}$, MCF-7. Effects against foetal lung fibroblasts (MRC-5) and normal breast epithelial cells (MCF10A) were assessed for preliminary examination of potential selectivity towards carcinoma cells. Mode of action studies led to the identification of ROS production as one possible mechanism of conofolidine's growth-inhibitory actions. Furthermore, increased DNA double strand breaks (DSBs) were detected, accompanied by perturbation of essential cell cycle regulatory proteins - CDK2, p21, cyclins A2 and D1 in a time-dependent fashion.




Figure 1. Structures of conofolidine(1), conophylline(2), leucophyllidine(3) and bipleiophylline(4).

2. Materials and Methods

2.1. Isolation and characterization of conofolidine

All 4 bisindoles were kindly provided by Prof. Kam, University of Malaya. Conofolidine was isolated from the EtOH extract of the leaves of *Tabernaemontana corymbosa* Roxb. ex Wall (Apocynaceae). Its two indoles correspond to taberhanine and 11-hydroxydeoxoapodine which are connected *via* a central dihydrofuran ring [14]. The structure and absolute configuration of conofolidine (Figure 1.1, $C_{44}H_{50}N_4O_{10}$, MW=794.89) were established by spectroscopic methods and X-ray diffraction analysis [14].

2.2. Cell culture

Cancer cell lines were obtained from the American Type Culture Collection (ATCC), grown and maintained in RPMI-1640 medium supplemented with 10% foetal bovine serum (FBS). Cell lines included MCF-7, MDA-MB-231, MDA-MB-468, SK-BR3, T-47D, ZR-75-B, Caco-2, HCT-116, HT-29, A549, MIAPaCa-2, PANC-1, DoHH2, Vallois. Cells were subcultured twice weekly to sustain their logarithmic growth and were allowed to grow in a humidified atmosphere at 37 °C, 5% CO₂ incubator. After ≤30 passages, and to reduce phenotypic or genotypic drift, new low passage cells were reintroduced from cell stocks that had been cryopreserved at -180 °C. Test compound top stocks were kept frozen at -80 °C as 10 mM/10 μL-aliquots in DMSO which were thawed promptly before use.

2.3. MTT assay

3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT, Alfa-Aesar, Lancashire, UK) colourimetric assays were used for testing the antiproliferative ability of test agents on the cancer cells. Cells were seeded into 96-well plates (3×10^3 cells/well) and allowed to attach overnight prior to test compound introduction (5 nM - 10 μM, $n \geq 4$). Serial dilutions were freshly prepared in culture medium before assay. Two measurements of viable cells (at the time of treatment introduction and

after 72 h exposure) were conducted following MTT addition. Plates were then incubated for 3 h allowing metabolic reduction of yellow MTT into insoluble purple formazan crystals which were solubilised in 150 μ L DMSO after supernatants' aspiration. Absorbance readings were recorded at 570 nm in a plate reader (PerkinElmer Life and Analytical Sciences, Buckinghamshire, UK). Dose-response profiles were constructed from mean absorbance values, and test agent concentrations that inhibited growth by 50% (GI_{50}) were calculated.

2.4. Clonogenic assay

Clonogenic assays were used to assess the ability of single cells to survive, proliferate and form colonies following brief exposure to conofolidine. Cells (~400/well) were seeded into 6-well plates in 2 mL medium. After 24 h incubation, conofolidine was introduced (0.1 and 0.5 μ M) while medium alone was added to controls. Following 24 h treatment, medium containing conofolidine was gently aspirated, cells were washed (2 \times sterile PBS, 37 $^{\circ}$ C) before fresh conofolidine-free medium was introduced, and plates were re-incubated for 7–9 days. Cells were examined daily until cells in controls had formed colonies consisting of >50 cells. Wells were washed with PBS, fixed with methanol (100%; 15 min), stained with methylene blue (0.5%; 15 min), washed (dH₂O), and air-dried, and stained colonies were then counted.

2.5. Cell cycle analysis

The method used is adapted from that described by Nicoletti *et al.* (1991) [15]. Cells were seeded and allowed to attach overnight in 6-well plates (1.0×10^5 and 0.5×10^5 cells/well for 24 h and 72 h exposures, respectively). Following treatment, floating cells were collected, attached cells trypsinised and cells were pooled together in FACS tubes. Samples were washed (PBS; 4 $^{\circ}$ C), and pelleted by centrifugation (1,200 rpm; 5 min; 4 $^{\circ}$ C). Cells were resuspended in fluorochrome solution (50 μ g/mL PI, 100 μ g/mL ribonuclease-A, 0.1% triton-X-100, 0.1% sodium citrate). Cell cycle analyses were performed using an FC-500 Beckman-Coulter flow cytometer. Data were collected and analysed using EXPO32 software.

2.6. Annexin-V/PI apoptosis assay

Cells were seeded at densities of 0.5 - 0.75×10^5 per well in a 6-well plate, incubated overnight and treated the following day. After 72 h $2 \times GI_{50}$ treatments, cells were harvested, washed (PBS; 4 $^{\circ}$ C) and transferred with the medium containing detached cells into FACS tubes. After centrifugation (1,200 rpm; 5 min; 4 $^{\circ}$ C) and aspiration of supernatants, cells were resuspended in 5 μ L annexin-V-FITC plus 100 μ L $1 \times$ annexin-V binding buffer (BD Pharmingen). Samples were briefly vortexed and incubated (protected from light for 15 min, 25 $^{\circ}$ C). PI (10 μ L) plus 400 μ L annexin-V binding buffer were added to each FACS tube followed by incubation for 10 min in the dark at 4 $^{\circ}$ C. Samples were analysed using Beckman Coulter FC-500 flow cytometer and EXPO32 software and percentages of early apoptotic (annexin-V positive (AV+), PI negative (PI-)) and late apoptotic cells (AV+, PI+) were recorded.

2.7. Caspase 3/7 assay

To measure executioner caspases' activity, the Caspase-Glo 3/7 (100 μ L; Promega) assay was conducted according to the manufacturer's protocol. Cells were seeded at a density of 5×10^3 in 100 μ L of medium/well in white-walled opaque 96-well plates and incubated overnight (37 $^{\circ}$ C, 5% CO₂). Conofolidine or medium alone was added for an additional 24 h. Caspase-Glo 3/7 reagents were then added to each well. After 30 min, relative luminescence was measured in a plate reader (PerkinElmer Life and Analytical Sciences, Buckinghamshire, UK).

2.8. Senescence β -galactosidase cell staining assay

Senescence was investigated using β -Gal staining (Cell Signaling Technology; CST (London, UK)). Cells were seeded at a density of 2.5×10^4 cell/well in 6-well plates, incubated overnight at 37 °C, 5% CO₂ and treated with conofolidine the following day. After 72 h, cells were washed with PBS and fixed before adding 1 mL β -Gal solution (1 mg/mL X-gal, pH=6.0). Plates were incubated at 37 °C \geq 16 h, before being inspected under a light inverted microscope for blue β -Gal staining; positive cells were counted in all wells.

2.9. Western Blotting

Following designated treatments, cells were collected and lysed using NP-40 lysis buffer containing protease and phosphatase inhibitors. Protein concentrations were measured using the Bradford assay [16]. Samples were then loaded (50 μ g protein each) and proteins were separated using sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE). By immunoblotting, proteins were transferred onto nitrocellulose membrane, probed against primary antibody (dilution 1:1000), and thereafter against secondary antibody (dilution 1:4000). Protein bands were detected using enhanced chemiluminescence (Amersham ECL Prime (Buckinghamshire, UK) and LI-COR Biosciences Ltd. C-DiGit[®] Blot Scanner, Cambridgeshire, UK). Densitometric analyses of western blots were performed using Image Studio. Primary antibodies were procured from CST (London, UK): whole and cleaved Poly (ADP-ribose) polymerase (PARP), CDK2, cyclin A2, cyclin D1, p21, NAD(P)H quinone dehydrogenase 1 (NQO1) and β -Actin. A secondary anti-mouse IgG/HRP antibody was purchased from Dako (Cambridgeshire, UK).

2.10. Confocal microscopy

Confocal microscopy was used to visualise the DNA-DSBs foci induced by conofolidine. Cells were seeded in eight-well μ -slides (Ibidi[®] Germany) and allowed to attach overnight in CO₂ incubator at 37 °C, then they were incubated with either medium alone or conofolidine or positive control – etoposide. Following 72 h exposure, cells were fixed with 3.7% formaldehyde in PBS, permeabilised with PBT (0.1% Triton-X-100 in PBS) and incubated with p-Histone γ -H2AX antibody (Sigma-Aldrich, UK) for DNA-DSBs foci staining and draq5 (Abcam, UK) for DNA staining. Images were captured using Zeiss LSM 510 META confocal microscope (Cambridgeshire, UK) and Zeiss LSM 510 image browser (version 4.2). The above protocol was used also for visual validation of mitotic abnormalities induced by conofolidine except that β -tubulin antibody TUBB1 (ThermoFisher-Scientific, UK) and vincristine - positive control - were used.

2.11. Monitoring of γ -H2AX phosphorylation induction by flow cytometry

Cells were seeded at a density of 1×10^6 in 100 mm² culture Petri dishes, incubated overnight and treated the following day with either conofolidine $2 \times \text{GI}_{50}$ or etoposide 2.0 μ M - positive control. After 72 h, cells were transferred into FACS tubes containing their respective medium. Cells were centrifuged (10 mins; 1500 RPM; 25 °C) and pelleted; supernatants were discarded and pellets were resuspended again. Thereafter, cells were fixed with 1% methanol-free formaldehyde in PBS, permeabilised with Triton-X-100 in PBS and incubated with p-Histone γ -H2AX antibody (Sigma-Aldrich, UK); dilution 1:3333 in 1% FBS in PBS; 1.5 h; 25 °C. For concurrent cell cycle analyses, cells were resuspended in PI (50 μ g/ml) and RNase (100 μ g/ml) in PBS solution (10 min.; 25 °C). The % of γ -H2AX (+) cells and cell cycle distribution were analysed using CyFlow[®] Space flow cytometer (Sysmex Partec, Germany) and Kaluza analysis software (Beckman Coulter, UK).

2.12. Reactive oxygen species determination

To measure ROS production, the ROS-Glo H₂O₂ (Promega) assay was conducted according to the manufacturer's protocol, which measures the most stable ROS-H₂O₂ level. Cells were plated at a density of $5 \times 10^3/70$ μ L medium in white-walled opaque bottom 96-well plates and incubated overnight (37 °C, 5% CO₂). Conofolidine or medium (10 μ L) was added to treated or control wells,

respectively, for an extra 24 h. After 18 h treatment, 20 μL H_2O_2 substrate was added to each well. Upon treatment period completion (24 h), 100 μL ROS-Glo solution was added to each well, and after 20 min, relative luminescence was measured in a plate reader (PerkinElmer Life and Analytical Sciences, Buckinghamshire, UK).

2.13. Statistical analysis

Experiments were repeated at least three times, internal repeats $n \geq 4$ (unless otherwise stated) and representative experiments are shown. Data are presented as mean \pm SD (or SEM). Statistical differences between study groups were analysed using two-way ANOVA (unless otherwise stated). Dunnett's multiple comparisons were used to test the significance, determined as p -value ≤ 0.05 .

3. Results

3.1. Conofolidine potently inhibits growth of cancer cells

The growth inhibitory properties of conofolidine (in addition to conophylline, leucophyllidine and bipleiophylline) were investigated by MTT assay. All cancer cell lines exhibited significant ($p \leq 0.05$) sensitivity to conofolidine (Table 1, Figure 2). Conofolidine remarkably suppressed oestrogen receptor-positive (ER+) MCF-7 growth (lowest $\text{GI}_{50} = 0.054 \mu\text{M}$). Triple-negative breast MDA-MB-468, colorectal HT-29 and pancreatic MIAPaCa-2 cells displayed similar GI_{50} values of $\sim 0.3 \mu\text{M}$. A549 lung and HCT-116 colorectal carcinoma cells were least sensitive ($\text{GI}_{50} = 0.405$ and $0.454 \mu\text{M}$ respectively). Conophylline, leucophyllidine and bipleiophylline were less potent yielding GI_{50} values of 0.07-1.7, 2.0-4.5 and 3.5-13.5 μM , respectively in the same carcinoma cell lines (Figure S1-3). DMSO vehicle did not affect cancer cell growth at the dilutions used (Figure S4).

To confirm conofolidine potency, and investigate its antitumour spectrum, additional cell lines were included in MTT assays (Figure S5, Table S1). Breast carcinoma cell lines were very sensitive to conofolidine (GI_{50} values $\leq 0.17 \mu\text{M}$); PANC-1 was similarly sensitive as MIAPaCa-2; while the lowest sensitivity was observed in Caco-2 cells ($\text{GI}_{50} = 0.417 \mu\text{M}$). Conofolidine's antiproliferative effect was also tested against two B-cell lymphoma cell lines (DoHH2 and Vallois), their GI_{50} values were $\leq 0.127 \mu\text{M}$ (Figure S5, Table S1). To study conofolidine's selectivity on cancerous compared to noncancerous cells, MRC-5 and MCF10A were used as *in vitro* models of foetal lung fibroblasts and proliferative benign fibrocystic epithelial breast cells respectively (Figure S5, Table S1). Conofolidine's GI_{50} in these cell lines were $\sim 1.0 \mu\text{M}$ for MRC-5 and $0.23 \mu\text{M}$ for MCF10A.

Table 1. Bisindoles' anti-proliferative effects on human carcinoma cell lines.

Human cell line		72 h exposure MTT GI_{50} (μM)			
Origin	Designation	Conofolidine	Conophylline	Leucophyllidine	Bipleiophylline
Breast carcinoma	MCF-7	0.054 \pm 0.08	0.066 \pm 0.03	2.79 \pm 0.98	3.49 \pm 0.47
	MDA-MB-468	0.292 \pm 0.03	0.557 \pm 0.08	1.99 \pm 0.03	5.81 \pm 1.11
Colon carcinoma	HCT-116	0.454 \pm 0.07	0.737 \pm 0.27	2.43 \pm 0.08	3.86 \pm 1.04
	HT-29	0.301 \pm 0.03	0.350 \pm 0.52	2.48 \pm 0.17	7.72 \pm 4.08
Pancreatic carcinoma	MIAPaCa-2	0.305 \pm 0.03	0.663 \pm 0.03	3.29 \pm 0.98	12.37 \pm 3.12
Lung carcinoma	A549	0.405 \pm 0.07	1.743 \pm 0.34	4.45 \pm 0.27	13.63 \pm 1.94

GI_{50} values were estimated from MTT data after 72 h exposure to bisindole alkaloids ($n=4$) and expressed as mean \pm SD of 3 independent trials.




Figure 2. Growth inhibitory effects of conofolidine in A549, HCT-116, HT-29, MCF-7, MDA-MB-468 and MIAPaCa-2. Each representative graph shows one independent MTT trial. Cells were seeded in 96-well plates (3×10^3 cells/well) and treated with TQ for 72 h. No. trials ≥ 3 ; n=4 per independent experiment.

3.2. Conofolidine potently inhibits colony formation

Clonogenic assays were performed to examine whether conofolidine can inhibit cancer cell colony formation. Guided by MTT assay results, two conofolidine concentrations were selected (0.1 and 0.5 μM). Conofolidine potently inhibited cancer cells' ability to form colonies (Figure 3, Figure S6). Breast MDA-MB-468 and pancreatic MIAPaCa-2 carcinoma colonies were dramatically decreased (99.9%) at 0.1 μM . At the same concentration, conofolidine suppressed MCF-7 (most sensitive cell line) colony formation by 91%, which, intriguingly, was similar to HCT-116 (less sensitive in MTT assay). However, comparable clonogenic inhibition profiles were seen in both A549 and HT-29 cells: at concentrations of 0.1 and 0.5 μM , conofolidine inhibited colony formation by ~30 and ~95%, respectively. Average plating efficiencies were A549 (68%), HCT-116 (45%), HT-29 (55%), MCF-7 (42%), MDA-MB-468 (28%) and MIAPaCa-2 (51%).




Figure 3. Representative photos of conofolidine's inhibition of MIAPaCa-2 colonies at 0.1 and 0.5 μ M. Mean \pm SD bars show mean survival fraction as % of control. Asterisk indicates significant inhibition ($p \leq 0.05$). Cells were seeded, treated with conofolidine (24 h) then medias replaced with fresh ones. When colonies contained ≥ 50 cells, colonies were fixed, stained and counted. Plating efficiencies ranged between $\sim 25\%$ and $\sim 65\%$. No. trials ≥ 3 ; $n=2$ per independent experiment.

3.3. Conofolidine caused potent perturbation of cell cycle progression

After confirming conofolidine's ability to inhibit cancer cell proliferation and colony formation, we sought to determine whether its mechanism of action involved cell cycle perturbation. Cell cycle analysis was conducted in A549, HCT-116, HT-29, MCF-7, MDA-MB-468 and MIAPaCa-2 after treatment with respective conofolidine $1\times$ and $2\times GI_{50}$ concentrations for 24 and 72 h (Figures 4 and 5). Conofolidine caused significant ($p \leq 0.05$) cell cycle perturbation including universal S-phase depletion, especially at $2\times GI_{50}$ treatment, declining to $\leq 10\%$ in HCT-116, HT-29, MCF-7 and MDA-MB-468 and $\leq 15\%$ in A549 and MIAPaCa-2 compared to 20-25% for untreated controls in all cell lines. Accompanying S-phase depletion, G1 arrest was observed, either sustained, in HT-29 and MCF-7 or temporarily in HCT-116 and MDA-MB-468; A549 exhibited decreased S-phase without G1 arrest. Interestingly, MIAPaCa-2 cells exhibited G2/M arrest while other cell lines exhibited significantly decreased G2/M events. After 24 h $1\times GI_{50}$ conofolidine treatment, G1 arrest accompanied S-phase reduction in HCT-116 ($\sim 48\%$ from control 40%), HT-29 (73% from control 55%), MCF-7 ($\sim 70\%$ from control 46%) and MDA-MB-468 ($\sim 65\%$ from control 41%). At $2\times GI_{50}$, 24 h exposure, comparable G1

increments were seen: HCT-116 (50%), HT-29 (73%), MCF-7 (67%) and MDA-MB-468 (62%). In MIAPaCa-2 cells, S-phase depletion was followed by accumulation of G2/M events after $2\times GI_{50}$, 24 and 72 h exposures: (~23% from control 14%). A less pronounced arrest in MIAPaCa-2 G2/M phase was seen at $1\times GI_{50}$, 24 h, however, it increased significantly after 72 h exposure reaching ~23% from control 14%. Concentration- and time-dependent increased pre-G1 (<2N-DNA) events were seen in all treatment groups of most cell lines except HT-29 and MCF-7 cells, in which significant increments were only observed at $2\times GI_{50}$, 72 h treatment: ~12% from control ~3%. Other cell lines exhibited significant pre-G1 at both $1\times$ and $2\times GI_{50}$ treatments. HCT-116 exhibited the most significant increments yielding ~20% at $1\times GI_{50}$, 24 and 72 h; and ~30% at $2\times GI_{50}$, 24 and 72 h exposures. In MDA-MB-468 and A549, comparable increases were observed after 72 h achieving ~17% and ~22% with $1\times$ and $2\times GI_{50}$ treatments, respectively. Intriguingly, MIAPaCa-2 cells showed the lowest significant ($p\leq 0.05$) pre-G1 events: 8% after $1\times GI_{50}$, 72 h and ~13% after $2\times GI_{50}$, 24 and 72 h exposures.




Figure 4. Mean \pm SD bars showing % distribution of cell cycle phases of A549, HCT-116, HT-29, MCF-7, MDA-MB-468, MIAPaCa-2 following 24, 72 h- $1\times$ and $2\times GI_{50}$ conofolidine treatments. S-phase depletion and pre-G1 increase are the most significantly encountered changes. Cells were treated; then stained with PI and at least 20,000 events/sample were measured. Asterisks indicate significant ($p\leq 0.05$) changes compared to control. No. trials ≥ 3 ; $n=2$ per independent experiment.




Figure 5. Representative cell cycle histograms showing conofolidine-induced perturbations in A549, HCT-116, HT-29 MCF-7, MDA-MB-468 and MIAPaCa-2 cell cycles after 24 and 72 h-treatments with their respective $1\times$ and $2\times GI_{50}$. The most prominent observations are depleted S-phase with occasional G1-arrest and increased pre-G1 (<2N) events. The G1/S arrest is mainly observed in HT-29 and MCF-7 treatment groups. An exceptionally increased G2/M can also be seen in MIAPaCa-2 cells. Cells were treated; then stained with PI and at least 20,000 events/sample were measured. No. trials ≥ 3 ; $n=2$ per independent experiment.

3.4. Conofolidine induced apoptosis

Considering conofolidine's ability to suppress proliferation, evoke cytotoxicity in carcinoma cells and generate pre-G1 populations; we hypothesised that conofolidine may induce cell death via apoptosis. The percentage of apoptotic cells was measured post-conofolidine treatment by flow cytometry following annexin-V-FITC/PI staining. Figures 6 and S7 show that conofolidine induced significant ($p < 0.05$) apoptosis when compared to controls in three out of six cancer cells: MDA-MB-468, MIAPaCa-2 and HCT-116. These cells displayed significant increases in both early apoptosis (AV+/PI-) and late apoptosis (AV+/PI+). Moreover, both MDA-MB-468 and MIAPaCa-2 exhibited the highest percentages of apoptotic events (AV+) following 72 h treatment with $2\times GI_{50}$, 54 ± 5.5 and $69 \pm 1.1\%$ respectively; while HCT-116 showed $30 \pm 4.7\%$ apoptosis. The percentage of late apoptotic cells (AV+/PI+) was comparable in these cell lines; however, MDA-MB-468 and MIAPaCa-2 displayed massive early apoptosis (AV+/PI-) compared to HCT-116. Other cell lines (A549, HT-29 and MCF-7) did not show significant apoptotic events after treatment with conofolidine $2\times GI_{50}$ values ($p > 0.05$).




Figure 6. (a) and (b) Representative dot-plots and mean \pm SD bars showing annexin V/PI results of carcinoma cells treated with conofolidine for 72 h with respective $2 \times GI_{50}$. Significantly increased apoptotic cells compared to control could only be seen in MDA-MB-468, MIAPaCa-2 and HCT-116 but not in HT-29. Like HT-29; A549 and MCF-7 did not show significant apoptotic events. Samples were stained with annexin V/PI and at least 10,000 events were detected. The percentage of apoptotic events was equal to sum of cells undergoing early-apoptosis (AV⁺/PI⁻) plus late-apoptosis (AV⁺/PI⁺). Asterisk indicate statistically significant ($p \leq 0.05$) increments compared to control. No. trials ≥ 3 ; $n=2$ per independent experiment. (c) Mean \pm SD bars showing changes in caspases 3/7 activities following 24 h conofolidine ($2 \times GI_{50}$) exposure in carcinoma cell lines. Conofolidine significantly increased caspases 3/7 activities (shown as fold change relative to control) in HCT-116, MDA-MB-468, and MIAPaCa-2. However, no significant changes were seen in A549, HT-29, and decreased activities were observed in MCF-7 cells. Assays were repeated 3 times ($n=2$). Asterisks indicate a statistically significant ($p \leq 0.05$) change compared to the control.

3.5. Conofolidine induced executioner caspases and PARP cleavage

To confirm annexin V-FITC/PI results, firstly we measured caspases 3/7 activities after 24 h, $2 \times GI_{50}$ treatment. As seen in Figure 6c, conofolidine increased significantly ($p \leq 0.05$) caspases 3/7 activities by ≥ 2 -fold in HCT-116, MDA-MB-468 and MIAPaCa-2 compared to control. No significant increase in caspases' activities was observed in A549, HT-29, and MCF-7. Thereafter, we carried out a series of western blots of whole and cleaved PARP in these cells following 24 and 72 h exposures to $2 \times GI_{50}$. HCT-116 displayed cleaved PARP (89 kDa) after 24 h exposure to conofolidine while less PARP cleavage can be seen in MDA-MB-468 and MIAPaCa-2. After 72 h, these three cell lines showed

extensive PARP cleavage. In contrast, A549 and HT-29 displayed faint cleaved PARP bands only after 72 h while no PARP cleavage can be detected in the caspase 3 deficient MCF-7 cells [17](Figure 7b).




Figure 7. Representative western blot bands showing expression of whole and cleaved PARP in protein lysates of carcinoma cell lines treated with conofolidine ($2 \times GI_{50}$, 24 h and 72 h). Western blots were performed using antibodies to detect whole and cleaved PARP and housekeeping gene β -Actin. **(a)** Cleaved PARP can be seen in HCT-116 after 24 h and in MDA-MB-468 and MIAPaCa-2 cells after 72 h exposures. **(b)** Faint bands of cleaved PARP can be seen in A549 and HT-29 cells after 72 h while no cleaved PARP can be seen in MCF-7. Assays were repeated 3 times.

3.6. Conofolidine promoted morphological changes associated with senescence induction

From our observations, conofolidine consistently induced distinctive changes in HT-29 cell morphology and size relative to control. These changes included cell boundary conversion from rounded, compact to polarised with an increased size and spread relative to control (Figure 8). The evolution of such phenotype was time-dependent emerging after 24 h treatment with conofolidine at $2 \times GI_{50}$ and continued throughout 72 h exposure. After this point, cells started to detach from the supporting matrix, suggesting cell death. When morphological changes accompany S-phase depletion, this could indicate induction of either differentiation or senescence [18,19]. Accordingly, two hypotheses - differentiation or senescence induction - were postulated. Differentiation was firstly tested (Figure S8) using sodium butyrate (a known differentiation inducer) and alkaline phosphatase activity (as a differentiation marker). However, the differentiation hypothesis was rejected since conofolidine failed to significantly increase HT-29 alkaline phosphatase activity as the positive control sodium butyrate did. The senescence hypothesis was then considered; thus β -Gal-histochemical staining was conducted in HT-29 populations in addition to MCF-7 and A549. Following 72 h exposure to conofolidine $2 \times GI_{50}$, these three cell lines developed blue β -Gal staining relative to control indicating senescence-induction (Figure 9). Statistical analyses showed a significantly increased senescence-associated β -Gal positive cells ($p \leq 0.05$) by 50-75% compared to 10-20% for untreated controls in all 3 cell lines.




Figure 8. Representative images of HT-29 cells obtained by light inverted microscope showing morphological and size alterations induced by conofolidine ($2\times GI_{50}$). (a) ~65% Confluent untreated control cells, (b) 24 h Conofolidine treated cells and (c) 72 h-conofolidine treated cells at 10 \times objective magnification. Examples of cells' morphology following 72 h incubation with vehicle control (d) and $2\times GI_{50}$ conofolidine (e); 40 \times objective magnification.




Figure 9. Representative images from light inverted microscope and mean \pm SD bars showing increased % of senescence-associated β -Gal positive cells induced by conofolidine ($2\times GI_{50}$, 72 h)

compared to untreated controls HT-29, MCF-7, and A549. Images of both HT-29 and MCF-7 cells were captured at 20x while A549 images were taken at 40x objective magnifications. Mean bars \pm SD showing significant ($p \leq 0.05$) increase in the percentage of senescence-associated β -Gal positive cells relative to untreated controls in HT-29, MCF-7 and A549 cells treated with conofolidine ($2 \times GI_{50}$, 72 h). Cells were either incubated for 72 h with medium alone or with conofolidine ($2 \times GI_{50}$); then stained with β -Gal after which images were captured from at least 4 different areas in each well of treatment groups. From captured images, senescence β -Gal positive cells were counted in both treated and untreated controls and at least 1000 cells were counted per each treatment group. Assays were repeated 3 times ($n=2$). Asterisks indicate a statistically significant ($p \leq 0.05$) change compared to the control.

3.7. Conofolidine downregulated CDK2, cyclins (A2, D1) and upregulated p21

To confirm cell cycle results, we investigated associated changes in protein expression by western blotting in selected cell lines (HT-29, HCT-116 and MDA-MB-468) depending on the death pathway triggered: senescence in HT-29 and apoptosis at different time points in HCT-116 and MDA-MB-468. The modulation of the expression of CDK2, cyclin A2, cyclin D1 and p21 was investigated following 24 and 72 h $2 \times GI_{50}$ conofolidine treatments (Figure 10). Consistent with cell cycle results, conofolidine induced time-dependent decreases in CDK2 and cyclin A2 expressions relative to control; CDK2 is required for G1/S-transition, and cyclin A2 upon binding with CDK2 is essential for orderly S-phase progression. Following 24 h, CDK2 expression decreased in these cell lines to $\leq 60\%$ of the control level. Further reductions were seen after 72 h treatment reaching $\sim 20\%$ in HT-29 and MDA-MB-468 and 40% in HCT-116 of control levels. The decreases in cyclin A2 were more pronounced reaching $\leq 30\%$ and $\leq 10\%$ of control levels after 24 and 72 h exposures, respectively in these cell lines. In addition to CDK2 and cyclin A2, G1 progression is controlled by cyclin D1 which activates CDK4/6 upon cell cycle initiation. Conofolidine markedly decreased cyclin D1 in HT-29 to 16% (compared to control levels) and MDA-MB-468 to 54% after 24 h and to $\leq 33\%$ in both of these cell lines following 72 h; these results corroborated accumulated events in the G1 phase. In contrast, in HCT-116 lysates, cyclin D1 expression was not significantly changed after 24 h exposure of cells to conofolidine, but increased to $>200\%$ (2-fold) after 72 h compared to control. The p21 is a known regulator of CDK/cyclin complexes' activities. Conofolidine significantly increased p21 expression compared to control in these cell lines. MDA-MB-468 exhibited remarkably increased p21 levels (>100 -fold) after 24 and 72 h exposures. Less enhanced p21 expression was seen in both HT-29 and HCT-116: ~ 3 -fold increases following 24 h in both cell lines; 2-fold and 4-fold increments in HT-29 and HCT-116, respectively after 72 h.




Figure 10. Representative western blot showing expression of and mean \pm SD error bars from densitometric analysis of CDK2, cyclin A2, cyclin D1 and p21 in protein lysates of HT-29, HCT-116 and MDA-MB-468 cells treated with conofolidine (24 h and 72 h, $2\times GI_{50}$). Western blots were done using antibodies against CDK2, cyclin A2, cyclin D1, p21 and housekeeping gene β -Actin. Mean \pm SD error bars presented as % decrease of the control value of CDK2, cyclin- A2 and D1; and fold increase of p21 relative to control. Conofolidine significantly decreased CDK2 and cyclin A2 and increased p21 expressions. Cyclin D1 is also decreased in HT-29 and MDA-MB-468 while its expression is increased in HCT-116 cells. No changed β -Actin expression was seen. Assays were repeated 2 times. Asterisks indicate a statistically significant ($p\leq 0.05$) change compared to the control.

3.8. Confocal images confirmed the induction of aberrant mitoses by conofolidine

After confirming conofolidine's effects on cell cycle regulatory proteins, we validated mitotic abnormalities in HCT-116 cells induced by conofolidine by confocal microscopy. Acquisition of images focused on the effects of conofolidine exposures (72 h; $1\times$ and $2\times GI_{50}$) on DNA, tubulin arrangement and mitoses compared with control and vincristine (72 h; 10 nM)-treated cells (Figure 11). While untreated cells (a-d) displayed the characteristics of cellular mitosis and cytokinesis, conofolidine-treated ones exhibited aberrant mitotic phenotypes: abundant multi-nucleation, mitotic slippage, chaotic mitosis (e-g), multipolar spindles (g, h), abnormal spindle, microtubules' disarrangement (i), membrane blebbing (j), DNA fragmentation (k) and unequal DNA content (m). Compared to vincristine-treated cells (n-p), both induced multipolar spindles and microtubules' disarrangement; however, vincristine evidently decreased microtubules' density to a greater extent than conofolidine.




Figure 11. Confocal images showing HCT-116 cells treated with conofolidine compared to untreated control and vincristine-treated cells. (a-d) Controls with medium only; (e-k) conofolidine 1×GI₅₀-treated cells; (l, m) conofolidine 2×GI₅₀-treated cells; (n-p) vincristine (10 nM) treated cells. Conofolidine induced variable morphological phenotypes including multi-nucleation with chaotic multiplication (arrows 1-3), multipolar large cells (arrows 4,5), loss of polarity with asymmetric division (arrow 6), membrane blebbing and/or DNA-fragmentation (arrows 7-9) and multinucleated asymmetric cells with varying DNA content (arrow 10). Vincristine-treated HCT-116 cells showed reduced microtubules density, microtubules irregularity with multipolar spindle.

3.9. Conofolidine increased the % γ -H2AX (+) events

Since DNA fragmentation and p21 upregulation were observed following exposure of cells to conofolidine, we examined by flow cytometry whether conofolidine (72 h; 2×GI₅₀) is able to induce DNA-DSBs in HCT-116, HT-29, MDA-MB-468 and MIAPaCa-2 cells. Our results showed that conofolidine induced significant ($p \leq 0.05$) DNA-DSBs following 72 h exposure in these cell lines (Figures 12 and S9–S11). In HCT-116, 21% of DSBs were observed with conofolidine-treated cells compared to 4% and 49% for control and etoposide (72 h; 2.0 μ M)-treated cells respectively. MDA-MB-468 cells showed 19% DSBs after conofolidine exposure compared to 3% and 34% for control and etoposide-treated cells respectively. In MIAPaCa-2 cells, 15% DSBs were seen in cells exposed to conofolidine (72 h; 2×GI₅₀) compared to 3% for control and 35% for etoposide (72 h; 2.0 μ M)-treated cells. However, <10% of DSBs were observed in HT-29 cells compared to 4% for control and 39% for etoposide-treated cells respectively. Concurrent cell cycle analyses were performed to determine in which cell cycle phase the DSBs emerged; interestingly most DNA-DSBs arose during the G1 phase; MIAPaCa-2 cells revealed some DSBs in G2/M phase as well.




Figure 12. (a) Representative histograms and dot-plots from independent trial showing HCT-116 cell cycle perturbation and γ -H2AX induction after 72 h, $2 \times GI_{50}$ conofolidine treatment. The majority of DSBs were seen in G1 phase of cell cycle compared to controls and etoposide (positive control)-treated cells. (b) Mean \pm SD bars showing changes in percentage of HCT-116 cell cycle phases and increased DNA-DSBs following 72 h, $2 \times GI_{50}$ conofolidine exposure compared to untreated control. Asterisks indicate a statistically significant ($p \leq 0.05$) change compared to the control. No. trials ≥ 3 ; $n = 2$ per independent experiment. At least 15000 events were obtained/sample. (c) Confocal images from one trial of HCT-116 treated with conofolidine showing γ -H2AX foci (green fluorescence) compared to control. Etoposide-treated cells also showed γ -H2AX foci formation. Arrows show examples of nuclei containing γ -H2AX foci saturation. Less DNA content was seen in each of conofolidine-treated cells compared to control. Assays were repeated 2 times and at least three images were obtained from each treatment.

3.10. Conofolidine induced γ -H2AX foci formation

To confirm DNA-DSBs flow cytometry results, confocal microscopy was performed in HCT-116 and MDA-MB-468 cells (Figures 12 and S9). Compared to untreated control cells, conofolidine-treated cells (72 h; $2\times GI_{50}$) consistently displayed green fluorescent γ -H2AX foci within HCT-116 and MDA-MB-468 nuclei. At least one DSBs' foci saturation per image (i.e. an uncountable number of detectable foci per nucleus leading to foci overlapping) was observed especially in HCT-116 nuclei. Etoposide (72 h; 2.0 μ M)-treated cells displayed increased formation of γ -H2AX foci within their nuclei. Reduced DNA content was seen in both conofolidine- and etoposide-treated cells compared to control.

3.11. Conofolidine induced ROS generation and NQO1 expression

ROS are chemically reactive molecules that can be generated as redox cycling by-products of quinone derivatives of oxygen-rich anticancer agents e.g. doxorubicin [20]. The conophylline family members - possibly including conofolidine - are converted to their more active quinone derivatives [21], hence they could similarly induce ROS generation. Elevated ROS levels can induce and maintain senescence in cancer cells through a mechanism involving sustained cell cycle arrest [22]. We hypothesised that conofolidine increased ROS production contributing to both apoptosis and senescence induction. To test this hypothesis, we measured ROS levels (following 24 h, $2\times GI_{50}$ exposure) in A549, HCT-116, HT-29, MCF-7, MDA-MB-468 and MIAPaCa-2 (Figure 13). ROS generation was modest but significant in colon, lung and pancreatic cancer cell lines. The most significant ($p\leq 0.05$) increase was observed in A549 (2-fold), HT-29 and HCT-116 (1.5-fold) relative to control. Small but significantly elevated ROS level was also observed in MIAPaCa-2 (1.3-fold) and MCF-7 (1.2-fold) relative to control while untreated and treated MDA-MB-468 cells were ROS-negative. Elevated ROS could induce cellular defense enzymes NQO1 which shows marked inducibility during oxidative stress [23]. Therefore, its expression was validated by western blot following conofolidine exposure. As expected, NQO1 expression was increased after 24 h in ROS-positive cells: HT-29, HCT-116, A549, MCF-7 and MIAPaCa-2. Similarly, after 72 h, NQO1 expression continued to increase suggesting that conofolidine efficiently maintained prolonged oxidative stress in these cells. Intriguingly, neither constitutive, nor induced NQO1 expression was seen in the MDA-MB-468 cells.




Figure 13. (a) Mean \pm SD error bars showing fold change relative to control of ROS level in carcinoma cells following conofolidine treatment ($2 \times GI_{50}$, 24 h). Significantly increased ROS is seen in A549, HCT 116, HT-29, MIAPaCa-2; MCF-7 showed slight increment while no significant change is observed in MDA-MB-468. Assays were repeated 3 times ($n=2$). Asterisks indicate a statistically significant ($p \leq 0.05$) change compared to the control. **(b)** Representative western blot showing expression of NQO1 in protein lysates of A549, HCT 116, HT-29, MCF-7, MDA-MB-468 and MIAPaCa-2 cells treated with conofolidine (24 h and 72 h, $2 \times GI_{50}$). Western blotting was performed using antibodies against NQO1 and housekeeping gene β -Actin. All cell lines (except MDA-MB-468) showed increased NQO1 expression following conofolidine treatment. No changed β -Actin expression was seen. Assays were repeated 2 times.

4. Discussion

Conofolidine, a novel bisindole alkaloid from *Tabernaemontana corymbosa* proved its broad-spectrum potency against carcinoma cells derived from breast (MCF-7, MDA-MB-468, MDA-MB-231, SK-BR3, T-47D, ZR-75-B), colon (HCT-116, HT-29, Caco-2), lung (A549), pancreas (MIAPaCa-2, PANC-1) and B-cell lymphoma (DoHH2, Vallois) cell lines (Table 1, Table S1). The breast carcinoma cells were predominantly the most sensitive to conofolidine. Compared to MRC-5, all carcinoma cell lines exhibited higher sensitivity $GI_{50} \leq 0.5 \mu M$. Although MCF10A - the proliferative benign fibrocystic epithelial breast - cell line showed $GI_{50} \sim 0.23 \mu M$, MCF-7, MDA-MB-231, T-47D, ZR-75-B, DoHH2 and Vallois cells yielded GI_{50} values $\leq 0.13 \mu M$. Indeed, the potency of conofolidine surpassed structurally-related conophylline (Table 1) against A549, MDA-MB-468 and MIAPaCa-2 cell lines. This could be a consequence of the structural variation: the second indole moiety *viz.* deoxoapodine in conofolidine instead of pachysiphine in conophylline (Figure 1). In addition to its antiproliferative actions, conofolidine efficiently suppressed the clonogenic potential of cancer cells (Figure 3). For example, MDA-MB-468, MIAPaCa-2 colony formation was significantly inhibited $>95\%$ by $1 \mu M$ conofolidine. Individual cells were unable to survive 24 h conofolidine challenge and form progeny colonies. Dose-dependent inhibition of A549 and HT-29 colony-forming ability was seen. Together, MTT and clonogenic assays demonstrated conofolidine's potential to compromise cancer cell proliferation and viability.

To explore conofolidine's cellular mechanisms of action, cell cycle analysis (using flow cytometry) was considered (Figures 4 and 5). Four distinctive perturbations (specifically at $2\times GI_{50}$) were observed: stark S-phase depletion, G1-phase arrest, G2/M arrest (seen only in MIAPaCa-2) and the emergence of sub-diploid population ($<2N$ -DNA) appearing as pre-G1 events. Following 24 h, conofolidine induced radical S-phase depletion. For example, $\leq 6\%$ HCT-116, HT-29, MCF-7, MDA-MB-468 cell population resided within S-phase using $1\times GI_{50}$ compared to 20-27% for untreated controls. Moreover, a concomitant increase of G1-populations started to emerge. After 72 h, conofolidine sustained statically significant G1/S-arrest, especially in HT-29 and MCF-7 cells. In contrast, A549 (less sensitive as indicated in previous assays) seemed to require a higher concentration of $2\times GI_{50}$ to produce profoundly equivalent depleted S-phase but without an obvious effect on G1 events. G1-arrest could be a consequence of inhibition of proteins' transcription and expression which take place during G1 and are required for DNA replication, leading ultimately to decreased numbers of cells escaping G1-block into S-phase. Similarly, structurally-related conophylline was reported to induce G1-arrest in endometrial cancer cells [24]. Thus, structurally-related conofolidine and conophylline may share similar anticancer mechanism(s) of action. However, conofolidine was able to decrease A549 S-phase population without affecting % of G1-cells suggesting that it may interfere directly with DNA replication machinery such as replicative helicase similar to anthracyclines [25]. In support of this suggestion, S-phase depletion was attained with minimal effect on G1 in MIAPaCa-2 cells which interestingly accompanied a G2/M arrest. These observations indicate putative anti-microtubule activity and resemble the effects of microtubule disrupting agents such as vinca bisindole alkaloids [26] and jerantinines A and B which share indole moiety structural similarities and are also isolated from *Tabernaemontana corymbosa* [8,9]. Concentration- and time-dependent increased pre-G1 events were evident in most cell lines indicating cell death by apoptosis. For instance, $\geq 20\%$ of HCT-116 cells accumulated at pre-G1 following 24 h exposure to $1\times GI_{50}$ which continued to increase to $\geq 30\%$ after 72 h. The depleted S-phase (with changed HT-29 morphology) led to a hypothesis that conofolidine might induce differentiation. Indeed, conophylline is known to differentiate pancreatic- β endocrine cells [27,28]. However, conofolidine, in contrast to the positive control (sodium butyrate), failed to impact HT-29 differentiation (Figure S8).

Considering the sub-diploid events seen in the cell cycle, an apoptotic mode of cell death was the first suggested death pathway to be investigated (Figures 6, 7 and S7). Annexin V-FITC/PI fluorescent staining and flow cytometric analyses of populations were undertaken, which distinguishes between early and late apoptosis. In early apoptosis, cell membrane asymmetry is lost exposing phosphatidylserine to the outer membrane surface. Thus annexin-V can bind and be measured by flow cytometry. PI can intercalate DNA and be quantified only upon the commencement of late apoptosis when membrane integrity deteriorates [29]. Significantly increased apoptotic events were only observed in MDA-MB-468, MIAPaCa-2, and HCT-116 populations. For instance, $>50\%$ of MIAPaCa-2 and MDA-MB-468 were AV+; AV+ events evolve in early apoptosis and are accompanied by PI+ events in late apoptosis when DNA becomes fragmented [30]. Indeed, pre-G1 populations are an indication of DNA cleavage into oligonucleosomal fragments [31]. DNA fragmentation is induced by caspase-activated DNase [4], thus caspases activities and emergence of caspase-mediated cleaved PARP were validated. Indeed, MDA-MB-468, MIAPaCa-2 and HCT-116 showed increased active caspases 3 and 7, which generated the 89 kDa-PARP fragment corroborating apoptosis. The timing of apoptosis-onset was different between these cells according to annexin-V/PI results which were validated by the increased caspase activity and appearance of cleaved PARP at different time points. Intriguingly, A549 cells which showed significant pre-G1 events, did not display significant AV+ events. These results are consistent with other studies that describe: i) the known apoptosis-reluctant phenotype of A549 cells [32]; and ii) the potential viability of pre-G1 A549 cells [33]. Indeed, conofolidine did not arrest the G1 but decreased S- and G2/M phases thereby delaying mitosis. Furthermore, these cells might be able to divide thereby producing two or three cells (tripolar mitosis) possessing sub-diploid DNA content - observed as pre-G1 events. Thus, conofolidine might cause postmitotic arrest in the second and even third cell cycles. Unsurprisingly,

the caspase 3 deficient MCF-7 cells did not show any significant AV+ events. This is likely because these cells were reported to have a base pair deletion of caspase 3 gene which ultimately abrogates mRNA translation [34]. Hence, no apparent cleaved PARP was observed in MCF-7 treated with conofolidine.

Inspired by conofolidine's potency, G1/S arrest and altered HT-29 cell morphology (Figure 8) without evident apoptosis or differentiation, we proposed senescence as an alternative pathway. Indeed, increased β -Gal positive HT-29, MCF-7 and A549 cells post-conofolidine exposure supported this proposal (Figure 9). For example, $\geq 60\%$ of HT-29 and MCF-7 cells were senescent compared to 15-20% of untreated cells; morphologically, these senescent cells appeared enlarged, flattened and prolonged. Senescent cells cease to divide - consistent with depleted S-phase, yet they remain metabolically active, staining positive for β -galactosidase, a lysosomal protein and putative senescence biomarker [3]. Cellular senescence may be, in principle, considered a cancer-preventive mechanism, especially in somatic cells that have undergone many divisions and accumulated multiple genetic mutations. By conversion to an immunogenic senescent phenotype, the cell can be eliminated *via* the immune system [35]. Senescence may also be considered a desired therapeutic outcome in cancer cells able to evade apoptosis, through for example p53 loss/mutation occurring in $>50\%$ human cancers [36] and exemplified by HT-29 cells. Controversially, persistent senescence may contribute to tumourigenesis [37] through acquisition of a senescence-associated secretory phenotype (SASP) that may promote tumour progression. However, corroborative evidence demonstrated herein reveals that the ultimate fate of conofolidine-induced senescence is death.

Interference with crucial cell cycle regulators - CDK2, cyclins A2, D1 and p21 - is expected with G1/S arrest inducers. Conofolidine affected the expression of these controllers in HT-29, HCT-116 and MDA-MB-468 cells (Figure 10). Downregulation of CDK2 and both cyclin A2 and D1 was observed; as a consequence, these cells were not able to progress G1 through S-phase. It may be postulated that there was i) less cyclin D1 available for CDK4/6 activation to phosphorylate the retinoblastoma protein (pRb) thereby triggering G1 initiation [38]; ii) decreased CDK2 expression required for both G1/S transition - upon binding to cyclin E2 - *via* phosphorylating both histone H1 and pRb [39], and S-phase completion - upon binding to cyclin A2 - *via* phosphorylating the replication machinery components e.g., DNA polymerase [40]. Diminished cyclin A2, which can activate CDK1 in addition to CDK2, leads to enhanced deterioration of S-phase progression through perturbation of DNA replication origin [41], and decreased G2/M transition [42] *via* disturbing nuclear envelope breakdown, nuclear accumulation of cyclin B1 [43] and anchoring of the mitotic spindle [44]. Thus, the observed G2 arrest accompanying S-phase depletion in MIA PaCa-2 cells may be explained. Our results demonstrating cyclin D1 downregulation in HT-29 and MDA-MB-468 are similar to previously described cyclin D1 suppression by conophylline in carcinoma cells [24]. Conversely, cyclin D1 in HCT-116 was exceptionally increased possibly because cyclin D1 is available in two isoforms (a and b); cyclin D1b upregulation induces G1 arrest with apoptosis in carcinoma cells [45]. Alternatively, HCT-116 cells might replicate their DNA without completing mitosis and/or cytokinesis, and re-enter G1 with tetraploid 4N DNA - driven by the increased cyclin D1- i.e. mitotic slippage. Indeed, confocal microscopy visually confirmed such aberrant division seen as frequent multi-nucleated cells (Figure 11). Upregulated p21 inhibits the kinase activity of CDK2 and CDK2/cyclin complexes [46,47]. In addition, p21 can suppress the proliferating cell nuclear antigen (PCNA) required for DNA replication thereby blocking S-phase progression [48]. Upregulation of p21 accompanied senescence-induction in HT-29 cells. Indeed, p21 mediates cellular senescence *via* p53-dependent and -independent pathways [49]. However, one could speculate whether p21 overexpression could contribute to apoptosis especially evident in MDA-MB-468. p21 may disrupt transcription factors e.g. STAT3 [50] which could inhibit the expression of antiapoptotic proteins as some authors reported [51]. Interestingly, increased p21 expression following conofolidine treatment seems to be p53-independent; this is because: i) MDA-MB-468 cells, which displayed >100 -fold p21 increase, harbour a gain of function (GOF)-R273H-mutation in their p53 gene; ii) HCT-116 cells which express WT-p53 exhibited only ~ 3 -fold p21 increase. However, recent studies have shown that p21 can be induced by ROS-mediated DNA-damage even when p53 function is compromised [52].

Confocal imaging of HCT-116 cells treated with conofolidine showed several aberrant mitotic phenotypes support our findings (Figure 11). Multinucleation and mitotic slippage (consistent with cyclin D1 upregulation); decreased DNA content, microtubules' disarrangement and changed polarity (consistent with cyclin A2 and CDK2 downregulation). Furthermore, DNA fragmentation and plasma membrane disintegration and/or blebbing confirmed apoptotic cells (as indicated in annexin V/FITC and cell cycle analyses). The multinucleated cells either succumbed directly to apoptosis, or continued another round of division. In support of the second fate, some cells exhibited aberrant, chaotic asymmetric multiplication with spindle disarrangement. Another form of mitotic abnormality appeared as multipolar giant cells with enlarged nuclei, which might be due to failure of nuclear envelope breakdown. The accumulation of DNA fragments might trigger genotoxic stress that leads to cell death. In comparison with the vincristine - a microtubule-targeting agent, both compounds disturbed microtubules' and mitotic spindle machinery. However, vincristine decreased microtubules' density more than conofolidine. DNA fragmentation is a likely consequence of lethal DNA-damage. Indeed, conofolidine caused DNA-DSBs as seen by intense γ -H2AX foci and increased percentage of γ -H2AX+ events with the majority of events emerging in G1- in addition to G2/M-phases of the cell cycle (Figures 12 and S9-S11). The histone protein H2AX is a robust reporter of unrepaired DNA-DSBs when phosphorylated to γ -H2AX at nascent DSB sites. Accumulation of γ -H2AX molecules in the chromatin around DSB sites creates foci at which DNA repair and chromatin remodeling proteins are recruited [53]. Confocal images revealed an increased number of DNA-DSBs foci with foci saturation was frequently seen in HCT-116 cells. In MDA-MB-468 cells, the DNA-damage detected likely precedes exacerbated conofolidine-induced apoptosis. Interestingly, these cells lack a functional pRB resulting in an attenuated response to DNA-damage [54]. Furthermore, conofolidine, by downregulating CDK2 and cyclin A2, halts the functions of both CDK2/cyclin A2 and also CDK2/cyclin A1 complexes. Considering the role of A-type cyclins in activating DNA-DSB repair mechanisms in collaboration with CDK2 and Ku DNA repair proteins [55], conofolidine rendered carcinoma cells more prone to DNA-damage leading to aggravated cytotoxicity and apoptosis. Compared to etoposide, a topoisomerase II inhibitor, conofolidine induced significant DSBs at lower concentrations, especially in MDA-MB-468 and MIAPaCa-2 cells: $\sim 0.6 \mu\text{M}$ conofolidine induced $\sim 20\%$ DSBs compared to $\sim 35\%$ for $2.0 \mu\text{M}$ etoposide. Thus, conofolidine demonstrated its potency as a DNA damaging agent comparable to etoposide.

We showed that conofolidine was capable of inducing ROS (H_2O_2) in our cancer cell line panel - with the exception of MDA-MB-468 (Figure 13). To cope with oxidative stress, cells increased NQO1 expression, an inducible quinone-detoxifying enzyme. In this regard, all ROS-positive cells exhibited NQO1 upregulation. It is noticeable that NQO1 expression is increased in these cells similarly although there is variation in their induced H_2O_2 level. One could speculate that conofolidine might induce other species of ROS e.g. hydroxyl radicals ($\text{OH}\cdot$) which may trigger NQO1 expression. Oxidative stress, by causing unrepairable DNA-damage, led to p21 upregulation and cell cycle arrest. Thus, ROS may act as mutagens and contribute to the induction and maintenance of cell senescence [56]. Interestingly, some studies reported that NQO1 stabilises p53 or causes p53 accumulation independent of MDM2, provoking senescence and apoptosis [57,58]. ROS are known triggers of apoptosis for plant-derived anticancer agents e.g. vinca alkaloids [59]. In the case of conofolidine, the oxidative stress elicited apoptosis in HCT-116 and MIAPaCa-2 cells through i) altering mitochondrial membrane permeability, cytochrome C release, caspase 3/7 activation and PARP-cleavage; ii) inducing DNA-damage which is aggravated by conofolidine's blockade of its repair mechanism *via* CDK2/cyclin A complex inhibition. Absence of NQO1 in MDA-MB-468 cells is consistent with other studies which failed to detect NQO1 expression in these cells [60,61]. Such absence might enhance conofolidine's anticancer action: conofolidine (like other conophylline family members) is converted to its quinone metabolite [14]; accordingly, the absence of the quinone-detoxifying NQO1 delayed its metabolic inactivation which enhanced its anticancer actions.

5. Conclusions

Taken together, we describe in this work for the first time the characterisation of anticancer activities of conofolidine - a novel bisindole alkaloid isolated from the plant *Tabernaemontana corymbosa*. Conofolidine potently suppresses (broad-spectrum) cancer cell growth and colony formation *in vitro*, inducing either senescence or apoptosis. To our knowledge, this is the first work investigating molecular mechanisms of conofolidine's antitumour action. Further characterisation of this promising anticancer candidate is justified.

Supplementary Materials: The following supporting information can be downloaded at the website of this paper posted on Preprints.org, Figure S1: Conophylline MTT; Figure S2: Leucophyllidine and bipleiophylline MTT, Figure S3: Comparison of conofolidine, conophylline, leucophyllidine and bipleiophylline GI₅₀s, Figure S4: DMSO MTT, Figure S5: Conofolidine MTT additional cell lines, Figure S6: Clonogenic photos, Figure S7: Annexin V/PI of A549 and MCF-7, Figure S8: HT-29 differentiation study, Figure S9: MDA-MB-468 γ -H2AX flow cytometry and confocal studies, Figure S10 and S11: HT-29 and MIAPaCa-2 γ -H2AX flow cytometry studies; Table S1: Conofolidine MTT additional cell lines.

Author Contributions: Conceptualisation M.A.H., C.E.N., K.H.L. and T.S.K.; data curation, investigation, validation, methodology, software, formal analysis, writing—original draft preparation, M.A.H.; resources, T.D.B. and T.S.K.; project administration, T.D.B.; writing—review and editing, K.H.L., T.S.K. and T.D.B.; supervision, T.D.B. and H.C.; funding acquisition, M.A.H. All authors have read and agreed to the published version of the manuscript.

Funding: The authors acknowledge the financial support from the Higher Committee of Education Development in Iraq (Ref. no. D-10-2013).

Data Availability Statement: The data presented in this study are available in the article and Supplementary Materials.

Acknowledgments: The authors of this work gratefully thank respectable lab colleagues at Division of Medicinal Chemistry and Structural Biology—Biodiscovery Institute and Flow Cytometry Facility—QMC, University of Nottingham, UK, for their technical supports and advices. Special acknowledgment should go to Prof Kam and his colleagues at Department of Chemistry, Faculty of Science, University of Malaya for generously providing conofolidine, conophylline, leucophyllidine and bipleiophylline.

Conflicts of Interest: The authors of this work declare that there is no conflict of interest.

References

1. da Rocha, A.B., R.M. Lopes, and G. Schwartzmann, *Natural products in anticancer therapy*. *Curr Opin Pharmacol*, 2001. **1**(4): p. 364-9.
2. Mann, J., *Natural products in cancer chemotherapy: past, present and future*. *Nat Rev Cancer*, 2002. **2**(2): p. 143-8.
3. Kim, R., M. Emi, and K. Tanabe, *The role of apoptosis in cancer cell survival and therapeutic outcome*. *Cancer Biology & Therapy*, 2006. **5**(11): p. 1429-1442.
4. Baig, S., et al., *Potential of apoptotic pathway-targeted cancer therapeutic research: Where do we stand?* *Cell death & disease*, 2016. **7**(1): p. e2058-e2058.
5. Hanahan, D. and R.A. Weinberg, *Hallmarks of cancer: the next generation*. *Cell*, 2011. **144**(5): p. 646-74.
6. Lee, M. and J.S. Lee, *Exploiting tumor cell senescence in anticancer therapy*. *BMB Rep*, 2014. **47**(2): p. 51-9.
7. Milczarek, M., *The Premature Senescence in Breast Cancer Treatment Strategy*. *Cancers (Basel)*, 2020. **12**(7).
8. Raja, V.J., et al., *Novel antitumour indole alkaloid, Jerantinine A, evokes potent G2/M cell cycle arrest targeting microtubules*. *Invest New Drugs*, 2014. **32**(5): p. 838-50.
9. Qazzaz, M.E., et al., *In vitro anticancer properties and biological evaluation of novel natural alkaloid jerantinine B*. *Cancer Lett*, 2016. **370**(2): p. 185-97.
10. Kam, T.-S., et al., *New alkaloids from the leaves of Tabernaemontana divaricata*. *Tetrahedron Letters*, 1992. **33**(7): p. 969-972.
11. Kam, T.-S., K.-Y. Loh, and C. Wei, *Conophylline and Conophyllidine: New Dimeric Alkaloids from Tabernaemontana divaricata*. *Journal of Natural Products*, 1993. **56**(11): p. 1865-1871.
12. Gan, C.-Y., et al., *Leucophyllidine, a Cytotoxic Bisindole Alkaloid Constituted From the Union of an Eburnan and a New Vinylquinoline Alkaloid*. *Organic Letters*, 2009. **11**(17): p. 3962-3965.
13. Kam, T.-S., et al., *Bipleiophylline, an Unprecedented Cytotoxic Bisindole Alkaloid Constituted from the Bridging of Two Indole Moieties by an Aromatic Spacer Unit*. *Organic Letters*, 2008. **10**(17): p. 3749-3752.

14. Nge, C.E., et al., *A Hexacyclic, Iboga-Derived Monoterpenoid Indole with a Contracted Tetrahydroazepine C-Ring and Incorporation of an Isoxazolidine Moiety, a Seco-Corynanthean, an Aspidosperma-Aspidosperma Bisindole with Anticancer Properties, and the Absolute Configuration of the Pyridopyrimidine Indole Alkaloid, Vernavosine*. J Nat Prod, 2016. **79**(10): p. 2709-2717.
15. Nicoletti, I., et al., *A rapid and simple method for measuring thymocyte apoptosis by propidium iodide staining and flow cytometry*. J Immunol Methods, 1991. **139**(2): p. 271-9.
16. Bradford, M.M., *A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding*. Analytical Biochemistry, 1976. **72**(1): p. 248-254.
17. Jänicke, R.U., *MCF-7 breast carcinoma cells do not express caspase-3*. Breast Cancer Research and Treatment, 2009. **117**(1): p. 219-221.
18. Liu, X., et al., *Induction of cell cycle arrest at G1 and S phases and cAMP-dependent differentiation in C6 glioma by low concentration of cycloheximide*. BMC Cancer, 2010. **10**(1): p. 684.
19. Maya-Mendoza, A., et al., *Immortalised breast epithelia survive prolonged DNA replication stress and return to cycle from a senescent-like state*. Cell death & disease, 2014. **5**(7): p. e1351-e1351.
20. Conklin, K.A., *Chemotherapy-associated oxidative stress: impact on chemotherapeutic effectiveness*. Integr Cancer Ther, 2004. **3**(4): p. 294-300.
21. Kam, T.S., H.S. Pang, and T.M. Lim, *Biologically active indole and bisindole alkaloids from Tabernaemontana divaricata*. Org Biomol Chem, 2003. **1**(8): p. 1292-7.
22. Kuilman, T., et al., *The essence of senescence*. Genes Dev, 2010. **24**(22): p. 2463-79.
23. Ross, D. and D. Siegel, *The diverse functionality of NQO1 and its roles in redox control*. Redox Biology, 2021. **41**: p. 101950.
24. Irie, T., et al., *Inhibition of attachment and chemotactic invasion of uterine endometrial cancer cells by a new vinca alkaloid, conophylline*. Anticancer Res, 1999. **19**(4B): p. 3061-6.
25. Bachur, N.R., et al., *Helicase inhibition by anthracycline anticancer agents*. Mol Pharmacol, 1992. **41**(6): p. 993-8.
26. Jordan, M.A., *Mechanism of action of antitumor drugs that interact with microtubules and tubulin*. Curr Med Chem Anticancer Agents, 2002. **2**(1): p. 1-17.
27. Kawakami, M., et al., *Promotion of beta-cell differentiation by the alkaloid conophylline in porcine pancreatic endocrine cells*. Biomed Pharmacother, 2010. **64**(3): p. 226-31.
28. Kojima, I. and K. Umezawa, *Conophylline: a novel differentiation inducer for pancreatic beta cells*. Int J Biochem Cell Biol, 2006. **38**(5-6): p. 923-30.
29. Giard, D.J., et al., *In vitro cultivation of human tumors: establishment of cell lines derived from a series of solid tumors*. J Natl Cancer Inst, 1973. **51**(5): p. 1417-23.
30. Bacso, Z., R.B. Everson, and J.F. Eliason, *The DNA of annexin V-binding apoptotic cells is highly fragmented*. Cancer Res, 2000. **60**(16): p. 4623-8.
31. Enari, M., et al., *A caspase-activated DNase that degrades DNA during apoptosis, and its inhibitor ICAD*. Nature, 1998. **391**: p. 43.
32. Demidenko, Z.N. and M.V. Blagosklonny, *Flavopiridol induces p53 via initial inhibition of Mdm2 and p21 and, independently of p53, sensitizes apoptosis-reluctant cells to tumor necrosis factor*. Cancer Res, 2004. **64**(10): p. 3653-60.
33. Demidenko, Z.N., et al., *Mechanism of G1-like arrest by low concentrations of paclitaxel: next cell cycle p53-dependent arrest with sub G1 DNA content mediated by prolonged mitosis*. Oncogene, 2008. **27**(32): p. 4402-4410.
34. Jänicke, R.U., et al., *Caspase-3 Is Required for DNA Fragmentation and Morphological Changes Associated with Apoptosis**. Journal of Biological Chemistry, 1998. **273**(16): p. 9357-9360.
35. Burton, D.G. and R.G. Faragher, *Cellular senescence: from growth arrest to immunogenic conversion*. Age (Dordr), 2015. **37**(2): p. 27.
36. Ozaki, T. and A. Nakagawara, *Role of p53 in Cell Death and Human Cancers*. Cancers, 2011. **3**(1): p. 994-1013.
37. Schmitt, C.A., B. Wang, and M. Demaria, *Senescence and cancer - role and therapeutic opportunities*. Nat Rev Clin Oncol, 2022. **19**(10): p. 619-636.
38. DeGregori, J., *The Rb network*. J Cell Sci, 2004. **117**(Pt 16): p. 3411-3.
39. Lauper, N., et al., *Cyclin E2: a novel CDK2 partner in the late G1 and S phases of the mammalian cell cycle*. Oncogene, 1998. **17**(20): p. 2637-43.
40. Woo, R.A. and R.Y. Poon, *Cyclin-dependent kinases and S phase control in mammalian cells*. Cell Cycle, 2003. **2**(4): p. 316-24.
41. Katsuno, Y., et al., *Cyclin A-Cdk1 regulates the origin firing program in mammalian cells*. Proceedings of the National Academy of Sciences, 2009. **106**(9): p. 3184-3189.
42. *Chapter 43 - G2 Phase, Responses to DNA Damage, and Control of Entry Into Mitosis*, in *Cell Biology (Third Edition)*, T.D. Pollard, et al., Editors. 2017, Elsevier. p. 743-754.
43. Gong, D., et al., *Cyclin A2 regulates nuclear-envelope breakdown and the nuclear accumulation of cyclin B1*. Curr Biol, 2007. **17**(1): p. 85-91.
44. Bendris, N., et al., *Cyclin A2, Rho GTPases and EMT*. Small GTPases, 2012. **3**(4): p. 225-8.

45. Wang, N., et al., *Cyclin D1b overexpression inhibits cell proliferation and induces cell apoptosis in cervical cancer cells in vitro and in vivo*. *Int J Clin Exp Pathol*, 2014. **7**(7): p. 4016-23.
46. Poon, R.Y., et al., *Cyclin-dependent kinases are inactivated by a combination of p21 and Thr-14/Tyr-15 phosphorylation after UV-induced DNA damage*. *J Biol Chem*, 1996. **271**(22): p. 13283-91.
47. Harper, J.W., et al., *Inhibition of cyclin-dependent kinases by p21*. *Molecular biology of the cell*, 1995. **6**(4): p. 387-400.
48. Funk, J.O., et al., *Inhibition of CDK activity and PCNA-dependent DNA replication by p21 is blocked by interaction with the HPV-16 E7 oncoprotein*. *Genes Dev*, 1997. **11**(16): p. 2090-100.
49. Georgakilas, A.G., O.A. Martin, and W.M. Bonner, *p21: A Two-Faced Genome Guardian*. *Trends Mol Med*, 2017. **23**(4): p. 310-319.
50. Coqueret, O. and H. Gascan, *Functional interaction of STAT3 transcription factor with the cell cycle inhibitor p21WAF1/CIP1/SDI1*. *J Biol Chem*, 2000. **275**(25): p. 18794-800.
51. Al Zaid Siddiquee, K. and J. Turkson, *STAT3 as a target for inducing apoptosis in solid and hematological tumors*. *Cell Res*, 2008. **18**(2): p. 254-67.
52. Masgras, I., et al., *Reactive oxygen species and mitochondrial sensitivity to oxidative stress determine induction of cancer cell death by p21*. *J Biol Chem*, 2012. **287**(13): p. 9845-9854.
53. Bonner, W.M., et al., *GammaH2AX and cancer*. *Nature reviews. Cancer*, 2008. **8**(12): p. 957-967.
54. Billecke, C.A., et al., *Lack of functional pRb results in attenuated recovery of mRNA synthesis and increased apoptosis following UV radiation in human breast cancer cells*. *Oncogene*, 2002. **21**(29): p. 4481-4489.
55. Müller-Tidow, C., et al., *The Cyclin A1-CDK2 Complex Regulates DNA Double-Strand Break Repair*. *Molecular and Cellular Biology*, 2004. **24**(20): p. 8917-8928.
56. Davalli, P., et al., *ROS, Cell Senescence, and Novel Molecular Mechanisms in Aging and Age-Related Diseases*. *Oxid Med Cell Longev*, 2016. **2016**: p. 3565127.
57. Liu, K., et al., *NQO1 Stabilizes p53 in Response to Oncogene-Induced Senescence*. *International journal of biological sciences*, 2015. **11**(7): p. 762-771.
58. Asher, G., et al., *NQO1 stabilizes p53 through a distinct pathway*. *Proceedings of the National Academy of Sciences*, 2002. **99**(5): p. 3099-3104.
59. Chiu, W.H., et al., *Vinca alkaloids cause aberrant ROS-mediated JNK activation, Mcl-1 downregulation, DNA damage, mitochondrial dysfunction, and apoptosis in lung adenocarcinoma cells*. *Biochem Pharmacol*, 2012. **83**(9): p. 1159-71.
60. Sutton, K.M., C.D. Doucette, and D.W. Hoskin, *NADPH quinone oxidoreductase 1 mediates breast cancer cell resistance to thymoquinone-induced apoptosis*. *Biochem Biophys Res Commun*, 2012. **426**(3): p. 421-6.
61. Pink, J.J., et al., *NAD(P)H:Quinone oxidoreductase activity is the principal determinant of beta-lapachone cytotoxicity*. *J Biol Chem*, 2000. **275**(8): p. 5416-24.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.