

Table 2: ITU characteristics of patients hospitalised with COVID-19 in English regions.

	London	North-East	North-West	Yorkshire	East Midlands	West Midlands	East of England	South-East	South-West	Total
Admissions to ITU [†]	13505 (10.3)	2010 (5.2)	7944 (6.5)	3966 (5.4)	3884 (5.8)	6200 (6.8)	5611 (7.2)	6719 (6.3)	2517 (6.0)	52,356 (7)
Number of ITU admissions [‡]	1.2 ± 0.5	1.1 ± 0.3	1.1 ± 0.3	1.1 ± 0.5	1.2 ± 0.6	1.1 ± 0.4	1.1 ± 0.4	1.1 ± 0.5	1.1 ± 0.4	1.1 ± 0.5
Number of organ support required [‡]	2.6 ± 1.8	2.4 ± 1.4	2.2 ± 1.3	2.4 ± 1.5	2.4 ± 1.6	2.7 ± 1.6	2.2 ± 1.4	2.3 ± 1.6	2.2 ± 1.7	2.4 ± 1.6
ITU length of stay (days) [‡]	10.7 ± 15.4	6.5 ± 10.3	7.6 ± 12.5	7.1 ± 11.8	8.2 ± 12.1	9.3 ± 11.3	7.4 ± 12.8	9.1 ± 13.4	7.3 ± 12.3	8.7 ± 13.2
Kidney support days [‡]	2.7 ± 7.9	1.4 ± 4.6	1.5 ± 4.9	1.5 ± 4.7	1.5 ± 5.1	1.9 ± 5.3	1.6 ± 5.7	1.8 ± 5.6	1.7 ± 5.7	1.9 ± 6.1
ITU mortality [†]	3747 (33.4)	619 (31.0)	1217 (39.5)	1420 (41.7)	684 (39.9)	1794 (37.7)	1542 (34.4)	1569 (32.2)	505 (25.0)	13,097 (34.9)

[†] Number (%), [‡] mean 2 standard deviations.