

Title: Mental health and violence against women in Afghanistan, India, and Sri Lanka: a situation analysis

Meaghen Quinlan-Davidson^{1 2 3}, Ayesha Ahmad⁴, Laura Asher⁵, Urvita Bhatia^{6 7}, Nayreen Daruwalla, Delan Devakumar³, Abhijit Nadkarni^{6 9}, Alexis Palfreyman³, Lamba Saboor¹⁰, T.H.Rasika Samanmalee¹¹, David Osrin

¹ The Margaret and Wallace McCain Centre for Child, Youth & Family Mental Health, Centre for Addiction and Mental Health, Toronto, Canada

² Dalla Lana School of Public Health, University of Toronto, Canada

³ Institute for Global Health, University College London, UK

⁴ St George's University of London, UK

⁵ Academic Unit of Lifespan and Population Health, School of Medicine, University of Nottingham, UK

⁶ Addictions Research Group, Sangath, Goa, India

⁷ Department of Psychology, Health and Professional Development, Oxford Brookes University, UK

⁸ Society for Nutrition, Education and Health Action, India

⁹ Centre for Global Mental Health, Department of Population Health, London School of Hygiene & Tropical Medicine, UK

¹⁰ Independent Consultant, Afghanistan

¹¹ Department of Psychiatry, Faculty of Medicine at the University of Colombo in Sri Lanka

Corresponding Author:
Meaghen Quinlan-Davidson
meaghen.quinlan@camh.ca
(416) 535-8501

Keywords. Violence against women; domestic violence; human trafficking; Afghanistan; India; Sri Lanka

DISCLOSURE OF COMPETING INTERESTS. None of the authors declares a conflict of interest.

ABSTRACT

Background. Globally, 10-53% of ever-partnered women have experienced physical or sexual intimate partner violence over their lifetime. Women survivors of violence are at high risk of poor mental health. In this study, we investigate women's exposure to violence and mental health conditions in Afghanistan, India, and Sri Lanka, while considering the policy and service contexts.

Methods. A situation analysis tool was developed for the study. We extracted information from grey and peer-reviewed literature and other publicly available data investigating the prevalence of violence against women and mental health conditions, policies addressing violence against women and mental health conditions in each country, and the services available to women exposed to violence and women with mental health conditions.

Results. Forty-six percent of women in Afghanistan, 21% of women in India and 5% of women in Sri Lanka reported experiencing physical violence within the last 12 months of the most recent survey. Meanwhile, 7% of ever-partnered women in Afghanistan, 6% of women in India and 7% of women in Sri Lanka reported experiencing sexual violence during their lifetime. In India, 6.9% of disability adjusted life years were attributed to childhood sexual abuse and 4.6% to intimate partner violence. In Sri Lanka, 14.6% of women exposed to physical or sexual violence by a partner had engaged in self-harm. We found no data on conflict-related sexual violence and trafficking. All three countries have made commitments to gender equality or preventing violence against women. Implementation of some of these policies, however, is unclear. The countries also have had mental health policies and services, but there is currently little intersection between mental health and violence against women.

Limitations. The situation analysis is limited by the data available and the generalizability of findings.

Conclusion. The three countries have limited data, policies, and legislation on the intersection between all forms of violence against women and poor mental health, as well as a paucity of mental health service provision.

Future Work. Future research should focus on integrating mental healthcare within social services; translating trauma-informed approaches into service provision; and addressing family violence, within VAW.

Study Registration. Not applicable

Funding Details. This research was funded by the National Institute for Health Care Research (NIHR) Global Health Research programme (17/63/47) using UK aid from the UK Government to support global health research. The views expressed in this publication are those of the authors and not necessarily those of the NIHR or the UK government.

Contents List.

Introduction	4
Methods	6
Results	8
Discussion	17
Appendix	26

List of Tables.

Table 1. Types of Violence Against Women in Sri Lanka, India, and Afghanistan	9-10
Table 2. Indicators of Poor Mental Health and Treatment in Sri Lanka, India and Afghanistan	12
Table 3. Policies, Legislation, Strategies, Plans and Programmes on Violence Against Women and Mental Health in Sri Lanka, India and Afghanistan	15-16
Table 4. Mental Health Resources in Sri Lanka, India and Afghanistan	20

Plain Language Summary. We describe a literature and database review on the prevalence of violence against women, mental health conditions, the policy context, and mental health services in Afghanistan, India, and Sri Lanka. The article provides information on the limited data and policies addressing the intersection between violence and poor mental health, as well as a lack of mental health service provision. Among ever-partnered women, 7% reported experiencing sexual violence in Afghanistan and Sri Lanka, respectively and 6% in India. The intersection between mental health and violence is limited and unclear. Future research on integrating mental healthcare within social services; translating trauma-informed approaches into service provision; and addressing family violence is proposed.

INTRODUCTION

Defined as public or private actions or threats made to intentionally hurt or harm women physically, sexually, or psychologically (such as coercion and loss of freedom),¹ violence against women (VAW) is a human rights violation and global public health problem that adversely affects women's health and wellbeing.² Global estimates suggest that between 10% and 53% of ever-partnered women have experienced physical or sexual intimate partner violence in their lifetime.^{3,4} These estimates are wide-ranging due to the challenges of collecting accurate data, given stigma, issues of non-response and safety, cultural and gender norms around violence, non-recognition of VAW, and issues of privacy and confidentiality, among others.⁵ Recurrent and chronic exposure to violence leads to persistent stress and results in poor mental health and wellbeing.³ This stress commonly manifests as anxiety, depression, post-traumatic stress disorder (PTSD), self-harming thoughts and/or behaviours, and/or substance misuse.^{1, 2, 3, 6, 7, 8, 9, 10, 11}

VAW is considered a gendered issue. The prevalence of domestic and sexual violence victimization is higher among women than men. Moreover, VAW exposure places women at higher risk of suffering from depression, anxiety, and suicide.^{12, 13 14}

Conflict further affects violence prevalence, mental health conditions, and the ability of services to respond. Conflict zones have been shown to have some of the highest prevalence of VAW globally,^{15, 16, 17} and individuals living in them have a greater likelihood of poor mental health than those in non-conflict settings.^{18, 19} Research on trafficking has shown similar findings. Trafficked individuals report comparable levels of poor mental health (anxiety, depression, and post-traumatic stress disorder (PTSD),^{20, 21} violence and coercion to survivors of domestic violence, prisoners of war, and survivors of torture.^{22, 23, 24, 25} Factors contributing to adverse mental health and violence among trafficked individuals include differential levels of power and control exerted by traffickers.^{22, 26, 27}

Despite the high prevalence of VAW and co-existing mental health problems in most parts of the world, access to care for women survivors of violence experiencing mental health problems is poor. Health systems have been slow to respond to VAW and provide targeted services.²⁸ Meanwhile, demand-

side barriers like stigma, sociocultural norms, and lack of financial resources affect uptake. These barriers could be attributed to the social acceptability of VAW in some settings, overburdened and underfunded mental health systems, and the consequent limited capacity of staff to respond adequately.^{29, 30} Additionally, health systems in conflict-affected settings are particularly fragile, with most of the population unable to obtain basic health services.³¹ In such settings, there is a lack of understanding of how mental health services respond (if at all) to women survivors of violence.

In light of these gaps, there is a need to develop effective mental health and VAW prevention and intervention strategies. The goal of our wider research programme was to develop a package of care to support the mental health of survivors of VAW in Afghanistan, India, and Sri Lanka. Importantly, Afghanistan has been in a state of conflict for more than four decades,³² with a recent escalation due to a change in government in August 2021.³³ Sri Lanka experienced 26 years of civil war (1983-2009)³⁴ and has recently experienced ethnic and religious tensions and political conflict.³⁵

Designing intervention strategies that are contextually relevant requires an in-depth understanding of epidemiology, as well as existing policies, legislation, and services.³⁶ Experiences of violence and its mental health consequences are shaped by the social, cultural, political, and economic context. For example, women living in countries without domestic violence legislation are estimated to be 7% more likely to experience violence than those in countries with legislation. In societies where violence is not criminalised, a perception that it is broadly acceptable tends to dominate public discourse. Survivors of violence may therefore be less likely to seek support from health services, police, or community resources, with physical and poor mental health repercussions.³⁷ In this situation analysis we aimed to quantify the forms of VAW and mental health conditions in Afghanistan, India, and Sri Lanka, to identify policies that integrate these two issues, as well as existing services for VAW and mental health and their integration.

Research group

Our group consisted of researchers based in non-government organisations (NGOs) and universities in Afghanistan, India, Sri Lanka and the United Kingdom. Partners in Afghanistan were based at Humanitarian Assistance for the Women and Children of Afghanistan (HAWCA), an NGO that supports women and children across Afghanistan including survivors of violence. Partners in India were based in Mumbai (Society for Nutrition, Education & Health Action (SNEHA)) and Goa (Sangath). SNEHA includes a group that works with communities to prevent and address VAW, while Sangath works to improve mental health by addressing the psychological and social needs of individuals and families. Partners in Sri Lanka were based at the Department of Psychiatry in the University of Colombo.

METHODS

As a detailed approach to identifying the barriers, facilitators, resources, challenges, and opportunities within settings,³⁸ a situation analysis was deemed to be the most appropriate method given the paucity of information available on mental health context in the context of VAW within these settings. The evidence was collected by mapping grey and peer-reviewed literature and other publicly available data. It did not include empirical data collection. Development of the package of care did, however, involve qualitative work with providers and survivors. This is detailed in the synopsis article in the NIHR Global Health Research journal. To guide data collection, we developed a situation analysis tool covering three domains and the following inclusion criteria: (i) the prevalence of violence and mental health conditions among women (prevalence of different forms of violence, help-seeking behaviours, prevalence of mental health conditions, such as depression, anxiety, PTSD, substance use, suicide and self-harm etc); (ii) policy context (policy and legislation for mental health, violence against women, and trafficking); and (iii) mental health services (mental health human resources, treatment coverage, violence and mental health screening). The situation analysis tool was designed by our transdisciplinary research team. The team also supported interpretation of the findings. We adapted the Programme for Improving Mental Health Care (PRIME) situation analysis tool created to guide a district-level situation analysis in

preparation for the implementation of mental health interventions in primary care in five countries.³⁹ Adaptations reflected our focus on mental healthcare for women survivors of violence and on national or regional indicators. Indicators were determined by consulting with members of the research group who had expertise in violence against women, trafficking, and mental health conditions. We made a pragmatic decision to focus on regional and country-level indicators due to the lack of locally-specific data and the possibility that as part of our research programme interventions would be implemented at more than one site within each country.

The situation analysis tool was initially completed by in-country investigators (AC, UB, THRS) in October 2018. The tool was used to search data between October and December 2018. A revision of the data in all countries took place in August and September 2022 (MQD). Data were primarily drawn from public domain health surveillance data, household survey collected through the Demographic and Health Surveys,⁴⁰ academic publications, policy documents, governmental and non-governmental reports, and the World Health Organization (WHO) Mental Health Atlas (see Appendix 1 for a list of main data sources by country). Data collected for Afghanistan was under the previous government. MQD, AA, AN, AP, LA and UB reviewed the data for completeness and accuracy. Although we did not use a systematic approach, we tried to establish the plausibility of data by triangulating with grey and unpublished literature where possible and checking with members of our research group with relevant expertise.

The situation analysis is part of a larger programme of work that included a transdisciplinary group with team members from Afghanistan, India, and Sri Lanka. These countries were included as team members aimed to develop a support package for the mental health of women survivors of violence and modern slavery. In addition, exploring evidence across select South Asian settings contributes to much-needed intervention based work in a high-VAW burden region. The idea was not to represent South Asia but to bring together a range of settings and stakeholders that work in the prevention of VAW, mental health, as well as academics conducting research on the topic.

RESULTS

The Prevalence of Violence and Mental Health Conditions

Women's Exposure to Violence

We found that multiple forms of violence against women were common (Table 1). Prevalence of violence was highest in Afghanistan, where 46% of women⁴¹ had experienced physical violence within 12 months preceding the survey (53% in their lifetime)⁴¹ and 7% of ever-married women had experienced sexual violence.⁴² There were high rates of marital control; for example, 35% of ever-partnered women reported psychological abuse (defined as jealousy or anger if they talked to other men), and 35% reported their partner insisted on knowing where they were at all times.⁴² The highest rates of physical violence perpetrated by family members were also reported in Afghanistan (40% of ever-married women 15-49 years).^{41, 42}

In India, 21% of women reported experiencing physical violence perpetrated by anyone within the last 12 months (lifetime estimates of 30%).⁴³ Sexual violence was prevalent, with 6% of women reporting it in their lifetime.⁴³ Meanwhile, 33% of ever-married women reported ever experiencing physical, emotional or sexual violence by their current partner.⁴³ Marital control among ever-married women was reported as the following: 27% experienced jealousy or anger if they talked to other men; 20% on knowing where they were at all times; and 9% were accused of infidelity.⁴³ Nearly one third of ever-married women (32%) in India also experienced family physical violence, which includes the husband's family and step-family members.⁴³

The COVID-19 pandemic appears to have increased VAW in India. In a rapid online survey across India in May 2020, 18% of women reported currently experiencing spousal violence (n=560). There was a 33% increase in spousal violence since lockdown. The rates of physical, sexual, verbal and emotional violence were estimated at 35%, 11%, 65% and 44%, respectively. Meanwhile, 76% of women

216 reported being sad or depressed due to violence and 37% reported having thoughts of harming
217 themselves. ⁴⁴ In a qualitative analysis of 586 women during lockdown between April and July 2020, 86%

Table 1. Types of Violence Against Women in Sri Lanka, India, and Afghanistan

Exposure to violence	Afghanistan	India	Sri Lanka
% women experienced physical violence from anyone	Last 12 months: 46% (35) Ever: 53% (35) Pregnant: unknown (35) Postnatal: unknown (35)	Last 12 months: 21% (37,82) Ever: 30% (37) Pregnant: 4% (37) Postnatal: unknown (37)	Last 12 months: 4.8% (30) Ever: 18.9% (30) Pregnant: unknown Postnatal: unknown
% ever-married women experienced sexual violence from anyone	7% (35)	6% (37)	7% (30)
% women experienced conflict-related sexual violence	No data found/identified	No data found/identified	No data found/identified
% of women trafficked	No data found/identified	No data found/identified	No data found/identified
% ever-partnered women experienced marital control	Type of controlling behaviour (36): 60% jealousy or anger 35% knowing where wife is at all times 25% accusations of infidelity	Type of controlling behaviour (37): 27% jealousy or anger 20% knowing where wife is at all times 9% accusations of infidelity	Type of controlling behaviour (30): 8% jealousy or anger 9% knowing where wife is at all times 6% accusations of infidelity
% ever-married women experienced physical / sexual intimate partner violence	56% (29)	33% (37)	20% (30)
% women experienced violence by family members (other than intimate partner)	94% (29)	Physical violence since age 15: 42% (37)	7% (30)
% women victims sought help in response to violence	Physical and sexual violence: 33% (36) Physical violence only: 18%	14% (37) Family: 65% Police: 3%	75% (73) Family: 27% Neighbours: 18%

219
220
221
222
223
224
225
226
227
228
229
230
231
232
233

	Sexual violence only: 9% Family: 34% Neighbours: 18% Police: Unknown Health Service: Unknown	Health Service: 1% Religious Leader: 2% Husband family: 29% Friend:15%	Police: 7% Health Service: 3% Government Institutions: 1.4% NGOs: 0.4% Other: 3%
--	--	---	--

of women at initial consultation reported experiencing emotional abuse, 76% economic violence, 70% intimate partner violence (IPV), 64% physical violence, 56% neglect, 56% controlling behaviours and 35% sexual violence; 86% reported surviving more than one of these forms, while 22% had experienced all of these forms of violence. 27% of women were still surviving violence during lockdown and 28% were surviving intimate partner or family violence.⁴⁵

In 2019, Sri Lanka conducted the first national Woman's Wellbeing Survey exploring VAW among ever-partnered women of reproductive age (15-49 years). The prevalence of physical violence among ever-partnered women within 12 months of the survey was 5% (lifetime estimate 19%).³⁴ Sexual violence was prevalent, with 7% of ever-partnered women (15-49) having experienced sexual violence.³⁴ Approximately 20% of women had experienced physical or sexual violence from a partner during their lifetime.³⁴ Approximately 19% of ever-partnered women reported experiencing controlling behaviours by a partner (during their lifetime), of whom 9% reported that their partner insisted on knowing where they were at all times; 8% experienced jealousy or anger if they spoke with another man and 6% were accused of infidelity.³⁴ Approximately 7% of ever-married women reported experiencing physical violence by a non-partner; of this, 42% was perpetrated by male family members, while 27% was perpetrated by female family members.³⁴ Concurrent to our situation analysis, WHO's 2018 country profile on VAW reported a wide range of 18-72% lifetime prevalence among women of reproductive age, indicating how prevalence data are informed, by what forms, and how VAW is researched in a given context⁴⁶.

Mental Health Problems

Table 2 illustrates the burden of mental health conditions across the three countries. Evidence on mental health conditions among survivors of violence in Afghanistan was scarce. A randomised-controlled trial evaluating women's empowerment in Afghanistan showed that, compared to unexposed women, women exposed to violence in the last 12 months were six times more likely to have suicidal thoughts.⁴⁷

260 **Table 2. Indicators of Poor Mental Health and Treatment in Sri Lanka, India and Afghanistan**

Prevalence of mental disorders	Afghanistan	India	Sri Lanka
Suicide rates (per 100,000 population)	5.7 per 100,000 (118)	Overall: 17.9 per 100,000 individuals (46) Females: 14.7 (13.1-16.2) (46) Males: 21.2 (14.6-23.6) (46)	12.9 per 100,000 (119) Males (49): 44.6 Females (49):16.8
PTSD (%)	No data found/identified	General population: 0.2% (83) Survivors of VAW: 14% (120,121) Survivors of conflict: 19% (121)	General population: 2% (49) Survivors of Conflict: 7% (50)
Common mental disorders (Anxiety and depression) (%)	No data found/identified	General population:10% (82) 3% (adults)	Survivors of conflict: 33% (50) (anxiety) 22% (50) (depression)
Alcohol abuse/dependence (%)	No data found/identified	General population: 4.6% (82)	General population: 5% (49)
Severe mental disorders (schizophrenia and other psychotic disorders, bipolar disorder, and moderate-to-severe depression) (per 100,000 population) ⁴⁸	General population: 47.0 (49)	Schizophrenia and other psychotic disorders: 1% (lifetime) and 0.4% (current prevalence)(81) Bipolar: 0.5% (lifetime) and 0.3% (current prevalence) (81)	No data found/identified
Treatment gap (%) (The proportion of people who need but have no access to mental health care and/or are not in treatment) ⁴⁹	No data found/identified	70% bipolar affective disorder (82) 75% psychotic disorders 85% major depressive disorder 83% Neurosis 86% alcohol use disorder	General population: 68% experience treatment gap (122)

261

262

263

In India, the Global Burden of Disease Study (2020) ⁵⁰ analysed mental health conditions between 1990 and 2017. The prevalence of depressive, anxiety and eating disorders was higher among females than males. The authors attributed these differences in prevalence to violence, sexual abuse, gender discrimination, negative sociocultural norms, and pregnancy and postpartum stress. ^{51, 52, 53, 54} The study also showed evidence for the contribution of risk factors such as childhood sexual abuse and intimate partner violence to mental health conditions. ⁵⁰ Estimates suggested that 7% and 5% of DALYs for depressive disorders were attributed to childhood sexual abuse and IPV, respectively, and were significantly higher among females than males. Women with depressive disorders were also more likely to die by suicide than males. ⁵⁰ Indeed, the suicide death rate among women in India was nearly three times that of countries with a similar Socio-Demographic Index, ⁵⁵ with married women representing the largest proportion of suicide deaths among all women. ^{56, 57} Indeed, prior cross-sectional studies conducted on suicide attempts in North India have showed a higher proportion of suicide attempts and psychiatric comorbidities among women. ^{58, 59, 60}

Estimates from the 2011 Mental Health Atlas in Sri Lanka showed that the suicide mortality rate was higher among males (44.6) than females (16.8 per 100,000 population). ⁶¹ Evidence from 2009 showed that approximately 2% experienced PTSD (7% of people affected by conflict). ⁶² In a cohort-analysis of police and hospital data on suicide, Knipe and colleagues (2014) showed that suicide rates were highest among women 17-25 years between 1975 and 2012. ⁶³ A more recent analysis indicates this trend towards younger female suicides has persisted through the latest national data 2022. ⁶⁴ In research on perinatal women, those with a history of intimate partner violence were four times more likely to report suicidal ideation and/or behaviour during their pregnancies compared to women without this violence history. ¹⁴ The Sri Lanka Women's Wellbeing Survey (2019) reported that women exposed to violence were more likely to exhibit emotional distress than those who were not exposed to violence. ³⁴ Estimates suggest that 15% of women who experienced physical or sexual violence by a partner engaged in self-harm, compared to 1% of women who had never experienced violence. At the same time, 36% of

women who experienced physical or sexual violence by a partner had suicidal thoughts, compared with 7% of women who had not.³⁴

Policy Context

Where there were policies on VAW and mental health, they focussed on prevention (Table 3). The countries varied in their policy, programme, and strategy responses to VAW and trafficking. In addition, mental healthcare policies for women survivors of violence were largely absent from the literature.

Afghanistan, India, and Sri Lanka have ratified the Convention on the Elimination of All Forms of Discrimination against Women (CEDAW),⁶⁵ as well as the Convention on the Rights of Persons with Disabilities.⁶⁶ Sri Lanka has ratified the United Nations Declaration on the Elimination of Violence Against Women,⁶⁷ while India has committed to it.⁶⁸ All three countries are signatories to the Sustainable Development Goals (SDG 5 aims to achieve gender equality)⁶⁹ and the Palermo Protocol (Table 3).⁷⁰

Afghanistan has shown some commitment to address violence against women. In 2009, a Presidential Decree was issued on the Law on the Elimination of Violence Against Women (EVAW), but it was opposed in Parliament in 2013 and its current status is unclear.⁷¹ In 2017, Afghanistan passed a law Prohibiting Human Trafficking and Migrant Smuggling. The law was expected to be scaled up across the country, but security, corruption, and warlord control have hindered its implementation.⁷² The Penal Code in Afghanistan recognises VAW, violence against individuals with mental health conditions, and trafficking as crimes.⁷³ The National Action Plan for the Women of Afghanistan (NAPWA) (2007-2017) aimed to pursue gender equality and women's empowerment through intersectoral strategies to prevent VAW. The country has experienced a shortage of lawyers to advocate for women's rights, but women's activists are working to prevent VAW and support improved mental health of women survivors of violence (Table 3).⁷⁴

314 **Table 3.** Policies, Legislation, Strategies, Plans and Programmes on Violence Against Women and Mental Health in Sri Lanka, India and
315 Afghanistan

Policies/ Legislation	Afghanistan	India	Sri Lanka
Policies/Legislation addressing VAW	<p>Convention on the Elimination of All Forms of Discrimination against Women (Ratified) (51)</p> <p>Convention on the Rights of Persons with Disabilities (Ratified) (52)</p> <p>Signatory on Sustainable Development Goals (55) and Palermo Protocol (56)</p> <p>Law on the Elimination of Violence Against Woman (59)</p> <p>Afghan Civil Law and the Penal Code (61)</p>	<p>Convention on the Elimination of All Forms of Discrimination against Women (Ratified) (51)</p> <p>Convention on the Rights of Persons with Disabilities (Ratified) (52)</p> <p>United Nations Declaration on the Elimination of Violence Against Women (Committed) (54)</p> <p>Signatory on Sustainable Development Goals (55) and Palermo Protocol (56)</p> <p>Protection of Women from Domestic Violence Act (2005) (66)</p> <p>National Policy for Women 2016 (Draft)</p>	<p>Convention on the Elimination of All Forms of Discrimination against Women (Ratified) (51)</p> <p>Convention on the Rights of Persons with Disabilities (Ratified) (52)</p> <p>United Nations Declaration on the Elimination of Violence Against Women (Ratified) (53)</p> <p>Signatory on Sustainable Development Goals (55) and Palermo Protocol (56)</p> <p>The Prevention of Domestic Violence Act (2005) (76)</p>
Plans/Programmes addressing VAW	National Action Plan for the Women of Afghanistan (2007-2017) (60)	No data found/identified	National Plan of Action to address Sexual and Gender-based Violence (2016-2020) (75)
Addressing trafficking	<p>Afghan Civil Law and the Penal Code (61)</p> <p>Law Prohibiting Human Trafficking and Migrant Smuggling (2017) (60)</p>	Anti-Trafficking Bill (2018) (71)	Convention on Preventing and Combatting Trafficking in Women and Children for Prostitution (Act No 30, 2005) (75)

Mental health strategy/ policy	National Mental Health Strategy 2011-2015 (NMHS) (78) National Mental Health Strategy (2019-2023) (78)	National Mental Health Policy (2014) (68) National Mental Healthcare Act (2017) (68) National Mental Health Programme (1982 – Present) (81)	Mental Health Policy of Sri Lanka (2005-2015) (96) Mental Health Policy of Sri Lanka 2020-2030
Supporting survivors of conflict	No data found/identified	No data found/identified	No data found/identified
Integration of VAW and Mental Health Strategies/services	National Mental Health Strategy 2019-2023 includes developing an integrated programme and package of mental health services and support to women survivors of violence (78).	One Stop Centres for women and children exposed to violence to obtain integrated services, including psychological, medical and legal services, in one location (74).	Gender and Women's Health Unit within the Family Health Bureau to address domestic violence within the health sector (76). Gender-Based Violence desks and Friendly Abode/Mithuru Piyasa/Natpu Nilayam centres provide safe havens for women exposed to domestic violence and operate in tertiary care hospitals (77).

Afghanistan made efforts to improve the mental health of its population. The National Mental Health Strategy (2011-2015) aimed to promote mental health, tackle stigma and discrimination associated with mental health conditions, reduce the impact of mental health conditions on individuals, families, and the community, prevent the development of mental health problems and mental health conditions, and provide quality, integrated, evidence- and rights-based care for people with mental health conditions. However, implementation of the Strategy was challenging due to chronic and multiple conflicts and an underdeveloped mental health infrastructure in an existing weak health system.⁷⁵ The country has recently introduced the National Mental Health Strategy 2019-2023, which aims to improve access to services, increase mental health resources, and strengthen the capacity of mental health care providers to respond to the population's mental health needs. It also includes developing an integrated programme and package of mental health services and support to women survivors of violence.⁷⁶ With the recent change in government in Afghanistan, the progress in its implementation is unclear.

In India, the government has committed to gender equality. In its Preamble, Fundamental Rights and Duties, the Constitution states that women have rights and are equal under law.⁷⁷ India passed The Protection of Women from Domestic Violence Act (PWDVA) in 2005. The Act articulates the legal rights, financial and other benefits to survivors of domestic violence. This Act has resulted in increased advocacy, as well as organisations working to ensure its implementation.⁷⁸ In 2016, the Ministry of Women & Child Development developed a National Policy for Women.⁷⁹ One of its objectives was to eliminate all forms of VAW in India. This Policy, however, has not yet been launched. India also has the Nirbhaya Fund, which provides financial resources to programmes that support the safety and dignity of women.⁸⁰ In 2018, India passed an Anti-Trafficking Bill to protect, rescue and rehabilitate victims of trafficking.⁸¹ The Bill takes a criminalisation approach to trafficking and sex work. However, due to poorly defined crimes and poor enforcement mechanisms, there have been low conviction rates (Table 3).⁸²

The National Mental Health Policy (NMHP) in India was implemented in 2014 and the National Mental Healthcare Act (NMHA) in 2017.⁸³ The NMHP aims to prevent mental health conditions and promote mental wellbeing and universal access to mental health services.⁸³ The Policy states that women should not be discriminated against in the provision of services. It also recognises the disproportionate burden of mental health conditions experienced by marginalised populations (e.g., victims of trafficking, those living in conflict, sex workers, children, and sexual minorities). The government provides shelters for homeless women under its social welfare scheme,⁸⁴ along with other third sector organisations which provide shelter for women exposed to violence. The country has also implemented One Stop Centres, state sponsored service centres for women and children exposed to violence. These provide an opportunity for survivors to obtain integrated services, including psychological, medical, and legal services, in one location (Table 3).⁸⁵

The Sri Lankan government has made several commitments to reduce women's exposure to violence and subsequent harm. These include the National Action Plan to Address Sexual and Gender-Based Violence (2016) to eradicate violence against women and children using a multisectoral approach,⁸⁶ and the Prevention of Domestic Violence Act (2005).⁸⁷ The country has also had some service developments, including a Gender and Women's Health Unit within the Family Health Bureau to address domestic violence within the health sector,⁸⁷ and the establishment of Women's and Children's Desks at selected police stations, which aim to be the contact point for survivors and those who wish to report violence, as well as, Gender-Based Violence desks and Friendly Abode/Mithuru Piyasa/Natpu Nilayam centres (56 centres in 21 districts) that provide support for women exposed to violence and operate in certain tertiary care hospitals.⁴⁶ Sri Lanka has also established the "1936" women's helpline and has been implementing gender mainstreaming programmes. The country also has field officers (Women's Development Officer, Child Rights Promotion Officer, Early Childhood Development Officer) working within communities to link people with available resources and select police stations have Women's and Children's Desks as focal points for violence reporting and survivor support, staffed by female officers (Table 3).⁴⁶

The government has shown commitment to address trafficking. Actions include the Convention on Preventing and Combatting Trafficking in Women and Children for Prostitution (Act No 30, 2005), standard operating procedures to identify and refer potential trafficking victims to protection services,⁸⁸ and training sessions to identify trafficking victims for members of civil society and local, district, and state officials, including health, probation, police, immigration, Criminal Investigation Department, and National Child Protection Authority officials.⁸⁸ Despite these advances, there is a lack of public awareness on the legislation and rights around VAW and trafficking; poor coordination between government and NGOs working on VAW and trafficking with mental health services; a lack of shelters for women survivors of violence and trafficking; unequal distribution of resources and services (mainly available in cities); a lack of monitoring and follow-up of existing programmes; and minimal funding allocated to the prevention of VAW, trafficking and mental health conditions (Table 3).⁸⁹

The Mental Health Policy of Sri Lanka (2005-2015) was passed in 2005 with the aim of establishing community-based, comprehensive mental health services. The policy focussed on several areas of mental health services, including service organisation, human resources, management at the national and provincial levels, research and ethics, stigma and mental wellbeing, the National Institute of Mental Health and mental health legislation.⁹⁰ Not all targets were reached under the previous policy and the recently launched Mental Health Policy (2020-2030) proposes to expand services by establishing at least one psychiatric centre or community psychiatric clinic in each Ministry of Health area around the country (Table 3).

Mental Health Services

Mental health infrastructure has evolved across the three countries through activist grassroots organisations and governmental responses, but mental health services and their accessibility are limited by a range of individual and systemic barriers. Significant treatment gaps were identified in Afghanistan, India, and Sri Lanka (Table 2) with limited mental health human resources (Table 4). There was limited

Table 4. Mental Health Resources in Sri Lanka, India and Afghanistan

	Afghanistan	India	Sri Lanka
<i>Human Resources (per 100,000)</i>			
Psychiatrists	0.34 (118)	0.30 (80)	0.58 (119)
Mental Health Workers	1.66 (80)	1.93 (80)	7.14(80)
Nursing Officers	0.07 (118)	0.80 (80)	2.93 (119)
Psychologists	0.35 (118)	0.07 (80)	0.29 (119)
<i>Facilities (N)</i>			
Community-based mental health outpatient facilities	2,622 (118)	1,217 (80)	20 (80)
Psychiatric hospitals	1 (80)	136 (80)	1 (80)
Psychiatric units in general hospitals	4 (80)	389 (80)	36 (119)
Mental hospital beds (per 100,000)	0.26 (118)	1.43 (80)	6.61 (119)

evidence on mental health care resources for women exposed to violence. It was not possible, therefore, to determine if women could safely disclose past or present experiences of violence within facilities, gain the care they needed, and if there were adequate diagnostic assessments to understand the psychological impact of violence.

The government of Afghanistan has made some progress with mental health services. Due to the National Mental Health Strategy (2010-2014),⁹¹ and its revision in 2015, the integration of mental health services within each level of the healthcare system led to a 75% increase in the number of provinces with an integrated Essential Package of Hospital Services based mental health services.⁹² However, access to mental healthcare remains a challenge, especially for women. This challenge is attributed to the socio-cultural context and ongoing conflict on women's ability to seek and afford healthcare, especially specialist healthcare. In addition, mental health stigma, limited awareness and low perceived needs, and reliance on traditional healers remain barriers to seeking mental healthcare.⁷⁶

Contributing to these challenges is the paucity of mental health human resources and facilities in Afghanistan. There were an estimated 0.34 psychiatrists, 0.07 nursing officers and 0.35 psychologists per 100,000 population. In addition to mental health outpatient facilities, there was one mental health hospital and four psychiatry units in general hospitals in the country. Afghanistan therefore relies on NGOs to provide mental healthcare, under the remit of the Ministry of Health. Most of the NGOs and large public mental health services with trained service providers are located in the major cities (Kabul, Herat and

Balkh) with other regions experiencing limited access to services.⁹³ According to a trans-sectional probability survey of the general population,⁹² results showed that 19% of participants who reported any mental health problem received help at some point, with 12% receiving help within the last 12-months. There was regional variation in mental health help-seeking, from 4% in the Central highland region to 22% in the South. Results indicated that mental health help providers included the health sector and non-health sector (e.g. healer or religious leader). Access to mental health care was adversely influenced by regional exposure to danger (i.e., number of attacks), traumatic events, and clinical needs.⁹²

India has demonstrated strong political and legal commitments to enhancing mental health services. Since 1982, the country has implemented the National Mental Health Programme,⁹⁴ which focuses on expanding access to care through capacity building and incorporating mental health into primary care. The District Mental Health Programme (DMHP) was created to focus on scaling up services at the community level. By 2014, approximately 20% of districts had implemented a DMHP.⁹⁴

Despite political and legal commitments, the implementation of mental health services has been challenging for India. The country faces a large treatment gap and a lack of accessibility. The most recent National Mental Health Survey of India⁹⁵ estimates that there is a large treatment gap for all mental disorders. For example, it is estimated that 85% of people with major depressive disorder do not access evidence-based care.⁹⁶ There is also a lack of evidence-based care, while women experience poorer mental health treatment compared to men.^{50, 97, 98, 99, 100, 101, 102} Indeed, women with mental health conditions may be institutionalised without consent.⁹⁶ Adding to the treatment gap is the shortage of mental health service providers. The most recent estimates suggest that there are less than one psychiatrist (0.3) and two mental health workers, 0.12 nurses and 0.07 psychologists per 100,000 population.⁹³ According to the literature, this gap is attributed to seeking mental health care from faith healers,⁹⁶ as well as a low perceived need for care, stigma^{103, 104, 105, 106} and discriminatory attitudes of healthcare providers.^{103, 107}

With the Mental Health Policy of Sri Lanka in 2005, community-based mental healthcare was prioritised. As a result, there was an increase in district hospital-based clinics, outreach clinics, home-

based care, community support centres, domiciliary care, helplines, and intermediate rehabilitation centres.¹⁰⁸ Although community services are available to all age groups, treatment services prioritise adults with severe mental disorders, including schizophrenia and substance use disorders.¹⁰⁹ In 2012, Sri Lanka initiated a screening programme for postnatal depression, embedded in the national pregnancy care programme, but implementation has been uneven and limited.¹⁴

Sri Lanka faces a shortage of mental health resources. In 2017, Sri Lanka had 0.52 psychiatrists, 1.47 Medical Officers of Mental Health, 3.28 nursing officers and 0.25 psychologists per 100,000 population.¹⁰⁹ The specific needs of women survivors of violence are overlooked, reducing accessibility to available mental health services.¹⁰⁹ Estimates suggest that the treatment gap for common and severe mental health disorders, as well as substance use disorders, is high, at 68%. This treatment gap is attributed to stigma and limitations in cognitive and other forms of access to mental health services.⁹³

DISCUSSION

This study is the first, to our knowledge, to investigate data from Afghanistan, India, and Sri Lanka on the mental health needs and context of women survivors of violence using a situation analysis tool. The evidence suggests that Afghanistan, India, and Sri Lanka share similarities in terms of a lack of data, implementation of policies and legislation on all forms of VAW, and limited recognition of women's mental health needs among survivors of violence. Further, there is a paucity of mental health service provision in these countries. These points should be considered in light of the scarcity of data on the topic.

The study showed that women in Afghanistan, India, and Sri Lanka experienced multiple forms of violence, with most of the data on violence focused on that which is committed by a partner or family member.^{12, 15} We were unable to identify data on the percentage of women who had experienced conflict-related sexual violence, or on women who had been trafficked, illustrating gaps in the literature. The lack

of data on conflict-related sexual violence and trafficking could be attributed to a fear of intimidation and retaliation against women survivors of conflict-related sexual violence, as well as a lack of collection.¹¹⁰ Similarly, the lack of data on trafficking could be attributed to the difficulty faced by community members, service providers, and law enforcement officers in identifying survivors, incomplete reporting, and a lack of services targeting trafficking survivors.¹¹¹ Without this information, it is difficult to develop prevention and intervention programmes for women survivors of conflict-related violence and trafficking with mental health conditions in these settings. Beyond being influenced by how violence research is designed and conducted (e.g., definitions, tools), the lower rates of VAW reported in Sri Lanka could be attributed to fear of not being believed, embarrassment, the normalisation of violence, protecting family reputation, and lack of knowledge of options and support services.^{34, 112} Meanwhile, limited evidence from India suggests that domestic violence increased threefold during the COVID-19 pandemic.^{113, 114, 115} There was a lack of data on the impact that COVID-19 has had on VAW in Afghanistan and Sri Lanka. Although we did not have specific age criteria, the data reported were based on estimates for women 15-49 years of age.

There was some evidence on mental health conditions, with limited data on the associations between women's exposure to different forms of VAW and symptoms of depression, suicidality, and self-harm in India and Sri Lanka.³⁴ Scarce evidence was shown in Afghanistan among women exposed to violence and suicidal thoughts.⁴⁷ This lack of evidence could be attributed to the paucity of mental health resources and cultural conceptualisations of mental health.¹¹⁶ Future research could investigate the bidirectional relationship between VAW and mental health conditions, with violence as a risk factor for, and consequence of, poor mental health. Prior evidence has shown a bidirectional, cumulative and dynamic interaction between mental health conditions and violence.¹¹⁷ Indeed, mental health conditions may influence a woman's decision to stay in a violent relationship, in choosing their partner and assessing their risk.¹¹⁸ At the same time, poor mental health status among women experiencing violence may adversely influence their ability to safely leave.¹¹⁹ These challenges are compounded by controlling

behaviour, wherein women with poor mental health may be less able to protect themselves against violence.¹¹⁹ The countries need to develop prevention policies and programmes to support women exposed to violence with mental health problems.

Although mental health policies were present in Afghanistan, India, and Sri Lanka, a lack of resources and individual barriers contributed to poor policy implementation. For example, there are challenges to developing a standardised mental health infrastructure that addresses the needs of women survivors of violence. Evidence has shown that mental health providers may not be trained on how to identify or screen for VAW.¹²⁰ There is also a lack of clinical reporting of women exposed to VAW.¹²¹ Due to limited health resources, the workload that mental health care providers have within these settings may prevent them from adequately and appropriately attending women exposed to violence.¹²² Further, women survivors of violence have reported discrimination by clinicians.¹²³ To address these challenges, mental health care services could screen for VAW, with referral mechanisms to specialised services and programmes. The services could ensure that mental health providers are appropriately trained to effectively treat women survivors of violence. In addition, there needs to be greater collaboration between community-based organisations and the mental health system to provide support networks for women survivors of violence.¹²³ Task-sharing, including through the use of alternative technologies, is another strategy whereby non-specialists and community health workers can be trained to deliver mental health services for women survivors of violence.^{124, 125} Safety and confidentiality would be important considerations.

Evidence from the situation analysis illustrates an opportunity to develop mental health interventions, policies and programmes for women survivors of violence in the three countries. A targeted, intersectoral approach to identify and treat mental health conditions among women survivors of violence is needed, with a particular focus on designing, testing, and refining interventions.^{57, 103, 126, 127} For example, there is a possibility of integrating mental and physical/sexual health and protection services with access to housing, legal, and financial services, as well clear referral pathways.^{30, 121} Integration implies ensuring the availability of adequate mental health services and appropriately trained service

providers, flexible service hours, seamless referral to other services, greater community awareness of these services, and the institutionalisation of a trauma-informed approach.¹²⁸

We have identified some limitations to the study. Given that the analysis included sources in the public domain, it is important to consider the rigor with which the data were collected. Our analysis was limited by the data available and their limited availability also precluded the direct comparison of findings across countries. For example, further research needs to be conducted on mental health resources in each country. In addition, some of the data collected and presented in this analysis are limited due to data collection and recruitment strategies, affecting the generalisability of findings. For example, approximately 69% of women respondents from the rapid online survey on spousal violence in India identified as being from upper and upper-middle income. Variability between regions, especially in a large country such as India, restricted our ability to assess implementation of VAW and mental health programmes. Future research needs to be conducted on women's economic and employment opportunities in relation to violence exposure and poor mental health. It should be noted that the analysis for Afghanistan covers up until the country came under Taliban control again in 2021. Where feasible and safe, new research should be conducted on VAW since the Taliban took over. This analysis did not include engagement with political and professional stakeholders, service users and communities in the co-design of the situation analysis and interpretation of findings but the wider work did involve qualitative data collection and engagement with survivors and service providers^{129, 130, 131}.

The utility of this knowledge is to provide a roadmap for how services could be updated in future work centred within these settings. In addition, and as part of the larger programme of work, additional research is required on integrating mental healthcare within social care; identifying ways to incorporate trauma-informed approaches into service provision; and addressing family violence.

CONCLUSION. This is the first study to investigate data from Afghanistan, India, and Sri Lanka on women's exposure to violence and mental health conditions. Based on the findings, the three countries experience a lack of data, prevention policies and legislation on different forms of VAW, and limited

recognition of women survivors' mental health needs. There is also a paucity of mental health service provision in these countries. These findings have implications for the development of effective VAW prevention and mental health promotion strategies.

ACKNOWLEDGMENTS.

We thank Alexander Catalano and Khawla Zainab for helping extract relevant data for India and Lida Ahmad from Afghanistan.

DATA SHARING. Due to the methods employed in this article, there is no data that can be shared.

ETHICS STATEMENT. Ethical permission for the research activities was granted by the UCL Research Ethics Committee (2744/007, 29 November 2018).

COMMUNITY ENGAGEMENT AND INVOLVEMENT. Community engagement and involvement was not included in the situation analysis. It was, however, part of the larger project, in which three different communities were involved, including survivors of domestic violence and human trafficking, counsellors, caseworkers, psychologists, psychiatrists, and lawyers.

EQUALITY, DIVERSITY, AND INCLUSION. Power differentials were considered across the entire, larger project in terms of knowledge and participant status. To address these power dynamics, one-to-one interviews were held with survivors of violence; we also engaged people with lived/living experience in workshops.

CONTRIBUTION OF EACH AUTHOR.

Meaghen Quinlan-Davidson: Formal analysis (lead), investigation (lead), writing – original draft; reviewing/editing (lead)

Ayesha Ahmed: Writing – reviewing and editing (equal)

Laura Asher: Writing – reviewing and editing (equal)

Nayreen Daruwall: Conceptualization (equal)

Delan Devakumar: Conceptualization (equal), project administration (lead); supervision (equal)

Abhijit Nadkarni: Writing – reviewing and editing (equal)

Alexis Palfreyman: Resources (equal); Writing – reviewing and editing (equal)

Lamba Saboor: Resources (equal); Writing – reviewing and editing (equal)

TH Rasika Samanmalee: Writing – reviewing and editing (equal)

David Osrin: Conceptualization (equal); funding acquisition (lead); supervision (equal)

INFORMATION GOVERNANCE. UCL is committed to handling all personal information in line with the UK Data Protection Act (2018) and the General Data Protection Regulation (EU GDPR) 2016/679. Under Data Protection legislation, UCL is the Data Processor, the partner organisations are Data Controllers, we process personal data in accordance with their instructions. You can find out more about how we handle personal data, including how to exercise your individual rights and the contact details for our Data Protection Officer here (<https://www.ucl.ac.uk/data-protection/data-protection-0>).

Appendix 1. Main Sources of Data by Country

Afghanistan

Websites from Ministry of Women's Affairs; Afghanistan Independent Human Rights Commission; WHO Mental Health Atlas (data is routinely collected every three years); US Department of State; Demographic and Health Survey (routinely collected household survey, every five years); and National Mental Health Strategy.

India

Websites from Demographic and Health Survey (routinely collected household survey, every five years); Ministry of Home Affairs, US State Department, Global Status Report on Alcohol, National Health Portal

Sri Lanka

Websites from the Ministry of Women and Child affairs; Ministry of Health and Indigenous Medicine; Director of Mental Health; Family Health Bureau; National Child protection Authority Sri Lanka; Sri Lanka Police Children and Women Bureau; Department of Social Services; WHO Mental Health Atlas (data is routinely collected every three years).

REFERENCES

1. Krantz G, Garcia-Moreno C. Violence against women. *J Epidemiol Community Health*. 2005;59(10):818-21. Access Date: 28/10/2024.
2. Global and regional estimates of violence against women: prevalence and health effects of intimate partner violence and non-partner sexual violence Geneva: WHO; 2013. Access Date: 28/10/2024.
3. Mannell J, Amaama SA, Boodoosingh R, Brown L, Calderon M, Cowley-Malcolm E, et al. Decolonising violence against women research: a study design for co-developing violence prevention interventions with communities in low and middle income countries (LMICs). *BMC Public Health*. 2021;21(1):1147. Access Date: 28/10/2024.
4. Ramsoomar L, Gibbs A, Chirwa ED, Dunkle K, Jewkes R. Pooled analysis of the association between alcohol use and violence against women: evidence from four violence prevention studies in Africa. *BMJ Open*. 2021;11(7):e049282. Access Date: 28/10/2024.
5. Guidelines for Producing Statistics on Violence against Women – Statistical Surveys. NYC: United Nations Department of Economic and Social Affairs; 2014. Access Date: 28/10/2024.
6. Chang EC, Kahle ER, Hirsch JK. Understanding how domestic abuse is associated with greater depressive symptoms in a community sample of female primary care patients: does loss of belongingness matter? *Violence Against Women*. 2015;21(6):700-11. Access Date: 28/10/2024.
7. Eberhard-Gran M, Schei B, Eskild A. Somatic symptoms and diseases are more common in women exposed to violence. *J Gen Intern Med*. 2007;22(12):1668-73. Access Date: 28/10/2024.
8. Jewkes R, Willan S, Heise L, Washington L, Shai N, Kerr-Wilson A, et al. Elements of the Design and Implementation of Interventions to Prevent Violence against Women and Girls Associated with Success: Reflections from the What Works to Prevent Violence against Women and Girls? Global Programme. *International Journal of Environmental Research and Public Health* [Internet]. 2021; 18(22).
9. Devries KM, Child JC, Bacchus LJ, Mak J, Falder G, Graham K, et al. Intimate partner violence victimization and alcohol consumption in women: a systematic review and meta-analysis. *Addiction*. 2014;109(3):379-91. Access Date: 28/10/2024.
10. Oram S, Khalifeh H, Howard LM. Violence against women and mental health. *Lancet Psychiatry*. 2017;4(2):159-70. Access Date: 28/10/2024.
11. Devries KM, Mak JY, Bacchus LJ, Child JC, Falder G, Petzold M, et al. Intimate partner violence and incident depressive symptoms and suicide attempts: a systematic review of longitudinal studies. *PLoS Med*. 2013;10(5):e1001439. Access Date: 28/10/2024.
12. Violence against women prevalence estimates, 2018: global, regional and national prevalence estimates for intimate partner violence against women and global and regional prevalence estimates for non-partner sexual violence against women. Geneva: WHO; 2021. Access Date: 28/10/2024.
13. Facts and figures: Ending violence against women: UN Women; 2024 [Available from: <https://www.unwomen.org/en/what-we-do/ending-violence-against-women/facts-and-figures#83915>]
14. Palfreyman A. Addressing Psychosocial Vulnerabilities Through Antenatal Care-Depression, Suicidal Ideation, and Behavior: A Study Among Urban Sri Lankan Women. *Front Psychiatry*. 2021;12:554808. Access Date: 28/10/2024.

15. Mannell J, Lowe H, Brown L, Mukerji R, Devakumar D, Gram L, et al. Risk factors for violence against women in high-prevalence settings: a mixed-methods systematic review and meta-synthesis. *BMJ Global Health*. 2022;7(3):e007704. Access Date: 28/10/2024.
16. Hossain M, Zimmerman C, Kiss L, Kone D, Bakayoko-Topolska M, Manan DK, et al. Men's and women's experiences of violence and traumatic events in rural Cote d'Ivoire before, during and after a period of armed conflict. *BMJ Open*. 2014;4(2):e003644. Access Date: 28/10/2024.
17. Devakumar D, Palfreyman A, Uthayakumar-Cumarasamy A, Ullah N, Ranasinghe C, Minckas N, et al. Mental health of women and children experiencing family violence in conflict settings: a mixed methods systematic review. *Confl Health*. 2021;15(1):74. Access Date: 28/10/2024.
18. Charlson F, van Ommeren M, Flaxman A, Cornett J, Whiteford H, Saxena S. New WHO prevalence estimates of mental disorders in conflict settings: a systematic review and meta-analysis. *Lancet*. 2019;394(10194):240-8. Access Date: 28/10/2024.
19. Dimitry L. A systematic review on the mental health of children and adolescents in areas of armed conflict in the Middle East. *Child Care Health Dev*. 2012;38(2):153-61. Access Date: 28/10/2024.
20. Ottisova L, Hemmings S, Howard LM, Zimmerman C, Oram S. Prevalence and risk of violence and the mental, physical and sexual health problems associated with human trafficking: an updated systematic review. *Epidemiol Psychiatr Sci*. 2016;25(4):317-41. Access Date: 28/10/2024.
21. Hossain M, Zimmerman C, Abas M, Light M, Watts C. The relationship of trauma to mental disorders among trafficked and sexually exploited girls and women. *Am J Public Health*. 2010;100(12):2442-9. Access Date: 28/10/2024.
22. Herman J. Trauma and recovery: The aftermath of violence—from domestic abuse to political terror. New York, NY, US: Basic Books; 2015. ix, 326-ix, p. Access Date: 28/10/2024.
23. Hopper E, Hidalgo, J. Invisible Chains: Psychological Coercion of Human Trafficking Victims. *Intercultural Human Rights Law Review* 2006;185. Access Date: 28/10/2024.
24. Johnson H, Thompson A. The development and maintenance of post-traumatic stress disorder (PTSD) in civilian adult survivors of war trauma and torture: a review. *Clin Psychol Rev*. 2008;28(1):36-47. Access Date: 28/10/2024.
25. Hougen HP. Physical and psychological sequelae to torture. A controlled clinical study of exiled asylum applicants. *Forensic Sci Int*. 1988;39(1):5-11. Access Date: 28/10/2024.
26. Iglesias-Rios L, Harlow SD, Burgard SA, Kiss L, Zimmerman C. Mental health, violence and psychological coercion among female and male trafficking survivors in the greater Mekong sub-region: a cross-sectional study. *BMC Psychol*. 2018;6(1):56. Access Date: 28/10/2024.
27. Muldoon KA, Deering KN, Feng CX, Shoveller JA, Shannon K. Sexual relationship power and intimate partner violence among sex workers with non-commercial intimate partners in a Canadian setting. *AIDS Care*. 2015;27(4):512-9. 28/10/2024 Access Date: 28/10/2024.
28. Preventing intimate partner violence improves mental health. Geneva: WHO; 2022. Access Date: 28/10/2024.
29. The crucial role of health services in responding to gender-based violence. Washington, DC: Population Reference Bureau; 2010. Access Date: 28/10/2024.
30. García-Moreno C, Hegarty K, d'Oliveira AF, Koziol-McLain J, Colombini M, Feder G. The health-systems response to violence against women. *Lancet*. 2015;385(9977):1567-79. Access Date: 28/10/2024.

31. Accessing essential health services in fragile, conflict-affected and vulnerable settings. Geneva: WHO; 2019. Access Date: 28/10/2024.
32. Mirzazada S, Padhani ZA, Jabeen S, Fatima M, Rizvi A, Ansari U, et al. Impact of conflict on maternal and child health service delivery: a country case study of Afghanistan. *Confl Health*. 2020;14:38. Access Date: 28/10/2024.
33. Afghanistan: Events of 2021: Human Rights Watch; 2021 [Available from: <https://www.hrw.org/world-report/2022/country-chapters/afghanistan>]
34. Women's Wellbeing Survey - 2019 Findings from Sri Lanka's first dedicated National Survey on Violence against Women and Girls. Colombo: Department of Census and Statistics SL; 2020. Access Date: 28/10/2024.
35. Salikuddin T. Five things to know about Sri Lanka's Crisis: Analysis and Commentary Washington, DC: United States Institute of Peace; 2022 [Available from: <https://www.usip.org/publications/2022/07/five-things-know-about-sri-lankas-crisis>]
36. Murphy J, Michalak E, Colquhoun H, Woo C, Ng C, Parikh S, et al. Methodological approaches to situational analysis in global mental health: a scoping review. *Global Mental Health*. 2019;6:e11. Access Date: 28/10/2024.
37. Mannell J, Hawkes S. Decriminalisation of gender-based violence is a global health problem. *BMJ Glob Health*. 2017;2(3):e000438. Access Date: 28/10/2024.
38. Weobong B, Ae-Ngibise KA, Sakyi L, Lund C. Towards implementation of context-specific integrated district mental healthcare plans: A situation analysis of mental health services in five districts in Ghana. *PLoS One*. 2023;18(5):e0285324. Access Date: 28/10/2024.
39. Hanlon C, Luitel NP, Kathree T, Murhar V, Shrivasta S, Medhin G, et al. Challenges and opportunities for implementing integrated mental health care: a district level situation analysis from five low- and middle-income countries. *PLoS One*. 2014;9(2):e88437. Access Date: 28/10/2024.
40. DHS Overview: The DHS Program website: ICF; 2012 [Available from: [https://dhsprogram.com/methodology/survey-Types/dHs.cfm#:~:text=Demographic%20and%20Health%20Surveys%20\(DHS,population%2C%20health%2C%20and%20nutrition](https://dhsprogram.com/methodology/survey-Types/dHs.cfm#:~:text=Demographic%20and%20Health%20Surveys%20(DHS,population%2C%20health%2C%20and%20nutrition))]
41. Press Release on Violence against Women - The causes, Grounds, and Situation of Violence against Women in Afghanistan during 10 months of the fiscal year 1396 [press release]. Afghanistan Independent Human Rights Commission 2017. Access Date: 28/10/2024.
42. Watt A. AFGHANISTAN DEMOGRAPHIC AND HEALTH SURVEY 2015 Ministry of Public Health Wazir Akbar Khan: Central Statistics Organization; 2017. Access Date: 28/10/2024.
43. NATIONAL FAMILY HEALTH SURVEY (NFHS-4) 2015-16 INDIA. Mumbai: Government of India: Ministry of Health and Family Welfare; 2015. Access Date: 28/10/2024.
44. Patojoshi A, Sidana A, Garg S, Mishra SN, Singh LK, Goyal N, et al. Staying home is NOT 'staying safe': A rapid 8-day online survey on spousal violence against women during the COVID-19 lockdown in India. *Psychiatry Clin Neurosci*. 2021;75(2):64-6. Access Date: 28/10/2024.
45. Huq M, Das T, Devakumar D, Daruwalla N, Osrin D. Intersectional tension: a qualitative study of the effects of the COVID-19 response on survivors of violence against women in urban India. *BMJ Open*. 2021;11(9):e050381. Access Date: 28/10/2024.
46. Country profile on gender-based violence in Sri Lanka. WHO; 2018. Contract No.: CC BY-NC-SA 3.0 IGO. Access Date: 28/10/2024.

47. Jewkes R, Corboz J, Gibbs A. Violence against Afghan women by husbands, mothers-in-law and siblings-in-law/siblings: Risk markers and health consequences in an analysis of the baseline of a randomised controlled trial. *PLoS One*. 2019;14(2):e0211361. Access Date: 28/10/2024.
48. Liu NH, Daumit GL, Dua T, Aquila R, Charlson F, Cuijpers P, et al. Excess mortality in persons with severe mental disorders: a multilevel intervention framework and priorities for clinical practice, policy and research agendas. *World Psychiatry*. 2017;16(1):30-40. Access Date: 28/10/2024.
49. Fernando N, Suveendran T, de Silva C. Decentralizing provision of mental health care in Sri Lanka. *WHO South East Asia J Public Health*. 2017;6(1):18-21. Access Date: 28/10/2024.
50. Sagar R, Pattanayak RD, Chandrasekaran R, Chaudhury PK, Deswal BS, Lenin Singh RK, et al. Twelve-month prevalence and treatment gap for common mental disorders: Findings from a large-scale epidemiological survey in India. *Indian J Psychiatry*. 2017;59(1):46-55. Access Date: 28/10/2024.
51. Beydoun HA, Beydoun MA, Kaufman JS, Lo B, Zonderman AB. Intimate partner violence against adult women and its association with major depressive disorder, depressive symptoms and postpartum depression: a systematic review and meta-analysis. *Soc Sci Med*. 2012;75(6):959-75. Access Date: 28/10/2024.
52. Varma D, Chandra PS, Thomas T, Carey MP. Intimate partner violence and sexual coercion among pregnant women in India: relationship with depression and post-traumatic stress disorder. *J Affect Disord*. 2007;102(1-3):227-35. Access Date: 28/10/2024.
53. Chen LP, Murad MH, Paras ML, Colbenson KM, Sattler AL, Goranson EN, et al. Sexual abuse and lifetime diagnosis of psychiatric disorders: systematic review and meta-analysis. *Mayo Clin Proc*. 2010;85(7):618-29. Access Date: 28/10/2024.
54. Albert PR. Why is depression more prevalent in women? *Journal of Psychiatry and Neuroscience*. 2015;40(4):219. Access Date: 28/10/2024.
55. Dandona R, Kumar GA, Dhaliwal RS, Naghavi M, Vos T, Shukla DK, et al. Gender differentials and state variations in suicide deaths in India: the Global Burden of Disease Study 1990-2016. *The Lancet Public Health*. 2018;3(10):e478-e89. 2024/10/29 Access Date: 28/10/2024.
56. Patel V, Ramasundarahettige C, Vijayakumar L, Thakur JS, Gajalakshmi V, Gururaj G, et al. Suicide mortality in India: a nationally representative survey. *Lancet*. 2012;379(9834):2343-51. Access Date: 28/10/2024.
57. Dandona R, Bertozzi-Villa A, Kumar GA, Dandona L. Lessons from a decade of suicide surveillance in India: who, why and how? *Int J Epidemiol*. 2017;46(3):983-93. Access Date: 28/10/2024.
58. Pandey A SH, Dalai P, Tripathi A, Dutt K, Sinha P. Psychiatric Morbidity in Suicide attempters attending University Hospital of North India. *Delhi Psychiatry Journal*. 2013;16(1):128-33. Access Date: 28/10/2024.
59. Bhagat S, Dar M, Ahmed I. Psychiatric Co-Morbidities of Suicide Attempters: A Cross Sectional Observation in a Tertiary Care Hospital of North India. *International Journal of Science and Healthcare Research*. 2021;6:40-6. Access Date: 28/10/2024.
60. Knipe D, Williams AJ, Hannam-Swain S, Upton S, Brown K, Bandara P, et al. Psychiatric morbidity and suicidal behaviour in low- and middle-income countries: A systematic review and meta-analysis. *PLoS Med*. 2019;16(10):e1002905. Access Date: 28/10/2024.
61. Mental Health Atlas: 2011. Geneva: WHO; 2011. Access Date: 28/10/2024.

62. Dorrington S, Zavos H, Ball H, McGuffin P, Rijdsdijk F, Siribaddana S, et al. Trauma, post-traumatic stress disorder and psychiatric disorders in a middle-income setting: prevalence and comorbidity. *Br J Psychiatry*. 2014;205(5):383-9. Access Date: 28/10/2024.
63. Knipe DW, Metcalfe C, Fernando R, Pearson M, Konradsen F, Eddleston M, et al. Suicide in Sri Lanka 1975–2012: age, period and cohort analysis of police and hospital data. *BMC Public Health*. 2014;14(1):839. Access Date: 28/10/2024.
64. Bandara P, Wickrama P, Sivayokan S, Knipe D, Rajapakse T. Reflections on the trends of suicide in Sri Lanka, 1997-2022: The need for continued vigilance. *PLOS Glob Public Health*. 2024;4(4):e0003054. Access Date: 28/10/2024.
65. Committee on the Elimination of Discrimination against Women OHCHR; [Available from: <https://www.ohchr.org/en/treaty-bodies/cedaw>]
66. Convention on the Rights of Persons with Disabilities New York: United Nations: Department of Economic and Social Affairs Social Inclusion; 2006 [Available from: <https://social.desa.un.org/issues/disability/crpd/convention-on-the-rights-of-persons-with-disabilities-crpd>]
67. Darj E, Wijewardena K, Lindmark G, Axemo P. 'Even though a man takes the major role, he has no right to abuse': future male leaders' views on gender-based violence in Sri Lanka. *Glob Health Action*. 2017;10(1):1348692. Access Date: 28/10/2024.
68. India reiterates commitment to UN Declaration on violence against women [press release]. India: Deccan Chronicle 2019. Access Date: 28/10/2024.
69. The Sustainable Development Agenda New York: United Nations; 2015 [Available from: <https://www.un.org/sustainabledevelopment/development-agenda/>]
70. United Nations Convention against Transnational Organized Crime and the Protocols Thereto New York: United Nations Office on Drugs and Crime; 2000 [Available from: <https://www.unodc.org/unodc/en/organized-crime/intro/UNTOC.html>]
71. Qazi Zada S. Legislative, institutional and policy reforms to combat violence against women in Afghanistan. *Indian Journal of International Law*. 2021;59(1):257-83. Access Date: 28/10/2024.
72. Law Prohibiting Human Trafficking and Migrant Smuggling (2017). Access Date: 28/10/2024.
73. “I Thought Our Life Might Get Better” Implementing Afghanistan’s Elimination of Violence against Women Law [press release]. Washington, DC: Human Rights Watch 2021. Access Date: 28/10/2024.
74. National Action Plan for the Women of Afghanistan (Eliminating Women-directed Violence in Public and Private Spaces) The Ministry of Women's Affairs 2007. Access Date: 28/10/2024.
75. National Strategy for Mental Health 2011-2015. Kabul: Ministry of Public Health; General Directorate of Preventive Medicines; Mental health and DDR Department 2010. Access Date: 28/10/2024.
76. National Strategy for Mental Health 2019-2023. Kabul: Ministry of Public Health General Directorate of Preventive Medicine Primary Health Care Directorate MENTAL HEALTH AND SUBSTANCE ABUSE DEPARTMENT; 2018. Access Date: 28/10/2024.
77. Fundamental Rights, Directive Principles, and Fundamental Duties. Government of India; 1950. Access Date: 28/10/2024.
78. Protection of Women from Domestic Violence Act, (2005). Access Date: 28/10/2024.

79. National Policy for Women: Articulating a Vision for Empowerment of Women Government of India Ministry of Women and Child Development; 2016. Access Date: 28/10/2024.
80. FRAMEWORK FOR NIRBHAYA FUND. New Delhi: Ministry of Women and Child Development; 2015. Access Date: 28/10/2024.
81. India Passes Anti-Trafficking Bill [press release]. Global March Against Child Labour 2018. Access Date: 28/10/2024.
82. India must bring its new anti-trafficking Bill in line with human rights law, urge UN experts [press release]. India: United Nations Human Rights Office of the High Commissioner 2018. Access Date: 28/10/2024.
83. Gupta S, Sagar R. National Mental Health Policy, India (2014): Where Have We Reached? Indian Journal of Psychological Medicine. 2021;44(5):510-5. 2024/10/28 Access Date: 28/10/2024.
84. Special Rapporteur on Violence against Women: Submission related to Shelters for Homeless Women from Housing and Land Rights Network (HLRN), India. Housing and Land Rights Network (India); 2015. Access Date: 28/10/2024.
85. One Stop Centre Scheme. New Delhi: Ministry of Women and Child Development; 2017. Access Date: 28/10/2024.
86. Policy Framework and National Plan of Action to address Sexual and Gender - Based Violence (SGBV) in Sri Lanka 2016 - 2020 Ministry of Women & Child Affairs and Social Development, United Nations Development Programme; 2021. Access Date: 28/10/2024.
87. Sri Lanka: Standard DHS, 2016. Ministry of Health and Indigenous Medicine; 2016. Access Date: 28/10/2024.
88. 2018 Trafficking in Persons Report. US Department of State; 2018. Access Date: 28/10/2024.
89. Optimizing Screening and Support Services for Gender-Based Violence and Trafficking in Person Victims: Sri Lanka. Colombo: The Asia Foundation; 2021. Access Date: 28/10/2024.
90. The Mental Health Policy of Sri Lanka 2005-2015, (2005). Access Date: 28/10/2024.
91. National Mental Health Strategy 2009-2014: For a Mentally Healthy Afghanistan. Kabul: Islamic Republic of Afghanistan Ministry of Public Health General Directorate of Preventive Medicines Mental Health & DDR Department; 2009. Access Date: 28/10/2024.
92. Kovess-Masfety V, Karam E, Keyes K, Sabawoon A, Sarwari BA. Access to Care for Mental Health Problems in Afghanistan: A National Challenge. Int J Health Policy Manag. 2022;11(8):1442-50. Access Date: 28/10/2024.
93. Mental health atlas 2017 Geneva: WHO; 2018. Access Date: 28/10/2024.
94. National Mental Health Programme (NMHP): Ministry of Health & Family Welfare Government of India; [Available from: <https://nhm.gov.in/index1.php?lang=1&level=2&sublinkid=1043&lid=359>]
95. National Mental Health Survey of India. National Institute of Mental Health and Neurosciences and Ministry of Health and Family Welfare Govt of India; 2017. Access Date: 28/10/2024.
96. Murthy RS. National Mental Health Survey of India 2015-2016. Indian J Psychiatry. 2017;59(1):21-6. Access Date: 28/10/2024.
97. Grover S, Raju VV, Sharma A, Shah R. Depression in Children and Adolescents: A Review of Indian studies. Indian J Psychol Med. 2019;41(3):216-27. Access Date: 28/10/2024.

98. Petersen I, Marais D, Abdulmalik J, Ahuja S, Alem A, Chisholm D, et al. Strengthening mental health system governance in six low- and middle-income countries in Africa and South Asia: challenges, needs and potential strategies. *Health Policy Plan*. 2017;32(5):699-709. Access Date: 28/10/2024.
99. Mugisha J, Abdulmalik J, Hanlon C, Petersen I, Lund C, Upadhaya N, et al. Health systems context(s) for integrating mental health into primary health care in six Emerald countries: a situation analysis. *Int J Ment Health Syst*. 2017;11:7. Access Date: 28/10/2024.
100. Shidhaye R, Raja A, Shrivastava S, Murhar V, Ramaswamy R, Patel V. Challenges for Transformation: A Situational Analysis of Mental Health Care Services in Sehore District, Madhya Pradesh. *Community Ment Health J*. 2015;51(8):903-12. Access Date: 28/10/2024.
101. Kaur R, Pathak RK. Treatment Gap in Mental Healthcare: Reflections from Policy and Research. *Economic and Political Weekly*. 2017;52(31):34-40.2024/10/28/ Access Date: 28/10/2024.
102. Arvind BA, Gururaj G, Loganathan S, Amudhan S, Varghese M, Benegal V, et al. Prevalence and socioeconomic impact of depressive disorders in India: multisite population-based cross-sectional study. *BMJ Open*. 2019;9(6):e027250. Access Date: 28/10/2024.
103. Firth J, Siddiqi N, Koyanagi A, Siskind D, Rosenbaum S, Galletly C, et al. The Lancet Psychiatry Commission: a blueprint for protecting physical health in people with mental illness. *Lancet Psychiatry*. 2019;6(8):675-712. Access Date: 28/10/2024.
104. Patel V, Saxena S, Lund C, Thornicroft G, Baingana F, Bolton P, et al. The Lancet Commission on global mental health and sustainable development. *The Lancet*. 2018;392(10157):1553-98.2024/10/28 Access Date: 28/10/2024.
105. 'HOW INDIA PERCEIVES MENTAL HEALTH': 2018 NATIONAL SURVEY REPORT: Live, Love, Laugh Foundation; 2018 [Available from: <https://www.thelivelovelaughfoundation.org/initiatives/research/how-india-perceives-mental-health>]
106. Dandona R. Mind and body go together: the need for integrated care. *Lancet Psychiatry*. 2019;6(8):638-9. Access Date: 28/10/2024.
107. Barik D, Thorat A. Issues of Unequal Access to Public Health in India. *Front Public Health*. 2015;3:245. Access Date: 28/10/2024.
108. Carpenter J. The history of mental health care in Sri Lanka. Sri Lanka: Marga Publications; 1988. Access Date: 28/10/2024.
109. Rajasuriya M, Hewagama M, Ruwanpriya S, Wijesundara H. Community-Based Psychiatric Services in Sri Lanka: a Model by WHO in the Making. *Consort Psychiatr*. 2021;2(4):40-52. Access Date: 28/10/2024.
110. Traunmüller R, Kijewski S, Freitag M. The Silent Victims of Sexual Violence during War: Evidence from a List Experiment in Sri Lanka. *Journal of Conflict Resolution*. 2019;63(9):2015-42.2024/10/28 Access Date: 28/10/2024.
111. Farrell A, Dank, M., Kafafian, M., Lockwood, S., Pfeffer, R. Capturing Human Trafficking Victimization Through Crime Reporting. Washington, DC: U.S. Department of Justice; 2019. Contract No.: 252520 Access Date: 28/10/2024.
112. Jayasuriya V, Wijewardena K, Axemo P. Intimate partner violence against women in the capital province of Sri Lanka: prevalence, risk factors, and help seeking. *Violence Against Women*. 2011;17(8):1086-102. Access Date: 28/10/2024.

113. Violence against women and girls data collection during COVID-19 NYC: UN Women; 2020 [Available from: <https://www.unwomen.org/en/digital-library/publications/2020/04/issue-brief-violence-against-women-and-girls-data-collection-during-covid-19>
114. Boserup B, McKenney M, Elkbuli A. Alarming trends in US domestic violence during the COVID-19 pandemic. *Am J Emerg Med*. 2020;38(12):2753-5. Access Date: 28/10/2024.
115. Roesch E, Amin A, Gupta J, García-Moreno C. Violence against women during covid-19 pandemic restrictions. *Bmj*. 2020;369:m1712. Access Date: 28/10/2024.
116. Global Mental Health: Principles and Practice: Oxford University Press; 2013. Available from: <https://doi.org/10.1093/med/9780199920181.001.0001>
117. Alegría M, NeMoyer A, Falgàs Bagué I, Wang Y, Alvarez K. Social Determinants of Mental Health: Where We Are and Where We Need to Go. *Curr Psychiatry Rep*. 2018;20(11):95. Access Date: 28/10/2024.
118. Tsai AC, Tomlinson M, Comulada WS, Rotheram-Borus MJ. Intimate Partner Violence and Depression Symptom Severity among South African Women during Pregnancy and Postpartum: Population-Based Prospective Cohort Study. *PLoS Med*. 2016;13(1):e1001943. Access Date: 28/10/2024.
119. Greene MC, Rees S, Likindikoki S, Bonz AG, Joscelyne A, Kaysen D, et al. Developing an integrated intervention to address intimate partner violence and psychological distress in Congolese refugee women in Tanzania. *Confl Health*. 2019;13:38. Access Date: 28/10/2024.
120. Plichta SB. Interactions between victims of intimate partner violence against women and the health care system: policy and practice implications. *Trauma Violence Abuse*. 2007;8(2):226-39. Access Date: 28/10/2024.
121. Jewkes R. Intimate partner violence: causes and prevention. *Lancet*. 2002;359(9315):1423-9. Access Date: 28/10/2024.
122. Kim HW, Salyers MP. Attitudes and perceived barriers to working with families of persons with severe mental illness: mental health professionals' perspectives. *Community Ment Health J*. 2008;44(5):337-45. Access Date: 28/10/2024.
123. Rodríguez M, Valentine JM, Son JB, Muhammad M. Intimate partner violence and barriers to mental health care for ethnically diverse populations of women. *Trauma Violence Abuse*. 2009;10(4):358-74. Access Date: 28/10/2024.
124. Armstrong G, Kermode M, Raja S, Suja S, Chandra P, Jorm AF. A mental health training program for community health workers in India: impact on knowledge and attitudes. *Int J Ment Health Syst*. 2011;5(1):17. Access Date: 28/10/2024.
125. Patel V, Weobong B, Weiss HA, Anand A, Bhat B, Katti B, et al. The Healthy Activity Program (HAP), a lay counsellor-delivered brief psychological treatment for severe depression, in primary care in India: a randomised controlled trial. *The Lancet*. 2017;389(10065):176-85. 2024/10/28 Access Date: 28/10/2024.
126. Lund C, Tomlinson M, Patel V. Integration of mental health into primary care in low- and middle-income countries: the PRIME mental healthcare plans. *Br J Psychiatry*. 2016;208 Suppl 56(Suppl 56):s1-3. Access Date: 28/10/2024.
127. Shidhaye R, Baron E, Murhar V, Rathod S, Khan A, Singh A, et al. Community, facility and individual level impact of integrating mental health screening and treatment into the primary healthcare system in Sehore district, Madhya Pradesh, India. *BMJ Glob Health*. 2019;4(3):e001344. Access Date: 28/10/2024.
128. Olson RM, García-Moreno C, Colombini M. The implementation and effectiveness of the one stop centre model for intimate partner and sexual violence in low- and middle-income

countries: a systematic review of barriers and enablers. *BMJ Glob Health*. 2020;5(3):e001883. Access Date: 28/10/2024.

129. Daruwalla N, Das T, Punekar S, Patil S, Manjrekar S, Pantvaidya S, et al. Global Health Research. Support needs of survivors of violence against women in urban India: a prospective analysis of client records. Southampton (UK): National Institute for Health and Care Research Copyright © 2024 Daruwalla et al.; 2024 Access Date: 28/10/2024.

130. Palfreyman A, Vijayaraj K, Riyaz S, Rizwan Z, Sivayokan S, Thenakoon THS, et al. What Women Want: Mental Health in the Context of Violence Against Women in Sri Lanka-A Qualitative Study of Priorities and Capacities for Care. *Violence Against Women*. 2024;10778012241230326. Access Date: 28/10/2024.

131. Kaul A, Saboor L, Ahmad A, Mannell J, Paphitis SA, Devakumar D. What are the experiences and psychosocial needs of female survivors of domestic violence in Afghanistan? A qualitative interview study in three Afghan provinces. *BMJ Open*. 2024;14(6):e079615. Access Date: 28/10/2024.