



Opinion paper

The importance of protein sources to support muscle anabolism in cancer: An expert group opinion



Katherine L. Ford^a, Jann Arends^b, Philip J. Atherton^c, Mariëlle P.K.J. Engelen^d,
Thiago J.M. Gonçalves^e, Alessandro Laviano^f, Dileep N. Lobo^{g,h}, Stuart M. Phillipsⁱ,
Paula Ravasco^{j,k}, Nicolaas E.P. Deutz^{d,**}, Carla M. Prado^{a,*}

^a Human Nutrition Research Unit, Department of Agricultural, Food & Nutritional Science, University of Alberta, Edmonton, Canada

^b Department of Medicine I, Medical Center - University of Freiburg, Faculty of Medicine, University of Freiburg, Germany

^c MRC Versus Arthritis Centre of Excellence for Musculoskeletal Ageing Research, Centre of Metabolism & Physiology (COMAP), University of Nottingham, Derby, UK

^d Center for Translational Research in Aging & Longevity, Department of Health and Kinesiology, Texas A&M University, College Station, TX, USA

^e Department of Nutrology and Clinical Nutrition, Sancta Maggiore Hospital, Prevent Senior Institute, São Paulo, Brazil

^f Department of Translational and Precision Medicine, Sapienza University of Rome, Rome, Italy

^g Gastrointestinal Surgery, Nottingham Digestive Diseases Centre, National Institute for Health Research (NIHR) Nottingham Biomedical Research Centre, Nottingham University Hospitals NHS Trust and University of Nottingham, Queen's Medical Centre, Nottingham, UK

^h MRC Versus Arthritis Centre for Musculoskeletal Ageing Research, School of Life Sciences, University of Nottingham, Queen's Medical Centre, Nottingham, UK

ⁱ Department of Kinesiology, McMaster University, Hamilton, Canada

^j Católica Medical School and Centre for Interdisciplinary Research in Health (CIIS), Universidade Católica Portuguesa, Lisbon, Portugal

^k Centre for Interdisciplinary Research Egas Moniz (CiEM), Egas Moniz Cooperativa de Ensino Superior, CRL, Almada, Portugal

ARTICLE INFO

Article history:

Received 19 October 2021

Accepted 24 November 2021

Keywords:

Cancer
Protein source
Animal protein
Muscle
Anabolism
Plant protein

SUMMARY

This opinion paper presents a short review of the potential impact of protein on muscle anabolism in cancer, which is associated with better patient outcomes. Protein source is a topic of interest for patients and clinicians, partly due to recent emphasis on the supposed non-beneficial effect of proteins; therefore, misconceptions involving animal-based (e.g., meat, fish, dairy) and plant-based (e.g., legumes) proteins in cancer are acknowledged and addressed. Although the optimal dietary amino acid composition to support muscle health in cancer is yet to be established, animal-based proteins have a composition that offers superior anabolic potential, compared to plant-derived proteins. Thus, animal-based foods should represent the majority (i.e., $\geq 65\%$) of protein intake during active cancer treatment. A diet rich in plant-derived proteins may support muscle anabolism in cancer, albeit requiring a larger quantity of protein to fulfill the optimal amino acid intake. We caution that translating dietary recommendations for cancer prevention to cancer treatment may be inadequate to support the pro-inflammatory and catabolic nature of the disease. We further caution against initiating an exclusively plant-based (i.e., vegan) diet upon a diagnosis of cancer, given the presence of elevated protein requirements and risk of inadequate protein intake to support muscle anabolism. Amino acid combination and the long-term sustainability of a dietary pattern void of animal-based foods requires careful and laborious management of protein intake for patients with cancer. Ultimately, a dietary amino acid composition that promotes muscle anabolism is optimally obtained through combination of animal- and plant-based protein sources.

© 2021 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Abbreviations: AICR, American Institute for Cancer Research; CI, confidence interval; DIAAS, digestible indispensable amino acid score; MPB, muscle protein breakdown; MPS, muscle protein synthesis; PDCAAS, protein digestibility corrected amino acid score; RDA, recommended dietary allowance; WCRF, World Cancer Research Fund.

* Corresponding author. 4-002 Li Ka Shing Center for Health Research Innovation, University of Alberta, Edmonton T6G 2P5, Canada.

** Corresponding author. Center for Translational Research in Aging & Longevity, Dept. of Health and Kinesiology, Texas A&M University, College Station, TX, USA.

E-mail addresses: nep.deutz@ctrnl.org (N.E.P. Deutz), carla.prado@ualberta.ca (C.M. Prado).

<https://doi.org/10.1016/j.clnu.2021.11.032>

0261-5614/© 2021 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Cancer is a leading cause of mortality worldwide and is the main cause of premature death in much of the Western Hemisphere and Western Europe [1]. Early and continued optimization of nutritional status, including elevated protein requirements [2,3], is crucial to prevent and minimize negative health outcomes (e.g., muscle loss) often observed in cancer. Optimal nutrition is a hallmark of successful cancer treatment as it can alleviate symptom burden, improve health and quality of life, and support survivorship [4–7].

Many people with cancer recognize the importance of nutrition and are often motivated to make lifestyle changes and improve their dietary choices [8–10]. However, access to regulated nutrition professionals in outpatient cancer centers is sparse, and only those with, or at significant risk for, depleted nutritional status are likely to be referred to a registered dietitian/nutritionist [11]. As such, the burden to seek dietary advice is often placed on the patient.

Self-guided dietary changes, including restricting or eliminating animal foods (e.g., meat and/or dairy) are common in cancer [9,12] and often used by patients in an attempt to “cure the cancer” or alleviate symptoms [13,14]. A Dutch study showed that people with cancer ($n = 239$) reported decreasing their meat intake and increasing intake of plant-based foods following a cancer diagnosis [15]. Similarly, a group of 1458 patients with stage I–IV colorectal cancer reported several dietary changes, including decreased meat and increased fruit, vegetables, fibers, wholegrains and fish consumption [14]. Decreased meat intake ($n = 376$) was more prevalent than increased fish consumption ($n = 342$), although these dietary changes were not quantified in relation to total protein intake [14], and thus the effect of habitual dietary change on protein intake was unclear. A study of the NutriNet-Santé cohort ($n = 696$) found that changes following a diagnosis of cancer included decreased vegetable, dairy, meat, soy, and alcohol consumption which cumulatively resulted in significantly lower total protein intake (-17.4 ± 12.5 g/day; $p < 0.0001$), compared with pre-diagnosis [16]. Dietary changes post-diagnosis resulting in decreased protein intake have also been observed in patients with breast cancer [17]. Although some of these changes are beneficial to overall health (e.g., decreased alcohol consumption), a diet containing exclusively (i.e., vegan diet) or predominantly (i.e., vegetarian diet) plant-based foods is concerning due primarily to the importance of animal-based protein for skeletal muscle health.

Acknowledging that healthy populations in the Americas, Europe, and Oceania achieve much of their protein intake from animal sources [18], the above-mentioned dietary changes should not be overlooked, nor the consequences underestimated. Aside from veganism, all diets contain a mixture of animal- and plant-based proteins. Initiating a dietary pattern that restricts or eliminates animal-based foods without careful consideration for increased protein needs may hinder optimal nutritional status following a diagnosis of cancer, especially related to the ability to sustain muscle mass. Dietary proteins provide essential amino acids required for muscle health. Muscle loss is a prominent problem experienced by people with cancer despite the increasing prevalence of obesity [19,20] and impacts prognosis and clinical outcomes [21–23]. Loss of muscle is a defining feature of malnutrition, which has several consequences ranging from economic [24] to physical burdens [23], and also of cancer cachexia, a debilitating multifactorial syndrome that results in functional debilitation and mortality [25]. Muscle loss occurs to varying degrees across stages and types of disease, body weight or weight stability/change, and leads to detrimental clinical outcomes, including lower tolerance to treatment and decreased survival [20–23,26]. In fact, low muscle mass is emerging as a clinical marker of biological age,

the latter of which has been proposed as an important consideration when prescribing anti-cancer therapies [27].

Given the importance of muscle health in the oncologic setting and the notion that cancer is a catalyst for dietary change, we aimed to address the potential impact of protein sources (i.e., animal- and plant-based) on muscle anabolism in cancer and suggest an optimal ratio of intake. Herein, animal-based proteins include beef, pork, chicken, fish, eggs, milk, cheese, etc. This opinion paper is a narrative review and expert group understanding of available data. Search term categories included ‘protein’, ‘muscle’, and ‘cancer’. Key concepts are summarized in Box 1.

2. Protein intake and muscle mass

Whole-body skeletal muscle mass is dependent on rates of muscle protein synthesis (MPS) and muscle protein breakdown (MPB), collectively termed muscle protein turnover [28–30]. In a healthy state, MPS and MPB are constantly changing in relation to food intake to maintain muscle mass [30]. To achieve muscle anabolism (i.e., growth), MPS on average must chronically exceed MPB to obtain a positive net protein balance. The homeostatic state of muscle protein turnover is disrupted in pro-inflammatory conditions such as cancer [30,31]. Upregulation of ubiquitin-proteasome/autophagy pathways [32] and a decline in MPS [33] results in increased degradation of intracellular proteins and subsequently loss of muscle mass [34]. Reduced protein intake because of inflammation-related anorexia and the adverse effects of cancer therapy further contribute to muscle loss [35,36]. In addition, muscle protein synthesis is greatly affected by protein intake and level of physical activity [29,37–40], which are often diminished in cancer [41–43], further exacerbating catabolism.

The nutritional value of protein is determined by the quantity and quality of constituent amino acids [44]. Amino acids are the dietary anabolic drivers of muscle mass accretion but vary in quality and do not equally promote anabolism [45,46]. The Protein Digestibility Corrected Amino Acid Score (PDCAAS) is indicative of essential amino acid content and digestibility of proteins [47]. Since the PDCAAS was developed, another measure of protein quality was introduced: The Digestible Indispensable Amino Acid Score (DIAAS) [48]. Notably, these scores do not suggest true skeletal muscle anabolic response to a particular amino acid but do provide a proxy method of quality comparison between proteins [28]. Dietary proteins have varying amino acid profiles whereby animal-based proteins offer greater anabolic stimuli when compared with plant-based alternatives [28,49,50]. Plant-based proteins are less digestible (i.e., lower PDCAAS) than animal-based proteins [47,51]. Other key differences between animal and plant proteins are highlighted in Fig. 1.

Soy protein isolate (i.e., a purified form of soy) is the plant-based protein that is an exception (high PDCAAS) as the absence of anti-nutritional factors in the isolated form of soy increases the digestibility of this product compared to soy protein in whole foods [51,52]. As with whole foods, animal-based protein supplements (e.g., whey) offer increased MPS and superior anabolic potential compared with plant-based protein powders, including soy protein isolate (PDCAAS 1.0) [28,53]. The decreased anabolic potential of plant-proteins (including soy) may be due to increased amino acid oxidation, decreased MPS, and lower leucine content compared to animal proteins [49,53]. The presence of anti-nutritional factors (e.g., trypsin inhibitors, tannins, phytates) may negatively impact digestibility and availability of amino acids in plant-proteins [52]. Plant-based proteins resist proteolysis in the gastrointestinal tract (i.e., decreasing their digestibility) due to their difference in structure when compared with animal-based proteins [54], which may be especially concerning for people suffering from gastrointestinal cancers (e.g., small bowel, pancreas, gastro-esophageal).

Box 1**Key Messages**

- People with cancer need more protein than healthy individuals.
- Animal proteins provide greater anabolic stimuli compared with plant-based protein foods and are, therefore, better for muscle health.
- A combination of animal ($\geq 65\%$ of protein intake) and plant proteins is likely to be optimal for supporting muscle health and avoiding malnutrition during the treatment of cancer.
- Eliminating animal proteins (i.e., vegan diet) is not a recommended dietary change to pursue during active treatment of cancer.
- During active treatment of cancer, the goals of nutritional intake shift and do not necessarily parallel the recommendations for cancer prevention and post-treatment.
- Theoretical arguments suggesting that nutrition feeds the tumor are not supported by evidence and should not be a reason to alter food choice.
- Individuals following well planned and balanced exclusively (i.e., vegan) or predominantly (i.e., vegetarian) plant-based diets for personal reasons (e.g., religious, ethical) may be able to support muscle health during treatment of cancer although a professional nutrition assessment is highly recommended.

Given the equally high protein quality (PDCAAS 1.0) of soy protein isolate as most animal proteins (e.g., whey), the role of soy for muscle health is a topic of interest [28,55]. Regardless, soy protein appears to be inferior to animal-based proteins at stimulating MPS and overall muscle anabolism in healthy older adults [49,53]. Leucine is a key regulator [45] and possibly the sole stimulator [56–59] of MPS. Leucine is also the most potent stimulator, compared with other essential amino acids, of the mTOR pathway – an essential anabolic pathway [46]. Animal-based proteins generally contain more leucine than plant-based proteins except for maize (corn), which in spite of its high leucine content is not a complete protein [18]. The role of collagenous animal proteins in muscle health has also been investigated. Collagen is an anomaly in the context of animal-based proteins because it completely lacks the essential amino acid tryptophan, rendering it incomplete and thus having a DIAAS of 0 [51,60]. A review of the literature suggests that MPS does not appear to be stimulated with collagen supplementation [51] although acute experiments with collagen also are difficult to interpret as tryptophan attached to albumin can temporarily provide tryptophan [61]. Future research in this area is needed; at this time, we do not recommend collagen supplementation for the specific purpose of muscle anabolism. Notably, despite the potential role of specific protein sources or nutrients, most dietary guidelines consider total protein intake rather than the amino acid composition of dietary protein.

3. Anabolic potential and protein source in cancer

The anabolic potential of skeletal muscle during cancer is controversial; studies indicate both anabolic resistance [62,63] and retained anabolic potential [19,62,64,65]. The etiology of cancer-associated muscle loss is complex and not fully understood despite its well-documented prevalence [20,63,64]. Cancer therapy often includes surgery, which can have a drastic and negative

impact on skeletal muscle. The loss of muscle in the perioperative period can stem from bedrest, the catabolic stress of the surgery itself, and diminished nutritional intake [63]. A small study in patients with colorectal cancer found that compared to healthy counterparts, those with cancer presented significantly less lean mass (which includes muscle mass) in their lower extremities before surgery [63]. Compared with the pre-operative period, those undergoing tumor resection had further lower lean leg mass loss post-operatively [63]. The benefits of nutrition therapy post-operatively and upon discharge continue to be doubted despite increasing acceptance of Enhanced Recovery After Surgery [66]. Although perioperative nutritional therapy alone may not alleviate all nutritional challenges in patients undergoing surgery for cancer, marginal gains achieved throughout the cancer trajectory have the capacity to improve clinically relevant outcomes [66]. Hence, small improvements in nutritional status should not be discounted if they do not amount to clinically meaningful changes in isolation as the additive effect of small improvements over time have the potential to bring about clinically significant positive change [66].

Animal-based proteins are of major importance during the active treatment of cancer to preclude detrimental loss of muscle and promote muscle anabolism. Only a few amino acid kinetic studies investigated whole-body protein synthesis and balance in cancer over the last decade [62,67–69], as reviewed by Antoun & Raynard [70]. Bozetti & Bozetti [71] previously reviewed the same topic and cited studies that suggested increased [72–74] or decreased/no change [73] in whole-body protein synthesis




Fig. 1. Visual comparison of select key nutritional differences between animal and plant proteins. Legend: ↑: higher; ↓: lower; ↔: variable; ∅: none. Images retrieved from smart.servier.com.

following amino acid infusions in persons with cancer. Although these studies forecast the effects of mimicking whole-food diets on muscle change, translating results from amino acid kinetics to whole-body anabolism is difficult, as whole-body protein turnover does not necessarily equate to skeletal muscle protein anabolism [70]. A study using a cachectic pig model that compared the effects of a dairy with dairy/plant combination diet on skeletal muscle anabolic response to feeding found that despite the diets having equal leucine content, the dairy/plant diet was not effective in inducing an anabolic response to feeding; in contrast, the whey protein diet was [75]. A study investigating the effects of leucine on tumor-bearing mice suggested that supplemental leucine may protect muscle from disease-induced wasting [76], although to our knowledge, a similar study is yet to be conducted in humans.

The role of the mTOR pathway in mediating amino acid-induced skeletal muscle anabolism is well-established. Nevertheless, the mTORC1 pathway is also involved in negative forms of anabolism, including tumor growth, such that some fear nutritionally-derived anabolic stimuli (e.g., protein) may also fuel or be associated with tumor growth [77]. Despite amino acids having heterogeneous effects on tumor growth in humans, the effect of protein intake on tumor growth has not been substantiated [78]. In general, international guidelines on nutrition in cancer acknowledge that theoretical arguments suggesting that nutrition feeds the tumor are not supported by evidence and should not be a reason to alter nutrition delivery [2].

4. Protein in nutrition oncology guidelines (during curative or palliative cancer treatment)

Protein intake in cancer is highly variable, and many patients do not meet the minimum recommended intake [41,42,79,80]. Nutritional oncology guidelines recommend a minimum intake of 1.0 g protein/kg body weight/d but suggest a target consumption of 1.2–2.0 g/kg/d [2]. These guidelines are similar to those for older adults, which recommend at least 1.0–1.2 g/kg/d, acknowledging that those with acute or chronic illness require more protein (1.2–1.5 g/kg/d) [81]. Given that targeting guideline-based protein levels with individualized nutrition support improves clinical outcomes in cancer [7] and that increased total protein intake in adults over the age of 65 years (similar to the median age of a cancer diagnosis) has a protective effect [82], it appears that total protein intake should be a co-primary consideration in addition to protein quality. Notably, historical concerns regarding the supposed negative impact of protein on kidney health are unfounded. Higher protein intakes (≥ 2.0 g/kg/d) are safe for people with healthy kidney function [2,44] and may reduce mortality in critically ill patients [83], although those with pre-existing kidney disease should maintain a lower intake [84,85].

Protein intake recommendations in cancer are notably higher than the recommended dietary allowance (RDA) of 0.8 g/kg/d for the healthy population, determined by nitrogen balance studies [86]. These studies primarily used high-quality proteins with a PDCAAS of 1.0 (e.g., animal-based proteins or soy protein isolate) [28,86–88]. Based on the methodology used to determine the RDA, this value should be considered a minimum amount needed to attain nitrogen balance rather than an amount sufficient to promote muscle maintenance or anabolism [44]. Conversely, the recommendations for patients with cancer are primarily based on expert opinions given the paucity of studies that investigated nitrogen balance or the impact of protein intake on clinical outcomes [2]. As oncology recommendations were derived from protein metabolism studies [2,62,89,90], the guidelines also acknowledge that the optimal amino acid composition for patients with cancer remains unknown [2].

The risk of malnutrition varies in people with cancer [91], especially between the curative and palliative setting, although the potential benefit of animal protein is present given the association of low muscle mass with malnutrition [23]. Studies employing isotopic tracer methods are needed to determine specific amino acid requirements in oncology settings [92]; for example, the indicator amino acid oxidation method is one technique that could be used to determine total protein requirements in a non-invasive manner [93]. Additionally, challenges of understanding optimal protein quantity and amino acid composition for muscle mass maintenance or anabolism are compounded by gut dysfunction, which is observed in patients with cancer, resulting in decreased protein digestion of oral food intake and absorption [94,95]. The reduction in protein digestion negatively impacts systemic amino acid availability and leads to increased quantity of undigested proteins in the colon [94,95]. The latter can alter microbial metabolism and generate harmful metabolites which may negatively affect muscle health [94,95].

5. Additional considerations and current evidence of animal- and plant-based protein intake during cancer treatment

One of the ten World Cancer Research Fund and the American Institute for Cancer Research (WCRF/AICR) recommendations for *prevention* of cancer is to follow the remaining nine recommendations for those *diagnosed* with cancer [96]. Translating dietary recommendations for prevention of cancer to patients with active cancer may provide insufficient nutritional targets and, therefore, suboptimal nutritional status. Some dietary strategies for prevention of cancer might result in worse outcomes once a diagnosis of cancer is made. Although controversial, red meat may be positively associated with the incidence of colon cancer, although the association may be inversely related to colon cancer mortality [97]. In a cohort of 992 patients with stage III colon cancer, restricted red and processed meat intake was associated with increased risk of death [97]. Additionally, diet quality was not associated with overall survival in a cohort of 1284 patients with metastatic colorectal cancer [98]. We propose that once a diagnosis of cancer is made, the goals of nutritional intake shift and do not necessarily parallel the recommendations for cancer prevention and post-treatment (survivors).

When considering protein sources, one concern regarding an exclusively, or even predominantly, plant-based diet during active treatment of cancer is the feasibility of obtaining adequate dietary amino acid intake to sustain functional muscle reserves, especially given the high risk of malnutrition in this population [91] (Fig. 3). Cancer therapy is frequently accompanied by nutrition-impact symptoms (e.g., nausea, anorexia, taste alterations) that can affect food intake and compound muscle catabolism [99,100]. Early satiation can also contribute to decreased oral intake and may be influenced by nutrients in the diet (e.g., protein, fiber). As discussed, protein intake is essential for muscle health in cancer, although caloric intake is also vital for optimizing nutrition in this vulnerable population. In older adults, essential amino acid supplementation has been proposed as a complementary measure, in addition to protein intake, that does not impact satiety, optimizes the ability to meet nutritional requirements, and promotes muscle health [101]. The efficacy of essential amino acid supplementation requires further investigation in clinical settings, including cancer. Similarly, although the satiating effect of protein has been studied in other populations, how a predominantly plant-vs. animal-based diet affects satiety is unknown in people with cancer or at risk of malnutrition [102]. Regardless of satiating effects, a larger volume of plant-based proteins than animal-based products is required to obtain adequate amino acid intake [103]. It follows that the higher

quality of animal-based proteins provides adequate protein intake from a smaller volume of food [28], as shown in Fig. 2.

Despite some insight into protein intake in this population, studies investigating protein quality or types of protein consumed are lacking [2]. In healthy middle-aged women, those consuming ~68% of their protein from animal sources (animal:plant protein intake ratio 2.09) had significantly higher muscle mass compared with vegetarians consuming ~55% of their protein from animal sources (animal:plant protein intake ratio 1.23) [104]. Given the paucity of research on optimal ratio of protein sources in cancer, interpretations can be drawn from studies in populations at similar risk of malnutrition as those undergoing treatment of cancer to provide insight and a starting point into appropriate nutrition needed to optimize health in cancer. In older adults with comorbidities, at least 65% of protein intake from high-quality protein (i.e., animal-based protein) is needed to avoid malnutrition [105]. Additionally, factors similar to those seen in people undergoing treatment of cancer (e.g., missing a meal, taste alterations) were independently associated with inadequate intake of ≥ 1 essential amino acid and subsequently greater risk of malnutrition [105]. Given the paucity of this type of research in cancer and the similarity in malnutrition risk and nutrition impact symptoms between populations, we propose that inferences be drawn and a minimum of 65% of protein intake from animal sources be considered as an optimal starting point to support muscle anabolism for people undergoing active cancer treatment. Future trials should seek to determine the optimal animal:plant protein ratio to support muscle mass in cancer.

Ongoing trials are investigating protein needs [106] and the clinical impact of increased protein [107,108] or amino acid [109,110] intake on muscle in people with cancer. The impact of

protein on muscle strength is also an important consideration in cancer given that muscle weakness can occur without loss of muscle mass [31]. A review that focused on nutritional interventions for muscle strength in cancer found no studies that compared plant-with animal-based diet interventions [111]. Clinical trials related to protein source and muscle anabolism in cancer primarily focused on MPS and MPB rather than whole-body lean mass response and its impact on clinical outcomes. A paradox exists in current research whereby humans consume predominately whole foods, yet research has focused on specific amino acids and their contribution to muscle protein turnover. Pragmatic studies employing a whole-food approach exploring the influence of protein sources on whole-body skeletal muscle anabolism are needed to guide future nutrition recommendations.

It is our opinion that in catabolic disease states such as cancer, the attributes of animal-based proteins contribute to optimal nutrition care and can be safely included in the diet. We acknowledge various reasons (ethical, religious, planetary, health, etc.) for choosing a plant-based diet. Those already consuming a balanced exclusively or predominantly plant-based diet may achieve adequate nutritional intake to support health, although appropriate knowledge of diet diversity is needed to ensure higher protein needs are met [112,113]. Initiating an unbalanced exclusively or predominantly plant-based dietary pattern during active treatment of cancer may impact negatively on the ability to achieve optimal protein intake. In contrast, a dietary pattern that combines protein from animal- and plant-based sources is, in our view, more likely to be the most suitable option for optimal health.

Lastly, although not the focus of this paper, and regardless of protein sources in the diet, exercise is a viable proponent of a multimodal approach to supporting muscle health in cancer. It is




Fig. 2. Select nutritional differences between animal and plant proteins are highlighted with food examples. Animal proteins include beef, pork, chicken, fish, eggs, milk, cheese, etc. Plant proteins include beans, lentils, soy, nuts, etc. Legend: ↑: higher; ↓: lower; ↔: variable; ∅: none. Images retrieved from [smart.servier.com](https://www.smart.servier.com). Canadian Nutrient File food codes: beef – 6112; beans – 7085.




Fig. 3. Flowchart of important nutrition-related considerations based on health status. Legend: ↑: increased; ↓: decreased; ↔: neutral; animal proteins: beef, pork, chicken, fish, eggs, milk, cheese, etc.; plant proteins: beans, lentils, soy, nuts, etc. Images retrieved from [smart.servier.com](https://www.smart.servier.com).

safe during and after treatment and is encouraged [2,114]. The role of resistance exercise and nutrition to mitigate muscle loss in pro-catabolic states has been extensively reviewed, emphasizing the importance of a multimodal approach to muscle health [115–117].

6. Dietary guidelines for healthy populations (for cancer prevention and post-treatment)

National dietary guidelines for healthy populations (including prevention of cancer and post-treatment) have increasingly emphasized plant-based foods for overall health, chronic disease prevention, and sustainability [118–121]. The WCRF/AICR has ten recommendations related to diet, nutrition, and physical activity for prevention of cancer. In line with select national dietary guidelines worldwide, the WCRF/AICR recommends limiting consumption of red and processed meat [96]. Although guidelines explicitly state not to avoid meat consumption entirely, the layperson may be inclined to adopt an exclusively or predominantly plant-based diet based on recommendations for cancer prevention. A review of the literature could not confirm that substituting animal-based proteins with plant-based options reduced the risk of developing cancer [122]. The limited and conflicting evidence of protein sources challenges the ability to draw confident conclusions; thus, the positive and negative effects of these types of foods must be weighed in relation to health priorities [123–126]. Elimination of specific foods or food groups without

supportive substitution is concerning as nutritional considerations of dietary intake are complex and should not be dichotomized as ‘good’ or ‘bad’. Protein needs for prevention of cancer and post-cancer treatment may not be elevated compared with protein needs during active cancer treatment, therefore the former can likely be reasonably achieved with predominantly plant-based foods given the lower risk for malnutrition and muscle loss in this population (Fig. 3). Thus, obtaining ≥65% of protein from animal sources may not be essential to sustain muscle mass in this healthy population. In general, protein intake from animal- and plant-based sources is highly variable. One study in healthy adults found that self-reported vegetarians ($n = 2370$) consumed 50.9% of their protein (g) from animal sources, whereas meat eaters ($n = 90,664$) ingested 70.8% of protein from animal sources [127]. Regardless of self-reported diet strategies, the definition of a predominantly plant-based diet remains controversial, compounding the challenge of determining optimal ratios of protein intake based on nutritional needs.

An important consideration is that the response of skeletal muscle to dietary anabolic stimuli may be blunted with age and, thus, older adults may have increased protein needs [81]. Moreover, increased protein intake is protective against mortality in adults over the age of 65 years [82]. Additional considerations for older adults with cancer include age-associated changes in appetite regulation, lack of hunger, a decline in taste and smell, reduced central and peripheral drive to eat, delayed gastric emptying and deteriorating dentition [128]. These are why animal protein intake

decreases with age, and the feasibility of adequate plant-protein intake in older adults may be compromised [49].

7. General benefits of plant-based proteins

Dietary guidelines for chronic disease prevention do not account for the source of protein, despite its importance given the diverse protein intake patterns observed worldwide [18]. As illustrated in Figs. 1 and 2, plant-based proteins are higher in essential fatty acids and fiber, whereas they are lower in protein, essential amino acids, and certain micronutrients (e.g., iron, vitamin B₁₂, zinc) [129].

Globally, plant-based options are the predominant source of protein [18,49]. Compared with animal counterparts, plant-based proteins are more environmentally sustainable, requiring fewer resources (e.g., land and water) and producing lower levels of greenhouse-gas emissions [119,130]. As such, many people are adopting an exclusively or predominantly plant-based diet, regardless of health status [119,131]. For example, national and international oncology groups support incorporating soy-based foods as a part of a healthy diet for prevention of cancer and during active treatment [132,133]. The high isoflavone content of soy continues to foster unfounded concerns regarding a potential link between soy intake and negative health outcomes relating to hormone production [122,132,134]. Soy intake has not been linked to cancer occurrence or recurrence; conversely, the isoflavones have anti-inflammatory properties that may decrease the risk of cancer occurrence [122,135].

In the absence of disease, it is possible to maintain a healthy diet without consuming animal products if eliminating the latter is substituted with alternatives that contain adequate essential nutrients [136]. Importantly, a regulated health care professional who is a nutrition expert (e.g., registered dietitian/nutritionist) should be consulted before making any drastic dietary changes. These professionals can counsel patients and provide strategies to achieve adequate intake, especially those who are already following an exclusively or predominantly plant-based diet while undergoing oncologic treatment.

8. Conclusion

We argue that the inclusion of anabolic-stimulating animal proteins like beef, pork, chicken, fish, eggs, milk, cheese, etc. is key to achieving clinical benefits related to muscle anabolism in cancer, especially given the higher protein needs due to the pro-inflammatory nature of the disease, low intake levels, and deconditioning of these patients. Drastic changes to food choice such as eliminating animal products from the diet may negatively interfere with the quality and quantity of protein intake if not properly substituted in the diet and do not align with international nutritional oncology guidelines. Without adequate substitution with other foods, nutritional status may be compromised if meat-derived proteins are reduced or eliminated. We support the increased incorporation of plant-based proteins. However, we caution that a dichotomous approach to eating that eliminates animal products may lead to inferior anabolic stimuli and greater quantity and variability of protein sources needed to achieve skeletal muscle anabolism, especially once diagnosed with cancer. Thus, we propose that in the oncologic setting, a minimum of 65% of protein intake should be derived from animal sources as an initial starting point although further research in this area is needed. Following successful cancer treatment, increased consumption of plant-based proteins may be adequate to sustain muscle health but requires careful planning if substituting animal products in the diet. Dietary changes can be optimized with the help of a registered dietitian/nutritionist.

Funding statement

This work is partially supported by CMP's Campus Alberta Innovation Program funding from the Government of Alberta.

Author contributions

All authors were responsible for conceptualization, writing and review/editing the manuscript. K.L.F wrote the first draft.

Conflict of interest

K.L.F has no conflicts to declare.

J.A. reports receiving honoraria from Baxter, B. Braun, Berg-Apotheke, Falk, Fresenius-Kabi, Helsinn, Nestlé and Nutricia.

P.J.A. reports consultancy fees and research income from Abbott Nutrition and Fresenius-Kabi.

M.P.K.J.E has no conflicts to declare.

T.J.M.G. reports receiving consulting fees for honoraria for lectures at industry-sponsored events; consulting fees from Nestlé Health Science and Nutricia.

A.L. reports receiving consulting fees for honoraria for lectures at industry-sponsored events; consulting fees from Abbott, Baxter, BBraun, Fresenius Kabi, Nestlé Health Science, Nutricia, and Smartfish; and research grant from Fresenius Kabi.

D.N.L. has no conflicts to declare.

S.M.P. reports grants from US National Dairy Council, during the conduct of the study; personal fees from the US National Dairy Council, non-financial support from Enhanced Recovery, outside the submitted work; In addition, Dr. Phillips has a patent Canadian 3052324 issued to Exerkine, and a patent US 20200230197 pending to Exerkine but reports no financial gains.

P.R. has no conflicts to declare.

N.E.P.D reports grants from Abbott Nutrition, Department of Defense, and National Institutes of Health, and honoraria from Abbott Nutrition.

C.M.P. reports receiving honoraria and/or paid consultancy from Abbott Nutrition, Nutricia, Nestle Health Science, Fresenius Kabi, Pfizer, and Helsinn.

Acknowledgements

We thank Camila Orsso for her assistance with the figures.

References

- [1] Bray F, Laversanne M, Weiderpass E, Soerjomataram I. The ever-increasing importance of cancer as a leading cause of premature death worldwide. *Cancer* 2021;127:3029–30. <https://doi.org/10.1002/cnrc.33587>.
- [2] Arends J, Bachmann P, Baracos V, Barthelemy N, Bertz H, Bozzetti F, et al. ESPEN guidelines on nutrition in cancer patients. *Clin Nutr* 2017;36:11–48. <https://doi.org/10.1016/j.clnu.2016.07.015>.
- [3] Roeland EJ, Bohlke K, Baracos VE, Bruera E, Fabbro E Del, Dixon S, et al. Management of cancer cachexia: ASCO guideline. *J Clin Oncol* 2020;38:2438–53. <https://doi.org/10.1200/JCO.20.00611>.
- [4] Prado CM, Anker SD, Coats AJS, Laviano A, von Haehling S. Nutrition in the spotlight in cachexia, sarcopenia and muscle: avoiding the wildfire. *J Cachexia Sarcopenia Muscle* 2021;12:3–8. <https://doi.org/10.1002/jcsm.12673>.
- [5] Ravasco P, Monteiro-Grillo I, Camilo ME. Does nutrition influence quality of life in cancer patients undergoing radiotherapy? *Radiother Oncol* 2003;67:213–20. [https://doi.org/10.1016/S0167-8140\(03\)00040-9](https://doi.org/10.1016/S0167-8140(03)00040-9).
- [6] de van der Schueren MAE, Laviano A, Blanchard H, Jourdan M, Arends J, Baracos VE. Systematic review and meta-analysis of the evidence for oral nutritional intervention on nutritional and clinical outcomes during chemo(radio)therapy: current evidence and guidance for design of future trials. *Ann Oncol* 2018;29:1141–53. <https://doi.org/10.1093/annonc/mdy114>.
- [7] Bargetzi L, Brack C, Herrmann J, Bargetzi A, Hersberger L, Bargetzi M, et al. Nutritional support during the hospital stay reduces mortality in patients with different types of cancers: secondary analysis of a prospective

- randomized trial. *Ann Oncol* 2021;32:1025–33. <https://doi.org/10.1016/j.annonc.2021.05.793>.
- [8] van Zutphen M, Boshuizen HC, Kok DE, van Baar H, Geijssen AJMR, Wesselink E, et al. Colorectal cancer survivors only marginally change their overall lifestyle in the first 2 years following diagnosis. *J Cancer Surviv* 2019;13:956–67. <https://doi.org/10.1007/s11764-019-00812-7>.
 - [9] Sullivan ES, Rice N, Kingston E, Kelly A, Reynolds JV, Feighan J, et al. A national survey of oncology survivors examining nutrition attitudes, problems and behaviours, and access to dietetic care throughout the cancer journey. *Clin Nutr ESPEN* 2021;41:331–9. <https://doi.org/10.1016/j.clnesp.2020.10.023>.
 - [10] Gavazzi C, Sieri S, Tracò F, Sproviero A, Vandoni G, Ricci R, et al. Changes in food habits in cancer patients in Italy: an AIOM-SINPE-FAVO survey. *Nutrition* 2018;55–56:140–5. <https://doi.org/10.1016/j.nut.2018.04.002>.
 - [11] Trujillo EB, Claghorn K, Dixon SW, Hill EB, Braun A, Lipinski E, et al. Inadequate nutrition coverage in outpatient cancer centers: results of a national survey. *J Oncol* 2019;2019:7462940. <https://doi.org/10.1155/2019/7462940>.
 - [12] Ghelfi F, Tieri M, Gori S, Nicolis F, Petrella MC, Filiberti A, et al. Do cancer patients change their diet in the e-health information era? A review of the literature and a survey as a proposal for the Italian population. *Food Res Int* 2018;104:59–68. <https://doi.org/10.1016/j.foodres.2017.10.021>.
 - [13] Salminen EK, Lagström HK, Heikkilä SP, Salminen SJ. Does breast cancer change patients' dietary habits? *Eur J Clin Nutr* 2000;54:844–8. <https://doi.org/10.1038/sj.ejcn.1601103>.
 - [14] Bours MJ, Beijer S, Winkels RM, Van Duijnhoven FJ, Mols F, Breedveld-Peters JJ, et al. Dietary changes and dietary supplement use, and underlying motives for these habits reported by colorectal cancer survivors of the Patient Reported Outcomes Following Initial Treatment and Long-Term Evaluation of Survivorship (PROFILES) registry. *Br J Nutr* 2015;114:286–96. <https://doi.org/10.1017/S0007114515001798>.
 - [15] van Veen MR, Winkels RM, Janssen SHM, Kampman E, Beijer S. Nutritional information provision to cancer patients and their relatives can promote dietary behavior changes independent of nutritional information needs. *Nutr Cancer* 2018;70:483–9. <https://doi.org/10.1080/01635581.2018.1446092>.
 - [16] Fassier P, Zelek L, Lécuyer L, Bachmann P, Touillaud M, Druenes-Pecollo N, et al. Modifications in dietary and alcohol intakes between before and after cancer diagnosis: results from the prospective population-based NutriNet-Santé cohort. *Int J Cancer* 2017;141:457–70. <https://doi.org/10.1002/ijc.30704>.
 - [17] Brunnvoll SH, Thune I, Bertheussen GF, Fjeldheim F, Flote VG, Frydenberg H, et al. Dietary changes in early-stage breast cancer patients from pre-surgery and over the 12 months post-surgery. *Br J Nutr* 2021;125:172–82. <https://doi.org/10.1017/S0007114520002627>.
 - [18] FAO UN Statistics Division. FAOSTAT food balance sheets. Available at: <http://www.fao.org/faostat/en/#search/protein>. [Accessed 17 October 2021].
 - [19] Engelen MPKJ, Van Der Meij BS, Deutz NEP. Protein anabolic resistance in cancer: does it really exist? *Curr Opin Clin Nutr Metab Care* 2016;19:39–47. <https://doi.org/10.1097/MCO.0000000000000236>.
 - [20] Prado CM, Lieffers JR, McCargar LJ, Reiman T, Sawyer MB, Martin L, et al. Prevalence and clinical implications of sarcopenic obesity in patients with solid tumours of the respiratory and gastrointestinal tracts: a population-based study. *Lancet Oncol* 2008;9:629–35. [https://doi.org/10.1016/S1470-2045\(08\)70153-0](https://doi.org/10.1016/S1470-2045(08)70153-0).
 - [21] Martin L, Birdsell L, MacDonald N, Reiman T, Clandinin MT, McCargar LJ, et al. Cancer cachexia in the age of obesity: skeletal muscle depletion is a powerful prognostic factor, independent of body mass index. *J Clin Oncol* 2013;31:1539–47. <https://doi.org/10.1200/JCO.2012.45.2722>.
 - [22] Xiao J, Caan BJ, Cespedes Feliciano EM, Meyerhardt JA, Kroenke CH, Baracos VE, et al. The association of medical and demographic characteristics with sarcopenia and low muscle radiodensity in patients with nonmetastatic colorectal cancer. *Am J Clin Nutr* 2019;109:615–25. <https://doi.org/10.1093/ajcn/nqy328>.
 - [23] Deutz NEP, Ashurst I, Ballesteros MD, Bear DE, Cruz-Jentoft AJ, Genton L, et al. The underappreciated role of low muscle mass in the management of malnutrition. *J Am Med Dir Assoc* 2019;20:22–7. <https://doi.org/10.1016/j.jamda.2018.11.021>.
 - [24] Snider JT, Linthicum MT, Wu Y, Lavallee C, Lakdawalla DN, Hegazi R, et al. Economic burden of community-based disease-associated malnutrition in the United States. *J Parenter Enteral Nutr* 2014;38:775–855. <https://doi.org/10.1177/0148607114550000>.
 - [25] Fearon K, Strasser F, Anker SD, Bosaeus I, Bruera E, Fainsinger RL, et al. Definition and classification of cancer cachexia: an international consensus. *Lancet Oncol* 2011;12:489–95. [https://doi.org/10.1016/S1470-2045\(10\)70218-7](https://doi.org/10.1016/S1470-2045(10)70218-7).
 - [26] Brown JC, Caan BJ, Cespedes Feliciano EM, Xiao J, Weltzien E, Prado CM, et al. Weight stability masks changes in body composition in colorectal cancer: a retrospective cohort study. *Am J Clin Nutr* 2021;113:1482–9. <https://doi.org/10.1093/ajcn/nqaa440>.
 - [27] Mandelblatt JS, Ahles TA, Lippman ME, Isaacs C, Adams-Campbell L, Saykin AJ, et al. Applying a life course biological age framework to improving the care of individuals with adult cancers: review and research recommendations. *JAMA Oncol* 2021. <https://doi.org/10.1001/jamaoncol.2021.1160>. Epub ahead of print.
 - [28] van Vliet S, Burd NA, van Loon LJC. The skeletal muscle anabolic response to plant- versus animal-based protein consumption. *J Nutr* 2015;145:1981–91. <https://doi.org/10.3945/jn.114.204305>.
 - [29] Wolfe RR. Regulation of muscle protein by amino acids. *J Nutr* 2002;132:3219S–24S. <https://doi.org/10.1093/jn/131.10.3219S>.
 - [30] Phillips SM, Glover EI, Rennie MJ. Alterations of protein turnover underlying disuse atrophy in human skeletal muscle. *J Appl Physiol* 2009;107:645–54. <https://doi.org/10.1152/japplphysiol.00452.2009>.
 - [31] van der Meij BS, Deutz NEP, Rodriguez RER, Engelen MPKJ. Increased amino acid turnover and myofibrillar protein breakdown in advanced cancer are associated with muscle weakness and impaired physical function. *Clin Nutr* 2019;38:2399–407. <https://doi.org/10.1016/j.clnu.2018.10.022>.
 - [32] Acharyya S, Guttridge DC. Cancer cachexia signaling pathways continue to emerge yet much still points to the proteasome. *Clin Cancer Res* 2007;13:1356–61. <https://doi.org/10.1158/1078-0432.CCR-06-2307>.
 - [33] Hardee JP, Montalvo RN, Carson JA. Linking cancer cachexia-induced anabolic resistance to skeletal muscle oxidative metabolism. *Oxid Med Cell Longev* 2017;2017:8018197. <https://doi.org/10.1155/2017/8018197>.
 - [34] Lecker SH, Goldberg AL, Mitch WE. Protein degradation by the ubiquitin-proteasome pathway in normal and disease states. *J Am Soc Nephrol* 2006;17:1807–19. <https://doi.org/10.1681/ASN.2006010083>.
 - [35] van der Meij BS, Teleni L, Engelen MPKJ, Deutz NEP. Amino acid kinetics and the response to nutrition in patients with cancer. *Int J Radiat Biol* 2019;95:480–92. <https://doi.org/10.1080/09553002.2018.1466209>.
 - [36] Dodson S, Baracos VE, Jatoi A, Evans WJ, Cella D, Dalton JT, et al. Muscle wasting in cancer cachexia: clinical implications, diagnosis, and emerging treatment strategies. *Annu Rev Med* 2011;62:265–79. <https://doi.org/10.1146/annurev-med-061509-131248>.
 - [37] Koopman R, Walrand S, Beelen M, Gijzen AP, Kies AK, Boirie Y, et al. Dietary protein digestion and absorption rates and the subsequent postprandial muscle protein synthetic response do not differ between young and elderly men. *J Nutr* 2009;139:1707–13. <https://doi.org/10.3945/jn.109.109173>.
 - [38] Paddon-Jones D, Short KR, Campbell WW, Volpi E, Wolfe RR. Role of dietary protein in the sarcopenia of aging. *Am J Clin Nutr* 2008;87:1562S–65. <https://doi.org/10.1093/ajcn/87.5.1562S>.
 - [39] Phillips SM. Nutrient-rich meat proteins in offsetting age-related muscle loss. *Meat Sci* 2012;92:174–8. <https://doi.org/10.1016/j.meatsci.2012.04.027>.
 - [40] Deutz NEP, Safar A, Schutzler S, Memelink R, Ferrando A, Spencer H, et al. Muscle protein synthesis in cancer patients can be stimulated with a specially formulated medical food. *Clin Nutr* 2011;30:759–68. <https://doi.org/10.1016/j.clnu.2011.05.008>.
 - [41] Hasegawa Y, Iijichi H, Saito K, Ishigaki K, Takami M, Sekine R, et al. Protein intake after the initiation of chemotherapy is an independent prognostic factor for overall survival in patients with unresectable pancreatic cancer: a prospective cohort study. *Clin Nutr* 2021;40:4792–8. <https://doi.org/10.1016/j.clnu.2021.06.011>.
 - [42] Lawson C, Ferreira V, Carli F, Chevalier S. Effects of multimodal prehabilitation on muscle size, myosteatosis, and dietary intake of surgical patients with lung cancer – a randomized feasibility study. *Appl Physiol Nutr Metab* 2021;46:1407–16. <https://doi.org/10.1139/apnm-2021-0249>.
 - [43] Smith WA, Nolan VG, Robison LL, Hudson MM, Ness KK. Physical activity among cancer survivors and those with no history of cancer – a report from the national health and nutrition examination survey 2003–2006. *Am J Transl Res* 2011;3:342–50.
 - [44] Wu G. Dietary protein intake and human health. *Food Funct* 2016;7:1251–65. <https://doi.org/10.1039/c5fo01530h>.
 - [45] Mitchell WK, Wilkinson DJ, Phillips BE, Lund JN, Smith K, Atherton PJ. Human skeletal muscle protein metabolism responses to amino acid nutrition. *Adv Nutr* 2016;7:828S–38S. <https://doi.org/10.3945/an.115.011650>.
 - [46] Atherton PJ, Smith K, Etheridge T, Rankin D, Rennie MJ. Distinct anabolic signalling responses to amino acids in C2C12 skeletal muscle cells. *Amino Acids* 2010;38:1533–9. <https://doi.org/10.1007/s00726-009-0377-x>.
 - [47] FAO/WHO. Protein quality evaluation: report of the joint FAO/WHO expert consultation 1989. Rome, Italy: FAO; 1991.
 - [48] FAO. Dietary protein evaluation in human nutrition: report of an FAO expert consultation. Rome, Italy: FAO; 2013. Available at: <http://www.fao.org/ag/humannutrition/35978-02317b979a686a57aa4593304ffc17f06.pdf>. [Accessed 26 September 2021].
 - [49] Berrazaga I, Micard V, Gueugneau M, Walrand S. The role of the anabolic properties of plant-versus animal-based protein sources in supporting muscle mass maintenance: a critical review. *Nutrients* 2019;11:1825. <https://doi.org/10.3390/nu11081825>.
 - [50] Park S, Church DD, Schutzler SE, Azhar G, Kim IY, Ferrando AA, et al. Metabolic evaluation of the dietary guidelines' ounce equivalents of protein food sources in young adults: a randomized controlled trial. *J Nutr* 2021;151:1190–6. <https://doi.org/10.1093/jn/nxaa401>.
 - [51] Deane CS, Bass JJ, Crossland H, Phillips BE, Atherton PJ. Animal, plant, collagen and blended dietary proteins: effects on musculoskeletal outcomes. *Nutrients* 2020;12:2670. <https://doi.org/10.3390/nu12092670>.
 - [52] Gilani GS, Xiao CW, Cockell KA. Impact of antinutritional factors in food proteins on the digestibility of protein and the bioavailability of amino acids and on protein quality. *Br J Nutr* 2012;108:S315–32. <https://doi.org/10.1017/S0007114512002371>.
 - [53] Yang Y. Myofibrillar protein synthesis following ingestion of soy protein isolate at rest and after resistance exercise in elderly men. *Nutr Metab* 2012;9:57–61. <https://doi.org/10.1186/1473-7075-9-57>.
 - [54] Carbonaro M, Maselli P, Nucara A. Relationship between digestibility and secondary structure of raw and thermally treated legume proteins: a Fourier

- transform infrared (FT-IR) spectroscopic study. *Amino Acids* 2012;43:911–21. <https://doi.org/10.1007/s00726-011-1151-4>.
- [55] Putra C, Konow N, Gage M, York CG, Mangano KM. Protein source and muscle health in older adults: a literature review. *Nutrients* 2021;13:373. <https://doi.org/10.3390/nu13030743>.
- [56] Devries MC, Mcglory C, Bolster DR, Kamil A, Rahn M, Harkness L, et al. Leucine, not total protein, content of a supplement is the primary determinant of muscle protein anabolic responses in healthy older women. *J Nutr* 2018;148:1088–95. <https://doi.org/10.1093/jn/nxy091>.
- [57] Devries MC, Mcglory C, Bolster DR, Kamil A, Rahn M, Harkness L. Protein leucine content is a determinant of shorter- and longer-term muscle protein synthetic responses at rest and following resistance exercise in healthy older women: a randomized, controlled trial. *Am J Clin Nutr* 2018;107:217–26. <https://doi.org/10.1093/ajcn/nqx028>.
- [58] Wilkinson DJ, Bukhari SSI, Phillips BE, Limb MC, Cegielski J, Brook MS, et al. Effects of leucine-enriched essential amino acid and whey protein bolus dosing upon skeletal muscle protein synthesis at rest and after exercise in older women. *Clin Nutr* 2018;37:2011–21. <https://doi.org/10.1016/j.clnu.2017.09.008>.
- [59] Bukhari SSI, Phillips BE, Wilkinson DJ, Limb MC, Rankin D, Mitchell WK, et al. Intake of low-dose leucine-rich essential amino acids stimulates muscle anabolism equivalently to bolus whey protein in older women at rest and after exercise. *Am J Physiol Endocrinol Metab* 2015;308:E1056–65. <https://doi.org/10.1152/ajpendo.00481.2014>.
- [60] Oikawa SY, Kamal MJ, Webb EK, McGlory C, Baker SK, Phillips SM. Whey protein but not collagen peptides stimulate acute and longer-term muscle protein synthesis with and without resistance exercise in healthy older women: a randomized controlled trial. *Am J Clin Nutr* 2020;111:708–18. <https://doi.org/10.1093/ajcn/nqz332>.
- [61] Ten Have GAM, Engelen MPKJ, Soeters PB, Deutz NEP. Absence of post-prandial gut anabolism after intake of a low quality protein meal. *Clin Nutr* 2012;31:273–82. <https://doi.org/10.1016/j.clnu.2011.09.008>.
- [62] Winter A, MacAdams J, Chevalier S. Normal protein anabolic response to hyperaminoacidemia in insulin-resistant patients with lung cancer cachexia. *Clin Nutr* 2012;31:765–73. <https://doi.org/10.1016/j.clnu.2012.05.003>.
- [63] Williams JP, Phillips BE, Smith K, Atherton PJ, Rankin D, Selby AL, et al. Effect of tumor burden and subsequent surgical resection on skeletal muscle mass and protein turnover in colorectal cancer patients. *Am J Clin Nutr* 2012;96:1064–70. <https://doi.org/10.3945/ajcn.112.045708>.
- [64] Prado CM, Sawyer MB, Ghosh S, Lieffers JR, Esfandiari N, Antoun S, et al. Central tenet of cancer cachexia therapy: do patients with advanced cancer have exploitable anabolic potential? *Am J Clin Nutr* 2013;98:1012–9. <https://doi.org/10.3945/ajcn.113.060228.1>.
- [65] Chevalier S, Winter A. Do patients with advanced cancer have any potential for protein anabolism in response to amino acid therapy? *Curr Opin Clin Nutr Metab Care* 2014;17:213–8. <https://doi.org/10.1097/MCO.0000000000000047>.
- [66] Adimah A, Lobo DN. Post-discharge oral nutritional supplementation after surgery for gastrointestinal cancer: real or marginal gains? *Clin Nutr* 2021;40:1–3. <https://doi.org/10.1016/j.clnu.2020.06.001>.
- [67] Engelen MPKJ, Safar AM, Barrant T, Koeman F, Deutz NEP. High anabolic potential of essential amino acid mixtures in advanced nonsmall cell lung cancer. *Ann Oncol* 2015;26:1960–6. <https://doi.org/10.1093/annonc/mdv271>.
- [68] van Dijk DPJ, van de Poll MCG, Moses AGW, Preston T, Olde Damink SWM, Rensen SS, et al. Effects of oral meal feeding on whole body protein breakdown and protein synthesis in cachectic pancreatic cancer patients. *J Cachexia Sarcopenia Muscle* 2015;6:212–21. <https://doi.org/10.1002/jcsm.12029>.
- [69] Engelen MPKJ, Klimberg VS, Allasia A, Deutz NEP. Presence of early stage cancer does not impair the early protein metabolic response to major surgery. *J Cachexia Sarcopenia Muscle* 2017;8:447–56. <https://doi.org/10.1002/jcsm.12173>.
- [70] Antoun S, Raynard B. Muscle protein anabolism in advanced cancer patients: response to protein and amino acids support, and to physical activity. *Ann Oncol* 2018;29(Suppl 2). <https://doi.org/10.1093/annonc/mdx809.ii10–7>.
- [71] Bozzetti F, Bozzetti V. Is the intravenous supplementation of amino acid to cancer patients adequate? A critical appraisal of literature. *Clin Nutr* 2013;32:142–6. <https://doi.org/10.1016/j.clnu.2012.10.017>.
- [72] Shaw J, Humberstone D, Douglas R, Koea J. Leucine kinetics in patients with benign disease, non-weight-losing cancer, and cancer cachexia: studies at the whole-body and tissue level and the response to nutritional support. *Surgery* 1991;109:37–50.
- [73] Hylander A, Warnold I, Edén E, Lundholm K. Effect on whole-body protein synthesis after institution of intravenous nutrition in cancer and non-cancer patients who lose weight. *Eur J Cancer* 1991;27:16–21. [https://doi.org/10.1016/0277-5379\(91\)90051-E](https://doi.org/10.1016/0277-5379(91)90051-E).
- [74] Bozzetti F, Gavazzi C, Ferrari P, Dworak F. Effect of total parenteral nutrition on the protein kinetics of patients with cancer cachexia. *Tumori* 2000;86:408–11. <https://doi.org/10.1177/030089160008600508>.
- [75] Revel A, Jarzagueta M, Peyron MA, Papet I, Hafnaoui N, Migné C, et al. At same leucine intake, a whey/plant protein blend is not as effective as whey to initiate a transient post prandial muscle anabolic response during a catabolic state in mini pigs. *PLoS One* 2017;12:e0186204. <https://doi.org/10.1371/journal.pone.0186204>.
- [76] Peters SJ, Van Helvoort A, Kegler D, Argilès JM, Luiking YC, Laviano A, et al. Dose-dependent effects of leucine supplementation on preservation of muscle mass in cancer cachectic mice. *Oncol Rep* 2011;26:247–54. <https://doi.org/10.3892/or.2011.1269>.
- [77] Ham DJ, Caldwell MK, Lynch GS. Leucine as a treatment for muscle wasting: a critical review. *Clin Nutr* 2014;33:937–45. <https://doi.org/10.1016/j.clnu.2014.09.016>.
- [78] Bossola M, Pacelli F, Rosa F, Tortorelli A, Doglietto GB. Does nutrition support stimulate tumor growth in humans? *Nutr Clin Pract* 2011;26:174–80. <https://doi.org/10.1177/0884533611399771>.
- [79] Hutton JL, Martin L, Field CJ, Wismer WV, Bruera ED, Watanabe SM, et al. Dietary patterns in patients with advanced cancer: implications for anorexia-cachexia therapy. *Am J Clin Nutr* 2006;84:1163–70. <https://doi.org/10.1093/ajcn/84.5.1163>.
- [80] Prado CMM, Lieffers JR, Bergsten G, Mourtzakis M, Baracos VE, Reiman T, et al. Dietary patterns of patients: with advanced lung or colorectal cancer. *Can J Diet Pract Res* 2012;73:e298–303. <https://doi.org/10.3148/73.4.2012.e298>.
- [81] Deutz NEP, Bauer JM, Barazzoni R, Biolo G, Boirie Y, Bosy-Westphal A, et al. Protein intake and exercise for optimal muscle function with aging: recommendations from the ESPEN Expert Group. *Clin Nutr* 2014;33:929–36. <https://doi.org/10.1016/j.clnu.2014.04.007>.
- [82] Levine M, Suarez J, Brandhorst S, Balasubramanian P, Cheng C, Madia F, et al. Low protein intake is associated with a major reduction in IGF-1, cancer, and overall mortality in the 65 and younger but not older population. *Cell Metabol* 2014;19:407–17. <https://doi.org/10.1016/j.cmet.2014.02.006>.
- [83] Zhu R, Allingstrup MJ, Perner A, Doig GS. The effect of IV amino acid supplementation on mortality in ICU patients may be dependent on kidney function: post hoc subgroup analyses of a multicenter randomized trial. *Crit Care Med* 2018;46:1293–301. <https://doi.org/10.1097/CCM.0000000000003221>.
- [84] Martin WF, Armstrong LE, Rodriguez NR. Dietary protein intake and renal function. *Nutr Metab* 2005;2:25. <https://doi.org/10.1186/1743-7075-2-25>.
- [85] Fiaccadori E, Sabatino A, Barazzoni R, Carrero JJ, Cupisti A, De Waele E, et al. ESPEN guideline on clinical nutrition in hospitalized patients with acute or chronic kidney disease. *Clin Nutr* 2021;40:1644–68. <https://doi.org/10.1016/j.clnu.2021.01.028>.
- [86] Rand WM, Pellett PL, Young VR. Meta-analysis of nitrogen balance studies for estimating protein requirements in healthy adults. *Am J Clin Nutr* 2003;77:109–27. <https://doi.org/10.1093/ajcn/77.1.109>.
- [87] Institute of Medicine. Dietary reference intakes for energy, carbohydrate, fiber, fat, fatty acids, cholesterol, protein, and amino acids (macronutrients). Washington: Institute of Medicine; 2005. <https://doi.org/10.17226/10490>.
- [88] World Health Organization. Protein and amino acid requirements in human nutrition: report of a joint WHO/FAO/UNU expert consultation. Geneva, Switzerland: World Health Organization; 2002. Available at: https://apps.who.int/iris/bitstream/handle/10665/43411/WHO_TRS_935_eng.pdf;jsessionid=F8E540F0997B2FC32EB290268D816461?sequence=1. [Accessed 26 September 2021].
- [89] Baracos VE. Skeletal muscle anabolism in patients with advanced cancer. *Lancet Oncol* 2015;16:13–4. [https://doi.org/10.1016/S1470-2045\(14\)71185-4](https://doi.org/10.1016/S1470-2045(14)71185-4).
- [90] Guadagni M, Biolo G. Effects of inflammation and/or inactivity on the need for dietary protein. *Curr Opin Clin Nutr Metab Care* 2009;12:617–22. <https://doi.org/10.1097/MCO.0b013e32833193bd>.
- [91] Arends J, Baracos V, Bertz H, Bozzetti F, Calder PC, Deutz NEP, et al. ESPEN expert group recommendations for action against cancer-related malnutrition. *Clin Nutr* 2017;36:1187–96. <https://doi.org/10.1016/j.clnu.2017.06.017>.
- [92] Prado CM, Purcell SA, Laviano A. Nutrition interventions to treat low muscle mass in cancer. *J Cachexia Sarcopenia Muscle* 2020;11:366–80. <https://doi.org/10.1002/jcsm.12525>.
- [93] Elango R, Ball RO, Pencharz PB. Indicator amino acid oxidation: concept and application. *J Nutr* 2008;138:243–6. <https://doi.org/10.1093/jn/138.2.243>.
- [94] Kirschner SK, Deutz NEP, Engelen MPKJ. Intestinal dysfunction in chronic disease. *Curr Opin Clin Nutr Metab Care* 2021;24:464–72. <https://doi.org/10.1097/mco.0000000000000780>.
- [95] van der Meij BS, Deutz NEP, Rodriguez RE, Engelen MPKJ. Early signs of impaired gut function affect daily functioning in patients with advanced cancer undergoing chemotherapy. *J Parenter Enteral Nutr* 2021;45:752–60. <https://doi.org/10.1002/jpen.1941>.
- [96] World Cancer Research Fund/American Institute for Cancer Research. Recommendations and public health and policy implications. London: World Cancer Research Fund International; 2018. Available at: <https://www.wcrf.org/wp-content/uploads/2021/01/Recommendations.pdf>. [Accessed 26 September 2021].
- [97] Van Blarigan EL, Fuchs CS, Niedzwiecki D, Zhang S, Saltz LB, Mayer RJ, et al. Association of survival with adherence to the American Cancer Society nutrition and physical activity guidelines for cancer survivors after colon cancer diagnosis the CALGB 89803/Alliance trial. *JAMA Oncol* 2018;4:783–90. <https://doi.org/10.1001/jamaoncol.2018.0126>.
- [98] Van Blarigan EL, Fuchs CS, Niedzwiecki D, Zhang S, Saltz LB, Mayer RJ, et al. Association of diet quality with survival among people with metastatic colorectal cancer in the Cancer and Leukemia B and Southwest Oncology Group 80405 Trial. *JAMA Netw Open* 2020;3:e2023500. <https://doi.org/10.1001/jamanetworkopen.2020.23500>.
- [99] Sanchez-Lara K, Sosa-Sanchez R, Green-Renner D, Rodríguez C, Laviano A, Motola-Kuba D, et al. Influence of taste disorders on dietary behaviors in

- cancer patients under chemotherapy. *Nutr J* 2010;9:15. <https://doi.org/10.1186/1475-2891-9-15>.
- [100] Coa KL, Epstein JB, Ettinger D, Jatoti A, McManus K, Platek ME, et al. The impact of cancer treatment on the diets and food preferences of patients receiving outpatient treatment. *Nutr Cancer* 2015;67:339–53. <https://doi.org/10.1080/01635581.2015.990577>.
 - [101] Ispoglou T, Witard OC, Duckworth LC, Lees MJ. The efficacy of essential amino acid supplementation for augmenting dietary protein intake in older adults: implications for skeletal muscle mass, strength and function. *Proc Nutr Soc* 2021;80:230–42. <https://doi.org/10.1017/S0029665120008010>.
 - [102] Lonnie M, Hooker E, Brunstrom JM, Corfe BM, Green MA, Watson AW, et al. Protein for life: review of optimal protein intake, sustainable dietary sources and the effect on appetite in ageing adults. *Nutrients* 2018;10:360. <https://doi.org/10.3390/nu10030360>.
 - [103] Pinckaers PJM, Trommelen J, Snijders T, van Loon LJC. The anabolic response to plant-based protein ingestion. *Sports Med* 2021;26–31. <https://doi.org/10.1007/s40279-021-01540-8>.
 - [104] Aubertin-Leheudre M, Adlercreutz H. Relationship between animal protein intake and muscle mass index in healthy women. *Br J Nutr* 2009;102:1803–10. <https://doi.org/10.1017/S0007114509991310>.
 - [105] Dasgupta M, Sharkey JR, Wu G. Inadequate intakes of indispensable amino acids among homebound older adults. *J Nutr Elder* 2005;24:85–99. https://doi.org/10.1300/J052v24n03_07.
 - [106] Clinicaltrials.gov. Identifier NCT04144907, protein needs in cancer [Internet]. Available at: <https://clinicaltrials.gov/ct2/show/NCT04144907>. [Accessed 17 October 2021].
 - [107] Ford KL, Sawyer MB, Trottier CF, Ghosh S, Deutz NEP, Siervo M, et al. Protein Recommendation to Increase Muscle (PRIME): study protocol for a randomized controlled pilot trial investigating the feasibility of a high protein diet to halt loss of muscle mass in patients with colorectal cancer. *Clin Nutr ESPEN* 2021;41:175–85. <https://doi.org/10.1016/j.clnesp.2020.11.016>.
 - [108] German Clinical Trials Register. Identifier DRKS00015744, Can oral protein intake of 1.5g/kg bw/day improve nutritional status, serum albumin levels, lean muscle mass and quality of life in patients under adjuvant chemotherapy after Kausch-Whipple surgery for pancreatic head cancer? [Internet]. Available at: <http://www.drks.de/DRKS00015744%0A>. [Accessed 17 October 2021].
 - [109] ANZCTR Australian New Zealand clinical trials registry. Identifier ACTRN12616001402437, Evaluation of the effects of a mixture of amino acids (Amixea) on lean body mass and muscle strength of patients with unresectable advanced non small cell lung cancer: a randomized, double blind, placebo controlled, multicentre study [Internet]. Available at: <https://anzctr.org.au/ACTRN12616001402437.aspx>. [Accessed 17 October 2021].
 - [110] Clinicaltrials.gov. Identifier: NCT04567459, the effect of nutrition for the colorectal cancer patients receiving chemotherapy-randomized controlled study [Internet]. Available at: <https://clinicaltrials.gov/ct2/show/NCT04567459>. [Accessed 17 October 2021].
 - [111] Zanetti M, Cappellari GG, Barazzoni R, Sanson G. The impact of protein supplementation targeted at improving muscle mass on strength in cancer patients: a scoping review. *Nutrients* 2020;12:2099. <https://doi.org/10.3390/nu12072099>.
 - [112] Mariotti F, Gardner CD. Dietary protein and amino acids in vegetarian diets—a review. *Nutrients* 2019;11:2661. <https://doi.org/10.3390/nu11122661>.
 - [113] Salomé M, De Gavelle E, Dufour A, Dubuisson C, Volatier JL, Fouillet H, et al. Plant-protein diversity is critical to ensuring the nutritional adequacy of diets when replacing animal with plant protein: observed and modeled diets of French adults (INCA3). *J Nutr* 2020;150:536–45. <https://doi.org/10.1093/jn/nxz252>.
 - [114] Segal R, Zwaal C, Green E, Tomasone JR, Loblaw A, Petrella T. Exercise for people with cancer: a clinical practice guideline. *Curr Oncol* 2017;24:40–6. <https://doi.org/10.3747/co.24.3376>.
 - [115] Glover EI, Phillips SM. Resistance exercise and appropriate nutrition to counteract muscle wasting and promote muscle hypertrophy. *Curr Opin Clin Nutr Metab Care* 2010;13:630–4. <https://doi.org/10.1097/MCO.0b013e32833f1ae5>.
 - [116] McGlory C, van Vliet S, Stokes T, Mittendorfer B, Phillips SM. The impact of exercise and nutrition on the regulation of skeletal muscle mass. *J Physiol* 2019;597:1251–8. <https://doi.org/10.1113/jp275443>.
 - [117] Phillips SM, Paddon-Jones D, Layman DK. Optimizing adult protein intake during catabolic health conditions. *Adv Nutr* 2020;11:S1058–69. <https://doi.org/10.1093/ADVANCES/NMAA047>.
 - [118] Fernandez MA, Bertolo RF, Duncan AM, Phillips SM, Elango R, Ma DWL, et al. Translating “protein foods” from the new Canada’s Food Guide to consumers: knowledge gaps and recommendations. *Appl Physiol Nutr Metabol* 2020;45:1311–23. <https://doi.org/10.1139/apnm-2020-0192>.
 - [119] Willett W, Rockström J, Loken B, Springmann M, Lang T, Vermeulen S, et al. Food in the Anthropocene: the EAT–Lancet Commission on healthy diets from sustainable food systems. *Lancet* 2019;393:447–92. [https://doi.org/10.1016/S0140-6736\(18\)31788-4](https://doi.org/10.1016/S0140-6736(18)31788-4).
 - [120] Santé Publique France. Recommandations relatives à l'alimentation, à l'activité physique et à la sédentarité pour les adultes. Saint-Maurice. 2019. Available at: <https://www.santepubliquefrance.fr/determinants-de-sante/nutrition-et-activite-physique/documents/rapport-synthese/recommandations-relatives-a-l-alimentation-a-l-activite-physique-et-a-la-sedentarite-pour-les-adultes>. [Accessed 26 September 2021].
 - [121] Herforth A, Arimond M, Álvarez-Sánchez C, Coates J, Christianson K, Muehlhoff E. A global review of food-based dietary guidelines. *Adv Nutr* 2019;10:590–605. <https://doi.org/10.1093/advances/nmy130>.
 - [122] Hertzler SR, Lieblein-Boff JC, Weiler M, Allgeier C. Plant proteins: assessing their nutritional quality and effects on health and physical function. *Nutrients* 2020;12:3704. <https://doi.org/10.3390/nu12123704>.
 - [123] Han MA, Zeraatkar D, Guyatt GH, Vernooij RWM, El Dib R, Zhang Y, et al. Reduction of red and processed meat intake and cancer mortality and incidence: a systematic review and meta-analysis of cohort studies. *Ann Intern Med* 2019;171:711–20. <https://doi.org/10.7326/M19-0699>.
 - [124] Rohrmann S, Linseisen J. Processed meat: the real villain? *Proc Nutr Soc* 2016;75:233–41. <https://doi.org/10.1017/S0029665115004255>.
 - [125] Kim SR, Kim K, Lee SA, Kwon SO, Lee J, Keum N, et al. Effect of red, processed, and white meat consumption on the risk of gastric cancer: an overall and dose–response meta-analysis. *Nutrients* 2019;11:826. <https://doi.org/10.3390/nu11040826>.
 - [126] Ramsden C, Domenichello A. PURE study challenges the definition of a healthy diet: but key questions remain. *Lancet* 2017;390:2018–9. [https://doi.org/10.1016/S0140-6736\(17\)32241-9](https://doi.org/10.1016/S0140-6736(17)32241-9).
 - [127] Allès B, Baudry J, Méjean C, Touvier M, Péneau S, Hercberg S, et al. Comparison of sociodemographic and nutritional characteristics between self-reported vegetarians, vegans, and meat-eaters from the nutritnet-santé study. *Nutrients* 2017;9:1023. <https://doi.org/10.3390/nu9091023>.
 - [128] Gonçalves T, Horie L, Gonçalves S, Bacchi M, Bailer M, Barbosa-Silva T, et al. Diretriz BRASPEN de terapia nutricional no envelhecimento. *Braspen J* 2019;34:2–58.
 - [129] Canadian nutrient file (CNF) – search by food. <https://food-nutrition.canada.ca/cnf-fce/index-eng.jsp>. [Accessed 17 October 2021].
 - [130] Springmann M, Spajic L, Clark MA, Poore J, Herforth A, Webb P, et al. The healthiness and sustainability of national and global food based dietary guidelines: modelling study. *BMJ* 2020;370:m2322. <https://doi.org/10.1136/bmj.m2322>.
 - [131] Marinangeli CPF, Fabek H, Ahmed M, Sanchez-Hernandez D, Foisy S, House JD. The effect of increasing intakes of plant protein on the protein quality of Canadian diets. *Appl Physiol Nutr Metabol* 2021;780:1–10. <https://doi.org/10.1139/apnm-2020-1027>.
 - [132] Rock CL, Doyle C, Demark-Wahnefried W, Meyerhardt J, Courneya KS, Schwartz AL, et al. Nutrition and physical activity guidelines for cancer survivors. *CA A Cancer J Clin* 2012;62:242–74. <https://doi.org/10.3322/caac.21142>.
 - [133] World Cancer Research Fund, American Institute for Cancer Research. Diet, nutrition, and physical activity and cancer: a global perspective. Continuous update project expert report 2018. London: World Cancer Research Fund International; 2018. <https://www.wcrf.org/diet-and-cancer/>. [Accessed 26 September 2021].
 - [134] Hüser S, Guth S, Joost HG, Soukup ST, Köhrle J, Kreienbrock L, et al. Effects of isoflavones on breast tissue and the thyroid hormone system in humans: a comprehensive safety evaluation. *Arch Toxicol* 2018;92:2703–48. <https://doi.org/10.1007/s00204-018-2279-8>.
 - [135] Messina M. Soy and health update: evaluation of the clinical and epidemiologic literature. *Nutrients* 2016;8:754. <https://doi.org/10.3390/nu8120754>.
 - [136] Davey GK, Spencer EA, Appleby PN, Allen NE, Knox KH, Key TJ. EPIC–Oxford: lifestyle characteristics and nutrient intakes in a cohort of 33 883 meat-eaters and 31 546 non meat-eaters in the UK. *Publ Health Nutr* 2003;6:259–68. <https://doi.org/10.1079/phn2002430>.