ADOLESCENCE

Dr. Naomi D. Harvey

The University of Nottingham, School of Veterinary Medicine & Science, Sutton Bonington, Leicestershire, LE12 5RD, UK

E-mail: Naomi.Harvey@nottingham.ac.uk

SYNONYMS

Pubescence; juvenile period

DEFINITION

The stage in human and non-human animal development through which a juvenile becomes a reproductively and behaviorally mature adult.

INTRODUCTION

Adolescence is a developmental stage in an animal's life history that is characterized by both reproductive and behavioral maturation. The terms 'puberty' and 'adolescence' are often used interchangeably as synonyms. However, it is more accurate to consider adolescence as the period in which an animal transitions into an adult that is reproductively, socially and cognitively mature (Sisk & Foster, 2004); encompassing puberty, which more precisely is the process through which an individual becomes reproductively mature. Specialists in this area argue that reproductive (gonadal) development and behavioral development are distinct processes, driven by separate neurobiological mechanisms, that are linked by interactions between gonadal steroid hormones and the nervous system (Sisk & Foster, 2004).

Adolescence is a period that is typically associated with marked changes in cognition and social behavior that include a weakened ability to regulate affect and behavior, increased reactivity to stressors, an increase in risk-taking behavior and a greater social awareness (Crone, 2009; Linda Patia Spear, 2009; Steinberg, 2005). These changes in behavior and cognition could be thought of as forming two distinct categories, primary and secondary. Primary changes contribute directly towards maturation, such as improvements in executive function, and development of sexual behavior. The reorganization of neural circuitry and increase in gonadal steroid hormones required to enact the primary changes, leads to temporary secondary changes associated with psychological vulnerability such as the increased sensitivity to stressors, increased risk-taking behavior and weaker regulatory control of behavior and affect (Schulz & Sisk, 2016; Steinberg, 2005).

Although all vertebrate species undergo a period of transition to become reproductively and behaviorally mature (Ball & Wade, 2013), mammalian species, particularly primates and rodents, are the dominant focus in scientific literature on this period of development.

THE ROLE OF HORMONES IN THE ONSET OF PUBERTY

In primates and sheep, the hypothalamic-pituitary-gonadal (HPG) axis (comprising the hypothalamus, anterior pituitary and gonads) is active during prenatal life and infancy, is suppressed during a pre-pubertal period of development then then reactivates and matures

throughout puberty resulting in reproductive maturation (Delemarre-Van De Waal, 2002; Plant, 2015; Plant & Zorub, 1982). The HPG axis reactivates when gonadotropin-releasing hormone (GnRH) is secreted by the hypothalamus following stimulation by the neuropeptide kisspeptin (Clarke, Dhillo, & Jayasena, 2015; Irwig et al., 2004). GnRH is released into the blood stream in pulses and stimulates the secretion of two gonadotrophins; luteinizing hormone (LH) and follicle stimulating hormone (FSH), from the anterior pituitary gland. Gonadal development is triggered by this release of LH and FSH, stimulation of ovarian follicles and spermatogenesis, signaling the ovaries to produce the steroid hormones estrogens and progesterone, and the Leydig cells in the testis to produce testosterone.

In contrast to primates and sheep, which experience an extended window of pre-pubertal development and reactivation of the HPG, spermatogenesis in rats and hamsters begins shortly after birth at days 5 and 9, respectively (van Haaster & de Rooij, 1993) and the rise in pubertal hypothalamic activity in rodents is not thought to be part of a reactivation of the HPG axis (Plant, 2015).

Whilst puberty is associated with clear physiological markers, adolescence is a period of gradual change, and there are no distinct physiological markers that indicate its beginning and end. In rats, the window within which adolescence occurs is widely considered to be days 28-42 after birth, with adulthood being reached by day 60, whilst some use a less conservative classification and consider adolescence to occur between days 21 (the typical onset of weaning) to 59 (McCormick & Mathews, 2010). Although adolescence in humans is typically thought of as occurring during the teenage years, hormonal and organizational brain changes begin between 9-10 years in females and 10-12 years in males (Peper & Dahl, 2013). The visible signs of puberty typically thought to signal its onset (such as onset of menarche in females, pubic hair growth and sexually dimorphic physical changes) actually occur towards the final stages of puberty. If we define human adolescence as beginning and ending according to the maturation of the brain, then this period of human development could be considered to begin around the age of 10 and cease in the mid to early twenties (Sisk & Zehr, 2005).

NEUROLOGICAL DEVELOPMENT IN ADOLESCENCE

Steroid hormones have two methods of action in the nervous system; activational or organizational. The activational effect of steroid hormones is temporary and typically associated with the adult response to its presence or absence (Sisk & Zehr, 2005). Of most relevance to adolescence is the organizational effect that steroid hormones play in reorganizing the nervous system, which is a permanent change. There is a large body of evidence indicating that neurological development continues throughout adolescence. In both human, and non-human animals, reorganization of neural circuitry occurs during adolescence, using many of the same processes involved in the original formation of the nervous system, including programmed cell death (apoptosis), neurogenesis, dendritic pruning and sexual differentiation (Schulz & Sisk, 2016). The structural changes associated with this adolescent neural reorganization are sex and brain-region specific (Sisk & Zehr, 2005).

In addition to reorganization and activation of the neural circuits associated with reproductive behavior, regions of the mammalian brain associated with emotional regulation, behavioral inhibition and risk/reward judgements are especially impacted by changes in brain structure and function (Steinberg, 2005). Areas of the medial and dorsal prefrontal cortex, and the ventral prefrontal cortex are integral to cognitive-regulation of goal-directed behavior and affective response (Nelson, Leibenluft, McClure, & Pine, 2005). Adolescent developmental changes in these areas involve pruning and increased myelination of synaptic networks responsible for social cognition, and appear to be driven by gonadal-independent development (Nelson et al., 2005).

The regions of the brain associated with affective responses to rewards and punishers include the nucleus accumbens, amygdala, and hypothalamus, and in contrast to those brain regions responsible for cognitive-regulation, these regions are heavily imbued with gonadal steroid receptors which moderate considerable reorganization during puberty as a result of gonadal development (Nelson et al., 2005). However, adolescent reorganization of the regions necessary for the expression of reproductive behavior is activated by both gonadal steroid hormones, and via unknown gonadal steroid-independent mechanisms (Sisk & Foster, 2004).

That some aspects of neurological remodeling of the adolescent brain can occur independently of gonadal maturation, highlights the fact that reproductive and behavioral maturation are distinct, though entwined, processes (Sisk & Foster, 2004). For example, the ability to express reproductive behavior in hamsters is reliant upon both gonadal maturation, and steroid-independent adolescent maturation; as copulatory behavior can be artificially triggered by testosterone in castrated hamsters after puberty (Schulz et al., 2004), but not before (Romeo, Richardson, & Sisk, 2002). Indeed, that puberty and adolescence are linked but not the same thing, has considerable consequences for long-term behavioral outcomes of hormone-brain interactions when the timing of pubertal and non-pubertal changes during adolescence differ.

The timing of gonadal maturation in mammals can be impacted by a number of extrinsic factors, including calorie intake, calcium levels, extreme exercise and even the quality of the infant-parent relationship and familial stress (Belsky, Houts, & Fearon, 2010). A mis-match in the timing of gonadal maturation and adolescent brain reorganization could have considerable effects on individual differences and risk of psycho-pathologies in adulthood (Sisk & Zehr, 2005) and further research on the behavioral outcomes of hormone-brain interactions during puberty and adolescence is warranted (Sisk & Foster, 2004). In human females, early gonadal maturation has been consistently associated with greater risks of clinical depression and eating disorders, in addition to problems with substance use (Golub et al., 2008), however, these results are confounded by problems of study design and are associations, not causation, so may be caused by correlations with unaccounted for confounding variable such as familial stress.

DIFFERENCES IN COGNITION AND BEHAVIOR DURING ADOLESCENCE

Adolescence triggers the emergence of considerable behavioral changes as the juvenile animal develops into the adult. Some of these changes are associated only with the adolescent phase, whilst others are permanent. Some of the most distinct changes are the emergence of reproductive behavior, agonistic and impulsive behavior, and stress-reactivity.

In rodents, the emergence of reproductive behavior occurs 1-2 weeks after pubertal testosterone increases in males, and 1-2 weeks after vaginal opening in females (Sisk & Zehr, 2005). To elicit reproductive behavior, the maturation of the gonadal axis alone isn't enough. Steroid-independent and steroid-dependent organizational changes in the brain are required during adolescence to enable these behaviors to be activated by gonadal steroid hormones and to function without impairment. Indeed, if gonadal hormones are absent during adolescent neurological development it can cause long lasting impairments to reproductive behavior (Sisk & Zehr, 2005). Examples of typical reproductive behaviors in mammals that are activated by the presence of gonadal hormones after adolescent remodeling include mounting, and lordosis (presenting to indicate sexual receptivity). The influence of pubertal gonadal hormones on the reproductive behavior of an animal depends on the species, sex and timing of exposure.

Development of other adult behavioral patterns, in addition to reproductive behavior, also rely on the organizational effect of gonadal hormones. A traditional behavioral test for measuring anxiety is the open-field test, in which lower levels of activity are interpreted as reflecting greater

levels of anxiety. Adult male rats are typically less active in an open field test than adult females, and they display less social behavior in novel environments. These differences emerge during puberty, can be prevented by castration and re-instated by replacement of testosterone during puberty. Such findings clearly demonstrate the organizational effect of gonadal hormones on these anxiety-related behaviors (Schulz, Molenda-Figueira, & Sisk, 2009). Comparatively, gonadal hormones expressed during puberty have also been shown to have a significant organizational impact on the development of spatial cognition via studies of people with naturally occurring disorders of the HPG axis (Sisk & Zehr, 2005). Specifically, a lack of gonadal-hormone expression during the normative puberty age window leads to impairments spatial cognition in adults that cannot be repaired by administration of gonadal hormone treatments after the age of twenty. Flank rubbing in Syrian hamsters and territorial scent marking in tree shrews during male-male interactions are further examples of behavior that emerges during adolescence due to gonadal-hormone dependent reorganization (Sisk & Zehr, 2005).

Adolescent rats show distinct adolescent-phase differences from adults in learning and memory that stem from extensive changes to the neural circuitry that mediate these processes during adolescence (McCormick & Mathews, 2010). The regions of the brain that mediate the majority of learning and memory are the medial prefrontal cortex, amygdala and hippocampus, all of which have dense corticosteroid receptors and are influenced by glucocorticoids. An example of the differences between adolescent and adult rats, is the higher thresholds for tolerating aversive motivational learning exhibited by adolescent rats when compared to adults (Pautassi, Myers, Spear, Molina, & Spear, 2008). Differences have also been demonstrated in the responses of adolescent rats to fear conditioning, that imply ongoing development of the amygdala and hippocampus (McCormick & Mathews, 2010). At least part of these differences in learning and memory in adolescents can be attributed to differences in the experience of stress by adolescents, impacting performance on tasks involving a stress component.

Pre-pubertal rats exhibit activation in response to anxiogenic drugs in the nucleus accumbens, part of the ventral striatum, whilst post-pubertal rats exhibited activation predominantly in the prefrontal cortex (Lyss, Andersen, LeBlanc, & Teicher, 1999). It is therefore, not surprising that stress-related behavior in rats differs significantly between adolescents and adults, given the developmental processes occurring in the limbic and prefrontal areas (Linda Patia Spear, 2009). Further discussion of differences in stress response of adolescents will be described below in reference to adolescence as a sensitive period of development.

ADOLESCENCE AS A SENSITIVE PERIOD OF DEVELOPMENT

As stated previously, adolescence involves considerable reorganization of the brain's neural circuitry, that can occur independently of gonadal reproductive maturation during puberty. Because the organizational effects of gonadal hormones on adult behavior are only possible during the adolescent period of development, adolescence is considered to be a sensitive period for neurological development. However, even if these changes occur at the correct times, the changes themselves are associated with additional psychological vulnerability in adolescents such as, increased risk-taking behavior, weaker regulatory control of behavior and affect, and increased sensitivity to stressors (Schulz & Sisk, 2016; Steinberg, 2005).

Spear (L P Spear, 2000) proposed a hypothesis that the organizational changes associated with adolescent neurological remodeling lead to increased stressor reactivity via the hypothalamic-pituitary-adrenal axis (HPA), which in turn increase the risk of developing psycho-pathologies in vulnerable individuals. In order to evaluate this hypothesis, it is vital to understand the role of adolescence in normally developing children, to determine whether changes in stress physiology and emotional reactivity are indeed characteristic of the transition from childhood to

adolescence (Linda Patia Spear, 2009). A set of seminal experiments published in a special edition of *Developmental Psychology* in 2009 focused on stress reactivity and emotional sensitivity during the transition into adolescence provide just such evidence.

Gunnar and colleagues (2009) examined 82 children between the ages of 9-15 years, recording baseline and stressor levels of salivary cortisol as a marker of HPA activation, pubertal development status, self-reported stress and cardiac measures of sympathetic and parasympathetic activity (Gunnar et al., 2009). The results of their study indicate clear changes in the activity of the HPA axis as children transition into adolescents, observing developmental increases cortisol under both baseline and stressor conditions. The increase in basal cortisol observed by Gunnar and colleagues was found to be associated with puberty, as opposed to simply being age-related. Their findings were strikingly similar to those of Stroud and colleagues (2009), who using similar psychophysiological methods, were also able to provide evidence for puberty related hyper-responsiveness to stressors (Stroud et al., 2009). Differences in baseline cortisol levels due to maturation in adolescents highlights the important of ascertaining pre-challenge baselines of any measures being used prior to implementing challenges in behavioral tests.

It has been suggested, that in humans, this normative developmental increase in reactivity in the HPA axis and autonomic nervous system could promote a shift towards dysregulation of the stress-response in at-risk individuals, leading to development of clinical psycho-pathologies such as depression (Linda Patia Spear, 2009; Stroud et al., 2009). This hypothesis promotes the consideration of adolescence as a sensitive period in which experiences during this time can have long-term impacts on an individual's behavior and cognition. Evidence for just such an effect can be found in studies using animal models, such as the work of Chaby and colleagues (Chaby, Cavigelli, White, Wang, & Braithwaite, 2013) who tested the impact of chronic unpredictable stress experienced by adolescent rats on their long-term cognitive bias, decision-making, coping response and exploratory behavior. Their results showed that unpredictable stress experienced during adolescence has lasting effects of cognition and behavior of rats, which exhibited greater sensitivity to reward loss and a negative cognitive bias lasting into adulthood. Such results support the notion that chronic unpredictable stress experienced during adolescence can induce long-term negative affective states, which is likely in turn to impact upon a suite of differences as adults in cognition and executive function inducing biases in decision-making, attention towards negative stimuli, judgement and memory (e.g. Paul, Harding, & Mendl, 2005; Shields, Moons, Tewell, & Yonelinas, 2016). Such negative cognitive biases could have adaptive ecological significance, if for example an animal inhabits an area with high-predation levels (Chaby et al., 2013) however, for humans and captive/companion animals that live outside of typical ecological niches these stress-induced alterations could have long-term detrimental impacts on their emotional and behavioral well-being.

CONCLUSION

Adolescence is a period of dramatic neurological development, that is organized by both gonadal and non-gonadal hormone pathways. This period encompasses puberty, with adolescent changes typically beginning long before physiological signs of puberty are apparent and ending long after reproductive maturity is reached. Significant remodeling of the brains neural circuitry during this period has both short-term and long-term impacts on behavior and cognition, that characterize adolescence as a sensitive period in development. Studies involving adolescent animals should take into account possible normative differences in basal levels for all response variables and consider the potential long-term impacts of testing on individual differences.

Furthermore, attention should be paid to this period of development by pet owners, as chronic stress experienced during adolescence could shape stress-reactivity later in life.

CROSS-REFERENCES

- Endocrine System
- Hormones and Behavior
- Life History
- Executive Function

REFERENCES

- Ball, G. F., & Wade, J. (2013). The value of comparative approaches to our understanding of puberty as illustrated by investigations in birds and reptiles. *Hormones and Behavior*, 64(2), 211–4. <http://doi.org/10.1016/j.yhbeh.2013.05.002>
- Belsky, J., Houts, R. M., & Fearon, R. M. P. (2010). Infant Attachment Security and the Timing of Puberty. *Psychological Science*, 21(9), 1195–1201. <http://doi.org/10.1177/0956797610379867>
- Chaby, L. E., Cavigelli, S. A., White, A., Wang, K., & Braithwaite, V. A. (2013). Long-term changes in cognitive bias and coping response as a result of chronic unpredictable stress during adolescence. *Frontiers in Human Neuroscience*, 7, 328. <http://doi.org/10.3389/fnhum.2013.00328>
- Clarke, H., Dhillo, W. S., & Jayasena, C. N. (2015). Comprehensive Review on Kisspeptin and Its Role in Reproductive Disorders. *Endocrinology and Metabolism (Seoul, Korea)*, 30(2), 124–41. <http://doi.org/10.3803/EnM.2015.30.2.124>
- Crone, E. A. (2009). Executive functions in adolescence: Inferences from brain and behavior. *Developmental Science*, 12(6), 825–830. <http://doi.org/10.1111/j.1467-7687.2009.00918.x>
- Delemarre-Van De Waal, H. A. (2002). Regulation of puberty. *Best Practice and Research: Clinical Endocrinology and Metabolism*, 16(1), 1–12. <http://doi.org/10.1053/beem.2001.0176>
- Golub, M. S., Collman, G. W., Foster, P. M. D., Kimmel, C. A., Meyts, E. R.-D., Reiter, E. O., ... Toppari, J. (2008, February 1). Pediatrics. *Pediatrics*. American Academy of Pediatrics. <http://doi.org/10.1542/peds.2007-1813f>
- Gunnar, M. R., Wewerka, S., Frenn, K., Long, J. D., & Griggs, C. (2009). Developmental changes in hypothalamus-pituitary-adrenal activity over the transition to adolescence: normative changes and associations with puberty. *Development and Psychopathology*, 21(1), 69–85. <http://doi.org/10.1017/S0954579409000054>
- Irwig, M. S., Fraley, G. S., Smith, J. T., Acohido, B. V., Popa, S. M., Cunningham, M. J., ... Steiner, R. A. (2004). Kisspeptin Activation of Gonadotropin Releasing Hormone Neurons and Regulation of KiSS-1 mRNA in the Male Rat. *Neuroendocrinology*, 80(4), 264–272. <http://doi.org/10.1159/000083140>
- Lyss, P. J., Andersen, S. L., LeBlanc, C. J., & Teicher, M. H. (1999). Degree of neuronal activation following FG-7142 changes across regions during development. *Brain Research. Developmental Brain Research*, 116(2), 201–3. [http://doi.org/10.1016/S0165-3806\(99\)00069-3](http://doi.org/10.1016/S0165-3806(99)00069-3)

- McCormick, C. M., & Mathews, I. Z. (2010). Adolescent development, hypothalamic-pituitary-adrenal function, and programming of adult learning and memory. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 34(5), 756–765. <http://doi.org/10.1016/j.pnpbp.2009.09.019>
- Nelson, E. E., Leibenluft, E., McClure, E. B., & Pine, D. S. (2005). The social re-orientation of adolescence: a neuroscience perspective on the process and its relation to psychopathology. *Psychological Medicine*, 35(2), 163–74. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/15841674>
- Paul, E. S., Harding, E. J., & Mendl, M. (2005). Measuring emotional processes in animals: the utility of a cognitive approach. *Neuroscience & Biobehavioral Reviews*, 29(3), 469–491. <http://doi.org/10.1016/J.NEUBIOREV.2005.01.002>
- Pautassi, R. M., Myers, M., Spear, L. P., Molina, J. C., & Spear, N. E. (2008). Adolescent but not adult rats exhibit ethanol-mediated appetitive second-order conditioning. *Alcoholism, Clinical and Experimental Research*, 32(11), 2016–27. <http://doi.org/10.1111/j.1530-0277.2008.00789.x>
- Peper, J. S., & Dahl, R. E. (2013). The Teenage Brain: Surging Hormones-Brain-Behavior Interactions During Puberty. *Current Directions in Psychological Science*, 22(2), 134–139. <http://doi.org/10.1177/0963721412473755>
- Plant, T. M. (2015). Neuroendocrine control of the onset of puberty. *Frontiers in Neuroendocrinology*, 38, 73–88. <http://doi.org/10.1016/j.yfrne.2015.04.002>
- Plant, T. M., & Zorub, D. S. (1982). The Role of Nongonadal Restraint of Gonadotropin Secretion in the Delay of the Onset of Puberty in the Rhesus Monkey (*Macaca mulatta*). *Journal of Animal Science*, 55, 43–55. Retrieved from http://jas.fass.org/content/55/Supplement_II/43
- Romeo, R. D., Richardson, H. N., & Sisk, C. L. (2002). Puberty and the maturation of the male brain and sexual behavior: recasting a behavioral potential. *Neuroscience and Biobehavioral Reviews*, 26(3), 381–91. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12034137>
- Schulz, K. M., Molenda-Figueira, H. A., & Sisk, C. L. (2009). Back to the Future: The Organizational-Activational Hypothesis Adapted to Puberty and Adolescence. *Hormones and Behavior*, 55(5), 597–604. <http://doi.org/10.1016/j.yhbeh.2009.03.010>.Back
- Schulz, K. M., Richardson, H. N., Zehr, J. L., Osetek, A. J., Menard, T. A., & Sisk, C. L. (2004). Gonadal hormones masculinize and defeminize reproductive behaviors during puberty in the male Syrian hamster. *Hormones and Behavior*, 45(4), 242–249. <http://doi.org/10.1016/j.yhbeh.2003.12.007>
- Schulz, K. M., & Sisk, C. L. (2016). The organizing actions of adolescent gonadal steroid hormones on brain and behavioral development. *Neuroscience & Biobehavioral Reviews*, 70, 148–158. <http://doi.org/10.1016/j.neubiorev.2016.07.036>
- Shields, G. S., Moons, W. G., Tewell, C. A., & Yonelinas, A. P. (2016). The effect of negative affect on cognition: Anxiety, not anger, impairs executive function. *Emotion (Washington, D.C.)*, 16(6), 792–7. <http://doi.org/10.1037/emo0000151>
- Sisk, C. L., & Foster, D. L. (2004). The neural basis of puberty and adolescence. *Nature Neuroscience*, 7(10), 1040–1047. <http://doi.org/10.1038/nn1326>
- Sisk, C. L., & Zehr, J. L. (2005). Pubertal hormones organize the adolescent brain and behavior.

Frontiers in Neuroendocrinology, 26(3–4), 163–174.

<http://doi.org/10.1016/j.yfrne.2005.10.003>

Spear, L. P. (2000). The adolescent brain and age-related behavioral manifestations. *Neuroscience and Biobehavioral Reviews*, 24(4), 417–63. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/10817843>

Spear, L. P. (2009). Heightened stress responsivity and emotional reactivity during pubertal maturation: Implications for psychopathology. *Development and Psychopathology*, 21(1), 87–97. <http://doi.org/10.1017/S0954579409000066>

Steinberg, L. (2005). Cognitive and affective development in adolescence. *Trends in Cognitive Sciences*, 9(2), 69–74. <http://doi.org/10.1016/j.tics.2004.12.005>

Stroud, L. R., Foster, E., Papandonatos, G. D., Handwerger, K., Granger, D. A., Kivlighan, K. T., & Niaura, R. (2009). Stress response and the adolescent transition: performance versus peer rejection stressors. *Development and Psychopathology*, 21(1), 47–68. <http://doi.org/10.1017/S0954579409000042>

van Haaster, L. H., & de Rooij, D. G. (1993). Spermatogenesis is accelerated in the immature Djungarian and Chinese hamster and rat. *Biology of Reproduction*, 49(6), 1229–35. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/8286605>