

COVID-19 infection and hospitalisation risk according to vaccination status and DMARD treatment in patients with rheumatoid arthritis

Authors:

René Cordtz ^{1,2}, Salome Kristensen ^{1,3}, Rasmus Westermann ¹, Kirsten Duch ^{1,4}, Fiona Pearce ⁵, Jesper Lindhardsen ⁶, Christian Torp-Pedersen ^{7,8}, Mikkel P. Andersen ⁷, Lene Dreyer ^{1,3,9}

Affiliations:

¹ Department of Rheumatology, Aalborg University Hospital, Aalborg, Denmark.

² Department of Rheumatology, Center for Rheumatology and Spine Diseases, Gentofte Hospital, Hellerup, Denmark.

³ Department of Clinical Medicine, Aalborg University, Aalborg, Denmark.

⁴ Unit of Clinical Biostatistics, Aalborg University Hospital, Aalborg, Denmark.

⁵ Division of Epidemiology and Public Health, Department of Rheumatology, University of Nottingham, Nottingham, United Kingdom.

⁶ Lupus and Vasculitis Clinic, Center for Rheumatology and Spine Diseases, Rigshospitalet, Copenhagen, Denmark.

⁷ Department of Cardiology, Nordsjællands Hospital, Hillerød, Denmark.

⁸ Department of Public Health, University of Copenhagen, Copenhagen, Denmark.

⁹ DANBIO, Denmark.

Correspondence:

Rene Cordtz

r.cordtz@rn.dk

Department of Rheumatology,

Aalborg University Hospital

Reberbansgade 15

9000-Aalborg

Denmark

ORCID: 0000-0002-5271-2574

ABSTRACT

Objectives: To investigate the incidence of COVID-19 hospitalisation in unvaccinated and vaccinated patients with rheumatoid arthritis (RA) compared with matched controls, and in patients with RA according to DMARD treatment.

Methods: Danish nationwide matched cohort study from January to October 2021. Patients with RA were identified in the DANBIO register and matched 1:20 with individuals from the general population on age, sex, and vaccination status. Primary and secondary outcomes were COVID-19 hospitalisation (Danish National Patient Register) and first-time positive SARS-CoV2 PCR test (Danish COVID-19 Surveillance Register), respectively. Stratified by vaccination status, incidence rates (IRs) per 1000 person years (PY) and comorbidity-adjusted hazard ratios (aHRs) in cause-specific Cox models were calculated with 95% confidence intervals.

Results: In total, 28 447 unvaccinated patients and 568 940 comparators had IRs for COVID-19 hospitalisation of 10.4 (8.0 to 13.4) and 4.7 (4.3 to 5.1) per 1000 PY, respectively (aHR 1.88, 1.44 to 2.46). When fully vaccinated, corresponding IRs were 0.9 (0.5 to 1.6) and 0.5 (0.4 to 0.6) per 1000 PY (aHR 1.94, 1.03 to 3.66). Unvaccinated RA patients had an aHR of 1.22 (1.09 to 1.57) for testing positive for SARS-CoV2 and 1.09 (0.92 to 1.14) among vaccinated. Vaccinated rituximab-treated patients had increased crude IR of COVID-19 hospitalisation compared with conventional DMARD treated patients.

Conclusion: The incidence of COVID-19 hospitalisation was increased for both unvaccinated and vaccinated patients with RA compared with controls. Importantly, the parallel decreasing risk for patients with RA suggests a comparable relative benefit of vaccination in most patients.

Key words: Rheumatoid arthritis, SARS-CoV2, epidemiology, COVID-19, biologics, rituximab, vaccination

KEY MESSAGES

- Regardless of vaccination, risk of COVID-19 hospitalisation was increased in patients with rheumatoid arthritis.
- The absolute risk of hospitalisation was, however, markedly lower among all vaccinated individuals.
- Less favourable outcomes were observed in rituximab-treated patients regardless of vaccination status but based on few events.

INTRODUCTION

Since the onset of the COVID-19 pandemic, researchers have focused on identifying particular risk groups in terms of contracting SARS-CoV2, being hospitalised with COVID-19 and of experiencing a severe outcome, e.g., death, during COVID-19 hospitalisation. In several studies, patients with rheumatoid arthritis (RA) have been associated with an increased risk of being hospitalised with COVID-19 when compared with the general population or non-RA individuals (1–7), but results have not been uniform when looking at the risk of contracting SARS-CoV2 (8–11) nor of having a severe outcome following hospitalisation (1,11,12).

In Denmark, a national vaccination strategy was implemented at the end of December 2020. The BNT162b2 (Pfizer BioNTech) and mRNA-1273 (Moderna) vaccines, both mRNA vaccines (13,14), were the first available vaccines in Denmark. Considering the immunologic dysregulation intrinsic to RA and the immunosuppressive treatments frequently adopted in these patients, some concerns have arisen regarding vaccine efficacy and safety in this population. In the phase three trials of BNT162b2 by Polack et al. (15) and mRNA-1273 by Baden et al. (14), treatment with immunosuppressive therapy was an exclusion criterion and only few participants suffered from rheumatic disease.

Whether the excess risk of COVID-19 hospitalisation associated with RA has disappeared after introduction of vaccinations, and if patients with RA in general or patients treated with certain disease-modifying antirheumatic drugs (DMARDs) should be prioritised for potential future booster shots is yet to be investigated on a nationwide scale. Thus, the aims of this study were to estimate the incidence of hospitalisation due to COVID-19 in patients with RA according to vaccination status compared with matched individuals from the general population; and as a secondary outcome, the risk of a first-time COVID-19 infection. Lastly, we estimated the

1
2
3 incidence of the primary and secondary outcomes in a subgroup analysis of patients with RA
4
5 treated with various DMARD regimens.
6
7
8
9

10 **PATIENTS AND METHODS**

11 **Study Design**

12
13 This was a population-based observational matched cohort study investigating the incidence of
14
15 COVID-19 hospitalisation in patients with RA from 1 January to 5 October 2021 via linkage of
16
17 several Danish nationwide registers. Secondly, we studied the incidence of a first-time positive
18
19 SARS-CoV2 PCR-test; and, for patients with RA, we explored the rate of COVID-19
20
21 hospitalisation and mortality according to vaccination status and type of DMARD treatment.
22
23 During the study period, the dominant SARS-CoV2 strains were Alpha (start of study period to
24
25 end of June 2021) and Delta (1 July to end of 5 October 2021)(16,17). According to Danish
26
27 legislation, no ethics approval is needed for register-based studies. The study has been approved
28
29 by the Data Protection Committee of Northern Jutland, Denmark (2020-032).
30
31
32
33
34
35
36

37 **Data Sources**

38
39 The Danish Civil Registration System contains information on all residents of Denmark
40
41 including the unique civil registration number, which enables linkage between registers (18). The
42
43 Civil Registration System was used to identify all Danish individuals above 18 years old and alive
44
45 on 1 January 2021, and to obtain information on age, sex, and vital status.
46
47
48
49

50 DANBIO is a nationwide clinical quality register for inflammatory arthritis, including RA, and
51
52 used in routine care by all departments of rheumatology in Denmark. The positive predictive
53
54 value (PPV) of an RA diagnosis in DANBIO has been estimated to 96% (19).
55
56
57
58
59
60

1
2
3 The Danish Vaccination Register (DDV) was used to obtain information on the exact type and
4 date of all first and second COVID-19 vaccine doses (20). From The National Surveillance at
5
6 The State Serum Institute, Denmark, information was obtained on the date of all first positive
7
8 PCR-tests for SARS-CoV2 in Denmark during the study period. The Danish National Patient
9
10 Register (DNPR) is a nationwide administrative register of contacts in the secondary and tertiary
11
12 health care sector (21). The Danish National Prescription Registry (NPR) holds information on
13
14 date of redemption and Anatomical Therapeutic Chemical Classification System (ATC) codes on
15
16 all prescriptions redeemed at pharmacies in Denmark (22).
17
18
19
20
21
22

23 **Study Population and Exposures**

24
25 Patients with RA were identified in DANBIO with ICD-10 codes M05 and M06 and included in
26
27 the study if registered before 1 May 2020 and alive on 1 January 2021. The vaccination history of
28
29 each patient with RA was obtained through linkage with the DDV, thereby obtaining
30
31 information on the date and exact type of first and second vaccination against COVID-19.
32
33
34
35
36

37 **Time-dependent vaccination status, matching ,and follow-up**

38
39 By 1 January 2021, all patients with RA were unvaccinated and, corresponding to this date, were
40
41 matched with up to 20 unvaccinated individuals from the general population with no history of
42
43 inflammatory rheumatic disease. Besides vaccination status, i.e., being unvaccinated, matching
44
45 parameters were year of birth and sex. Unvaccinated patients and matched comparators were
46
47 then followed from 1 January 2021 to the date of first COVID-19 hospitalisation, death, date of
48
49 first vaccination, or 5 October 2021, whichever occurred first.
50
51
52
53
54

55 Those patients who received a first vaccination and had, at that point, not been hospitalised with
56
57 COVID-19, were then matched with 20 individuals from the general population that also had no
58
59 history of COVID-19 hospitalisation. The matching parameters were once again vaccination
60

1
2
3 status (now partially vaccinated), year of birth, sex, and, additionally, the month and year of
4
5 receiving their first vaccination. The latter matching parameter was included to indirectly
6
7 minimise temporal variations in the pressure of the epidemic between patients and their matched
8
9 controls. This stratum of partially vaccinated patients and matched comparators were followed
10
11 from the date of their first vaccination to the date of first COVID-19 hospitalisation, death, date
12
13 of second vaccination, or 5 October 2021, whichever occurred first.
14
15
16
17
18

19 Similarly, patients with RA who received a second vaccination and both during their follow-up in
20
21 the unvaccinated and partially vaccinated strata had not been hospitalised with COVID-19, were
22
23 matched with 20 individuals with no history of COVID-19 hospitalisation nor inflammatory
24
25 rheumatic disease. Matching parameters were vaccination status (i.e. fully vaccinated), year of
26
27 birth, sex, and month and year of receiving their first and second vaccination. Patients and
28
29 matched comparators were followed-up from the date of their second vaccination to the date of
30
31 first COVID-19 hospitalisation, death, or 5 October 2021, whichever occurred first.
32
33
34

35 In all vaccination status strata, individuals were excluded for further follow-up at time of their
36
37 first COVID-19 hospitalisation. Thus, a patient or comparator could not contribute multiple
38
39 outcomes in one vaccination stratum, nor could they progress to and contribute with follow-up
40
41 to the next vaccination status if they had been hospitalised.
42
43
44

45 **Outcome**

46
47
48 The primary outcome of COVID-19 hospitalisation was identified through linkage with DNPR
49
50 and ICD-10 codes created specifically for the pandemic by the Danish Ministry of Health in
51
52 accordance with the definition established by the World Health Organization (ICD-10 codes
53
54 B34.2A, B97.2 and B97.2A)(23). COVID-19 hospitalisation was defined as registration with one
55
56 of the abovementioned ICD-10 codes listed as the primary diagnosis and a hospital-stay lasting
57
58 > 24 hours. For descriptive purposes, the proportion who had a severe outcome during their
59
60

1
2
3 hospitalisation is presented (see Supplementary Table S1, available at *Rheumatology* online, for
4 definition). The secondary outcome was a first-time positive PCR-test for SARS-CoV2 according
5 to the National Surveillance.
6
7
8
9

10 11 12 **Other covariates**

13
14 Chronic lung disease, cardiovascular disease, diabetes mellitus type I and II, chronic kidney
15 disease, and cancer (except non-melanoma skin cancer) were a priori comorbidities of interest
16 and were identified using ICD-10 codes in the DNPR and/or redeemed prescriptions registered
17 in the NPR of available at *Rheumatology* online). For descriptive purposes, information on
18 redeemed prescriptions of conventional synthetic DMARD (csDMARD) and prednisolone
19 within 1 year prior to start of follow-up, was obtained from NPR using ATC-codes
20 (Supplementary Table S1).
21
22
23
24
25
26
27
28
29

30
31
32 For biological DMARDs (bDMARD) including TNF-inhibitors, rituximab, interleukin-6
33 inhibitors, and abatacept, as well as certain csDMARDs, SKS-procedure codes in the DNPR
34 were used to obtain information on date of administration for these drugs in the year leading up
35 to start of follow-up (Supplementary Table S1). Unfortunately, neither DNPR nor NPR held
36 information on administration of leflunomide or JAK-inhibitor treatment during the study
37 period.
38
39
40
41
42
43
44
45
46
47

48 **Statistical Analysis**

49
50 Age- and sex standardised incidence rates (IRs) of hospitalisation per 1000 person years (PY)
51 were estimated with 95% confidence intervals (95%CI). In each vaccination status stratum, the
52 incidence of hospitalisation with COVID-19 in patients with RA was compared with their
53 matched comparators, using a stratified cause-specific Cox-regression taking into account the
54 competing risk of death (24). RA was considered the main exposure variable and stratification
55
56
57
58
59
60

1
2
3 was done by matched pairs. Additionally, a comorbidity-adjusted model was made
4
5 (comorbidities: chronic lung disease, diabetes mellitus, cardiovascular disease, chronic kidney
6
7 disease, and cancer) and hazard ratios (HRs) for COVID-19 hospitalisation and the competing
8
9 event of death were presented, both with their corresponding 95% CI. The proportional hazards
10
11 assumption was evaluated by plotting the scaled Schoenfeld residuals against time.
12
13
14

15
16 To illustrate the overall absolute risk of first COVID-19 hospitalisation for the RA and
17
18 comparator groups in the respective vaccination states while taking into account the competing
19
20 risk of death, the Aalen-Johansen estimated cumulative incidence (in %) was plotted as a
21
22 function of follow-up time. The x-axis was from time 0 to the 3rd quartile of follow-up within
23
24 each vaccination stratum.
25
26
27

28
29 Following exclusion of patients with a history of COVID-19 prior to start of follow-up, IRs, and
30
31 cause-specific HR for a first-time positive SARS-CoV2 PCR test were estimated for patients with
32
33 RA and comparators. Data management and statistical analyses were performed in R 4.0.3.
34
35 Cause-specific Cox models were fitted using the ‘riskRegression’ package, and the Aalen-
36
37 Johansen cumulative incidence functions were estimated using the ‘prodlm’ package (25,26).
38
39 The model specifications are shown in the Supplementary Data S1, available at *Rheumatology*
40
41 online.
42
43
44
45
46
47

48 **Subgroup and exploratory analyses**

49
50 Considering only patients with RA, the three vaccination strata were split by exposure to
51
52 DMARDs as (‘Any bDMARD treatment except rituximab’, ‘rituximab’, ‘csDMARD treatment
53
54 only’ and ‘no DMARD treatment’). Patients that had received multiple types of bDMARDs in
55
56 the year leading up to start of follow-up was allocated to ‘Any bDMARD’. The IRs and HRs for
57
58 first-time infection and hospitalisation with COVID-19, respectively, were estimated The Cox-
59
60

1
2
3 regression was adjusted for age (continuous variable) and sex. Within each treatment-stratum we
4 presented both the proportion of patients that had redeemed at least 2 prescriptions of
5 prednisolone in the year leading up to start of follow-up and the proportion of patients with a
6 history of cancer.
7
8
9
10

11
12 Lastly, for exploratory purposes, among rituximab treated patients with RA, we calculated the
13 mean time since last infusion at start of follow-up (date of vaccination) in those with and without
14 COVID-19 infection.
15
16
17
18
19
20

21 **Sensitivity analyses**

22
23 A sufficient vaccine response is not expected until 7-14 days post-vaccination. Thus, a potential
24 interaction between exposure group (RA vs matched comparators) and time since follow-up was
25 investigated, by calculating HRs for the days 0 to 10 and 10+ post-vaccination. This analysis was
26 repeated using cut-off values for follow-up with days 0 to 14 and 14+.
27
28
29
30
31
32
33

34 **RESULTS**

35
36 A total of 28 447 SARS-CoV2 unvaccinated patients with RA started follow-up on 1 January
37 2021. The median age was 67.7 years, 71.3% were women, 68% were treated with csDMARD
38 and 23.7% treated with a bDMARD (Table 1). During follow-up, 65 unvaccinated patients with
39 RA were hospitalised with COVID-19 corresponding to an IR of 10.4 per 1000 PY (95%CI 8.0
40 to 13.4), 171 patients died and 96.2% went on to receive the first vaccination (Table 2).
41
42
43
44
45
46
47

48 Following adjustment for comorbidities, the HR for COVID-19 hospitalisation in the
49 unvaccinated RA group was 1.88 (95%CI 1.44 to 2.46) compared with the matched group (Table
50 2).
51
52
53
54
55
56

57 In patients with RA, and thus, also in matched individuals, the first vaccination was
58 predominantly an mRNA vaccine: 89.2% BNT162b2 and 8.4% mRNA-1273. Upon receipt of
59
60

1
2
3 the first vaccination, the groups were followed for a median of 28 and 30 days and had IRs of
4 COVID-19 hospitalisation of 5.5 (95%CI 3.0 to 10.0) and 2.2 (95%CI 1.8 to 2.7) per 1000 PY,
5
6 respectively (Table 2).
7
8

9
10 In Denmark, non-mRNA vaccines were excluded from the state-sponsored COVID-19
11 vaccination program by mid-April 2021. Thus, 99.9% of patients with RA received an mRNA
12 vaccination as their second dose (91.2% BNT162b2 and 8.7% mRNA-1273). The IR of
13 COVID-19 hospitalisation in patients with RA had decreased to 0.9 (95%CI 0.5 to 1.6) and to
14 0.5 (95%CI 0.4 to 0.6) per 1000 PY for matched comparators. The adjusted HR for COVID-19
15 hospitalisation remained increased at 1.94 for fully vaccinated individuals with RA compared to
16 general population comparators (95%CI 1.03 to 3.66).
17
18
19
20
21
22
23
24
25
26
27

28 The absolute risk of hospitalisation was 0.20 % for unvaccinated patients with RA at 60 days and
29 0.08 % for matched individuals, whereas the absolute risk remained below 0.05 % in both fully
30 vaccinated groups after 180 days of follow-up (Figure 1).
31
32
33
34
35
36

37 Sensitivity analyses revealed an interaction between exposure and time since vaccination in the
38 partially vaccinated stratum, where the excess risk was exclusively seen from day 10+
39 (Supplementary Table S2, available at *Rheumatology* online, and Figure 1 Panel B). These patterns
40 remained when changing the cut-off to 14 days.
41
42
43
44
45
46
47

48 Among individuals with no history of COVID-19 infection, unvaccinated patients with RA had
49 an adjusted HR of 1.22 (95%CI 1.09 to 1.57) for a first-time positive SARS-CoV2 PCR test
50 (Table 3). Among vaccinated, there was no difference in incidence of first-time SARS-CoV2
51 infection (Table 3). In patients with RA, bDMARD treated were younger and a higher
52 proportion was women (Table 4). The rituximab treated group had higher proportions of
53 patients with a history of cancer and prednisolone users. For unvaccinated, patients treated with
54
55
56
57
58
59
60

1
2
3 rituximab had a HR of 4.71 (95%CI 1.98 to 11.18) for hospitalisation with csDMARD treated
4 patients as reference (Table 4). For fully vaccinated, the HR for COVID-19 hospitalisation
5
6 remained increased for rituximab treated although based on very few (≤ 3) events. The IRs and
7
8 HRs for a first-time positive PCR-test were also numerically increased for patients receiving
9
10 rituximab (Supplementary Table S3, available at *Rheumatology* online).
11
12
13
14
15

16
17 Among unvaccinated rituximab treated patients not hospitalised with COVID-19, the mean
18
19 number of days since last infusion at start of follow-up was 105 with the corresponding number
20
21 of days at 124 for those hospitalised during follow-up. Of the latter, none received rituximab
22
23 infusion between start of follow-up and time of COVID-19 hospital admission, whereas 20% of
24
25 non-hospitalised patients treated with rituximab received an infusion between start and end of
26
27 follow-up. The mean number of days since last rituximab infusion at date of second vaccination
28
29 dose was 120 and 175 for those hospitalised and not hospitalised with COVID-19, respectively.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

DISCUSSION

This nationwide matched cohort study showed that the incidence of COVID-19 infection and hospitalisation was markedly lower both among vaccinated patients with RA and matched controls. However, patients with RA had an excess risk of hospitalisation with COVID-19 even when vaccinated, whereas the incidence of SARS-CoV2 infection was only increased in patients with RA up until receiving their first vaccination dose. For rituximab treated patients, the risks of first-time SARS-CoV2 infection and COVID-19 hospitalisation were increased compared with other DMARD regimens, regardless of vaccination status.

Numerous studies from the pre-vaccine era of COVID-19 have shown increased risk of hospitalisation with or death from COVID-19 in patients with RA compared with non-RA individuals (1–7). However, to our knowledge, this study represents the first nationwide study specifically investigating if this excess risk of hospitalisation has remained in the era of the COVID-19 vaccines.

In a pre-vaccine era of COVID-19 study, England et al. used the Veterans Affairs in the US to assess the risk of COVID-19 infection found a HR of 1.25 for patients with RA, remarkably similar to the HR of 1.22 for unvaccinated patients with RA in the present study (6). Papagoras et al. compared severity of COVID-19 infection in vaccinated and unvaccinated patients with systemic rheumatic disease, and found that a higher proportion of unvaccinated patients were hospitalised (29% vs 10%) for their COVID-19 (27). In our study, the IR of hospitalisation decreased from 10.4 in unvaccinated to 0.9 per 1000 PY in vaccinated patients with RA. Importantly, the parallel decreasing IR for patients with RA and comparators suggests a comparable relative benefit of vaccination in general. However, other factors than the vaccination contributed to the decrease in both groups: follow-up started at different calendar months for each stratum, i.e., unvaccinated started follow-up in January, whereas vaccinated

1
2
3 strata started follow-up late winter and early spring for partially vaccinated. Even with few
4
5 calendar months separating the start of follow-up, variations in factors such as which virus strain
6
7 was dominating; weather conditions; and, importantly, the measures of general restrictions issued
8
9 by the authorities, surely play a part. Disentangling the effect of vaccination from these factors is
10
11 impossible in the present study, and thus, the 10-fold decreased in the RA group does not
12
13 correspond to the effect of vaccination. However, the decreased rate of hospitalisation in
14
15 vaccinated patients with RA does suggest that the vaccines were effective in most patients with
16
17 RA. In line with this, studies investigating the immunogenicity of mRNA vaccines in patients
18
19 with RA found that most patients amounted an antibody response following vaccination, but
20
21 also, that this response was potentially inferior compared with that of healthy controls (28–32).
22
23 In addition to the intrinsic dysregulated immune system in RA, there are other potential
24
25 contributors to this lower antibody response: studies have shown that the RA cornerstone drug,
26
27 methotrexate, hampers the immunogenicity of the BNT162b2 vaccine (33,34). Also, the case of
28
29 decreased antibody production is even stronger for patients treated with rituximab (34–36).
30
31 Among fully vaccinated rituximab treated patients in the present study, although based on few
32
33 events, there was a numerically increased risk of breakthrough infections and hospitalisation with
34
35 COVID-19. However, the proportion of rituximab treated patients suffering from previous or
36
37 ongoing cancer and receiving concomitant glucocorticoids were higher compared to the other
38
39 DMARD-groups; and both are important confounding factors for which we could not adjust
40
41 without overfitting the Cox model. There was no clear pattern with regards to time since last
42
43 rituximab infusion and time to vaccination nor time to hospitalisation; this is in line with what
44
45 was observed in a Swedish study of rituximab-treated patients with multiple sclerosis (37).
46
47 With regards to the specific absolute and relative incidence rates in the present study, it is
48
49 important to remember two factors putting the results in context: first, the absolute risk of
50
51 hospitalisation may be considered low, but it is worth noting that Denmark has had one of the
52
53 lowest SARS-CoV2 infection rates in Europe during the pandemic, and the study period
54
55
56
57
58
59
60

1
2
3 specifically covered here was when the infection rates were at their lowest from 2020 to
4
5 2021. Thus, the absolute risks do not transfer to other countries with different, and likely higher
6
7 background rates, whereas the relative rates are relevant and of interest to different RA
8
9 populations in other countries. Second, a Danish study showed that patients with RA were more
10
11 likely to self-isolate and take stronger measures of social distancing even when federal general
12
13 restrictions were abandoned. Thus, the HRs for both infection and hospitalisation may be
14
15 underestimating the true biological difference (38).
16
17
18
19
20

21 We believe that the totality of findings in the existing literature and the results in the present
22
23 study, constitute a strong argument for prioritising patients with RA high for future booster
24
25 shots.
26
27
28
29

30 There are important limitations that needs mentioning: First, we did not have access to
31
32 information on incident patients with RA for the period of May 2020 to January 2021. In
33
34 Denmark, the incidence of RA is roughly 1700 per year, and therefore about 900 patients may
35
36 have been misclassified as non-RA. The impact of this in a 1:20 general population matching
37
38 setting is expected to be minimal. Another limitation is that we did not have access to register
39
40 information on leflunomide and JAK-inhibitor treatment, of which namely the latter could have
41
42 been of interest to study (7,39). Some of our subgroup analyses were hampered by a low number
43
44 of events. Lastly, the use of diagnoses in an administrative register could lead to misclassification.
45
46 However, the high validity of the COVID-19 outcomes is very reassuring, and although we can
47
48 only speculate, we have very little reason not to think that the sensitivity of these ICD-10 codes
49
50 is also high given the registration process and incentive to register in DNPR (23).
51
52
53
54
55
56

57 There are also strengths to our study. Importantly, the validity of the RA diagnosis in DANBIO
58
59 is very high (19), thus ensuring that the risk estimates provided are in fact for a true RA
60

1
2
3 population. Also, the vaccination data from DDV has a high validity and carries detailed
4
5 information, enabling us to match on both temporality as well as exact type of vaccine at each
6
7 shot.
8
9

10
11
12 In conclusion, the results of this study confirm the observed increased risk of hospitalisation
13
14 with COVID-19 in the pre-vaccination era in patients with RA compared with non-RA
15
16 individuals. Further, although the absolute risk was low among fully vaccinated individuals, the
17
18 relative incidence rate has remained increased in patients with RA compared with matched
19
20 individuals from the general population, but nonetheless underlines the importance of
21
22 vaccination.
23
24
25
26
27

28 **ACKNOWLEDGEMENTS**

29
30 The authors wish to acknowledge The Danish Departments of Clinical Microbiology and Statens
31
32 Serum Institut for carrying out laboratory analysis, registration, and release of the national SARS-
33
34 CoV-2 surveillance data use in the present study. Further, the authors wish to thank all the
35
36 Danish departments of rheumatology for reporting to the DANBIO register.
37
38
39
40

41 **AUTHOR CONTRIBUTIONS:** Cordtz, Westermann, Lindhardsen, Andersen, and Torp-
42
43 Pedersen had full access to the data and take responsibility of the integrity of the data. Cordtz
44
45 takes responsibility for the accuracy of the data analyses.
46
47

48 *Concept and design:* Cordtz, Pearce, Dreyer.

49
50 *Acquisition, analyses, or interpretation of data:* All authors

51
52 *Drafting of the manuscript:* Cordtz, Westermann, Kristensen, Dreyer

53
54 *Critical revision of the manuscript for important intellectual content:* All authors

55
56 *Statistical analyses:* Cordtz and Duch

57
58
59 *Obtained funding:* Dreyer and Torp-Pedersen
60

1
2
3 *Administrative, technical, or material support:* Andersen, Torp-Pedersen, Dreyer
4

5 *Supervision:* Dreyer, Torp-Pedersen, Kristensen
6
7
8
9

10 **FUNDING:** The study was financially supported by Aalborg University Hospital. Dr. Pearce is
11 funded by an NIHR advanced fellowship NIHR300863. The funder had no impact on the
12 planning, design, analyses nor manuscript writing of the study.
13
14
15

16
17
18 **CONFLICTS OF INTEREST:** F.P. reports a grant from Vifor Pharma outside the submitted
19 work. Torp-Pedersen reports grants from Bayer, grants from Novo Nordisk outside the
20 submitted work. L.D has received research grant/research support from BMS, and speakers'
21 bureau from Eli Lilly and Galderma. The remaining authors have declared no conflicts of
22 interest.
23
24
25
26
27
28
29

30
31 **DATA AVAILABILITY:** The dataset generated during the current study is not publicly
32 available. All data were accessed in the research environment of Statistics Denmark where
33 multiple registries can be combined with the limitation that individual data cannot be exported
34 from the re- search environment. Further, individual information is encrypted. Thus, datasets
35 cannot be made available.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REFERENCES

1. Cordtz R, Lindhardsen J, Soussi BG, Vela J, Uhrenholt L, Westermann R, et al. Incidence and severeness of COVID-19 hospitalization in patients with inflammatory rheumatic disease: A nationwide cohort study from Denmark. *Rheumatology*. 2021;60(SI):SI59–67.
2. Bower H, Frisell T, Di Giuseppe D, Delcoigne B, Ahlenius GM, Baecklund E, et al. Impact of the COVID-19 pandemic on morbidity and mortality in patients with inflammatory joint diseases and in the general population: A nationwide Swedish cohort study. *Ann Rheum Dis*. 2021;80(8):1086–93.
3. MacKenna B, Kennedy NA, Mehkar A, Rowan A, Galloway J, Mansfield KE, et al. Risk of severe COVID-19 outcomes associated with immune-mediated inflammatory diseases and immune modifying therapies: a nationwide cohort study in the OpenSAFELY platform. *medRxiv*. 2021;2021.09.03.21262888.
4. Curtis JR, Zhou X, Rubin DT, Reinisch W, Yazdany J, Robinson PC, et al. Characteristics, Comorbidities, and Outcomes of SARS-CoV-2 Infection in Patients With Autoimmune Conditions Treated With Systemic Therapies: A Population-based Study. *The Journal of Rheumatology*. 2021. jrheum.210888.
5. Akiyama S, Hamdeh S, Micic D, Sakuraba A. Prevalence and clinical outcomes of COVID-19 in patients with autoimmune diseases: a systematic review and meta-analysis. *Ann Rheum Dis*. 2020 Oct;
6. England BR, Roul P, Yang Y, Kalil AC, Michaud K, Thiele GM, et al. Risk of COVID-19 in Rheumatoid Arthritis: A National Veterans Affairs Matched Cohort Study in At-Risk Individuals. *Arthritis Rheumatol*. 2021;73(12):2179–88.
7. Bower H, Frisell T, Di Giuseppe D, Delcoigne B, Ahlenius GM, Baecklund E, et al. Effects of the COVID-19 pandemic on patients with inflammatory joint diseases in Sweden: From infection severity to impact on care provision. *RMD Open*. 2021;7(3):1–10.

- 1
2
3 8. Jung Y, Kwon M, Choi HG. Association between previous rheumatoid arthritis and
4 COVID-19 and its severity: a nationwide cohort study in South Korea. *BMJ Open*. 2021
5 Oct;11(10):e054753.
6
7
- 8
9
10 9. Zhong J, Shen G, Yang H, Huang A, Chen X, Dong L, et al. COVID-19 in patients with
11 rheumatic disease in Hubei province, China: a multicentre retrospective observational
12 study. *Lancet Rheumatol*. 2020 Sep;2(9):e557–64.
13
14
- 15
16 10. Salvarani C, Bajocchi G, Mancuso P, Galli E, Muratore F, Boiardi L, et al. Susceptibility
17 and severity of COVID-19 in patients treated with bDMARDs and tsDMARDs: A
18 population-based study. *Ann Rheum Dis*. 2020;79(7):986–8.
19
20
- 21
22 11. Conway R, Grimshaw AA, Konig MF, Putman M, García AD-, Tseng LY, et al. SARS-
23 CoV-2 Infection and COVID-19 Outcomes in Rheumatic Disease: A Systematic
24 Literature Review And Meta-Analysis. *Arthritis Rheumatol*. 2021;(Nov 21):1–29.
25
26
- 27
28 12. Xu C, Yi Z, Cai R, Chen R, Thong BY-H, Mu R. Clinical outcomes of COVID-19 in
29 patients with rheumatic diseases: A systematic review and meta-analysis of global data.
30 *Autoimmun Rev*. 2021 Apr;20(4):102778.
31
32
- 33
34 13. Walsh EE, Frenck RWJ, Falsey AR, Kitchin N, Absalon J, Gurtman A, et al. Safety and
35 Immunogenicity of Two RNA-Based Covid-19 Vaccine Candidates. *N Engl J Med*. 2020
36 Dec;383(25):2439–50.
37
38
- 39
40 14. Baden LR, El Sahly HM, Essink B, Kotloff K, Frey S, Novak R, et al. Efficacy and Safety
41 of the mRNA-1273 SARS-CoV-2 Vaccine. *N Engl J Med*. 2021;384(5):403–16.
42
43
- 44
45 15. Polack FP, Thomas SJ, Kitchin N, Absalon J, Gurtman A, Lockhart S, et al. Safety and
46 Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N Engl J Med*. 2020;383(27):2603–
47 15.
48
49
- 50
51 16. The Danish Ministry of Health. Status på virusvarianter [Internet]. 2021 [cited 2022 Feb
52 15]. Available from: [https://sum.dk/ministeren/sociale-medier-
53 arkiv/2021/oktober/2910-status-paa-virusvarianter-facebook](https://sum.dk/ministeren/sociale-medier-arkiv/2021/oktober/2910-status-paa-virusvarianter-facebook)
54
55
56
57
58
59
60

- 1
2
3 17. Danish COVID-19 Genome Consortium. Nextstrain [Internet]. 2022 [cited 2022 Feb 15].
4 Available from: <https://www.covid19genomics.dk/nextstrain>
5
6
- 7
8 18. Schmidt M, Pedersen L, Sørensen HT. The Danish Civil Registration System as a tool in
9 epidemiology. *Eur J Epidemiol*. 2014;29(8):541–9.
10
11
- 12 19. Ibfelt EH, Sørensen J, Jensen D V., Dreyer L, Schiøtz-Christensen B, Thygesen PH, et
13 al. Validity and completeness of rheumatoid arthritis diagnoses in the nationwide
14 DANBIO clinical register and the Danish national patient registry. *Clin Epidemiol*.
15 2017;9:627–32.
16
17
- 18 20. Grove Krause T, Jakobsen S, Haarh M, Mølbak K. The Danish vaccination register.
19 *Eurosurveillance*. 2012;17(17):2.
20
21
- 22 21. Schmidt M, Schmidt SAJ, Sandegaard JL, Ehrenstein V, Pedersen L, Sørensen HT. The
23 Danish National patient registry: A review of content, data quality, and research potential.
24 *Clin Epidemiol*. 2015;7:449–90.
25
26
- 27 22. Pottegård A, Schmidt SAJ, Wallach-Kildemoes H, Sørensen HT, Hallas J, Schmidt M.
28 Data resource profile: The Danish national prescription registry. *Int J Epidemiol*.
29 2017;46(3):798.
30
31
- 32 23. Bodilsen J, Leth S, Nielsen SL, Holler JG, Benfield T, Omland LH. Positive predictive
33 value of ICD-10 diagnosis codes for COVID-19. *Clin Epidemiol*. 2021;13:367–72.
34
35
- 36 24. Austin PC, Lee DS, Fine JP. Introduction to the Analysis of Survival Data in the Presence
37 of Competing Risks. *Circulation*. 2016;133(6):601–9.
38
39
- 40 25. Gerds TA, Ozenne B. riskRegression: Risk Regression Models and Prediction Scores for
41 Survival Analysis with Competing Risks. 2020.
42
43
- 44 26. Gerds TA. prodlim: Product-Limit Estimation for Censored Event History Analysis
45 [Internet]. 2019. Available from: <https://cran.r-project.org/package=prodlim>
46
47
- 48 27. Papagoras C, Fragoulis GE, Zioga N, Simopoulou T, Deftereou K, Kalavri E, et al. Better
49 outcomes of COVID-19 in vaccinated compared to unvaccinated patients with systemic
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 rheumatic diseases. *Ann Rheum Dis.* 2021;annrheumdis-2021-221539.
4
5 28. Furer V, Eviatar T, Zisman D, Peleg H, Paran D, Levartovsky D, et al. Immunogenicity
6 and safety of the BNT162b2 mRNA COVID-19 vaccine in adult patients with
7 autoimmune inflammatory rheumatic diseases and in the general population: a multicentre
8 study. *Ann Rheum Dis.* 2021 Oct;80(10):1330–8.
9
10 29. Simon D, Tascilar K, Fagni F, Krönke G, Kleyer A, Meder C, et al. SARS-CoV-2
11 vaccination responses in untreated, conventionally treated and anticytokine-treated
12 patients with immune-mediated inflammatory diseases. *Ann Rheum Dis.* 2021
13 Oct;80(10):1312–6.
14
15 30. Ferri C. Impaired immunogenicity to COVID-19 vaccines in autoimmune systemic
16 diseases. High prevalence of non-response in different patients' subgroups _ Elsevier
17 Enhanced Reader.pdf.
18
19 31. Geisen UM, Berner DK, Tran F, Sümbül M, Vullriede L, Ciripoi M, et al.
20 Immunogenicity and safety of anti-SARS-CoV-2 mRNA vaccines in patients with chronic
21 inflammatory conditions and immunosuppressive therapy in a monocentric cohort. *Ann*
22 *Rheum Dis.* 2021;80(10):1306–11.
23
24 32. Simader E, Tobudic S, Mandl P, Haslacher H, Perkmann T, Nothnagl T, et al.
25 Importance of the second SARS-CoV-2 vaccination dose for achieving serological
26 response in patients with rheumatoid arthritis and seronegative spondyloarthritis. *Ann*
27 *Rheum Dis.* 2021;annrheumdis-2021-221347.
28
29 33. Haberman RH, Herati RS, Simon D, Samanovic M, Blank RB, Tuen M, et al.
30 Methotrexate hampers immunogenicity to BNT162B2 mRNA covid-19 vaccine in
31 immune-mediated inflammatory disease. *Ann Rheum Dis.* 2020;1339–44.
32
33 34. Ammitzbøll C, Bartels LE, Bøgh Andersen J, Risbøl Vils S, Elbæk Mistegård C, Dahl
34 Johannsen A, et al. Impaired Antibody Response to the BNT162b2 Messenger RNA
35 Coronavirus Disease 2019 Vaccine in Patients With Systemic Lupus Erythematosus and
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 Rheumatoid Arthritis. *ACR Open Rheumatol.* 2021;3(9):622–8.
4
5
6 35. Deepak P, Kim W, Paley MA, Yang M, Carvidi AB, Demissie EG, et al. Effect of
7 immunosuppression on the immunogenicity of mRNA vaccines to SARS-CoV-2: a prospective
8 cohort study. *Ann Intern Med.* 2021;174(11):1572–85.
9
10
11
12 36. Deepak P, Kim W, Paley MA, Yang M, Carvidi AB, El-Qunni AA, et al. Glucocorticoids
13 and B Cell Depleting Agents Substantially Impair Immunogenicity of mRNA Vaccines to
14 SARS-CoV-2. *medRxiv Prepr Serv Heal Sci.* 2021;
15
16
17
18 37. McKay KA, Piehl F, Englund S, He A, Langer-Gould A, Hillert J, et al. Rituximab
19 Infusion Timing, Cumulative Dose, and Hospitalization for COVID-19 in Persons with
20 Multiple Sclerosis in Sweden. *JAMA Netw Open.* 2021;4(12):10–3.
21
22
23
24 38. Glintborg B, Jensen DV, Engel S, Terslev L, Pfeiffer Jensen M, Hendricks O, et al. Self-
25 protection strategies and health behaviour in patients with inflammatory rheumatic
26 diseases during the COVID-19 pandemic: Results and predictors in more than 12 000
27 patients with inflammatory rheumatic diseases followed in the Danish DANBIO registry.
28 *RMD Open.* 2021;7(1):1–13.
29
30
31
32 39. Strangfeld A, Schäfer M, Gianfrancesco MA, Lawson-Tovey S, Liew JW, Ljung L, et al.
33 Factors associated with COVID-19-related death in people with rheumatic diseases:
34 Results from the COVID-19 Global Rheumatology Alliance physician-reported registry.
35 *Ann Rheum Dis.* 2021;80(7):930–42.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Characteristics of patients with rheumatoid arthritis and matched comparators stratified by vaccination status.

	Unvaccinated		Partially vaccinated		Fully vaccinated	
	Rheumatoid arthritis	General population	Rheumatoid arthritis	General population	Rheumatoid arthritis	General population
N	28 447	568 940	27 154	542 610	26 217	523 826
Women, N (%)	20 286 (71.3%)	405 720 (71.3%)	19 336 (71.2%)	386 384 (71.2%)	18 611 (71.0%)	371 851 (71.0%)
Age in years, median (IQR)	67.7 (34.2-88.3)	67.8 (34.2-88.4)	68.4 (36.4-88.6)	68.4 (36.5-88.6)	68.9 (40.9-88.7)	68.9 (41.0-88.7)
Median (IQR) follow-up time	102 (62-137)	115 (88-146)	28 (22-35)	30 (21-39)	150 (111-189)	150 (111-189)
History of COVID-19, N (%)	534 (1.9%)	11 421 (2.0%)	708 (2.6%)	15 772 (2.9%)	700 (2.7%)	15 890 (3.0%)
History of hospitalisation with COVID-19, N (%)	57 (0.2%)	870 (0.2%)	97 (0.4%)	1567 (0.3%)	96 (0.4%)	1585 (0.3%)
Chronic kidney disease, N (%)	660 (2.3%)	10 890 (1.9%)	662 (2.4%)	15 144 (2.8%)	641 (2.4%)	15 299 (2.9%)
Cardiovascular disease, N (%)	5847 (20.6%)	96 644 (17.0%)	5770 (21.2%)	108 789 (20%)	5692 (21.7%)	107 759 (20.6%)
Use of cardiovascular drugs, N (%)	14 694 (51.7%)	270 220 (47.5%)	14 431 (53.1%)	280 652 (51.7%)	14 281 (54.5%)	278 675 (53.2%)
Hospital-diagnosis of diabetes mellitus (type I or II), N (%)	2266 (8%)	38 267 (6.7%)	2205 (8.1%)	46 636 (8.6%)	2168 (8.3%)	46 493 (8.9%)
Use of anti-diabetic drugs, N (%)	2741 (9.6%)	52 509 (9.2%)	2725 (10.0%)	60 495 (11.1%)	2679 (10.2%)	60 128 (11.5%)
Chronic lung disease, N (%)	2431 (8.5%)	29 188 (5.1%)	2385 (8.8%)	37 909 (7.0%)	2341 (8.9%)	37 844 (7.2%)
Use of inhalation-medicine, N (%)	4545 (16%)	68 846 (12.1%)	4368 (16.1%)	77 422 (14.3%)	4288 (16.4%)	76 669 (14.6%)
History of cancer (except NMSC), N (%)	3608 (12.7%)	72 501 (12.7%)	3563 (13.1%)	88 832 (16.4%)	3534 (13.5%)	89 743 (17.1%)
Methotrexate, N (%)	15 801 (55.5%)	2572 (0.5%)	15 043 (55.4%)	6430 (1.2%)	14 603 (55.7%)	6728 (1.3%)
Sulfasalazine, N (%)	4042 (14.2%)	593 (0.1%)	3726 (13.7%)	1410 (0.3%)	3545 (13.5%)	1407 (0.3%)
Hydroxychloroquine, N (%)	2961 (10.4%)	690 (0.1%)	2787 (10.3%)	4 (0%)	2698 (10.3%)	5 (0%)

1						
2						
3						
4	Other csDMARD, N (%)	3143 (11.0%)	1019 (0.2%)	2899 (10.7%)	1504 (0.3%)	2771 (10.6%)
5	Prednisolone, N (%)	3566 (12.5%)	11 464 (2.0%)	3311 (12.2%)	2775 (0.5%)	3202 (12.2%)
6						
7	TNF-inhibitor, N (%)	4799 (16.9%)	612 (0.1%)	4672 (17.2%)	15 974 (2.9%)	4480 (17.1%)
8						
9	Abatacept, N (%)	423 (1.5%)	13 (0%)	410 (1.5%)	2563 (0.5%)	399 (1.5%)
10						
11	Tocilizumab, N (%)	843 (3.0%)	66 (0%)	803 (3.0%)	100 (0%)	765 (2.9%)
12						
13	Rituximab, N (%)	630 (2.2%)	581 (0.1%)	572 (2.1%)	303 (0.1%)	544 (2.1%)
14						
15						
16						
17						
18						
19						
20						
21						
22						
23						
24						
25						
26						
27						
28						
29						
30						
31						
32						
33						
34						
35						
36						
37						
38						
39						
40						
41						
42						
43						
44						
45						
46						

Abbreviations: IQR, interquartile range; NMSC, non-melanoma skin cancer; csDMARD, conventional synthetic DMARD; TNF, tumour necrosis factor;

Table 2. Incidence rates and hazard ratios for COVID-19 hospitalisation and death stratified by vaccination status.

	Unvaccinated		Partially vaccinated		Fully vaccinated	
	Rheumatoid arthritis	General population	Rheumatoid arthritis	General population	Rheumatoid arthritis	General population
N	28 447	568 940	27 154	542 610	26 217	523 826
PY of observation	8297	186 493	2370	49 051	10 846	216 529
Median [IQR] days of follow-up	102 (62-137)	115 (88-146)	28 (22-35)	30 (21-39)	150 (111-189)	150 (111-189)
Hospitalised with COVID-19, N	65	727	11	95	11	131
Age and sex standardised COVID-19 hospitalisation rate per 1000 PY, (IQR)	10.4 (8.0-13.4)	4.7 (4.3-5.1)	5.5 (3.0-10.0)	2.2 (1.8-2.7)	0.9 (0.5-1.6)	0.5 (0.4-0.6)
HR for COVID-19 hospitalisation – crude, (IQR)	2.01 (1.54-2.61)	1 (Ref.)	2.36 (1.2-4.66)	1 (Ref.)	1.63 (0.88-3.01)	1 (Ref.)
HR for COVID-19 hospitalisation - adjusted for comorbidities ^a, (IQR)	1.88 (1.44-2.46)	1 (Ref.)	2.47 (1.25-4.89)	1 (Ref.)	1.94 (1.03-3.66)	1 (Ref.)
N (%) of COVID-19 hospitalisations with a severe outcome ^b	11 (16.9%)	209 (28.7%)	≤3	26 (27.4%)	4 (36.4%)	42 (32.1%)
Died during follow-up, N	171	2912	69	1273	391	7926
Age and sex standardised mortality rate per 1000 PY, (IQR)	27.7 (23.7-32.4)	21.4 (20.6-22.2)	33.7 (26.6-42.8)	29.2 (27.7-30.9)	29.1 (26.3-32.2)	30.1 (29.4-30.7)
HR for death – crude, (IQR)	1.37 (1.15-1.62)	1 (Ref.)	1.17 (0.88-1.56)	1 (Ref.)	0.98 (0.89-1.09)	1 (Ref.)
HR for death - adjusted for comorbidities ^a, (IQR)	1.31 (1.10-1.56)	1 (Ref.)	1.25 (0.94-1.68)	1 (Ref.)	1.05 (0.94-1.16)	1 (Ref.)

Abbreviations: PY, person years; IQR, interquartile range; HR, hazard ratio

^a Adjusted for history of cancer, cardiovascular disease, diabetes mellitus, chronic kidney disease, and chronic lung disease.

^b A severe outcome was defined as the composite of either acute respiratory distress syndrome, admission to an intensive care unit, and/or death

Table 3. Incidence rates and hazard ratios for first-time positive SARS-CoV2 PCR test stratified by vaccination status.

	Unvaccinated		Partially vaccinated		Completed vaccination	
	Rheumatoid arthritis	General population	Rheumatoid arthritis	General population	Rheumatoid arthritis	General population
Number of individuals, N	27 903	557 415	26 444	526 815	25 517	507 926
Median age (IQR)	67.8 (57.4-76.2)	67.9 (57.5-76.2)	68.5 (58.4-76.7)	68.6 (58.5-76.7)	69 (59.4-76.9)	69.1 (59.5-76.9)
Female, N (%)	19 885 (71.3%)	397 314 (71.3%)	18 805 (71.1%)	374 962 (71.2%)	18 088 (70.9%)	360 304 (70.9%)
History of cancer, N (%)	3551 (12.7%)	71 336 (12.8%)	3491 (13.2%)	86 829 (16.5%)	3456 (13.5%)	87 629 (17.3%)
Person years of observation, N	8053	180 804	2290	45 965	10 558	209 456
Median days of follow-up, N (IQR)	102 (62-137)	115 (88-145)	28 (22-35)	29 (21-39)	150 (112-189)	150 (112-189)
First-time COVID-19 infections, N	309	6690	58	1345	107	1942
Age and sex standardised COVID-19 infection rate per 1000 person years (IQR)	37.8 (33.6-42.6)	33.9 (33.1-34.8)	27 (20.7-35.1)	28.5 (27-30.2)	11.3 (9.2-13.9)	10.4 (9.9-10.9)
HR – crude (IQR)	1.24 (1.09-1.41)	1 (Ref.)	0.87 (0.59-1.27)	1 (Ref.)	1.09 (0.9-1.32)	1 (Ref.)
HR - adjusted for comorbidities ^a (IQR)	1.22 (1.09-1.57)	1 (Ref.)	0.87 (0.95-1.74)	1 (Ref.)	1.09 (0.92-1.14)	1 (Ref.)

Died, N	161	2739	62	1187	369	7637
Mortality rate per 1000 PY, N (IQR)	26.8 (22.8-31.4)	20.6 (19.8-21.4)	31.5 (24.5-40.5)	29.3 (27.7-31.1)	28.3 (25.5-31.4)	30 (29.3-30.7)
HR for death – crude (IQR)	1.38 (1.16-1.64)	1 (Ref.)	1.18 (0.88-1.6)	1 (Ref.)	0.96 (0.86-1.06)	1 (Ref.)
HR for death - adjusted for comorbidities^a (IQR)	1.22 (1.07-1.38)	1 (Ref.)	0.87 (0.59-1.27)	1 (Ref.)	1.09 (0.89-1.32)	1 (Ref.)

Abbreviations: PY, person years; IQR, interquartile range; HR, hazard ratio

^a Adjusted for history of cancer, cardiovascular disease, diabetes mellitus, chronic kidney disease, and chronic lung disease.

Table 4. Characteristics and rates of COVID-19 hospitalisation in patients with rheumatoid arthritis stratified by DMARD treatment.

	Unvaccinated				Partially vaccinated				Fully vaccinated			
	TNFi, abatacept, or IL6	Rituximab	csDMARD	No DMARD	TNFi, abatacept, or IL6	Rituximab	csDMARD	No DMARD	TNFi, abatacept, or IL6	Rituximab	csDMARD	No DMARD
N	5768	630	14969	7080	5625	572	14062	6895	5420	544	13626	6627
Median (IQR) age in years	63.3 (53.3-71.7)	66.6 (57-73.1)	69 (58.7-76.8)	69.4 (58.2-78)	63.7 (54.1-71.9)	67 (58-73.7)	69.8 (59.8-77.2)	70.3 (59.4-78.6)	64.3 (55.1-72.2)	67.5 (58.6-74)	70.2 (60.7-77.3)	71 (60.8-78.9)
Female, N (%)	4336 (75%)	485 (77%)	10492 (70%)	4973 (70%)	4214 (75%)	440 (77%)	9856 (70%)	4826 (70%)	4046 (75%)	413 (76%)	9523 (70%)	4629 (70%)
History of cancer, N (%)	337 (6%)	221 (35%)	2011 (13%)	1039 (15%)	342 (6%)	205 (36%)	1952 (14%)	1064 (15%)	349 (6%)	199 (37%)	1924 (14%)	1062 (16%)
Glucocorticoid use, N (%)	769 (13%)	143 (23%)	1719 (11%)	935 (13%)	720 (13%)	136 (24%)	1526 (11%)	929 (13%)	677 (12%)	124 (23%)	1479 (11%)	922 (14%)
PY of observation	1712.6	163.6	4261.6	2159.6	504.9	46.2	1200.1	618.9	2214.4	248.2	5664	2719
Median (IQR) days of follow-up	107 (62-140)	75 (56-122)	101 (62-133)	106.5 (63-143)	28 (22-35)	25 (22-35)	28 (22-35)	28 (22-35)	147 (108-189)	179.5 (135.2-197)	151 (115-189)	148 (108-188)
N hospitalised with COVID-19	5	6	39	15	0	0	5	6	≤3	≤3	4	4

1													
2													
3	Standardised												
4	COVID-19												
5	hospitalisation	8.3 (2.8-24.4)	44.4 (17.7-	12.1 (8.6-	8.0 (4.8-	-	-	4.8 (2.0-	8.3 (3.7-	-	-	0.6 (0.2-	0.9 (0.3-
6	rate per 1000 PY,		111.7)	17.2)	13.4)			11.6)	18.7)			1.7)	2.4)
7	N (IQR)												
8	N (%) of COVID-												
9	19 hospitalisations												
10	with a severe	0	≤3	6 (15.4%)	4 (26.7%)	0	0	≤3	≤3	0	≤3	≤3	≤3
11	outcome ^b												
12													
13	HR for COVID-19												
14	hospitalisation –	0.47 (0.19-	4.71 (1.98-	1 (Ref.)	0.74 (0.41-	-	-	1 (Ref.)	1.78 (0.53-	0.80 (0.09-	11.69	1 (Ref.)	1.86 (0.46-
15	crude, (IQR)	1.21)	11.18)		1.34)				5.96)	7.41)	(2.07-		7.51)
16											66.06)		
17	Died, N (%)	25	5	73	68	≤3	≤3	33	32	41	7	178	165
18													
19													
20	Mortality rate per	24.9 (15.4-	40.9 (15.5-	20.9 (16.5-	38.7 (30.4-	23.3 (4.8-	38.2 (5.4-	29.4 (20.9-	52.0 (36.5-	22.0 (15.8-	33.5 (15.6-	24.3 (20.9-	42.2 (35.8-
21	1000 PY, N (IQR)	40.2)	108.4)	26.5)	49.2)	112.7)	271.2)	41.4)	74.1)	30.7)	71.8)	28.2)	49.7)
22													
23													
24	HR for death –	1.48 (0.93-	2.39 (0.96-	1 (Ref.)	1.63 (1.17-	0.39 (0.12-	0.95 (0.13-	1 (Ref.)	1.33 (0.81-	0.95 (0.67-	1.29 (0.60-	1 (Ref.)	1.57 (1.27-
25	crude, (IQR)	2.34)	5.92)		2.27)	1.29)	7.00)		2.18)	1.34)	2.75)		1.95)
26													

Abbreviations: TNFi, tumour necrosis factor inhibitor; IL6, interleukin 6; csDMARD, conventional synthetic DMARD; IQR, interquartile range; PY, person years; HR, hazard ratio. ^b Severe outcome defined as the composite of either acute respiratory distress syndrome, admission to an intensive care unit, and/or death




Figure 1. Cumulative incidence (%) of first COVID-19 hospitalisation according to vaccination status. Aalen-Johansen estimated cumulative incidence for (A) unvaccinated, (B) partially vaccinated and (C) fully vaccinated patients with rheumatoid arthritis and matched controls.