

- "This document is the Accepted Manuscript version of a Published Work that appeared in final form in [Organic Letters], copyright © American Chemical Society after peer review and technical editing by the publisher. To access the final edited and published work see [<http://pubs.acs.org/articlesonrequest/AOR-SrYtcTQyy2REUxcMIckA>]."

# Iron catalyzed indolizine synthesis from pyridines, diazo compounds and alkynes

Tim Douglas,<sup>†,‡</sup> Anca Pordea<sup>‡</sup> and James Dowden<sup>†,\*</sup>

<sup>†</sup>School of Chemistry and <sup>‡</sup>Department of Chemical and Environmental Engineering, University of Nottingham, University Park, Nottingham, NG7 2RD, United Kingdom.


**KEYWORDS:** indolizines, diazo reagents, ylides, metal carbenes, dehalogenation.

**ABSTRACT:** Iron (III) catalyzed synthesis of indolizines from commercially available alkyne, pyridine and diazo precursors is reported. This mild, expedient method is tolerant of various solvents and proceeds with as little as 0.25 mol % of [Fe(TPP)Cl]. Significantly, this multicomponent reaction is compatible with electrophilic alkynes; control experiments demonstrate the importance of catalyst in promoting pyridinium ylide formation over background reactivity.

Catalytic, multicomponent reactions that give direct access to indolizines from commercially available precursors are highly desirable. Strong demand for this heterocycle reflects its versatile photochemical properties,<sup>1,5</sup> and biological activity.<sup>6-7</sup> In turn, this has stimulated much innovation in indolizine synthesis.<sup>8-11</sup> Exploiting the abundance of commercially available pyridines would be attractive, especially if they are used directly without the need for prior functionalization. An initial report of rhodium(II) catalyzed indolizine synthesis involved transformation of pyridine-tethered diazo precursors *via* metal carbenoids and ylides in the presence of electrophilic alkynes; just one example from this report involved three-component transformation from a simple pyridine.<sup>12</sup> Synthesis of indolizines by combination of alkenes, diazo reagents and quinolines or pyridines was described very recently, although a large amount of catalyst (20 mol % CuF<sub>2</sub> and PPh<sub>3</sub>) was required and poor yields were obtained from reactions involving pyridines.<sup>13</sup> A general method involving alkynes and catalytically generated pyridinium ylides is yet to be reported.

Recently, we described efficient synthesis of alkaloid-inspired spirotetrahydroindolizines by combination of pyridine, diazo compound, alkenes and an iron (III) catalyst.<sup>14</sup> We were keen to investigate whether indolizines could be accessed by performing the related reaction with electrophilic alkynes. Marshalling the greater reactivity of these dipolarophiles is a potential challenge, however. In the ideal reaction pathway (Scheme 1), the diazo precursor **A** is converted to an electrophilic metal carbenoid **B** which is intercepted by pyridine to give an ylide **C**. Subsequent addition to the electrophilic alkyne and cyclisation give a dihydroindolizine **D**, which undergoes aromatization to generate the target indolizine **E**. Clearly, direct addition of pyridine to alkyne is a potential competing


**Scheme 1.** Envisaged reaction pathway for indolizine synthesis *via* catalytically generated pyridinium ylides.



pathway but, once formed, pyridinium ylides are expected to be significantly more nucleophilic than pyridines.<sup>15</sup>

Herein, we report catalytic synthesis of indolizines by one-pot condensation of commercially available pyridines, alkynes, and diazo ester precursors. The commercially available [Fe(TPP)Cl] catalyst is derived from cheap and abundant iron and successfully operates with as little as 0.25 mol % equivalents at mild temperatures in various solvents. Characterization of background reaction products underlines the remarkable capacity of this catalyst to promote a pathway *via* highly reactive metal carbenoid and pyridinium ylides.

Initially, reaction of pyridine (3 equiv), ethyl diazoacetate (EDA) (1.5 equiv) and ethyl propiolate (1 equiv) using [Fe(TPP)Cl] as catalyst (1 mol %) at ambient temperature was investigated. Reagents were combined in a vessel open to air, without additional oxidant. To our satisfaction, these reaction conditions returned an excellent yield of the target indolizine **1** (89%, Table 1, entry 1). The same reaction was performed in a variety of solvents (e.g. acetone, ethyl acetate, toluene, etc.) from which

**Table 1.** Initial reaction optimization.

entry	pyridine (equiv)	Fe(PPP)Cl (mol %)	solvent	yield (%) <sup>a</sup>
1	3	1	CH <sub>2</sub> Cl <sub>2</sub>	89
2	3	1	CH <sub>3</sub> CN	80
3	3	1	acetone	82
4	3	1	toluene	79
5	3	1	EtOAc	70
6	3	1	H <sub>2</sub> O	13
7	3	1	H <sub>2</sub> O:EtOH(2:8)	82
8	1	1	CH <sub>2</sub> Cl <sub>2</sub>	87
9	1	0.25	CH <sub>2</sub> Cl <sub>2</sub>	82
10	1	0.1	CH <sub>2</sub> Cl <sub>2</sub>	27


<sup>a</sup>Isolated yield after column chromatography.

consistently good yields of indolizine **1** were obtained (70–82%, entries 2–5).<sup>16</sup>

The yield of indolizine **1** was not significantly reduced when pyridine was used in equal ratio to alkyne (87%, entry 8). Subsequently, reducing the amount of catalyst to 0.25 mol % also gave **1** in good yield (82%, entry 9) but a poor yield (27%, entry 10) was obtained if the catalyst was used at 0.1 mol %. For the rest of this study reactions were compared using the most productive conditions, specifically 1 mol % catalyst in dichloromethane.

In the absence of catalyst, the same combination of reagents furnished 3*H*-pyrazole derivative **2** in significant yield (49%, Table 2, entry 1). Presumably, cycloaddition between EDA and the alkyne gives an initial 1*H*-pyrazole product,<sup>17</sup> that undergoes subsequent 1,4-conjugate addition with a further equivalent of ethyl propiolate. The lack of product, **2**, when [Fe(PPP)Cl] was present (1 mol %) reinforces the notion of a catalytic pathway *via* metal carbenoid and pyridinium ylide.

When ethyl propiolate was replaced with more electrophilic dimethyl acetylenedicarboxylate an exothermic reaction occurred prior to addition of EDA, which may be attributed to deleterious reaction between alkyne and pyridine. Specifically, 4*H*-quinolizine **3** (66%, entry 2) was isolated as the main product from this reaction mixture, which is consistent with reported preparation of this molecule.<sup>18</sup> Indolizine **4** was obtained in 4% yield from the same reaction. The yield of **4** improved to 53% by co-addition of pyridine and EDA to a mixture of catalyst and alkyne at -20 °C and subsequent warming to room temperature (entry 3). Methyl 2-butynoate, pyridine and EDA gave indolizine **5** in 34% yield using these conditions (entry 4).


**Table 2.** Further optimization of indolizine synthesis.

entry	alkyne	change from std. conditions	product (yield %)
1		no catalyst, (1.3 equiv pyridine) <sup>a</sup>	 <b>2</b> (49%)
2		(3 equiv pyridine) <sup>a</sup>	 <b>3</b> (66%)
3		-20 °C <sup>b</sup>	 <b>4</b> (53%)
4		-20 °C <sup>b</sup>	 <b>5</b> (34%)

<sup>a</sup>Reactions performed by addition of EDA (1.5 equiv) to a solution of pyridine (as specified), alkyne (1.0 equiv), [Fe(PPP)Cl] 1 mol % in CH<sub>2</sub>Cl<sub>2</sub> at room temperature. <sup>b</sup>EDA (1.5 equiv) and pyridine (1.0 equiv, 0.6 mmol) added a solution of alkyne (1.0 equiv), [Fe(PPP)Cl] 1 mol % in CH<sub>2</sub>Cl<sub>2</sub> at the specified temperature.

Indolizines were readily obtained using these improved reaction conditions and a variety of commercially available pyridines. For example, reaction of 4-methoxypyridine gave indolizine **6** in excellent yield (77%, Scheme 2). Pyridines with various 4-substituents also gave indolizine products. 4-Acetylpiperidine gave the corresponding indolizine **7** in excellent yield (75%). Reaction of 4-piperidines with other electron withdrawing substituents (CF<sub>3</sub>, CN) gave the corresponding indolizines **8** (57%) and **9** (45%) respectively. Indolizines **10** (74%) and **11** (59%) were obtained from 4-methylpiperidine and 4-phenylpiperidine precursors, respectively. 2-Methylpiperidine also reacted well to give indolizine **12** in 58% yield.

**Scheme 2.** Indolizine products incorporating various pyridines.<sup>a</sup>




<sup>a</sup>Reactions were performed using pyridines (1.0 equiv, 0.6 mmol), ethyl propiolate (1.0 equiv), EDA (1.5 equiv), [Fe(PPP)Cl] 1 mol % in CH<sub>2</sub>Cl<sub>2</sub> at -20 °C then warming to room temperature for 16 h; all yields after chromatography. <sup>b</sup>Product ratios estimated by relative integration of peaks corresponding to 6- or 8- indolizine position by <sup>1</sup>H NMR spectroscopy. <sup>c</sup>5 equivalents of 2-methylpyridine were used.

3-Substituted pyridines also reacted in good yields, the resulting indolizine products were obtained as a mixture of regioisomers corresponding to substitution at either the 6- or 8-position. 3-Formylpyridine gave a (1:1) mixture of indolizine regioisomers **13a** and **13b** in 51% yield. Indolizines were produced with a slight preference for substitution at the 8-position from reaction of either 3-methylpyridine (69% yield, **14a** 5:3 **14b**) or 3-fluoropyridine (**15a** 33% and **15b** 19%). A more significant preference was observed in the reaction of 3-methoxypyridine, from which the 8-methoxyindolizine isomer **16a** was obtained as the major product (68%), whereas 6-methoxyindolizine **16b** was isolated from the same reaction in 4% yield. The influence of electron donating substituents on the products of addition to pyridinium species has been noted previously.<sup>19</sup> Interestingly, reaction of 3-bromopyridine gave the indolizine product arising from proto-debromination, **1** (60%) Scheme 3A.

We sought to further explore this dehalogenation. The same indolizine product **1** was obtained in the non-catalysed route using K<sub>2</sub>CO<sub>3</sub> to generate ylide by deprotonation of a pyridinium salt (40%, Scheme S1A),<sup>20,21</sup> absolving the iron catalyst from involvement in this process. Performing the catalytic reaction in a biphasic mixture of CH<sub>2</sub>Cl<sub>2</sub> and D<sub>2</sub>O (3:1) gave indolizine products with bromine at the 6- or 8-position (**17a**, 37%, **17b**, 25% respectively, Scheme 3B) along with a mixture of proto- and deuterio-debrominated products (**1** and **18ab**, 14%); deuterium was incorporated in similar ratio (**18a** 5:3 **18b**) to brominated co-products (**17a** and **17b**). Based on these observations, we speculate that the initial dihydropyridine cycloadduct undergoes tautomerization and deuterium exchange to form an intermediate (e.g. **[19a]**) from which bromine is collected by an external nucleo-


phile, e.g. pyridinium ylide to furnish **1** or deuterated analogue **18a** (Scheme 3C). Formation of **18b** from the dihydropyridine precursor of **17b** via an analogous pathway is envisaged (Scheme S2).<sup>20</sup> A related dehalogenation mechanism has been noted for other *N*-heterocycles.<sup>22,23</sup>

**Scheme 3.** Dehalogenation observed during reaction of 3-bromopyridine A) in CH<sub>2</sub>Cl<sub>2</sub>; B) in CH<sub>2</sub>Cl<sub>2</sub>:D<sub>2</sub>O (1:1). C) proposed debromination mechanism to **1** or **18a**.



<sup>a</sup>Reaction conditions: 3-bromopyridine (1.0 equiv, 1.74 mmol), ethyl propiolate (1.0 equiv), EDA (1.5 equiv), [Fe(PPP)Cl] 1 mol % in CH<sub>2</sub>Cl<sub>2</sub>:D<sub>2</sub>O (3:1) at room temperature for 16 h; combined yield for inseparable mixture of **1** and **18ab**. <sup>b</sup>Product ratio estimated by relative integration of protons at 5-, 6- or 8- indolizine position by <sup>1</sup>H NMR spectroscopy.

#### Scheme 4. Indolizine products of various alkynes.<sup>a</sup>



<sup>a</sup>Reactions were performed using 4-acetylpyridine (1.0 equiv, 0.6 mmol), alkynes (1.0 equiv), EDA (1.5 equiv), [Fe(PPP)Cl] 1 mol % in CH<sub>2</sub>Cl<sub>2</sub> at -20 °C then warming to room temperature for 16 h; all yields after chromatography.

Generally, electron deficient alkynes were required to produce indolizines in acceptable yield. For example, methyl propiolate and phenylsulfonylacetylene reacted with 4-acetylpyridine to give indolizines **20** and **21** in 85% and 41% yield respectively (Scheme 4), but target heterocycles (**22**, **23**) were not obtained from the reaction with trimethylsilylacetylene or phenylacetylene.

Disubstituted alkynes such as dimethyl acetylenedicarboxylate or ethyl 3-trifluoromethylpropiolate gave the corresponding indolizines **24** in 60% yield and **25** in 65% yield respectively. Alternative benzyl diazoacetate ester performed equally well in reaction with ethylpropiolate to produce indolizine **26** in 64% yield.

In conclusion, a general and expedient, one pot synthesis of indolizines has been developed. The reaction uses commercially available pyridine, alkyne and diazo ester precursors, while the catalyst is derived from cheap and abundant iron. Significantly, the iron catalyzed route via pyridinium ylides is compatible with electrophilic alkynes by out-competing with potentially significant background reactions.

#### ASSOCIATED CONTENT

**Supporting Information.** The Supporting Information is available free of charge on the ACS Publications website at <http://pubs.acs.org>.

Supplementary tables, schemes, detailed experimental and compound characterization data (PDF).

#### AUTHOR INFORMATION

##### Corresponding Author

\* E-mail: [james.dowden@nottingham.ac.uk](mailto:james.dowden@nottingham.ac.uk);

##### Author Contributions

All authors have given approval to the final version of the manuscript.

#### Funding Sources

EPSRC Centre for Doctoral Training in Sustainable Chemistry (grant number EP/L015633/1) and EPSRC (EP/J021008/1).

#### Notes

The authors declare no competing financial interest.

#### ABBREVIATIONS

EDA, ethyl diazoacetate; TPP, tetraphenylporphyrin.

#### REFERENCES

- Huckaba, A. J.; Yella, A.; Brogdon, P.; Scott Murphy, J.; Nazeeruddin, M. K.; Grätzel, M.; Delcamp, J. H., *Chem. Commun.* **2016**, 52, 8424-8427.
- Huckaba, A. J.; Yella, A.; McNamara, L. E.; Steen, A. E.; Murphy, J. S.; Carpenter, C. A.; Punekey, G. D.; Hammer, N. I.; Nazeeruddin, M. K.; Grätzel, M.; Delcamp, J. H., *Chem. Eur. J.* **2016**, 22, 15536-15542.
- Huckaba, A. J.; Giordano, F.; McNamara, L. E.; Dreux, K. M.; Hammer, N. I.; Tschumper, G. S.; Zakeeruddin, S. M.; Grätzel, M.; Nazeeruddin, M. K.; Delcamp, J. H., *Adv. Energy Mater.* **2015**, 5, 1401629.
- Choi, E. J.; Kim, E.; Lee, Y.; Jo, A.; Park, S. B., *Angew. Chem. Int. Edit.* **2014**, 53, 1346-1350 DOI: 10.1002/anie.201308826.
- Song, Y. R.; Lim, C. W.; Kim, T. W., *Luminescence* **2016**, 31, 364-371.
- Chen, P.; Chaikuad, A.; Bamborough, P.; Bantscheff, M.; Bountra, C.; Chung, C.-w.; Fedorov, O.; Grandi, P.; Jung, D.; Lesniak, R.; Lindon, M.; Mueller, S.; Philpott, M.; Prinjha, R.; Rogers, C.; Selenski, C.; Tallant, C.; Werner, T.; Willson, T. M.; Knapp, S.; Drewry, D. H., *J. Med. Chem.* **2016**, 59, 1410-1424.
- Hay, D. A.; Rogers, C. M.; Fedorov, O.; Tallant, C.; Martin, S.; Monteiro, O. P.; Muller, S.; Knapp, S.; Schofield, C. J.; Brennan, P. E., *MedChemComm* **2015**, 6, 1381-1386.
- Zhu, C.; Zhao, P.; Qiao, Y.; Xiao, K.; Song, C.; Chang, J., *J. Org. Chem.* **2017**, 82, 7045-7049.
- Meazza, M.; Leth, L. A.; Erickson, J. D.; Jorgensen, K. A., *Chem. - Eur. J.* **2017**, 23, 7905-7909.
- Erguven, H.; Leitch, D. C.; Keyzer, E. N.; Arndtsen, B. A., *Angew. Chem. Int. Ed.* **2017**, 56, 6078-6082.
- Review Sadowski, B.; Klajn, J.; Gryko, D. T., *Org. Biomol. Chem.* **2016**, 14, 7804-7828.
- Padwa, A.; Austin, D. J.; Precedo, L.; Zhi, L., *J. Org. Chem.* **1993**, 58, 1144-1150.
- Chen, R.; Zhao, Y.; Sun, H.; Shao, Y.; Xu, Y.; Ma, M.; Ma, L.; Wan, X., *J. Org. Chem.* **2017**, 82, 9291-9304.
- Day, J.; McKeever-Abbas, B.; Dowden, J., *Angew. Chem. Int. Edit.* **2016**, 55, 5809-5813.
- Allgaeuer, D. S.; Mayer, P.; Mayr, H., *J. Am. Chem. Soc.* **2013**, 135, 15216-15224.
- Other catalysts were evaluated but gave inferior yields compared to [Fe(PPP)Cl] (Table S1).<sup>20</sup>
- Vuluga, D.; Legros, J.; Crousse, B.; Bonnet-Delpon, D., *Green Chem.* **2009**, 11, 156-159.
- Acheson, R. M.; Taylor, G. A., *J. Chem. Soc.* **1960**, 1691-1701.
- Sundberg, R. J.; Hamilton, G.; Trindle, C., *J. Org. Chem.* **1986**, 51, 3672-3679.
- Supplementary Tables and Schemes are provided in the Supporting Information
- Matsumoto, K.; Ikemi, Y.; Konishi, H.; Shi, X.; Uchida, T., *Heterocycles* **1988**, 27, 2557-2562.
- Jedinák, L.; Zátoková, R.; Zemánková, H.; Šustková, A.; Cankař, P., *J. Org. Chem.* **2017**, 82, 157-169.
- Gorringer, A. M.; Lloyd, D.; Wasson, F. I.; Marshall, D. R.; Duffield, P. A., *J. Chem. Soc. C*, **1969**, 1449-1456.

TOC graphic:

