Abstract

The development of well-constrained palaeo-proxies that enable the reconstruction of past climate change is becoming an ever more important field of scientific enquiry within the palaeobotanical community, with the potential to deliver broader impacts linked to understanding of future anthropogenic climate change. One of the major uncertainties in predicting climate change is how the hydrological cycle will respond to future warming. Griener and Warny (2015, Review of Palaeobotany and Palynology 221, 138-143) suggested that pollen size might be a useful proxy for tracking moisture availability, as pollen size appears to be negatively correlated with moisture. Given the long fossil record of pollen and spores such a proxy would have broad scope and the potential to deliver much needed information. Here we set out to fully evaluate and test the robustness of this proxy. We focus on a number of a key issues: controls on pollen size, data analysis, and finally proxy validation. Using this approach we find that there is little theoretical or empirical support for the original relationship proposed by Griener and Warny. Consequently it is currently premature to use pollen size as a moisture availability proxy in the fossil record. However, we recognise that the technique may have potential and conclude by offering a series of recommendations that would rigorously assess and test for a relationship between pollen size and moisture availability.

Keywords

Pollen size, moisture availability, *Nothofagus*, phylogeny, palaeoclimate proxy

1. Introduction

In the absence of direct measurements of climate in Earth's past, palaeoclimate proxies have become essential for reconstructing climatic trends and ground-truthing climate models (Masson-Delmotte et al., 2013). However before these proxies can be deployed they need to be fully tested to assess both accuracy and the precession that underlie their predictive elements (Lomax and Fraser, 2015; Lomax et al., 2012). While fossil pollen and spores have traditionally been used in a passive manner (Jardine et al., 2016) to infer past climates based on the climatic tolerances of their nearest living extant relatives (e.g. Mosbrugger and Utescher, 1997; Utescher et al., 2014), there have also been attempts to use palynological morphological (Griener and Warny, 2015; Kürschner et al., 2013) and chemical data (Fraser et al., 2014; Fraser et al., 2011; Jardine et al., 2017; Jardine et al., 2016; Lomax and Fraser, 2015; Lomax et al., 2008; Rozema et al., 2009; Rozema et al., 2001a; Rozema et al., 2001b; Watson et al., 2007) to reconstruct climatic parameters more directly.

Griener and Warny (2015), hereafter G&W15, introduced a proxy for moisture availability based on the size of pollen grains. This proxy is centered on the idea that larger pollen grains will be relatively more resistant to desiccation stress, because of a decreased surface area to volume ratio (Ejsmond et al., 2011; Griener and Warny, 2015). There should therefore be a negative correlation between moisture availability and pollen size at certain taxonomic levels (i.e. within species, genera or families). Moisture availability is a key climatic parameter (Wilf et al., 1998) and an important control on plant distributions (Engelbrecht et al., 2007; Gentry, 1988). A new, robust proxy for moisture availability would therefore be a particularly valuable tool for the scientific community. G&W15 demonstrated this proxy by relating the size of modern *Nothofagus* pollen grains to mean annual precipitation (MAP), and then reconstructing size trends in Antarctic *Nothofagidites*

lachlaniae-complex pollen grains from the Eocene to the Miocene. Partial validation of the fossil pollen size data was carried out via a descriptive (i.e. non-statistical) comparison to the intrinsic water use efficiency (WUE_i) of Eocene *Nothofagus*, derived from the carbon isotope discrimination (Δ) of *Nothofagidites* sporopollenin (Griener et al., 2013).

Here, we focus on a number of key issues that we have identified in the theoretical basis and validation of the moisture availability proxy of G&W15. Specifically, we focus on three aspects: first, known controls on pollen size; second, the data analysis techniques employed by G&W15 for validating their modern pollen size data, including a lack of accounting for phylogenetic autocorrelation; and third, the lack of supporting evidence for the palaeo-moisture availability reconstruction developed by G&W15 in coeval fossil data. Finally, we suggest some ways in which this proxy may be more fully developed and validated.

2. Controls on pollen size

2.1 Genome size

There is extensive literature showing that pollen size in extant plants is related to genome size and that this relationship scales with the level of ploidy (Bennett, 1972; De Storme et al., 2013). For example Bennett (1972) showed that there is a highly significant positive relationship between pollen volume and genome size in the grasses ($y = 2.1 + 0.643 \cdot x$, $r^2 = 0.91$, $p < 0.001$, $n = 15$). Although these data are not corrected for the effects of phylogeny (see section 3, below) it does highlight that factors other than moisture availability have the capacity to drive changes in pollen size and that these factors can explain a greater proportion of the variance. A full investigation of the relationship between pollen diameter and genome size indicates that relationship appears to hold within disparate plant groups

88 especially when looking at ploidy but does not scale across broad groups when phylogeny is
89 accounted for (Knight et al., 2010). These findings suggest that there is a strong phylogenetic
90 signal (see section 3) that links pollen size to genome size and that is independent of
91 environment.

92
93 There is some evidence for polyploidy being a selective advantage in dryer
94 environments. The larger cells of polyploids can lead to higher xylem hydraulic conductance,
95 for example (Thompson et al., 2015), as well as differences in stomatal apparatus
96 (Manzaneda et al., 2012). Manzaneda et al. (2012) found that aridity is a strong predictor of
97 ploidy level in the temperate grass *Brachypodium distachyon*, with tetraploid individuals
98 having higher water use efficiency and increased tolerance to drought relative to diploid
99 individuals. Further support for polyploidy conferring a selective advantage with regard to
100 water stress comes from Garbutt and Bazzaz (1983), Watanabe (1986) and Li et al. (1996),
101 all of whom demonstrated that polyploid plants are more tolerant of water stress than
102 diploids. However, there is also evidence for polyploid plants being less well adapted to
103 drought conditions (Baldwin, 1941). Similar dichotomous results are also found when
104 comparing polyploid plants to diploids when looking for trends that confer cold tolerance and
105 shade tolerance (see Maherali et al., 2009 for details). Taken together these findings suggest
106 that while there is some tentative support for a general relationship between pollen size and
107 moisture availability (Ejzmond et al., 2011; Griener and Warny, 2015), it is highly likely that
108 this response would be mediated through genome size, rather than being a direct cause and
109 effect relationship. Consequently, any other factors influencing the relationship between
110 ploidy/genome size and habitat preference would bias the G&W15 moisture availability
111 proxy, limiting the inferences that could be made from it.

2.2 *Palynological processing methods, mounting media and taphonomy*

It has long been known that different palynological processing methods can modify pollen size (Christensen, 1946; Faegri and Iversen, 1975; Moore et al., 1991; Reitsma, 1969), although this was not directly discussed by G&W15, either in relation to their own samples or as implications for future moisture availability reconstructions. This is particularly important because G&W15 treated their modern and fossil samples differently, with no processing for the modern samples and a standard processing protocol of HF and HCl for the fossil samples. HF has been shown to decrease the size of pollen grains (Faegri and Iversen, 1975; Moore et al., 1991; Schüler and Behling, 2011), meaning that the size changes demonstrated by G&W15 may not be directly comparable across datasets. Since no pollen size-MAP calibration or quantitative precipitation estimates were attempted by G&W15 this in itself does not compromise their results, but it will need to be considered if this proxy is developed further as a quantitative technique. Other processing methods not carried out by G&W15, such as treatment with KOH and acetolysis, have been reported to cause size changes in palynomorphs (Christensen, 1946; Faegri and Iversen, 1975; Moore et al., 1991; Reitsma, 1969), so as with HF care will need to be taken when analysing grain size trends across sample sets that have undergone different processing protocols.

The choice of storage and mounting media is also known to have an impact on pollen size (Andersen, 1960; Christensen, 1946; Cushing, 1961; Faegri and Iversen, 1975; Moore et al., 1991; Reitsma, 1969; Sluyter, 1997). G&W15 mounted their modern and fossil samples in glycerine jelly, which can cause swelling of pollen grains, either through absorption of water from the atmosphere (Christensen, 1946; Moore et al., 1991) or softening of the exine and subsequent deformation caused by pressure from the microscope coverslip (Cushing, 1961). G&W15 did not state how long their samples were stored in glycerine jelly prior to

analysis, so it is not possible to say how much of an impact it may have had on their reported measurements. Clearly any pollen size analyses that make use of samples that have been stored in glycerine jelly will need to take possible size changes into account, especially if different batches of samples were processed at different times. An alternative storage and mounting medium such as silicone oil (Andersen, 1960; Sluyter, 1997) may be a better option for pollen size measurements, or using coverslip supports (e.g. sand grains or splints from other coverslips) to limit the downward pressure on the pollen grains (Cushing, 1961).

Finally, taphonomic processes will impact upon the size and shape of fossil palynomorphs. Damage to pollen grains and spores through folding, pinching or breaking is a common occurrence in fossil palynological samples (Havinga, 1967; Mander et al, 2012; Tweddle and Edwards, 2010; Twiddie and Bunting, 2010), and makes consistent measurements challenging, especially across changes in taphonomic regimes or in comparison to modern specimens. Careful quality control will therefore be needed when selecting specimens for measurement, which may limit the broader utility of the proxy unless the target taxon is abundant and well-preserved through the time period of interest. Taken together, these various factors show that care needs to be taken when selecting and processing samples for pollen size analysis, and that all processing protocols need to be fully reported in the literature. While these specific issues may not have compromised the analysis in G&W15, they will need to be carefully considered and taken into account in any further research on pollen size-based proxies, whether these relate to moisture availability or genome size in the fossil record.

3. Analysis of data

The modern validation dataset of G&W15 comprised 157 measurements of *Nothofagus* pollen diameters taken from 19 herbarium specimens representing 12 species (not 13, as stated in G&W15), with between 1 and 30 grains measured per specimen. G&W15 tested the relationship between pollen size and mean annual precipitation using ordinary least squares (OLS) linear regression, fitting a linear model through the mean values for each sample. OLS regressions were carried out for the whole dataset together (Fig. 1 in G&W15) and separately for the subgenera *Brassospora*, *Fuscospora* and *Nothofagus* (Fig. S1 in G&W15). The subgenus *Lophozonia* is only represented by one measurement in the G&W15 dataset, from the species *N. obliqua*, and so could not be modelled separately.

One of the underlying assumptions of OLS regression is that the values in the response variable are independent and identically distributed (i.i.d.) (Rohlf, 2006; Zuur et al., 2009). Data from a range of taxa are never truly independent, because the taxa will be in some way descended from a common ancestor, with more closely related species being more similar than distantly related species (Garland and Ives, 2000). This phylogenetic signal in the response variable violates the assumption of independence (Garland and Ives, 2000; Martins and Hansen, 1997; Rohlf, 2006), inflating the Type 1 error rate (i.e. incorrectly rejecting a true null hypothesis, and finding a statistically significant result where none is present) (Rohlf, 2006).

Several methods have been developed to account for the phylogenetic signal within biological trait data (Cooper et al., 2016). One of the most commonly used is phylogenetic independent contrasts (PIC), which calculates differences (contrasts) between the character/trait values of pairs of sister taxa across the tips and nodes of a phylogeny (Felsenstein, 1985; Garland et al., 1992; Garland and Ives, 2000). PIC implicitly assumes a

Brownian motion model of trait evolution, i.e. an evolutionary random walk through trait space (Cooper et al., 2016; Garland and Ives, 2000). The calculated contrasts are phylogenetically independent and can be used in standard statistical analyses such as correlations, allowing for the role of phylogeny in driving the relationship to be assessed. For example from a palaeobotanical perspective PIC has been used to explore if carbon isotope fractionation is driven by phylogeny (Lomax et al., 2012) and to determine that within angiosperms the relationship between genome size and guard cell length is independent of phylogeny (Beaulieu et al., 2008; Lomax et al., 2014), enabling broad scale reconstruction of the genome size of fossil plants (Lomax et al., 2014).

More recently, phylogenetic generalised least squares (P-GLS) regression has been developed, which allows for phylogenetic relatedness to be directly modelled as a correlation structure within the linear regression framework (Blomberg et al., 2012; Garland and Ives, 2000; Grafen, 1989; Martins and Hansen, 1997). P-GLS regression has the advantages that (i) it is more flexible than PIC, allowing for different underlying models of trait evolution (Rohlf, 2006), (ii) the variables can be modeled directly, rather than as contrasts (Garland and Ives, 2000), and (iii) it can be readily extended to more complex mixed effects models (Blomberg et al., 2012). However when a Brownian motion model of trait evolution is assumed in P-GLS regression the two techniques give identical results (Blomberg et al., 2012; Garland and Ives, 2000; Rohlf, 2006).

Here, we re-analyse the G&W15 dataset using P-GLS. G&W15 made their dataset (specimen pollen size mean and standard deviation, and associated MAP) available in Table S1 of their paper. We use the *Nothofagus* molecular phylogeny of Sauquet et al. (2012), downloaded from TreeBASE (www.treebase.org) on 18/01/2017, to represent the

phylogenetic relatedness among taxa. The phylogeny comprises 27 *Nothofagus* species (Fig 1), as well as 21 outgroup species from the core Fagales, and so was first trimmed down to the taxa represented in the G&W15 dataset. *Nothofagus rubra* and *Nothofagus starkenborghii* are not present in the molecular phylogeny, and so were removed from the G&W15 dataset. The pollen size and MAP means for the remaining 10 taxa were then calculated for use in the P-GLS regression. We used a simple Brownian motion model of trait evolution across the trimmed *Nothofagus* phylogeny as a correlation structure in the P-GLS regression. Data analysis was carried out in R v. 3.3.1 (Team, 2016) using the packages ape v. 4.1 (Paradis et al., 2004), phytools v. 0.5-64 (Revell, 2012), nlme v. 3.1-131 (Pinheiro et al., 2017) and astrochron v. 0.6.5 (Meyers, 2014).

Mapping pollen size directly onto the molecular phylogeny using ancestral character estimation (Revell, 2013) shows a degree of phylogenetic structuring (i.e. similar values in the sister taxa *N. carrii* and *N. grandis*, and *N. dombeyi* and *N. betuloides*), confirming the importance of correcting for phylogenetic non-independence in the regression model (Fig 2a; see also Fernández et al., 2016). Regressing pollen size onto MAP (pollen size is the response or dependent variable, MAP is the explanatory or independent variable) using P-GLS regression gives the model $y = 35.2 - 0.003 \cdot x$, $p = 0.003$, $n = 10$, whereas an OLS regression of the same version of the dataset gives the model $y = 38.4 - 0.005 \cdot x$, $p = 0.0006$, $n = 10$ (Fig 2b). While both model fits are statistically significant, the p value is an order of magnitude lower in the P-GLS model, and the slope is also shallower. The different slopes between the two models shows the importance of including phylogeny for any regression models being used for calibration and quantitative moisture availability reconstructions.

There is also the more general point that the G&W15 dataset itself is quite small, with limited replication of individual plants within each species. Seven of the species in the dataset are represented by just one herbarium specimen, three by two specimens, and two by three specimens. This not only makes it challenging to look at within-species variation, to investigate whether populations at different MAP levels exhibit differences in pollen size, but also makes it impossible to assess whether non-significant relationships at the subgenus level are due to a genuine lack of a response to moisture availability or insufficient data. G&W15 carried out regressions for the three subgenera with more than one species represented, and only found a significant relationship for one of them, *Brassospora* (Fig. 3, note that the subgenus *Nothofagus* was reported as significant but had a p value of <0.17 in G&W15). The significant relationship for *Brassospora* is largely driven by one outlier, the species *N. discoidea*, which with a mean grain size of $32.1\ \mu\text{m}$ is considerably larger than the other *Brassospora* species sampled (Figs. 3 and 2a). Removing the *N. discoidea* data point from the dataset and re-running the regression reduces the slope of the fitted model from -0.006 to -0.003 , the r^2 from 0.73 to 0.28 , and the p value from 0.003 to 0.18 (i.e. the relationship is no longer statistically significant without this one species). There is therefore limited evidence for a consistent, statistically robust relationship between pollen size and MAP below the level of genus. While this is very possibly down to the small sample sizes involved, this in itself suggests that the dataset used by G&W15 is too limited to robustly validate this proxy, and that further replication across taxonomic ranks is needed if it is to be confidently deployed as a palaeoclimate proxy.

4. Lack of evidence in fossil data

Past moisture availability reconstructions based on fossil pollen size should show good agreement with other moisture availability or precipitation proxies (assuming these

themselves are broadly reliable). G&W15 measured *Nothofagidites lachlaniae*-complex pollen size from the Eocene to the Miocene of the Antarctic Peninsula (data available in Table S2 of G&W15), and interpreted changes in the context of moisture availability. Griener et al. (2013) generated a moisture availability proxy record for the late Eocene portion of the same cores, based on *Nothofagidites* pollen $\delta^{13}\text{C}$ discrimination. The $\Delta^{13}\text{C}$ record shows an overall decrease through time, suggesting a decrease in moisture availability, and thus we would expect this to be reflected in a trend towards larger pollen sizes (i.e. the two records should show a negative correlation).

The decrease in moisture availability suggested by the $\Delta^{13}\text{C}$ dataset is less clear in the pollen size dataset, however, especially given the large errors on the pollen size means (Fig. 4a). Interpolating the $\Delta^{13}\text{C}$ data to the same depth levels as the size data, and regressing size onto $\Delta^{13}\text{C}$ with OSL regression gives the model $y = 33.40 - 0.48 \cdot x$, $r^2 = 0.08$, $p = 0.37$, $n = 12$ (Fig. 4b). Therefore the regression reveals the expected negative correlation, but the relationship is not statistically significant and the $\Delta^{13}\text{C}$ data only explains a small proportion of the variance in the pollen size data. The negative correlation between pollen size and $\Delta^{13}\text{C}$ is also largely driven by one extreme value (Fig. 4b). Given that the size and discrimination data have been developed on the same plant group from the same sedimentary record this result offers no support for a link between moisture availability and pollen size. This proxy therefore cannot be successfully validated in the fossil record, although we acknowledge that other factors that drive stomatal closure (such as elevated CO_2) which control discrimination could also influence $\Delta^{13}\text{C}$. However experimental work has shown that water availability does act as a primary control on $\Delta^{13}\text{C}$ even over a range of CO_2 concentrations relevant to the geological record (Lomax et al., 2012).

5. Suggestions for further proxy development

For the reasons outlined above, we feel that it is currently premature to use pollen size as a moisture availability proxy in the fossil record. In order to fully test and validate this as a robust proxy, we suggest that the following measures are needed:

1. A better understanding of the relationship(s) between genome size, abiotic factors including moisture availability and pollen size.
2. Considerably more modern pollen size data, with greater replication across taxonomic ranks, to robustly quantify the relationship between pollen size and moisture availability. Analysis of these data will require a proper accounting for phylogenetic dependency, and also any spatial autocorrelation between sampling localities (Zuur et al., 2009).
3. Data from controlled growth experiments, with pollen harvested from plants grown under a range of environmental conditions, to better understand the external controls on pollen size within and among species.
4. Further palaeo-palynological datasets where there are existing precipitation indicators, to allow for validation in the fossil record. Quaternary pollen records may be best suited to this, because of better-constrained environmental conditions compared to the deep time fossil record.

Acknowledgements

We would like to thank Surangi Punyasena and one anonymous reviewer, whose comments greatly improved this paper. We would also like to thank Kathryn Griener and Sophie Warny for making their dataset freely available alongside their paper, which allowed for this reanalysis to be carried out. B.H.L.'s research on palaeogenome size was funded via NERC

New Investigators grant NE/J004855/1. P.E.J. was funded by ERC grant 649081 (MAGIC) and NERC standard grant NE/K005294/1.

References

Andersen, S.T., 1960. Silicone oil as a mounting medium for pollen grains. *Danmarks Geologiske Undersøgelse (Series) IV 1*, 1-24.

Baldwin, T.J., 1941. *Galax*: the genus and its chromosomes. *Journal of Heredity* 32, 249-254.

Beaulieu, J.M., Leitch, I.J., Patel, S., Pendharkar, A., Knight, C.A., 2008. Genome size is a strong predictor of cell size and stomatal density in angiosperms. *New Phytologist* 179, 975–986.

Bennett, M.D., 1972. Nuclear DNA Content and Minimum Generation Time in Herbaceous Plants. *Proceedings of the Royal Society of London B: Biological Sciences* 181, 109-135.

Blomberg, S.P., Lefevre, J.G., Wells, J.A., Waterhouse, M., 2012. Independent contrasts and PGLS regression estimators are equivalent. *Systematic Biology* 61, 382-391.

Christensen, B.B., 1946. Measurement as a means of identifying fossil pollen. *Danmarks Geologiske Undersøgelse (Series) IV 3(2)*, 1-22.

Cooper, N., Thomas, G.H., FitzJohn, R.G., 2016. Shedding light on the 'dark side' of phylogenetic comparative methods. *Methods in Ecology and Evolution* 7, 693-699.

335 Cushing, E.J., 1961. Size increase in pollen grains mounted in thin slides. *Pollen et Spores*
336 3(2), 265-274.
337

338 De Storme, N., Zamariola, L., Mau, M., Sharbel, T.F., Geelen, D., 2013. Volume-based
339 pollen size analysis: an advanced method to assess somatic and gametophytic ploidy in
340 flowering plants. *Plant Reproduction* 26, 65-81.
341

342 Ejsmond, M.J., Wrońska-Pilarek, D., Ejsmond, A., Dragosz-Kluska, D., Karpińska-Kołaczek,
343 M., Kołaczek, P., Kozłowski, J., 2011. Does climate affect pollen morphology? Optimal size
344 and shape of pollen grains under various desiccation intensity. *Ecosphere* 2, 1-15.
345

346 Engelbrecht, B.M., Comita, L.S., Condit, R., Kursar, T.A., Tyree, M.T., Turner, B.L.,
347 Hubbell, S.P., 2007. Drought sensitivity shapes species distribution patterns in tropical
348 forests. *Nature* 447, 80-82.
349

350 Faegri, K., Iversen, J., 1975. Textbook of pollen analysis, 3rd edition. Munksgaard,
351 Copenhagen.
352

353 Felsenstein, J., 1985. Phylogenies and the comparative method. *American Naturalist* 125, 1-
354 15.
355

356 Fernández, D.A., Santamarina, P.E., Tellería, M.C., Palazzesi, L., Barreda, V.D., 2016.
357 Pollen morphology of *Nothofagus* (Nothofagaceae, Fagales) and its phylogenetic
358 significance. *Acta Palaeobotanica* 56, 223-245.
359

360 Fraser, W.T., Lomax, B.H., Jardine, P.E., Gosling, W.D., Sephton, M.A., 2014. Pollen and
 361 spores as a passive monitor of ultraviolet radiation. *Frontiers in Ecology and Evolution* 2:12.
 362 <http://dx.doi.org/10.3389/fevo.2014.00012>
 363
 364 Fraser, W.T., Sephton, M.A., Watson, J.S., Self, S., Lomax, B.H., James, D.I., Wellman,
 365 C.H., Callaghan, T.V., Beerling, D.J., 2011. UV-B absorbing pigments in spores:
 366 biochemical responses to shade in a high-latitude birch forest and implications for
 367 sporopollenin-based proxies of past environmental change. *Polar Research* 30, 8312.
 368
 369 Garbutt, K., Bazzaz, F.A., 1983. Leaf demography, flower production and biomass of diploid
 370 and tetraploid populations of *Phlox drummondii* Hook. on a soil moisture gradient. *New*
 371 *Phytologist* 93, 129-141.
 372
 373 Garland, T.J., Harvey, P.H., Ives, A.R., 1992. Procedures for the analysis of comparative data
 374 using phylogenetically independent contrasts. *Systematic Biology* 41, 18-32.
 375
 376 Garland, T.J., Ives, A.R., 2000. Using the Past to Predict the Present: Confidence Intervals
 377 for Regression Equations in Phylogenetic Comparative Methods. *The American Naturalist*
 378 155, 346-364.
 379
 380 Gentry, A.H., 1988. Changes in plant community diversity and floristic composition on
 381 environmental and geographical gradients. *Annals of the Missouri Botanical Garden* 75, 1-
 382 34.
 383

384 Grafen, A., 1989. The phylogenetic regression. Philosophical Transactions of the Royal
 385 Society of London B: Biological Sciences 326, 119-157.
 386
 387 Griener, K.W., Nelson, D.M., Warny, S., 2013. Declining moisture availability on the
 388 Antarctic Peninsula during the Late Eocene. Palaeogeography, Palaeoclimatology,
 389 Palaeoecology 383-384, 72-78.
 390
 391 Griener, K.W., Warny, S., 2015. *Nothofagus* pollen grain size as a proxy for long-term
 392 climate change: An applied study on Eocene, Oligocene, and Miocene sediments from
 393 Antarctica. Review of Palaeobotany and Palynology 221, 138-143.
 394
 395 Havinga, A.J., 1967. Palynology and pollen preservation. Review of Palaeobotany and
 396 Palynology 2, 81-98.
 397
 398 Jardine, P.E., Abernethy, F.A.J., Lomax, B.H., Gosling, W.D., Fraser, W.T., 2017. Shedding
 399 light on sporopollenin chemistry, with reference to UV reconstructions. Review of
 400 Palaeobotany and Palynology 238, 1-6.
 401
 402 Jardine, P.E., Fraser, W.T., Lomax, B.H., Sephton, M.A., Shanahan, T.M., Miller, C.S.,
 403 Gosling, W.D., 2016. Pollen and spores as biological recorders of past ultraviolet irradiance.
 404 Scientific Reports 6, 1-8.
 405
 406 Knight, C.A., Clancy, R.B., Götzenberger, L., Dann, L., Beaulieu, J.M., 2010. On the
 407 Relationship between Pollen Size and Genome Size. Journal of Botany 2010, 612017.
 408

409 Kürschner, W.M., Batenburg, S.J., Mander, L., 2013. Aberrant *Classopollis* pollen reveals
 410 evidence for unreduced (2n) pollen in the conifer family Cheirolepidiaceae during the
 411 Triassic–Jurassic transition. Proceedings of the Royal Society of London B: Biological
 412 Sciences 280, 20131708.
 413
 414 Li, W.-L., Berlyn, G.P., Ashton, P.M.S., 1996. Polyploids and their structural and
 415 physiological characteristics relative to water deficit in *Betula papyrifera* (Betulaceae).
 416 American Journal of Botany 83, 15-20.
 417
 418 Lomax, B.H., Fraser, W.T., 2015. Palaeoproxies: Botanical monitors and recorders of
 419 atmospheric change. Palaeontology 58, 759-768.
 420
 421 Lomax, B.H., Fraser, W.T., Sephton, M.A., Callaghan, T.V., Self, S., Harfoot, M., Pyle, J.A.,
 422 Wellman, C.H., Beerling, D.J., 2008. Plant spore walls as a record of long-term changes in
 423 ultraviolet-B radiation. Nature Geoscience 1, 592-596.
 424
 425 Lomax, B.H., Hilton, J., Bateman, R.M., Upchurch, G.R., Lake, J.A., Leitch, I.J., Cromwell,
 426 A., Knight, C.A., 2014. Reconstructing relative genome size of vascular plants through
 427 geological time. New Phytologist 201, 636-644.
 428
 429 Lomax, B.H., Knight, C.A., Lake, J.A., 2012. An experimental evaluation of the use of C₃ δ
 430 ¹³C plant tissue as a proxy for the paleoatmospheric δ ¹³CO₂ signature of air. Geochemistry,
 431 Geophysics, Geosystems 13, Q0AI03.
 432

433 Maherali, H., Walden, A.E., Husband, B.C., 2009. Genome duplication and the evolution of
 434 physiological responses to water stress. *New Phytologist* 184, 721-731.
 435
 436 Mander, L., Wesseln, C.J., McElwain, J.C., Punyasena, S.W., 2012. Tracking taphonomic
 437 regimes using chemical and mechanical damage of pollen and spores: an example from the
 438 Triassic–Jurassic mass extinction. *PloS ONE*, 7(11), e49153.
 439
 440 Manzaneda, A.J., Rey, P.J., Bastida, J.M., Weiss-Lehman, C., Raskin, E., Mitchell-Olds, T.,
 441 2012. Environmental aridity is associated with cytotype segregation and polyploidy
 442 occurrence in *Brachypodium distachyon* (Poaceae). *New Phytologist* 193, 797-805.
 443
 444 Martins, E.P., Hansen, T.F., 1997. Phylogenies and the comparative method: a general
 445 approach to incorporating phylogenetic information into the analysis of interspecific data.
 446 *The American Naturalist* 149, 646-667.
 447
 448 Masson-Delmotte, V., Schulz, M., Abe-Ouchi, A., Beer, J., Ganopolski, A., González Rouco,
 449 J.F., Jansen, E., Lambeck, K., Luterbacher, J., Naish, T., Osborn, T., Otto-Bliesner, B.,
 450 Quinn, T., Ramesh, R., Rojas, M., Shao, X., Timmermann, A., 2013. Information from
 451 Paleoclimate Archives, in: Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K.,
 452 Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M. (Eds.), *Climate Change 2013: The*
 453 *Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of*
 454 *the Intergovernmental Panel on Climate Change*. Cambridge, United Kingdom and New
 455 York, NY, USA.: Cambridge University Press, 383-464.
 456

457 Meyers, S.R., 2014. Astrochron: An R Package for Astrochronology. [http://cran.r-](http://cran.r-project.org/package=astrochron)
458 [project.org/package=astrochron](http://cran.r-project.org/package=astrochron)
459
460 Moore, P.D., Webb, J.A., Collinson, M.E., 1991. Pollen Analysis, 2nd edition. Blackwell
461 Science Ltd, Oxford.
462
463 Mosbrugger, V., Utescher, T., 1997. The coexistence approach – a method for quantitative
464 reconstructions of Tertiary terrestrial palaeoclimate data using plant fossils. *Palaeogeography,*
465 *Palaeoclimatology, Palaeoecology* 134, 61-86.
466
467 Paradis, E., Claude, J., Strimmer, K., 2004. APE: analyses of phylogenetics and evolution in
468 R language. *Bioinformatics* 20, 289-290. <http://cran.r-project.org/package=ape>
469
470 Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., Team, R.C., 2017. nlme: Linear and
471 Nonlinear Mixed Effects Models, 3.1-128 ed. <http://cran.r-project.org/package=nlme>
472
473 R Core Team 2016. R: A language and environment for statistical computing. R Foundation
474 for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>
475
476 Reitsma, T.J., 1969. Size modification of recent pollen grains under different treatments.
477 *Review of Palaeobotany and Palynology* 9(3), 175-202.
478
479 Revell, L.J., 2012. phytools: An R package for phylogenetic comparative biology (and other
480 things). *Methods in Ecology and Evolution* 3, 217-223. [http://cran.r-](http://cran.r-project.org/package=phytools)
481 [project.org/package=phytools](http://cran.r-project.org/package=phytools)

482

483 Revell, L.J., 2013. Two new graphical methods for mapping trait evolution on phylogenies.

484 *Methods in Ecology and Evolution* 4, 754-759.

485

486 Rohlf, F.J., 2006. A comment on phylogenetic correction. *Evolution* 60, 1509-1515.

487

488 Rozema, J., Blokker, P., Mayoral Fuertes, M.A., Broekman, R., 2009. UV-B absorbing

489 compounds in present-day and fossil pollen, spores, cuticles, seed coats and wood: evaluation

490 of a proxy for solar UV radiation. *Photochemical and Photobiological Sciences* 8, 1233-1243.

491

492 Rozema, J., Broekman, R.A., Blokker, P., Meijkamp, B., de Bakker, N., van de Staaij, J., van

493 Beem, A., Ariese, F., Kars, S.M., 2001a. UV-B absorbance and UV-B absorbing compounds

494 (*para*-coumaric acid) in pollen and sporopollenin: the perspective to track historic UV-B

495 levels. *Journal of Photochemistry and Photobiology B: Biology* 62, 108-117.

496

497 Rozema, J., Noordjik, A.J., Broekman, R.A., van Beem, A., Meijkamp, B.M., de Bakker,

498 N.V.J., van de Staaij, J.W.M., Stroetenga, M., Bohncke, S.J.P., Konert, M., Kars, S., Peat, H.,

499 Smith, R.I.L., Convey, P., 2001b. (Poly)phenolic compounds in pollen and spores of

500 Antarctic plants as indicators of UV-B: A new proxy for the reconstruction of past solar UV-

501 B? *Plant Ecology* 154, 11-26.

502

503 Sauquet, H., Ho, S.Y., Gandolfo, M.A., Jordan, G.J., Wilf, P., Cantrill, D.J., Bayly, M.J.,

504 Bromham, L., Brown, G.K., Carpenter, R.J., Lee, D.M., Murphy, D.J., Sniderman, J.M.,

505 Udovicic, F., 2012. Testing the impact of calibration on molecular divergence times using a

506 fossil-rich group: the case of *Nothofagus* (Fagales). *Systematic Biology* 61, 289-313.

507

508 Schüler, L., Behling, H., 2011. Poaceae pollen grain size as a tool to distinguish past
509 grasslands in South America: a new methodological approach. *Vegetation History and*
510 *Archaeobotany* 20, 83-96.

511

512 Sluyter, A., 1997. Analysis of maize (*Zea mays* subsp. *mays*) pollen: normalizing the effects
513 of microscope-slide mounting media on diameter determinations. *Palynology* 21, 35-39.

514

515 Thompson, K.A., Husband, B.C., Maherali, H., Bonser, S., 2015. No influence of water
516 limitation on the outcome of competition between diploid and tetraploid *Chamerion*
517 *angustifolium* (Onagraceae). *Journal of Ecology* 103, 733-741.

518

519 Tweddle, J.C., Edwards, K.J., 2010. Pollen preservation zones as an interpretive tool in
520 Holocene palynology. *Review of Palaeobotany and Palynology* 161, 59–76.

521

522 Twiddle, C.L., Bunting, M.J., 2010. Experimental investigations into the preservation of
523 pollen grains: A pilot study of four pollen types. *Review of Palaeobotany and Palynology*
524 162, 621-630.

525

526 Utescher, T., Bruch, A.A., Erdei, B., François, I., Ivanov, D., Jacques, F.M.B., Kern, A.K.,
527 Liu, Y.-S.C., Mosbrugger, V., Spicer, R.A., 2014. The Coexistence Approach – Theoretical
528 background and practical considerations of using plant fossils for climate quantification.
529 *Palaeogeography, Palaeoclimatology, Palaeoecology* 410, 58-73.

530

531 Watanabe, K., 1986. The cytogeography of the genus *Eupatorium* (Compositae) – a review.
 532 Plant Species Biology 1, 99-116.
 533
 534 Watson, J.S., Septhon, M.A., Septhon, S.V., Self, S., Fraser, W.T., Lomax, B.H., Gilmour, I.,
 535 Wellman, C.H., Beerling, D.J., 2007. Rapid determination of spore chemistry using
 536 thermochemolysis gas chromatography-mass spectrometry and micro-Fourier transform
 537 infrared spectroscopy. Photochemistry and Photobiology 6, 689-694.
 538
 539 Wilf, P., Wing, S.L., Greenwood, D.R., Greenwood, C.L., 1998. Using fossil leaves as
 540 paleoprecipitation indicators: An Eocene example. Geology 26, 203-206.
 541
 542 Zuur, A.F., Ieno, E.N., Walker, N.J., Saveliev, A.A., Smith, G.M., 2009. Mixed effects
 543 models and extensions in ecology with R. Springer, New York.
 544

Figure legends

Fig. 1. *Nothofagus* phylogeny of Sauquet et al. (2012), with the taxa in the G&W15 dataset in bold type.

Fig. 2. A) *Nothofagus* phylogeny of Sauquet et al. (2012) limited to the taxa in the G&W15 dataset, with pollen size mapped on using ancestral character estimation (Revell, 2013). B) Pollen size plotted against mean annual precipitation, showing the mean values for each species. Dashed line = ordinary least squares (OSL) model, solid line = phylogenetic generalised least squares (P-GLS) model.

Fig. 3. *Nothofagus* pollen size from G&W15 plotted against mean annual precipitation, showing the mean values for each specimen and OSL regression fits for each subgenus (solid lines are fitted models, dashed lines are 95% confidence intervals). Individual regression models are: *Brassospora*, $y = 39.90 - 0.006 * x$, $r^2 = 0.73$, $p = 0.003$, $n = 9$; *Fuscospora*, $y = 31.55 - 0.002 * x$, $r^2 = 0.37$, $p = 0.28$, $n = 5$; *Nothofagus*, $y = 35.47 - 0.005 * x$, $r^2 = 0.89$, $p = 0.06$, $n = 4$.

Fig. 4. A) Eocene *Nothofagidites* pollen size and $\Delta^{13}\text{C}$ data for the SHALDRIL 3C core. Black points and line = $\Delta^{13}\text{C}$ data from Griener et al. (2013). Grey points with error bars = pollen size data from G&W15, points are sample means and error bars are 1 standard deviation. B) Eocene *Nothofagidites* pollen size regressed against *Nothofagidites* $\Delta^{13}\text{C}$, with the $\Delta^{13}\text{C}$ data interpolated to the same sampling depths as the pollen size data. Black solid line is the fitted regression model, dashed lines are 95% confidence intervals.

Jardine and Lomax

Is pollen size a robust proxy for moisture availability?

Figure 1

571
572

Jardine and Lomax
Is pollen size a robust proxy for moisture availability?
Figure 2

573

574

575

Jardine and Lomax
Is pollen size a robust proxy for moisture availability?
Figure 3

Jardine and Lomax
Is pollen size a robust proxy for moisture availability?
Figure 4

590

591